

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)

2. REPORT TYPE
Technical Papers

3. DATES COVERED (From - To)

4. TITLE AND SUBTITLE

5a. CONTRACT NUMBER

5b. GRANT NUMBER

5c. PROGRAM ELEMENT NUMBER

6. AUTHOR(S)

5d. PROJECT NUMBER

2303

5e. TASK NUMBER

m208

5f. WORK UNIT NUMBER

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)
AFRL/PRS
5 Pollux Drive
Edwards AFB CA 93524-7048

8. PERFORMING ORGANIZATION
REPORT

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory (AFMC)
AFRL/PRS
5 Pollux Drive
Edwards AFB CA 93524-7048

10. SPONSOR/MONITOR'S
ACRONYM(S)

11. SPONSOR/MONITOR'S
NUMBER(S)

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for public release; distribution unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT

1121 041

15. SUBJECT TERMS

16. SECURITY CLASSIFICATION OF:

17. LIMITATION
OF ABSTRACT

18. NUMBER
OF PAGES

19a. NAME OF RESPONSIBLE
PERSON

Leilani Richardson

a. REPORT

b. ABSTRACT

c. THIS PAGE

Unclassified

Unclassified

Unclassified

A

19b. TELEPHONE NUMBER

(include area code)

(661) 275-5015

Standard Form 298 (Rev. 8-98)
Prescribed by ANSI Std. Z39.18

62

separate items are enclosed

TP-1998-090

C8

MEMORANDUM FOR IN-HOUSE PUBLICATIONS

FROM: PROI (TI) (STINFO)

30 Apr 98

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-TP-1998-090
H.Y. Yoo and Jerry Boatz "Theoretical Study of the Mechanism of the Decomposition Process of High
Energy Density Materials" **HEDM Conference Presentation** (Statement A)

Theoretical Study of the Mechanism of the Decomposition Process of High Energy Density Materials

Hi Young Yoo and Jerry Boatz*

National Research Council, Washington, DC 20418

*Air Force Research Laboratory, Edwards AFB, CA 93524-7680

AFOSR HEDM CONTRACTOR'S MEETING
5/18-20, 1998
MONTEREY, CA

DISTRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited



20021121 041

Theoretical Study of the Mechanism of the Decomposition Process of High Energy Density Materials

Hi Young Yoo and Jerry Boatz

National Research Council, Washington, DC 20418
Air Force Research Laboratory, Edwards AFB, CA 93524-7680

One of the goals of the HEDM program is the development of high performance monopropellants which are also less toxic than currently used systems such as hydrazine. Included in this effort is the characterization of decomposition mechanisms, as an initial step toward identification of a suitable catalyst. *Ab initio* quantum mechanical calculations are performed on the decomposition mechanisms of $[\text{NH}_2\text{Me}_2]^+[\text{NO}_3]^-$, a potential monopropellant replacement for hydrazine. The potential energy surfaces of two gas-phase decomposition processes have been explored: (1) proton transfer and (2) methyl cation transfer reactions. These reaction pathways have been examined for both the isolated cation $[\text{NH}_2\text{Me}_2]^+$ and in the presence of a counter anion, $\text{X}=\text{Cl}^-$, $[\text{NO}_3]^-$. For $\text{X}=\text{Cl}^-$, transition states for both pathways have been located at the RHF/6-31G* level and the corresponding intrinsic reaction coordinates (IRCs) have been traced. Comparison of activation barriers and reaction enthalpies for these gas-phase decomposition pathways will be presented here.



Hydrazine is the state of the art monopropellant currently. However, it has several disadvantages including toxicity, volatility, and handling.

Researchers are continually looking for new monopropellant candidates which include energetic materials such as substituted ammonium salts.

Our research effort has concentrated on the decomposition routes of some HEDM materials. *Ab initio* quantum mechanical calculations have been carried out by our group to explore possible decomposition mechanisms.

The model compounds under study are $[\text{NH}_2\text{Me}_2]^+ [\text{X}]^-$, where $\text{X} = \text{NO}_3$ and Cl , and $[\text{N}(\text{NH}_2)_2\text{Me}_2]^+ [\text{NO}_3]^-$.

- To map out the potential energy surface of these model compounds
- To determine the energetics of decomposing species
- To design a catalyst that stabilizes the transition state

Decomposition Mechanisms of $[\text{NH}_2\text{Me}_2]^+[\text{X}]^-$,

$\text{X} = \text{Cl}$ or NO_3


Proton Transfer



Methyl Cation Transfer




A Proton Transfer Transition Structure of $[\text{NH}_2\text{Me}_2]^+[\text{Cl}]^-$ At RHF/6-31G*(6d)



Intermediate Complex

Energy
(kcal/mol)


0.0



Transition Structure

4.6


A Methyl Cation Transfer Transition Structure of $[\text{NH}_2\text{Me}_2]^+[\text{Cl}]^-$ At RHF/6-31G*(6d)



Intermediate Complex

Energy
(kcal/mol)

0.0



Transition Structure

41.6

Calculated Reaction Energies of $[\text{NH}_2\text{Me}_2]^+[\text{Cl}]^-$

Proton Transfer



Methyl Cation Transfer



CONCLUSION

Based on the *ab initio* calculation studies of $[\text{NH}_2\text{Me}_2]^+ [\text{Cl}]^-$

- The activation energy of the proton transfer decomposition process is much lower than that of the methyl cation transfer pathway.
- The transition structure of proton transfer is much tighter than that of methyl cation transfer.
- The methyl cation transfer pathway is 2.6 kcal/mol less endothermic than the proton transfer process.
- Our calculations imply that the proton transfer is most likely the first step to occur in the decomposition process of $[\text{NH}_2\text{Me}_2]^+ [\text{Cl}]^-$.