

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY)			2. REPORT TYPE Technical Papers		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE					5a. CONTRACT NUMBER	
					5b. GRANT NUMBER	
					5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)					5d. PROJECT NUMBER 2303	
					5e. TASK NUMBER M208	
					5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048					8. PERFORMING ORGANIZATION REPORT	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048					10. SPONSOR/MONITOR'S ACRONYM(S)	
					11. SPONSOR/MONITOR'S NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.						
13. SUPPLEMENTARY NOTES						
14. ABSTRACT						
15. SUBJECT TERMS						
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	
a. REPORT	b. ABSTRACT	c. THIS PAGE			Leilani Richardson	
Unclassified	Unclassified	Unclassified	A		19b. TELEPHONE NUMBER <i>(include area code)</i> (661) 275-5015	

62

separate items are enclosed

TP-FY99-0122
✓ Spreadsheet
✓ DTS

ERC # 99-021 ?

MEMORANDUM FOR PRS (In-House)

FROM: PROI (TD) (STINFO)

28 May 1999

SUBJECT: Authorization for Release of Technical Information, Control Number: AFRL-PR-ED-TP-FY99-0122
Suri and Tinnirello, "Bicyclopropylidene and 1,5-Hexadiyne from Bench Scale to Pilot Scale: Problems and Solutions"

Presentation HEDM Conference (Statement A)

Bicyclopropylidene and 1,5-Hexadiyne from Bench Scale to Pilot Scale: Problems and Solutions

Suresh C. Suri and Michael Tinnirello

Air Force Research Laboratory/PRS; ERC Inc.

10 East Saturn Blvd., Edwards AFB, CA 93524

E-Mail: suresh_suri@ple.af.mil

20021122 014

DISTRIBUTION STATEMENT A:
Approved for Public Release -
Distribution Unlimited

Presentation Outline

- Goal
- Criteria for Fuel Selection
- Structural Requirements and Selection for hydrocarbons
- Synthetic Results and Scale Up Challenges
- Future Efforts

Goal

- To come up with a fuel with 2-5% increase of Isp over LOX/RP-1
 - LOX/RP-1 (del.) = 263 sec*
 - LOX/RP-1 (calc.) = 300 sec*

* Determined at sea level and 1000 psi chamber pressure

Task Objective


- Survey of energetic hydrocarbons
- Selection of hydrocarbons based on improved theoretical performance
- Synthesis of target hydrocarbons at bench scale
 - Easy preparation, ^{or} Cost effective and safe
- Translate bench-scale synthesis to pilot scale

Criteria for Fuel Selection

- Predicts Better Performance (Isp) Over LOX/RP-1 System
- Most Desirable Physical Properties
 - Lower Vapor Pressure Compared to RP-1
 - Higher Density (\geq RP-1 = 0.801 g/mL)
 - Freezing Point ($\leq -10^{\circ}\text{C}$; RP-1 = -41.4°C)
 - Boiling Point \geq B. P. of RP-1
- Thermally Stable
- Compatible with the Current System

Structural Requirement for High Energy Contents

- The Energy Contents Can be Increased by Adding Unsaturation in the Molecule



Heat of Formation of Saturated

Hydrocarbons

• Compound	Structure	ΔH_f (Obs)
• Ethane	CH_3CH_3	-20.04
• Propane	$\text{CH}_3\text{CH}_2\text{CH}_3$	-25.02
• Butane	$\text{CH}_3(\text{CH}_2)_2\text{CH}_3$	-30.03
• Pentane	$\text{CH}_3(\text{CH}_2)_3\text{CH}_3$	-35.08

• $\Delta H_f/C = \sim -5 \text{ Kcal/mole}$

Heat of Formation of Unsaturated

Hydrocarbons

• Compound	Structure	$\Delta H_f(\text{Obs})$
• Ethylene	$\text{CH}_2=\text{CH}_2$	+12.5
• 1,3-Butadiene	$\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$	+26.11
	$\Delta H_f/\text{C} = \sim +6.25 \text{ Kcal/mole}$	
• Acetylene	$\text{HC}\equiv\text{CH}$	+54.36
	$\Delta H_f/\text{C} = \sim +27.1 \text{ Kcal/mole}$	

Structural Requirement for High Energy Contents (Cont...)

- The Energy Contents is Also Increased by Incorporating Strain in the Molecule

	ΔH_f
– Ring Compound	
– Cyclopropane	+ 12.73 Kcal/mole
– Cyclobutane	+ 6.78 Kcal/mole
– Cyclopentane	- 18.44 Kcal/mole

Survey of Hydrocarbons



Cyclopropane
 $\Delta H_f = 12.7 \text{ Kcal/mole}$
 $= 0.3 \text{ Kcal/g}$
 $I_{sp} = 312 \text{ Sec.}$



[2.2] Spiropentane
 $\Delta H_f = 44.4 \text{ Kcal/mole}$
 $= 0.65 \text{ Kcal/g}$
 $I_{sp} = 311 \text{ Sec.}$




Bicyclopropylidene
 $\Delta H_f = 76.1 \text{ Kcal/mole}$
 $= 0.95 \text{ Kcal/g}$
 $I_{sp} = 312.5 \text{ Sec.}$



Cyclopropylacetylene
 $\Delta H_f = 64.0 \text{ Kcal/mole}$
 $= 0.97 \text{ Kcal/g}$
 $I_{sp} = 311.3 \text{ Sec.}$




Bicyclopropylacetylene
 $\Delta H_f = 73.4 \text{ Kcal/mole}$
 $= 0.69 \text{ Kcal/g}$
 $I_{sp} = 307.2 \text{ Sec.}$



Dicyclopropylidene
 $\Delta H_f = 104.6 \text{ Kcal/mole}$
 $= 1.13 \text{ Kcal/g}$
 $I_{sp} = 313.4 \text{ Sec.}$


Survey of Hydrocarbons



1,5-Hexadiyne

$$\Delta H_f = 91.8 \text{ Kcal/mole}$$
$$= 1.18 \text{ Kcal/g}$$


$$I_{sp} = 311.8 \text{ Sec}$$



1,7-Octadiyne

$$\Delta H_f = 79.9 \text{ Kcal/mole}$$
$$= 0.75 \text{ Kcal/g}$$


$$I_{sp} = 308.2 \text{ Sec}$$



Quadricyclane

$$\Delta H_f = 72.2 \text{ Kcal/mole}$$
$$= 0.78 \text{ Kcal/g}$$

$$I_{sp} = 307 \text{ Sec}$$




[3]-Triangulane

$$\Delta H_f = 72.3 \text{ Kcal/mole}$$
$$= 0.77 \text{ Kcal/g}$$


$$I_{sp} = 311.4 \text{ sec}$$

Survey of Hydrocarbons


[1.1.1]Propellane and its Derivatives




$$\begin{aligned}\Delta H_f &= 83.0 \text{ Kcal/mole} \\ &= 1.25 \text{ Kcal/g} \\ \text{Isp} &= 316.6 \text{ sec}\end{aligned}$$




$$\begin{aligned}\Delta H_f &= 51.0 \text{ Kcal/mole} \\ &= 0.75 \text{ Kcal/g} \\ \text{Isp} &= 313.9 \text{ sec}\end{aligned}$$



$$\begin{aligned}\Delta H_f &= 45.0 \text{ Kcal/mole} \\ &= 0.54 \text{ Kcal/g} \\ \text{Isp} &= 311.2 \text{ sec}\end{aligned}$$




$$\begin{aligned}\Delta H_f &= 26.0 \text{ Kcal/mole} \\ &= 0.21 \text{ Kcal/g} \\ \text{Isp} &= 308.0 \text{ sec}\end{aligned}$$




$$\begin{aligned}\Delta H_f &= 95.0 \text{ Kcal/mole} \\ &= 0.70 \text{ Kcal/g} \\ \text{Isp} &= 309.9 \text{ sec}\end{aligned}$$


Selection of Target Molecules




Bicyclopropylidene
 $I_{sp} = 312.5 \text{ Sec}$



1,5-Hexadiyne
 $I_{sp} = 311.8 \text{ Sec}$




Cyclopropylacetylene
 $I_{sp} = 311.3 \text{ Sec}$




1,7-Octadiyne
 $I_{sp} = 308.2 \text{ Sec}$


Literature Methodology



P. LePerchec and J. M. Conia, *Tetrahedron Lett.* 1970, 1587




Literature Methodology




1. A.H. Schmidt, U. Schirmer and J.-M. Conia; *Chem. Ber.* **1976**, 109, 258
2. W. Weber and A.de Meijere; *Syn. Comm.* **1986**, 16, 837

Kulinkovich Reaction




O. G. Kulinkovich, S. V. Sviridov, D. A. Vasilievskii; *Synthesis* 1991, 234



1. A. de Meijere, S. I. Kozhushkov, T. Spaeth and N. S. Zefirov; *J. Org. Chem.* 1993, **58**, 502
2. S.C. Suri; *Technical Report PL-TR-97-3057*, 1997, p 26






Literature Methodology for 1,5-Hexadiyne



AFRL/PRS Methodology

- Eliminated Use of Free Halogen
- Eliminated Use of Methylene Chloride
- Eliminated Use of Liquid Ammonia/Sodium

1,5-hexadiene
1,5-hexadiene




Hazard Characteristics of Hydrocarbons

Compound	Olin Matheson Liquid Impact* <i>cc</i> Kg/cm	Julius Peters Sliding Friction* <i>cc</i> N	NOL Card GAP At Zero Card
RP-1	>200 <i>cc</i> Kg/cm	>371N	Negative
Bicyclopropylidene	>200 <i>cc</i> Kg/cm	133N	Negative
Cyclopropylacetylene	>200 <i>cc</i> Kg/cm	78N	Negative
1,5-Hexadiyne	56 <i>cc</i> Kg/cm	112N	Negative
1,7-Octadiyne	148 <i>cc</i> Kg/cm	100N	Negative

* Obtained five negative results

Proposed Mechanism of Kulinkovich Reaction



Problems	Consequences	Solution
<ul style="list-style-type: none"> Rise in temperature (Exothermic reaction) 	<ul style="list-style-type: none"> Loss of flammable solvent ($F_p = -45^\circ\text{C}$) Product rearranges to cyclopropyl ethyl ketone 	<ul style="list-style-type: none"> Perform addition of Grignard reagent below 0°C Operation is done below 30°C
<ul style="list-style-type: none"> Water contamination 	<ul style="list-style-type: none"> Decreases the concentration of Grignard reagent 	<ul style="list-style-type: none"> Purge the reactor with nitrogen gas all the time to reduce the condensation of water vapors in the reactor. Use anhydrous ether
<ul style="list-style-type: none"> High acid concentration while quenching 	<ul style="list-style-type: none"> Probability of formation of rearranged product 	<ul style="list-style-type: none"> Use of low concentration of acid
<ul style="list-style-type: none"> Gummy deposit on the wall of reactor and around cooling coil 	<ul style="list-style-type: none"> Methylcyclopropyl carboxylate entraps in the gummy material. 	<ul style="list-style-type: none"> Decrease the size of the batch. Try Continuous Process
<ul style="list-style-type: none"> By Products (Isopropanol and Methanol) 	<ul style="list-style-type: none"> Reacts with brominating reagent in the second step. 	<ul style="list-style-type: none"> Azeotrope removal of Isopropanol & methanol using ethylacetate at $\leq 50^\circ\text{C}$


Problems	Consequences	Solution
Contamination of Isopropanol/methanol	Consumption of brominating agent to form 2-bromopropane/bromomethane	Try to minimize IPA/methanol contamination in step 1. After checking GC, compensate for IPA/Methanol by adding excess of reagent
Contamination of Pyridine	Carried over to next step	Wash the product in pmethylene chloride with aqueous HCl
Distill off solvent directly from reactor	Resulted in thick solid triphenyl phosphine oxide in the reactor.	Transfer to rotary evaporator directly and remove 2/3 of dichloromethane followed by treatment with pentane to form free flowing solid

Problems	Consequences	Solution
<ul style="list-style-type: none"> • Exothermic Reaction 	<ul style="list-style-type: none"> • Loss of Product 	<ul style="list-style-type: none"> • Reaction vessel is equipped with condensor hooked to chiller at $\leq -10^{\circ}\text{C}$.
<ul style="list-style-type: none"> • Direct Distillation under high vacuum at room temperature 	<ul style="list-style-type: none"> • Loss of Product 	<ul style="list-style-type: none"> • Quenching by adding the reaction mixture into ice-water and extracted with pentane • Distilling off pentane under vacuum using water aspirator at dry ice-acetone temperature. • Putting multiple cold-traps in series
<ul style="list-style-type: none"> • Purification 	<ul style="list-style-type: none"> • 	<ul style="list-style-type: none"> • Using packed column • It further removes traces of pentane

Future Target Molecules



Bicyclopropylacetylene
Isp = 307.2_{Sec}



Bicyclopropylidenemethane
Isp = 313.4

Summary

- The synthesis of 1-cyclopropylcyclopropan-1-ol was developed by generating Grignard reagent in situ, thus avoiding handling of moisture sensitive and flammable preformed ethylmagnesium bromide.
- Three steps synthesis was used to prepare 7-8 lbs of bicyclopropylidene. There is a need to find an alternative synthetic route (maximum 2 steps) for it.
- About 200 g of 1,5-hexadiyne was synthesized using environmentally friendly process that eliminates the use of free bromine, controlled solvent dichloromethane and liquid ammonia, was worked out.
- Collected hazard data on bicyclopropylidene and 1,5-hexadiyne