

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE Technical Paper		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE				5a. CONTRACT NUMBER F04011-77-0-0045	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER DARP	
				5e. TASK NUMBER A205	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)				8. PERFORMING ORGANIZATION REPORT	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/PRS 5 Pollux Drive Edwards AFB CA 93524-7048				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
20021212 128					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Leilani Richardson
Unclassified	Unclassified	Unclassified	(A)		19b. TELEPHONE NUMBER (include area code) (661) 275-5015

5 Separate items enclosed

DAK/A

99 4 1 5

MEMORANDUM FOR PRS (In-House Contractor Publication)

FROM: PROI (STINFO)

01 April 2002

SUBJECT: Authorization for Release of Technical Information, Control Number: **AFRL-PR-ED-TP-2002-073**
Ashwani Vij (ERC) et al., "Experimental Detection of the Pentazole Anion, N₅"

Publication in Science (magazine)
(Deadline: ASAP)

(Statement A)

1. This request has been reviewed by the Foreign Disclosure Office for: a.) appropriateness of distribution statement, b.) military/national critical technology, c.) export controls or distribution restrictions, d.) appropriateness for release to a foreign nation, and e.) technical sensitivity and/or economic sensitivity.

Comments: _____

Signature _____ Date _____

2. This request has been reviewed by the Public Affairs Office for: a.) appropriateness for public release and/or b) possible higher headquarters review.

Comments: _____

Signature _____ Date _____

3. This request has been reviewed by the STINFO for: a.) changes if approved as amended, b) appropriateness of references, if applicable; and c.) format and completion of meeting clearance form if required

Comments: _____

Signature _____ Date _____

4. This request has been reviewed by PR for: a.) technical accuracy, b.) appropriateness for audience, c.) appropriateness of distribution statement, d.) technical sensitivity and economic sensitivity, e.) military/national critical technology, and f.) data rights and patentability

Comments: _____

APPROVED/APPROVED AS AMENDED/DISAPPROVED

PHILIP A. KESSEL Date
Technical Advisor
Space and Missile Propulsion Division

Experimental Detection of the Pentazole Anion, N_5^-

Ashwani Vij,^{1*} James G. Pavlovich,² William W. Wilson,¹ Karl O. Christe^{1,3*}

The pentazole anion has been generated from *para*-hydroxyphenylpentazole and identified by electrospray ionization mass spectrometry. Whereas at low collision voltages the *para*-phenoxyphenylpentazole anion undergoes stepwise N_2 elimination generating the corresponding azide and nitrene, at high collision voltages the N_5^- anion is formed. Fragmentation of the pentazole anion produces the N_3^- anion as the principal negative ion. These experiments provide the first experimental proof for the existence of the pentazole anion. They also demonstrate that under suitable reaction conditions the C-N bond in a phenylpentazole can selectively be broken with conservation of the pentazole ring, thus providing a potential synthetic route to the pentazole anion.

Nitrogen and oxygen are unique among the chemical elements. In contrast to the other elements, their homonuclear single bond energies are significantly less than one half of their double bond or one third of their triple bond energies. Consequently, homonuclear polynitrogen and polyoxygen species are thermodynamically highly unstable and the

¹ERC, Inc., Air Force Research Laboratory, Edwards Air Force Base, CA 93524, USA.

²Department of Chemistry, University of California, Santa Barbara, CA 93106, USA.

³Loker Hydrocarbon Research Institute, University of Southern California, Los Angeles, CA 90089-1661, USA.

*To whom correspondence should be addressed. E-mail: karl.christe@Edwards.af.mil

number of known compounds is very limited. Due to the highly endothermic heats of formation, their syntheses and handling present great challenges. It is therefore no surprise that for oxygen only one metastable allotrope, i. e., ozone, is known and for nitrogen none are known that can be isolated in bulk, while most other elements can exist in the form of many stable allotropes.

Polynitrogen compounds have been studied extensively for the last two decades. In view of the great experimental difficulties, most of the efforts have been limited to theoretical studies (1-9). The first major breakthrough in the synthesis area was achieved in 1999 with the synthesis of the N_5^+ cation in the form of a marginally stable AsF_6^- salt (10). Subsequently, the thermally more stable $N_5^+SbF_6^-$ was synthesized, and the crystal structure of $N_5^+Sb_2F_{11}^-$ was determined (11).

Based on Born-Haber cycle considerations, the stability of an ionic salt is governed by four factors: the lattice energy, the electron affinity of the cation, the first ionization potential of the anion, and the activation energy barriers of the ions toward decomposition. Theoretical calculations from our and other (1,5,8,13-19) research groups predict that the unknown pentazole anion (see Figure 1) has a first ionization potential and activation energy barrier toward decomposition that might be high enough to provide a stable salt. As a result, a worldwide effort is underway to synthesize this anion. Although the existence and stability of substituted pentazole ring compounds have been demonstrated successfully more than 40 years ago by Huisgen and Ugi (20-23) and substituted pentazoles have been well characterized (24-29), all attempts to prepare either the parent HN_5 molecule (30,31) or its anion, N_5^- , have so far been unsuccessful. In this paper, we wish to report the first experimental detection of this important anion.

In our pursuit of the N_5^- anion, the following strategy was employed: (i) the use of Ugi-Huisgen-type, substituted phenylpentazoles as starting materials; (ii) the transfer of maximum negative charge to the pentazole ring by the use of highly electron donating substituents on the phenyl ring in *para*-position to the pentazole substituent to increase the aromaticity and stability of the pentazole ring, while at the same time weakening the connecting C-N bond; (iii) the selective cleavage of the C-N bond while keeping the N-N bonds of the pentazole ring intact; and (iv) the use of an analytical method that is ideally suited for the generation and detection of anions. A similar approach for steps (i) – (iii) has recently been published, but attempts to cleave the C-N bond by ozonolysis were unsuccessful (32). The reasons, outlined above, prompted us to choose *para*-hydroxy (32) or *para*-dimethylamino (22) substituted phenylpentazoles as starting materials and negative ion electron spray ionization mass spectrometry (ESIMS) (33-35) as the analytical tool.

The *para*-dimethylaminophenylpentazole and *para*-hydroxyphenylpentazole were dissolved in strongly polar solvents, such as CH_3CN or a mixture of CH_3OH and CH_2Cl_2 , and infused into the spectrometer with a syringe pump. Desired negative ion peaks were mass-selected and subjected to secondary negative ion mass spectroscopy (MS/MS) at variable collision voltages.

The most interesting results were obtained with *para*-hydroxyphenylpentazole in CH_3CN solution. An intense, parent – H, peak was observed at $m/e = 162$ and mass-selected for subsequent MS/MS studies. Using a low collision voltage of –10 Volt, the $m/e = 162$ peak ($OC_6H_4N_5^-$) underwent stepwise N_2 , N_2 , and CO loss, giving rise to intense peaks with m/e values of 134 ($OC_6H_4N_3^-$), 106 ($OC_6H_4N^-$), and 78 ($C_5H_4N^-$), respectively. The loss of the first N_2 molecule is due to the opening of the pentazole ring

and produces the phenoxyazide ion. The second N_2 loss occurs from the azido group and generates a nitrene. The nitrene nitrogen can readily insert into the phenyl ring, and the resulting seven-membered ring can undergo a facile CO extrusion giving a pyridine anion. Secondary fragmentation of the $m/e = 134$ peak at a collision voltage of -30 Volt gave rise to intense peaks at $m/e = 78$ ($C_5H_4N^-$), 52 ($C_3H_2N^-$), and 50 (C_3N^-). The secondary fragmentation patterns of the $m/e = 106$ and 78 peaks at collision voltages of -75 Volt gave only a very intense m/e peak at 50 (C_3N^-).

Using high collision voltages of about -75 Volt for the secondary MS of the $m/e = 162$ peak, however, gave a very different fragmentation pattern. The only peaks observed were $m/e = 70$, 52 , 50 , and 42 (see Figure 2). The $m/e = 70$ peak can only be due to N_5^- , and the $m/e = 52$ and 50 peaks are due to $C_3H_2N^-$ and C_3N^- , respectively, and, as shown above, result from the fragmentation of the pyridine anion. The $m/e = 42$ peak is due to the azide anion, N_3^- . These results clearly demonstrate that at high collision voltages the pentazole anion is formed. The fragmentation of the $m/e = 70$ peak to N_3^- (and neutral N_2 that is not observable in the negative ion spectrum) is in accord with the theoretically predicted decomposition pathway of N_5^- (8) and further supports its identification as N_5^- . The N_3^- and N_2 fragments have been calculated at the CCSD(T)/aug-cc-pVTZ level of theory to be 60 kJ/mol lower in energy than N_5^- , with an energy barrier of 116 kJ/mol to the cycloreversion (8). These values indicate that bulk N_5^- salts should be manageable on a preparative scale. Furthermore, the formation of the N_5^- peak from a starting material containing a pentazole ring and the vibrational instability of open-chain N_5^- (8), establish beyond doubt that the observed N_5^- species must be the long-sought pentazole anion.

In summary, our results constitute the first experimental detection of the pentazole anion and demonstrate that in suitably substituted phenyl-pentazoles the C-N bond can be cleaved while leaving the pentazole ring intact. Since the substituted phenylpentazoles are easily accessible, this approach holds great promise for the bulk synthesis of N_5^- salts, and experiments in this direction are in progress in our laboratories. Our results on N_5^+ (10,11) and N_5^- , together with the recent observations of the N_4 molecule as a metastable species with a lifetime exceeding 1 microsecond (36), the observation of a new but ill-characterized poly-nitrogen species from a discharge generated nitrogen plasma (37), and exciting progress in high nitrogen compounds (29,38), indicate a bright future for experimental polynitrogen chemistry.

References and Notes

1. R. J. Bartlett, Chem. Ind. 140 (2000), and references cited therein; a compilation of data for N_2 to N_{10} can be found at <http://www.qtp.ufl.edu/~bartlett/polynitrogen.pdf>.
2. S. Fau, R. J. Bartlett, J. Phys. Chem. A **105**, 4096 (2001).
3. M. Tobita, R. J. Bartlett, J. Phys. Chem. A **105**, 4107 (2001).
4. T. M. Klapoetke, Angew. Chem. Int. Ed. **38**, 2536 (1999), and references cited therein.
5. M. N. Glukhovtsev, H. Jiao, P. v. Rague Schleyer, Inorg. Chem. **35**, 7124 (1996).
6. H. H. Michels, J. A. Montgomery, Jr., K. O. Christe, D. A. Dixon, J. Phys. Chem. **99**, 187 (1995).
7. M. W. Schmidt, M. S. Gordon, J. A. Boatz, Int. J. Quant. Chem. **76**, 434 (2000); G. Chung, M. W. Schmidt, M. S. Gordon, J. Phys. Chem. A **104**, 5647 (2000).

8. M. T. Nguyen, T. K. Ha, *Chem. Phys. Lett.* **335**, 311 (2001).
9. X. Wang, H. R. Hu, A. Tian, N. B. Wong, S. H. Chien, W. K. Li, *Chem. Phys. Lett.* **329**, 483 (2000).
10. K. O. Christe, W. W. Wilson, J. A. Sheehy, J. A. Boatz, *Angew. Chem. Int. Ed.* **38**, 2004 (1999).
11. A. Vij, W. W. Wilson, V. Vij, F. S. Tham, J. A. Sheehy, K. O. Christe, *J. Am. Chem. Soc.* **123**, 6308 (2001).
12. W. W. Wilson, A. Vij, V. Vij, M. Gerken, S. Schneider, T. Schroer, K. O. Christe, unpublished results.
13. L. Gagliardi, G. Orlandi, S. Evangelisti, B. O. Roos, *J. Chem. Phys.* **114**, 10733 (2001).
14. M. Lein, J. Frunzke, A. Timoshkin, G. Frenking, *Chem. Eur. J.* **7**, 4155 (2001).
15. S. Fau, K. J. Wilson, R. J. Bartlett, *J. Phys. Chem.*, in press.
16. M. T. Nguyen, M. Sana, G. Leroy, J. Elguero, *Can. J. Chem.* **61**, 1435 (1983).
17. M. T. Nguyen, M. A. McGinn, A.F. Hegarty, J. Elguero, *Polyhedron*, **4**, 1721 (1985).
18. V. A. Ostrovskii, G. B. Erusalimskii, M. B. Shcherbinin, *Russ. J. Org. Chem.* **31**, 1284 (1995).
19. M. N. Glukhovtsev, P. v. R. Schleyer, C. Maerker, *J. Phys. Chem.* **97**, 8200 (1993).
20. R. Huisgen, I. Ugi, *Angew. Chem.* **68**, 705 (1956); *Chem. Ber.* **90**, 2914 (1957).
21. I. Ugi, R. Huisgen, *Chem. Ber.* **91**, 531 (1958).
22. I. Ugi, H. Perlinger, L. Behringer, *Chem. Ber.* **91**, 2324 (1958).
23. I. Ugi, *Angew. Chem.* **73**, 172 (1961).
24. J. D. Wallis, J. D. Dunitz, *J. Chem. Soc., Chem. Commun.* 910 (1983).


25. M. Witanowski, L. Stefaniak, H. Januszewski, K. Bahadur, G. A. Webb, *J. Cryst. Mol. Struct.* **5**, 137 (1975).
26. R. Mueller, J. D. Wallis, W. v. Philipsborn, *Angew. Chem. Int. Ed.* **24**, 513 (1985).
27. R. N. Butler, S. Collier, A. F. M. Fleming, *J. Chem. Soc., Perkin Trans. 2*, 801 (1996).
28. R. N. Butler, A. Fox, S. Collier, L. A. Burke, *J. Chem. Soc., Perkin Trans. 2*, 2243 (1998).
29. A. Hammerl, T. M. Klapoetke, *Inorg. Chem.* **41**, 906 (2002).
30. R. Janoschek, *Angew. Chem. Int. Ed.* **32**, 230 (1993).
31. K. F. Ferris, R. J. Bartlett, *J. Am. Chem. Soc.* **114**, 8302 (1992).
32. V. Benin, P. Kaszynski, J. G. Radziszewski, *J. Org. Chem.* **67**, 1354 (2002).
33. M. Yamashita, J. B. Fenn, *J. Chem. Phys.* **88**, 4451 (1984); C. M. Whitehouse, R. N. Dreyer, M. Yamashita, J. B. Fenn, *Anal. Chem.* **57**, 675 (1985).
34. R. B. Cole (ed), *Electrospray Ionization Mass Spectrometry* (Wiley-Interscience, New York, NY, 1997).
35. B. H. Lipshutz, K. L. Stevens, B. James, J. G. Pavlovich, J. P. Snyder, *J. Am. Chem. Soc.* **118**, 6796 (1996).
36. F. Cacace, G. de Petris, A. Troiani, *Science* **295**, 480 (2002).
37. J. P. Zheng, J. Waluk, J. Spanget-Larsen, D. M. Blake, J. G. Radziszewski, *Chem. Phys. Lett.* **328**, 227 (2000).
38. D. E. Chavez, M. A. Hiskey, R. D. Gilardi, *Angew. Chem. Int. Ed.* **39**, 1791 (2000).
39. This work was financially supported by the Defense Advanced Research Project Agency, the Air Force Office of Scientific Research, and the National Science Foundation. The authors are grateful to Drs. Robert Corley, Arthur Morrish, Don

Woodbury, and Michael Berman for their steady support, to Prof. W. Kaska for the use of the ESIMS, and to Drs. Michael Gerken, Thorsten Schroer, Stefan Schneider, Ralf Haiges, and Ross Wagner for support and stimulating discussions.

FIGURE CAPTIONS

1. Minimum energy structure of the planar D_{5h} pentazole anion from ref.1, calculated at the CCSD(T)/aug-cc-pVTZ level of theory, bond length in Å.

2. Negative ion mass spectrum of the mass-selected $m/e = 162$ peak due to $\text{OC}_6\text{H}_4\text{N}_5^-$, recorded at a collision voltage of -75 Volt. The peaks at $m/e = 70$ and 42 are due to the N_5^- anion and its decomposition product N_3^- , respectively, while the peaks at 52 and 50 are the $\text{C}_3\text{H}_2\text{N}^-$ and C_3N^- fragments, respectively, resulting from the breakdown of the phenoxyazide anion.



TOF Product (162.1): 0.017 to 4.984 min from Sample 2 (pOH, AcCN) of kas032102qp.wiff, Smoothed

Max. 0.416666

MS/MS
162
-75V

