

A NEURAL NETWORK FOR ESTIMATION OF AORTIC PRESSURE FROM THE RADIAL ARTERY PRESSURE PULSE

A. Qasem, A. Avolio, G. Frangakis

Graduate School of Biomedical Engineering, New South Wales University, Sydney, NSW, Australia

Abstract. A neural network is developed to estimate aortic pressure from the radial artery pressure pulse waveform. Invasively measured aortic and radial artery pressure in 51 adult subjects were used to train the network. Tests in a separate group of 21 subjects of similar age range showed a high correlation ($r > 0.93$) between measured and estimated systolic, diastolic and pulse pressure, with mean absolute errors (%) of 2.5 ± 0.3 , 3.5 ± 0.6 , 4.8 ± 0.7 respectively. This method has potential applications in obtaining accurate estimates of central aortic pressure values from non-invasive radial artery pulse measurements. Such neural networks can be trained in specific subgroups (eg diabetics) to improve the estimation of central aortic pressure from the peripheral pulse.

Keywords - Neural Network, Radial and Aortic Pressure, Transfer Function.




Fig 1 Amplitude of transfer function between aorta and radial artery.

I. INTRODUCTION

There is increasing interest in stratification of cardiovascular risk factors in subgroups of the population with emphasis on pulse pressure and central aortic pressure values [1]. The conventional measures of systolic and diastolic pressure in the arm do not give an accurate measure of pressure in the aorta because of the amplification of the pressure pulse as it propagates from the heart to the peripheral arteries [2,3]. However, since aortic pressure can only be measured essentially by invasive means, methods have been developed to estimate aortic pressure from non-invasive signals such as peripheral pressure waveforms, or simultaneous pressure and flow signals [4, 5-10]. One of these methods, the general transfer function method (GTF), relates aortic to radial pressure waveforms in the frequency domain (Fig. 1). This was obtained using invasive measurements and has been shown to be applicable in adults to estimate central aortic pressure from the radial pulse with reasonable accuracy [4,10].

While these general techniques take into account most of the morphological waveform changes, such as frequency depended amplification (Fig. 1), they do not account for additional effects of age or other conditions that may be specific to different risk groups [11]. This study attempts to apply neural network techniques to develop strategies for accurate estimation of central aortic pressure so that individual parameters are taken into account

II. METHODOLOGY

An artificial neural network is a mathematical representation of a real network in terms of neurons, function and connections. A neural network consists of neurons which relate multiple inputs to a single output using a mathematically defined function. All the inputs to the neuron are multiplied by variables called weights. Neurones can be arranged vertically in one layer and a neural network can have more than one layer (Fig. 2). In this case we will use a three layer neural network. The method involves a learning or training process where weights are changed according to a set of inputs and outputs using iterative methods. As a result, the weights will be such that it will produce the lowest error in the learning set. This type of neural network is described as a back-propagation network [12,13]

The inputs to the neural network in this study were age, height, weight, gender, heart period, radial systolic, mean and pulse pressure. These inputs were chosen since they are known influence blood pressure [3]. The outputs were aortic systolic and diastolic pressure.

Simultaneous invasive recordings of aortic and radial pressure waveforms were obtained for 72 subjects. A learning set was obtained in 51 subjects (16F, 35M, Age 38-84, Ht 150-189cm, Wt 54-119Kg), and the test set in 21 subjects (2F, 19M, Age 39-76, Ht 156-185cm, Wt 60-119 Kg). Each recording was of about 10 to 12 pulses from which an averaged pressure waveform was calculated.

Report Documentation Page

Report Date 25 Oct 2001	Report Type N/A	Dates Covered (from... to) -
Title and Subtitle A Neural Network for Estimation of Aortic Pressure From The Radial Artery Pressure Pulse		Contract Number
		Grant Number
		Program Element Number
Author(s)	Project Number	
	Task Number	
	Work Unit Number	
Performing Organization Name(s) and Address(es) Graduate School of Biomedical Engineering New South Wales University Sydney, NSW, Australia		Performing Organization Report Number
Sponsoring/Monitoring Agency Name(s) and Address(es) US Army Research, Development & Standardization Group (UK) PSC 802 Box 15 FPO AE 09499-1500		Sponsor/Monitor's Acronym(s)
		Sponsor/Monitor's Report Number(s)
Distribution/Availability Statement Approved for public release, distribution unlimited		
Supplementary Notes Papers from 23rd Annual International Conference of the IEEE Engineering in Medicine and Biology Society, October 25-28, 2001, held in Istanbul, Turkey. See also ADM001351 for entire conference on cd-rom., The original document contains color images.		
Abstract		
Subject Terms		
Report Classification unclassified	Classification of this page unclassified	
Classification of Abstract unclassified	Limitation of Abstract UU	
Number of Pages 3		

From an investigation of different neural network structures ($n=26$), the final structure selected was one with 15 neurons in layer 1, 5 neurones in layer 2 and 2 neurones in layer 3, since this structure produced the lowest error of all those investigated.




Fig. 2. Neural network with 3 layers and n input X and m outputs Y

III. RESULTS

Table 1 shows the low average error in aortic systolic, diastolic and pulse pressure produced by the neural network when used on the test set. Figs. 3,4,5 show the high correlation between the estimated and measured aortic pressure. R^2 for systolic, diastolic and pulse pressure are 0.94, 0.93 and 0.95 respectively.

TABLE 1

Error in % in estimating aortic pressure using neural network.

%	Systolic	Diastolic	Pulse
Mean $\pm$ SEM	2.5 $\pm$ 0.3	3.5 $\pm$ 0.6	4.8 $\pm$ 0.7
Max	5	10.3	11.2




Fig 3. Estimated and measured aortic systolic pressure




Fig 4. Estimated and measured aortic diastolic pressure




Fig 5. Estimated and measured aortic pulse pressure

IV. DISCUSSION

Neural networks can be trained to solve complex and non-linear problems where it is difficult to do so with conventional mathematical methods. Hence, because of the non-linear and complex inter-relationship between aortic and radial pressure, neural networks may be applied to estimations of central aortic pressure using a single input signal from a peripheral accessible site such as the radial artery.

This investigation has shown that a neural network can be trained in a group of subjects to derive aortic systolic and diastolic pressure values from a calibrated peripheral pulse. The neurons utilised a normalized sigmoid function of the form $y = \log\text{Sig}(n)$, where n is the input and y the output with limits between 0 and 1. It has been shown that this function is efficient in converging to a minimum error [13].

As more invasive pressure measurements become available in subgroups of subjects with specific cardiovascular risk factors or disorders, the method can be refined to estimate weighting coefficients for that particular group. It will provide a complimentary method to the current

methods employing the GTF Furthermore, this method can be developed to predict other aortic parameters related to the pressure waveform to provide a better non-invasive assessment of the cardiovascular system.

V. CONCLUSION

The results of this investigation show that neural networks have potential applications in obtaining accurate estimates of central aortic pressure values from non-invasive radial artery pulse measurements. Such neural networks can be trained in specific subgroups (eg diabetics) to improve the estimation of central aortic pressure from the peripheral pulse.

ACKNOWLEDGMENT

We acknowledge the contribution of Dr. Alfredo Pauca of Wakeforest University in North Carolina USA, who provided us with the invasive recordings of radial and aortic pressure waveform.

REFERENCES

- [1] Franklin SS, Khan SA, Wong ND, Larson MG, Levy D. "Is pulse pressure useful in predicting risk for coronary heart Disease?" The Framingham heart study. *Circulation*. 1999 27;100(4):354-60.
- [2] Karamanoglu M, Gallagher D, Avolio AP, O'Rourke MF, "Pressure wave propagation in a multibranched model of the human upper limb", *American Journal of Physiology*, 269, 38: H1363-H1369, 1995
- [3] M O'Rourke, R. Kelly, A. Avolio, *The Arterial Pulse*, Philadelphia, Lea & Febiger, 1992
- [4] M. Karamanoglu, M. O'Rourke, A. Avolio, R. Kelly, An Analysis of the relationship between central aortic and peripheral pressure waves in man, *European Heart Journal*, 1993; 14, pp 160-7.
- [5] M. O'Rourke, *Method for ascertaining the pressure pulse and related parameters in the ascending aorta from the contour of the pressure pulse in the peripheral arteries*, U.S Patent No. 5265011, Nov. 23, 1993 .
- [6] A. Pauca, S Wallenhaupt, N Kon, W Tucker, Does radial artery pressure accurately reflect aortic pressure ?, *Chest*, 1992, 102, pp 1193-98.
- [7] T. Sharir, A. Marmor, C. Ting, W Chen, C Liu, M Chang, F. Yin, D. Kass., Validation of a method for non-invasive measurement of central arterial pressure, *Hypertension*, 1993,21(1):74-82.
- [8] N Stergiopoulos, B Westerhof, N Westerhof, Physical basis of pressure transfer from periphery to artery: a model based study, *American Journal of Physiology*, 1998,274:43, pp H1386-92
- [9] C. Chen, E. Nevo, B Fetics, P Pak, F, Yin, L Maughan, D Kass, Estimation of central aortic pressure waveform by mathematical transformation of radial tonometry pressure, *Circulation*, 95(7):1827-36.
- [10] B Fetics, E Nevo, C. Chen, D Kass, Parametric model derivation of transfer function for noninvasive estimation of aortic pressure by radial tonometry, *IEEE Transactions on Biomedical Engineering*, 1999; 46:6: pp698-706.
- [11] R Kelly, C Hayward, A Avolio, M O'Rourke, Noninvasive determination of age-related changes in the human arterial pulse, *Circulation*, 1989; 80:pp1652-1659
- [12] M Caudill, C. Butler, *Understanding Neural Networks: Computer Explorations*, Vol 1 and 2, Cambridge, MA: the MIT Press, 1992.
- [13] Demuth H, Beale M, *MATLAB Neural Network Toolbox User's Guide*, The Math Works Inc. Natick MA, USA 1994