

AD \_\_\_\_\_

Award Number: DAMD17-99-1-9123

TITLE: Molecular Markers in Hereditary Breast Cancer

PRINCIPAL INVESTIGATOR: Olufunmilayo I. Olopade, M.D.

CONTRACTING ORGANIZATION: The University of Chicago  
Chicago, Illinois 60637

REPORT DATE: October 2002

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20030411 042

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE October 2002	3. REPORT TYPE AND DATES COVERED Final (15 Sep 99 - 14 Sep 02)
----------------------------------	--------------------------------	---

4. TITLE AND SUBTITLE Molecular Markers in Hereditary Breast Cancer	5. FUNDING NUMBERS DAMD17-99-1-9123
--	--

6. AUTHOR(S) : Olufunmilayo I. Olopade, M.D.
---

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The University of Chicago Chicago, Illinois 60637  E-Mail: folopade@medicine.bsd.uchicago.edu	8. PERFORMING ORGANIZATION REPORT NUMBER
--	---

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012	10. SPONSORING / MONITORING AGENCY REPORT NUMBER
---	---

11. SUPPLEMENTARY NOTES

12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited	12b. DISTRIBUTION CODE
---	------------------------

13. Abstract (*Maximum 200 Words*) (*abstract should contain no proprietary or confidential information*)

We are entering a new era of medicine where genetic markers are going to be used to make clinical management decisions. My long term career goal is to further our understanding of the genetic alterations which characterize human breast cancer in a way that will eventually lead to early diagnosis, more effective treatment or prevention of the disease. The promise of research into breast cancer genetics is that it will provide us with new insights into the etiology of breast cancer that can be translated to strategies for early diagnosis and treatment for the larger population of women who develop breast cancer without having a genetic predisposition.

14. SUBJECT TERMS: breast cancer	15. NUMBER OF PAGES 8
	16. PRICE CODE

17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited
--	---	--	---

## Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4-6
Key Research Accomplishments.....	6
Reportable Outcomes.....	6-7
Conclusions.....	7
References.....	7-8
Appendices.....	NA

## INTRODUCTION

As a physician-scientist, I have had extensive training in clinical oncology and in molecular biology and genetics; I am ideally positioned to bridge the gap between the two. The academic award has represented an outstanding opportunity for me to critically appraise the emerging role of genetics in clinical breast cancer care and forge new avenues of research. Toward this goal, I plan to accomplish the following during the period of my academic award.

1) perform a thorough review of the cytogenetic and molecular genetics literature to identify potential chromosomal regions that may harbor genes whose abnormal function is critically involved in the development of breast cancer.

2) develop a robust panel of markers that can be used for clinical correlative studies of hereditary breast cancers.

3) develop a tissue repository composed of biological specimens from 500 patients with inherited breast cancer (e.g fresh frozen tumor specimens, or paraffin embedded tumor specimens and normal blood lymphocytes, DNA and sera whenever possible).

Using these unique resources, my future studies will characterize the molecular pathways that allow a normal breast cell to become cancerous in individuals who are genetically predisposed. I will also develop longitudinal follow up studies to correlate clinical outcomes with molecular characterization and epidemiologic risk factors. These studies will no doubt lead to an improved understanding of the biology of breast cancer which will ultimately translate into more effective therapies.

### Task I

**perform a thorough review of the cytogenetic and molecular genetics literature to identify potential chromosomal regions that may harbor genes whose abnormal function is critically involved in the development of breast cancer.**

This year we published two reviews on the genetics of breast cancer. In the next year, we are completing two manuscripts that will focus on the chromosomal abnormalities and genetic alterations in breast cancer.

### **Publications**

**Olopade OI, Pichert G.** Cancer genetics in oncology practice. *Ann Oncol* 2001 Jul;12(7):895-908

**White, M, Bradbury, A, Olopade OI.** Breast and Ovarian Cancer Risk Assessment and Risk Reduction: Strategies for the Primary Care Physician. In Press *Women's Health*

### Task II

**develop a robust panel of markers that can be used for clinical correlative studies of hereditary breast cancers.**

We have developed several probes for fluorescent in situ hybridization and have begun to apply these probes to a panel of breast tumors in our tumor bank.

**a) Dissection of Cooperating Oncogenes involved in *BRCA1* tumor progression**

To examine whether amplification of *HER-2/neu* contributes to the aggressive biology of *BRCA1*-associated tumors, we performed fluorescence *in situ* hybridization (FISH) on formalin-fixed paraffin-embedded breast tumor tissue sections from 53 *BRCA1* mutation carriers and 41 randomly selected age-matched sporadic breast cancer cases. Although *BRCA1*-associated and sporadic tumors were equally likely (19% versus 22%) to exhibit *HER-2/neu* amplification (defined as a ratio of *HER-2/neu* copies to chromosome 17 centromere (*CEP17*) copies  $\geq 2$ ), 6 (15%) of the sporadic tumors were highly amplified (defined as a ratio  $\geq 5$ ) versus none of the *BRCA1*-associated tumors ( $p = 0.048$ ). *HER-2* protein overexpression as measured by immunohistochemical analysis (IHC) was not observed among the *BRCA1*-associated cases ( $p = 0.042$ ). Four out of 21 (19%) sporadic tumors exhibited strong membranous staining of *HER-2* (intensity level of 3+) as compared to 0/39 *BRCA1*-associated tumors. Our data suggest that a germ line mutation in the *BRCA1* tumor suppressor gene is associated with a significantly lower level of *HER-2/neu* amplification. Thus, it is possible that *BRCA1*-associated and *HER-2/neu*-highly amplified tumors progress through distinct molecular pathways and the aggressive pathologic features of *BRCA1*-associated tumors appear unrelated to amplification of the adjacent *HER-2/neu* oncogene (Grushko et al. 2002).

We further examined whether *MYC* amplification contributes to tumor progression in *BRCA1*-associated human breast cancer, and analyzed tumors using a *MYC/CEP8* assay on formalin-fixed paraffin-embedded tumor tissues from 28 women with known deleterious germ line *BRCA1* mutations and 49 sporadic cases, including 18 with hypermethylation of the *BRCA1* gene promoter. We observed a *MYC/CEP8* amplification ratio  $\geq 2$  in 17 of 28 (61%) *BRCA1*-mutated and in 13 of 49 (27%) sporadic tumors ( $P = 0.009$ ). Of the 13 sporadic cases with *MYC* amplification, 7 (54%) were *BRCA1*-methylated. In total, *MYC* amplification was found in the majority of tumors with inactivated *BRCA1* (24/46, 52% versus 6/31, 19%;  $p = 0.01$ ). We concluded that *MYC* oncogene amplification is associated with multi-step tumor progression in both *BRCA1*-associated hereditary and methylated sporadic tumors, which suggests that *BRCA1* promoter methylation may be an early step in the development of some sporadic cancers. We hypothesize that *BRCA1* may function as a tumor suppressor gene in part by regulating *MYC* oncogenic activity.

#### **b) . Hypermethylation of *BRCA1* and ER promoter**

We assessed *BRCA1* and estrogen receptor (ER) promoter methylation in 5 breast cancer cell lines and 132 primary breast tissues by Methylation-Specific (M-PCR). *BRCA1* and ER expression were determined in breast tumor cell lines and primary tissues by RT-PCR. In addition, we performed FISH using *BRCA1* and *CEP17* probes on both sporadic and *BRCA1*-associated hereditary breast cancer. We observed *BRCA1* methylation in the UACC-3199 positive control cell line and in 39 of 132 sporadic (29.5%) tumors. *BRCA1* methylation was correlated with chromosome 17 aneusomy and down-regulation or complete absence of the transcript. *BRCA1* methylation correlated inversely with age of onset: 40% of tumors from cases under 55 years old were methylated vs. 25% of cases over 55 years old (Table 4). The methylated cases were equally distributed among all histological types and there was no difference in the proportion of African American women (27.5%) vs. non-Hispanic White women with methylated tumors (28.6%). The majority of *BRCA1*-methylated cases (79%) were ER (-) and/or ER methylated. *MYC* and *HER2/neu* amplification in methylated tumors were intermediate in values between hereditary *BRCA1*-associated and sporadic unmethylated tumors, suggesting that *BRCA1* methylation might be incomplete in some tumors. These results suggest that silencing of the *BRCA1* gene by methylation occurs in a significant proportion of sporadic breast cancers and may be an early event during tumor progression (Wei et al. in preparation). There may be a slight difference in the proportion of black women with methylated tumors.

#### **Publications**

Grushko TA, Blackwood MA, Schumm PL, Hagos FG, Adeyanju MO, Feldman MD, Sanders MO, Weber BL, Olopade OI. Molecular-cytogenetic analysis of HER-2/neu gene in BRCA1-associated breast cancers. *Cancer Res.* 2002 Mar 1;62(5):1481-8.

Min-Jie Wei<sup>1</sup>, Tatyana Grushko<sup>1</sup>, Soma Das<sup>2</sup>, James Dignam<sup>3</sup>, Fitsum Hagos<sup>1</sup>, Lise Sveen<sup>1</sup>, James Fackenthal<sup>1</sup> and Olufunmilayo I Olopade<sup>1</sup>. Methylation of the *BRCA1* Promoter in Sporadic Breast Cancer Related to *BRCA1* Copy Number and Pathologic Features. Manuscript in preparation

Tatyana A. Grushko,, James J. Dignam, , Soma Das, Anne Blackwood, Charles M. Perou, , April J. Adams, , Fitsum G. Hagos, Lise Sveen , Karin K. Ridderstråle, Kristin Anderson, Barbara L. Weber Olufunmilayo I. Olopade, M.D. *MYC* is amplified in *BRCA1*-associated breast cancers. Manuscript in preparation

### **Task III**

**develop a tissue repository composed of biological specimens from 500 patients with familial or hereditary breast cancer (e.g fresh frozen tumor specimens, or paraffin embedded tumor specimens and normal blood lymphocytes, DNA and sera whenever possible).**

We have developed a clinical protocol for the tumor bank. The protocol has not yet been approved by the DOD Human Subjects Review Panel. Hence we have not enrolled any patients specifically to this study. However, we have identified collaborators and other sources of tumor materials that will be ready and available for recruitment once our study is approved. Our protocol is still awaiting approval by the DOD.

### **KEY RESEARCH ACCOMPLISHMENTS:**

We are defining important pathways in BRCA1 tumor progression.

### **REPORTABLE OUTCOMES:**

Academic Productivity in 2002.

1. Olopade OI, Fackenthal JD, Dunston G, Tainsky MA, Collins F, Whitfield-Broome C. Breast cancer genetics in African Americans. *Cancer.* 2003 Jan 1;97(1 Suppl):236-45. PMID: 12491487

2: Narod SA, Dube MP, Klijn J, Lubinski J, Lynch HT, Ghadirian P, Provencher D, Heimdal K, Moller P, Robson M, Offit K, Isaacs C, Weber B, Friedman E, Gershoni-Baruch R, Rennert G, Pasini B, Wagner T, Daly M, Garber JE, Neuhausen SL, Ainsworth P, Olsson H, Evans G, Osborne M, Couch F, Foulkes WD, Warner E, Kim-Sing C, Olopade O, Tung N, Saal HM, Weitzel J, Merajver S, Gauthier-Villars M, Jernstrom H, Sun P, Brunet JS. Oral Contraceptives and the Risk of Breast Cancer in BRCA1 and BRCA2 Mutation Carriers. *J Natl Cancer Inst.* 2002 Dec 4;94(23):1773-9.

3: Huo Z, Giger ML, Olopade OI, Wolverton DE, Weber BL, Metz CE, Zhong W, Cummings SA. Computerized analysis of digitized mammograms of BRCA1 and BRCA2 gene mutation carriers. *Radiology.* 2002 Nov;225(2):519-26. PMID: 12409590 [PubMed - indexed for MEDLINE]

4: Antman K, Abraido-Lanza AF, Blum D, Brownfield E, Cicatelli B, Debor MD, Emmons K, Fitzgibbon M, Gapstur SM, Gradishar W, Hiatt RA, Hubbell FA, Joe AK, Klassen AC, Lee NC, Linden HM, McMullin J, Mishra SI, Neuhaus C, Olopade FI, Walas K. Reducing disparities in breast cancer

survival: a Columbia University and Avon Breast Cancer Research and Care Network Symposium. *Breast Cancer Res Treat.* 2002 Oct;75(3):269-80.  
PMID: 12353816

6: Fackenthal JD, Cartegni L, Krainer AR, Olopade OI. BRCA2 T2722R is a deleterious allele that causes exon skipping. *Am J Hum Genet.* 2002 Sep;71(3):625-31.  
PMID: 12145750

7: Euhus DM, Smith KC, Robinson L, Stucky A, Olopade OI, Cummings S, Garber JE, Chittenden A, Mills GB, Rieger P, Esserman L, Crawford B, Hughes KS, Roche CA, Ganz PA, Seldon J, Fabian CJ, Klemp J, Tomlinson G. Pretest prediction of BRCA1 or BRCA2 mutation by risk counselors and the computer model BRCAPRO. *J Natl Cancer Inst.* 2002 Jun 5;94(11):844-51.

8: Rebbeck TR, Lynch HT, Neuhausen SL, Narod SA, Van't Veer L, Garber JE, Evans G, Isaacs C, Daly MB, Matloff E, Olopade OI, Weber BL. Prophylactic oophorectomy in carriers of BRCA1 or BRCA2 mutations. *N Engl J Med.* 2002 May 23;346(21):1616-22.

9: Grushko TA, Blackwood MA, Schumm PL, Hagos FG, Adeyanju MO, Feldman MD, Sanders MO, Weber BL, Olopade OI. Molecular-cytogenetic analysis of HER-2/neu gene in BRCA1-associated breast cancers. *Cancer Res.* 2002 Mar 1;62(5):1481-8.

## CONCLUSIONS:

The observed similarities between *BRCA1*-mutated and *BRCA1*-methylated tumors led us to propose a tumor progression model in which early loss of *BRCA1* causes defects in chromosome structure, cell division, and viability, so that a *BRCA1*-deficient cell must acquire additional alterations that overcome these problems and presumably force tumor evolution down a limited set of pathways. Our FISH results are consistent with data from DNA microarray studies that suggest that breast cancers arising in the setting of germ line *BRCA1* mutations have unique gene expression profiles, and sporadic tumors with methylated *BRCA1* may be misclassified with the *BRCA1*-mutation-positive group. A review of the set of genes published by Heldenfalk et al. (Heldenfalk et al. 2001) and in the recent paper by van 't Veer et al. (van 't Veer et al. 2002) demonstrated that *MYC* on 8q was overexpressed in *BRCA1* mutation carriers. In addition, the *BRCA1* mutant tumors we have studied appear to have a profile that is most consistent with the basal-like subtype suggested by Perou et al. (Perou et al. 2000), based upon the following observations. First, both (meaning sporadic basal-like tumors and *BRCA1* mutant tumors) tend to be high grade, ER/PR negative and *HER2/neu*-negative, and both show the high expression and/or amplification of *MYC*. In fact, *MYC* emerged as one of the most relevant genes that defined the basal-like group and was expressed more than 2-4 fold above background in the majority of cases (Perou, unpublished results). Moreover, we have previously shown that *BRCA1*-mutated tumors express specific basal cytokeratins in a manner suggestive of an ER-negative basal-like epithelial cell of origin (Olopade and Grushko 2001) and are never associated with high levels of *HER-2/neu* amplification (Grushko et al. 2002). Therefore, it is reasonable to suggest that *BRCA1*-mutated tumors are mostly basal-like (ER-, HER2-) and that *MYC* amplification/overexpression further defines *BRCA1*-deficient tumor cells.

## REFERENCES:

van 't Veer LJ, Dai H, van de Vijver MJ, He YD, Hart AA, Mao M, Peterse HL, van der Kooy K, Marton MJ, Witteveen AT, Schreiber GJ, Kerkhoven RM, Roberts C, Linsley PS, Bernards R, Friend SH. Gene expression profiling predicts clinical outcome of breast cancer.

Nature. 2002 Jan 31;415(6871):530-6.

Hedenfalk I, Duggan D, Chen Y, Radmacher M, Bittner M, Simon R, Meltzer P, Gusterson B, Esteller M, Kallioniemi OP, Wilfond B, Borg A, Trent J.

Gene-expression profiles in hereditary breast cancer.

N Engl J Med. 2001 Feb 22;344(8):539-48.

Perou CM, Sorlie T, Eisen MB, van de Rijn M, Jeffrey SS, Rees CA, Pollack JR, Ross DT, Johnsen H, Akslen LA, Fluge O, Pergamenschikov A, Williams C, Zhu SX, Lonning PE, Borresen-Dale AL, Brown PO, Botstein D.

Molecular portraits of human breast tumours.

Nature. 2000 Aug 17;406(6797):747-52.

**APPENDICES:**

N/A