

Award Number: DAMD17-00-1-0095

TITLE: Characterization of SIRPs in Prostate Cancer Cells

PRINCIPAL INVESTIGATOR: William E. Seaman, M.D.

CONTRACTING ORGANIZATION: Northern California Institute for Research
and Education
San Francisco, California 94121

REPORT DATE: March 2003

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20031029 074

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE March 2003	3. REPORT TYPE AND DATES COVERED Annual (15 Feb 02 - 14 Feb 03)	
4. TITLE AND SUBTITLE Characterization of SIRPs in Prostate Cancer Cells		5. FUNDING NUMBERS DAMD17-00-1-0095	
6. AUTHOR(S) William E. Seaman, M.D.			
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Northern California Institute for Research and Education San Francisco, California 94121 E-Mail: bseaman@medicine.ucsf.edu		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES			
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited			12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) <p>Signal regulatory proteins (SIRPs include SIRPβ1, which activates cells, and SIRPα1, which inhibits the cellular response to several growth factors, and which regulates cell adhesion and spreading.</p> <p>We demonstrated by PCR that 3 of 3 prostate cancer cell lines (PC-3, DU-145 and LNCaP) express transcripts for SIRPs. Under this contract, we generated a monoclonal antibody that recognizes both SIRPβ1 and SIRPα1, thereby confirming the expression of SIRPs on PC-3 cells and, to a lesser extent on DU-145 cells. The receptor could not be detected on LNCaP cells. We have since shown by PCR, Western blotting, and by surface staining that PC-3 and DU-145 cells express SIRPα1 but not SIRPβ. We find that they also express the tyrosine phosphatase, SHP-2, and that SHP-2 binds to SIRPα1 when it is phosphorylated, demonstrating that this pathway for the function of SIRPα1 is intact. We have created constructs of epitope-tagged SIRPα1, either intact or with mutations that would alter SHP-2 binding, in order to study its function in PC-3 cells.</p> <p>We have also worked in particular on the characterization of the SIRPα1 protein in prostate cancer cells. Is there more than one form, due either to alternate splicing or to post-translational modification? These studies have proved challenging, but we expect to complete them, along with all of the objectives of the contract, over the coming year (no-cost extension).</p>			
14. SUBJECT TERMS signaling, Prostate Cancer, Growth Factor, Signal Regulatory Protein SHPS-1			15. NUMBER OF PAGES 7
			16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited

Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	5
Reportable Outcomes.....	6
Conclusions.....	6
References.....	6
Appendices.....	

INTRODUCTION

Our Studies are based on our identification by PCR of transcripts for signal regulator protein $\alpha 1$ (SIRP $\alpha 1$) in three prostate cancer cell lines, PC-3, LNCaP, and DU-145.

We proposed six objectives:

1. Examine SIRP transcript size and expression level in different prostate cancer cell lines by Northern blotting.
2. Use RT-PCR to obtain and sequence full-length SIRP transcripts from the prostate cancer cell lines PC-3, DU-145, and LNCaP.
3. Use hybridization to screen cDNA libraries from these prostate cancer cell lines to identify additional SIRP cDNA clones.
4. Overexpress wild-type and mutated SIRPs in prostate cancer lines and, as a control, in NIH3T3 fibroblasts, to assess their effect on the cellular growth response to EGF. These will include studies of the effect of cross-linking SIRPs in cell growth. Mutational studies will include: for alpha SIRPs, mutate the cytoplasmic tyrosine to phenylalanine, for beta SIRPs, mutate the transmembrane lysine to alanine.
5. Assess the effect of EGF on the phosphorylation of SIRPs in prostate cancer cells and of associated proteins obtained by co-immunoprecipitation.
6. Produce monoclonal antibodies (mAbs) against SIRPs.

BODY

Signal regulatory proteins (SIRPs, also known as SHPS-1, BIT, p84, and Myd-1) are normally expressed on certain hematopoietic cells and some brain cells (1-3). SIRP $\beta 1$ activates cells and is expressed on cells of monocyte/macrophage lineage. Its ligand is unknown. SIRP $\alpha 1$ inhibits the response of several cell types to growth factors (1), and it regulates integrin-mediated cell adhesion and spreading (4,5). Its ligand is CD47 (integrin associated protein)(6,7). SIRP $\beta 1$ and SIRP $\alpha 1$ are highly homologous in their extracellular portions, which include three immunoglobulin (Ig)-like domains (one V and two C domains). By alternative splicing, SIRP $\alpha 1$ can also be expressed with a single Ig-like (V) domain (8). Phosphorylated SIRP $\alpha 1$ binds SHP-2, a tyrosine phosphatase that is widely distributed (1). Thus, expression of SIRPs on tumor cells might be functional and could regulate the response to growth factors and/or the capacity of tumors to invade.

This report is for the third year of our studies. During the first year, we: (i) produced monoclonal antibodies to SIRPs (cross-reactive with both SIRP $\alpha 1$ and SIRP $\beta 1$), (ii) used the antibodies to confirm surface expression of SIRPs on PC-3 and DU-145 prostate cancer cells (LNCaP did not stain with mAb, but had only low levels of transcripts by PCR), and (iii) stably overexpressed SIRP $\alpha 1$ and SIRP $\beta 1$ on PC-3 cells. These findings were published as an abstract for the Annual Meeting of the American Association for Cancer Research, March, 2001 (attached).

During the second year, we: (i) completed objective one by performing Northern blotting of RNA from PC-3, DU-145, and LNCaP cells using, as a probe, a PCR product covering most of the extracellular domain, which revealed (for PC-3 and DU-

145) a dominant band at ~3.5kb and a secondary band at ~2.2kb, similar to transcripts in the U373 glioblastoma cell line, which expresses SIRP α 1 (unpublished); (ii) used specific PCR primers to demonstrate transcripts for SIRP α but not SIRP β in PC-3 cells; (iii) conformed by Western blotting that SIRP α is expressed in PC-3 cells, (iv) performed hybridization screening of a PC-3 DNA library from Drs. Shutsung Liao and John Kokontis at the University of Chicago, which unfortunately led us to find that their subclone of PC-3 lacks SIRP α ; (v), probed the NCBI human genome and the Celera human genome with each exon of SIRP β 1 and SIRP α 1, by which we identified a only single gene for SIRP α within the SIRP family locus on chromosome 20, but also a second potential SIRP α gene on chromosome 22, where SIRP α is encoded as a single exon, evidently a retrotransposon (one of our PCR products correlates with this gene suggesting that genes both may be expressed in PC-3 cells), (vi) by the same methods, identified the known genes for SIRP β as well as several loci that may encode other SIRP β proteins (but as confirmed in this year's work, these are not expressed in the prostate cancer cells), (vii) used PC-3 cells treated with pervanadate (to increase tyrosine phosphorylation of all proteins) to demonstrate that PC-3 cells express SHP-2 tyrosine phosphatase, (viii) demonstrated association of SIRP α with SHP-2 in PC-3 prostate cancer cells, (ix) initiated studies using protein deglycosylation to confirm the size of SIRP α in prostate cancer cells.

During the third year, we spent much of our effort on the characterization of intact and deglycosylated SIRP α protein in PC-3 cells. Although we are experienced in these area, we went through a prolonged period in which we obtained inconsistent results in these studies. We believe we have resolved these issues with the finding that the PC-3 cells express some full-length SIRP α protein, but that there may be an additional, smaller form. Because of these problems, we requested and received a no-cost extension, and we expect to complete our studies within this time.

Additional results obtained during the third year include (i) confirmation that SIRP β is not expressed on the prostate cancer cell lines by using a monoclonal antibody that recognizes SIRP β 1 but not SIRP α 1; (ii) construction of transcripts encoding SIRP α 1 mutated at the cytoplasmic tyrosine required for the recruitment of SHP-2; (iii) production of additional monoclonal anti-SIRP antibodies.

KEY RESEARCH ACCOMPLISHMENTS

Year 1

1. The production of monoclonal antibodies to SIRPs
2. The use of monoclonal antibodies to confirm the expression of SIRPs on prostate cancer cells.
3. Stable overexpression of SIRP α 1 and of SIRP β 1 in PC-3 cells.

Year 2

1. Confirmation of SIRP transcripts in prostate cancer cells by Northern blotting.
2. Confirmation of SIRP α 1 transcripts in PC-3 cells by PCR (no evidence for SIRP β).
3. Confirmation by Western blotting that SIRP α is expressed in PC-3 cells
4. Demonstration that PC-3 cells express the SIRP substrate SHP-2
5. Demonstration in PC-3 cells of the interaction of SIRP with SHP-2.

Year 3

1. Resolution of SIRP α protein size, expressed in prostate cancer cells in both glycosylated and deglycosylated forms (this work is still in progress).
2. Demonstration by flow cytometry that prostate cancer cells do not express SIRP β .
3. Construction of mutant SIRP α 1, lacking the cytoplasmic tyrosine required for the recruitment of SHP-2.

REPORTABLE OUTCOMES

PC-3 cells express SIRP α , and phosphorylation of this receptor leads to its association with the tyrosine phosphatase, SHP-2. We wish to resolve the exact form of SIRP α before we report this.

CONCLUSIONS

Our studies have confirmed the hypothesis that prostate cancer cell lines express transcripts for SIRP α and that SIRP α is expressed on the cell surface. Further, they express SHP-2, and this phosphatase associates with phosphorylated SIRP α in PC-3 prostate cancer cells, supporting the hypothesis that this receptor is functional. Studies with Western blotting suggest that PC-3 cells may express both full-length SIRP α and a smaller form, as yet uncharacterized.

REFERENCES

1. Kharitonov A, Chen Z, Sures I, Wang H, Schilling J, Ullrich A. A family of proteins that inhibit signalling through tyrosine kinase receptors. *Nature*. 1997 Mar 13;386(6621):181-6.
2. Chuang W, Lagenaur CF. Central nervous system antigen P84 can serve as a substrate for neurite outgrowth. *Dev Biol*. 1990 Feb;137(2):219-32.
3. Cant CA, Ullrich A. Signal regulation by family conspiracy. *Cell Mol Life Sci*. 2001 Jan;58(1):117-24.
4. Inagaki K, Yamao T, Noguchi T, Matozaki T, Fukunaga K, Takada T, Hosooka T, Akira S, Kasuga M. SHPS-1 regulates integrin-mediated cytoskeletal reorganization and cell motility. *EMBO J*. 2000 Dec 15;19(24):6721-31.
5. Liu Y, Buhning HJ, Zen K, Schnell FJ, Williams IR, Parkos CA. Signal regulatory protein (SIRP{alpha}), a cellular ligand for CD47, regulates PMN transmigration. *J Biol Chem*. 2002, Jan 15 (online)

6. Jiang P, Lagenaur CF, Narayanan V. Integrin-associated protein is a ligand for the P84 neural adhesion molecule. *J Biol Chem.* 1999 Jan 8;274(2):559-62.
7. Seiffert M, Cant C, Chen Z, Rappold I, Brugger W, Kanz L, Brown EJ, Ullrich A, Buhring HJ. Human signal-regulatory protein is expressed on normal, but not on subsets of leukemic myeloid cells and mediates cellular adhesion involving its counterreceptor CD47. *Blood.* 1999 Dec 1;94(11):3633-43.
8. Veillette A, Thibaudeau E, Latour S. High expression of inhibitory receptor SHPS-1 and its association with protein-tyrosine phosphatase SHP-1 in macrophages. *J Biol Chem.* 1998 Aug 28;273(35):22719-28.