

REPORT DOCUMENTATION PAGE

AFRL-SR-AR-TR-03-

0436

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, gathering existing data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE October 17, 2003	3. REPORT TYPE AND DATES COVERED Final Report 6/15/99 - 6/14/02	
4. TITLE AND SUBTITLE Probing 2D Physics with Microwave			5. FUNDING NUMBERS F49620-99-1-0284.	
6. AUTHOR(S) Daniel Tsui				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Princeton University, Electrical Engineering Department Princeton, NJ 08544			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Office of Scientific Research (AFOSR) 4015 Wilson Blvd, Rm 713 Arlington, VA 22203-1954			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES				
12a. DISTRIBUTION / AVAILABILITY STATEMENT APPROVED FOR PUBLIC RELEASE, DISTRIBUTION UNLIMITED			12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 Words) Microwave absorption, using the co-planar waveguide configuration that we developed earlier, was employed to investigate electron dynamics of the high mobility 2D charge carriers in GaAs/Al _x Ga _{1-x} As heterostructures. Three experiments were carried out: <ul style="list-style-type: none"> (1) Dynamic response of composite fermions in an anti-dot lattice (2) Observation of microwave induced cyclotron harmonics, and (3) Investigation of pinning of the Wigner crystal and measurement of the crystal correlation lengths as a function of carrier density. 				
14. SUBJECT TERMS			15. NUMBER OF PAGES 3	
			16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	

20031112 088

Final Report Submitted to Air Force Office of Scientific Research (AFOSR)**for Award #F49620-99-1-0284****“Probing 2D Physics with Microwave”****Daniel Tsui****Electrical Engineering Department,
Princeton University, Princeton NJ, 08544****October 17, 2003****1. WORK ACCOMPLISHED – Three experiments were completed****1.1 Dynamic response of composite fermions in an anti-dot lattice**

The ground state of the strongly interacting 2D electron gas in a B field at half-filling of the lowest Landau level (i.e. filling factor $\nu = 1/2$) is a Fermi liquid of composite fermions (CF's), each can be thought of as an electron bound to two flux quanta. These bizarre particles feel no net B field at $\nu = 1/2$ and the two series of Fractional Quantum Hall Effect (FQHE) states at $\nu = p/(2p \pm 1)$, with $p = \text{integers}$, can be viewed as the integer Quantum Hall Effect of the CF's resulting from the Landau quantization of their kinetic energy by the effective B they feel away from half-filling. We have studied the microwave response of the CF's in uniform samples and in samples fabricated with anti-dot arrays. We find that, in samples with anti-dot arrays, the ac conductivity increases strongly with increasing microwave frequency, strongest at $\nu = 1/2$ and clearly observed in the entire CF Landau quantization region of the $\nu = p/(2p \pm 1)$ FQHE series.

This frequency dependence is contrary to expectation for ordinary metallic systems and is not observable for $T \geq 600$ mK. It is attributed to microwave excitation of the chiral Luttinger liquids of the edge states of the $p/(2p \pm 1)$ FQHE series. However, what are the electronic processes? how to understand the excitations? and how to think through in terms of the bizarre particles of CF's? are questions still under intense current theoretical debate.

1.2 Microwave induced cyclotron harmonics

It is well known from Kohn's Theorem that excitation by the electromagnetic radiation at the cyclotron frequency harmonics is not allowed in a translationally invariant electron system, and no harmonics of microwave cyclotron resonance has been observed. It is therefore a great surprise that giant microwave induced change in the dc resistance, $\Delta R/R$ of up to 250% were observed at the 2D electron cyclotron resonance frequency and its harmonics in our high

mobility samples in the 4-40GHz frequency range. This change in resistance is proportional to the square root of the microwave power. Investigations focusing on the physical origin of this effect and the potential for its use in detector application are being continued.

1.3 Pinning mode of the Wigner crystal

The sharp microwave resonance observed in the high B field induced insulating phase of the 2D electrons in high quality GaAs/Al_xGa_{1-x}As heterostructures, which we reported a few years ago, was a great surprise and an enigma. The insulating phase has long been expected to be a pinned Wigner crystal, and it would be natural to attribute the resonance to its pinning mode. However, the B field dependence of the resonance frequency is opposite to that expected and pinning by random disorder should not give rise to a sharp resonance.

This mystery is now solved by our new experiments on higher quality and lower density 2D electron samples, grown by Dr. Loren Pfeiffer of Bell Labs, which have made it clear that the insulator is indeed a pinned Wigner solid, showing the expected behavior of decreasing resonance frequency, f_{pk} , with increasing B ($B = (n_s/\nu) * 4.14 \times 10^{-7} \text{G-cm}^2$ in Fig. 2).




Fig. 1 Microwave resonance in $\text{Re} \sigma_{xx}$ vs f




Fig. 2 Resonance frequency, f_{pk} vs $1/\nu$

The resonance is the pinning mode of the electron solid. But, the physics is much richer than having been expected. It turns out that the expected B field dependence is that of the classical limit, only now accessible in the lowest density samples. In the earlier experiment, the classical

limit is inaccessible and the observed B field dependence of the resonance frequency reflects the pinning of the quantum crystal where exchange interaction is important. The sharpness of the pinning is now understood as a characteristic of the long range Coulomb force by the electrons, which correlates the guiding center motion of the electron lattice in the randomly pinned domains.

Fig. 1 shows the resonance at different densities varied by the gate voltage on the device, and Fig. 2 summarized data on the B field dependence of the resonance frequency. For B higher than that indicated by the dotted line in Fig. 2, the system approaches that of a classical 2D electron solid.

2. JOURNAL PUBLICATIONS

2.1) "Giant Microwave Photo Resistance of Two-dimensional Electron Gas," P.D. Ye, L.W. Engel, D.C. Tsui, J.A. Simmons, J.R. Wendt, G.A. Vawter, and J.L. Reno, *Applied Phss. Lett.*, **79**, p. 2193 (2001).

2.2) "High Magnetic Field Microwave Conductivity of 2D Electrons in an Array of Antidots," P.D. Ye, L.W. Engel, D.C. Tsui, J.A. Simmons, J.R. Wendt, G.A. Vawter, and J.L. Reno., *Phys. Rev. B*, **65**, 121305 (R), (2002)

2.3) "Correlation lengths of Wigner crystal order in a 2D electron system at high B fields," P.D. Ye, L.W. Engel, D.C. Tsui, R.M. Lewis, L.N. Pfeiffer, and K. West, *Phys. Rev. Lett.*, **89**, 176802 (2002).

3. PERSONNEL

3.1 P.D. Ye (now member of technical staff at Agere Systems),

3.2 R.M. Lewis

3.3 Yong Chen