

AD_____

Award Number: DAMD17-00-1-0295

TITLE: Role of DNA-Dependent Protein Kinase in Breast Cancer
Development/Progression

PRINCIPAL INVESTIGATOR: Suk-Hee Lee, Ph.D.

CONTRACTING ORGANIZATION: Indiana University
Indianapolis, Indiana 46202-5167

REPORT DATE: July 2003

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20031112 162

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE July 2003	3. REPORT TYPE AND DATES COVERED Final (15 Jul 00-14 Jun 03)	
4. TITLE AND SUBTITLE Role of DNA-Dependent Protein Kinase in Breast Cancer Development/Progression			5. FUNDING NUMBERS DAMD17-00-1-0295	
6. AUTHOR(S) Suk-Hee Lee, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Indiana University Indianapolis, Indiana 46202-5167 E-Mail: slee@iupui.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Original contains color plates. All DTIC reproductions will be in black and white.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) DNA-dependent protein kinase (DNA-PK) is a nuclear serine/threonine protein kinase involved in various DNA metabolism and damage signaling pathways. DNA damage activates DNA-PK, which in turn phosphorylates a number of key proteins involved in replication, repair, and transcription. Accordingly, DNA-PK has long been suspected as a factor involved in sensing and transmitting DNA damage signals to the downstream target, which eventually contributes to the genomic stability and prevention of cancer. The overall goal of this proposal is to explore the role of DNA-PK in the development and progression of breast cancer. Since DNA-PK is a DNA repair factor as well as involved in damage signaling pathway, levels of DNA-PK activity among breast cancer would contribute to their drug resistance and also provide the basis for selection of patients for treatment with chemotherapy drugs. From the three-year study funded by DOD, we concluded that a peptide-based inhibitor preventing DNA-PKcs from forming a complex with Ku70/Ku80 significantly lowered DNA-PK activity. Furthermore, treatment of these breast cancer cells with target peptide significantly lowered the cell growth only in the presence of ionizing radiation, indicating that the peptide-based inhibitor exhibited a positive effect of on lowering drug resistance by specifically targeting DNA-PK <i>in vivo</i>. This result also validates a physiologic role for DNA-PK in chemotherapy drug resistance of breast cancers. Funding from the DOD breast cancer research also allowed us to study a potential role for DNA-PK in damage-induced cell cycle arrest, which was published recently.				
14. SUBJECT TERMS Damage-induced cell cycle checkpoint, DNA repair, DNA replication, S-phase arrest				15. NUMBER OF PAGES 68
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

NSN 7540-01-280-5500

Standard Form 298 (Rev. 2-89)
Prescribed by ANSI Std. Z39-18
298-102

Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	10
Reportable Outcomes.....	10
Conclusions.....	11
References.....	11
Appendices.....	13

INTRODUCTION:

The overall goal of this proposal is to explore the role of DNA-dependent protein kinase (DNA-PK) in the development and progression of breast cancer. DNA-PK is a nuclear serine/threonine protein kinase involved in various DNA metabolism and damage signaling pathways. DNA damage activates DNA-PK, which in turn phosphorylates a number of key proteins involved in replication, repair, and transcription. Accordingly, DNA-PK has long been suspected as a factor involved in sensing and transmitting DNA damage signals to the downstream target, which eventually contributes to the genomic stability and prevention of cancer. To see a correlation between DNA-PK activity and different stages of cancer, DNA-PK activity levels of various stages of breast cancer will be analyzed. Outcome of this study will provide us the information as to whether DNA-PK can be used as a prognostic measure of breast cancer progression. We will also analyze breast cancer cells grown in culture and grown as tumor grafts in mice to see whether DNA-PK plays a role in chemotherapy resistance. A strong correlation between DNA-PK activity and drug resistance of breast cancers would provide the basis for selection of patients for treatment with chemotherapy drugs. Furthermore, the information regarding the role of DNA-PK in drug resistance would be very useful for drug discovery aimed at increasing the sensitivity of tumors to chemotherapy.

BODY:**Task 1. To characterize the relationship between DNA-PK and breast cancer development/progression (months 1-36)**

In order to assess DNA-PK activity and its expression in various stages of breast cancer cells, we obtained several tissue samples from the Indiana University Cancer Center (IUCC) Tumor Bank and carried out a preliminary study. The number of breast cancer tissues we tested, however, were only 5 samples that were not enough to make any conclusion on the relationship between DNA-PK and breast cancer development/progression. We had some difficulty in obtaining enough number of specimens for different stages of breast cancer by the FIGO (Federation of International Gynecologic Organization) guideline, however the IU Cancer Center strongly supports our study and provided enough number of specimens to carry out our proposed study. This study is currently in progress. Once obtained, tissue samples will be grown in tissue culture dish (25 x 150 mm) and cell extracts will be prepared for measurement of DNA-PK activity. Also, molecular analysis of the alteration of DNA-PK activity in breast cancers will follow to evaluate the relationship between DNA-PK activity and the degree status of breast cancer.


Task 2. To determine whether DNA-PK plays a role in DNA repair and/or the chemotherapy drug resistance among breast cancers (months 1-36)**DNA-PK activity and chemotherapy drug resistance of breast cancer cells (months 6-24):**

We initially proposed to determine whether altered DNA-PK activity in breast cancers is associated with chemotherapy drug. To see a relationship between DNA-PK and drug resistance of breast cancer cells, we prepared a peptide-based inhibitor that binds to the C-terminus of Ku80 (Gell & Jackson, 1999) and competitively inhibits DNA-PKs from binding to Ku70/Ku80. We tested these peptides for their effects on DNA-PK kinase activity *in vitro* using a specific substrate peptide (EPPLSQEAFADLWKK) (Lees-Miller *et al*, 1990). The target peptide (HNI-38) effectively inhibited DNA-PK activity under the conditions where the control peptide (HN-26) had no effect, indicating that HNI-38 prevented DNA-PKs from forming a complex with Ku70/Ku80 (Figure 2).

A previous study indicates that DNA-PK mutant cells exhibit sensitivity to irradiation and cisplatin treatment (Britten *et al*, 1999; Frit *et al*, 1999). Also, studies with drug-resistant and sensitive cells indicate that higher levels of DNA-PK expression lead to drug-resistant cells, whereas the low DNA-PK activity was associated with cells with drug-sensitive phenotype (Shen *et al*, 1997). Since the target peptide effectively interferes with DNA-PK activity, we attempted to analyze whether alteration of DNA-PK activity induced by treatment with peptide-based inhibitor affected drug resistance of breast cancer cells (Figures 3-5).

Targeted Inhibition of DNA-PK

Ku70 and Ku80 form a heterodimeric complex that is important for DNA-termini binding; neither Ku70 nor Ku80 alone is active in DNA binding activity (Wu and Lieber, 1996; Gell and Jackson, 1999). The C-terminus of both Ku70 and Ku80 are necessary for heterodimer assembly as well as for DNA-termini binding (Wu and Lieber, 1996; Gell and Jackson, 1999). A recent protein interaction study indicated that DNA-PKcs interacting domain is localized at the extreme C-terminus of Ku80 (amino acids 720-732) (Gell and Jackson, 1999). Since the C-terminus of Ku80 is also likely involved in heterodimer assembly and DNA termini binding, this region (amino acids 720-732 of Ku80) was selected to synthesize a target peptide that would prevent DNA-PKcs from binding to Ku70/Ku80 regulatory subunits (see Figure 1A). To deliver a peptide to the cancer cells, a cell-permeable peptide import domain and the nuclear localization domain were added to the target peptide to obviate the need for permeabilization or microinjection of individual cells (Lin *et al*, 1995; Figure 1B).



HNI-38: AAVALLPAVLLALLAPVQRKRQKLMY—
 HN-26: AAVALLPAVLLALLAPVQRKRQKLMY
 HI-29: AAVALLPAVLLALLAPY—
 NI-22: VQRKRQKLMY—

Figure 1A (upper). A peptide co-therapy strategy for targeted inhibition of DNA-PK in cancer cell co-therapy. Treatment of cells with ionizing radiation (or chemotherapy drug) induces strand-break DNA

damage. To repair DNA damage, DNA-PK heterotrimeric complex (Ku70, Ku80, and DNA-PKcs) needs to be assembled at the ends of DNA. Target peptide representing amino acids 720-732 of Ku80 not only interacts with DNA-PKcs, but may also be involved in Ku heterodimer assembly and DNA termini binding. As a result, cells treated with target peptide will exhibit poor or no DNA repair and become highly sensitive to ionizing radiation or chemotherapy drug. **1B (lower)**. The Sequences of synthetic peptide-based inhibitor and control peptides (single-letter amino acid code). Membrane-translocating hydrophobic signal sequence is underlined and the nuclear localization sequence is shown in **bold face**. Twelve residue of peptide inhibitor region is indicated as bar (▬) at the C-terminus. The tyrosine residue (Y) is included for ^{125}I -labeling to determine the import efficiency of the peptide into the cells. Peptide-based inhibitor contains the hydrophobic region localization sequence, so-called membrane-translocating carrier, which not only facilitates secretion of proteins, but also is important for importing synthetic peptides into the cell (Lin YZ *et al*, 1995). This localization peptide is capable of carrying a functional domain such as nuclear localization signal (NLS) (Boulikas T, 1994). We therefore synthesized a 38-residue peptide (HNI-38) comprising the signal peptide sequence (AAVALLPAVLLALLAP) and NLS (VQRKRQKLM) followed by a tyrosine (Y) residue and 12-residue of peptide inhibitor sequence (EGGDVDDLLDMI) representing the C-terminus of Ku80 (amino acids #721-732). The tyrosine residue was used for ^{125}I -labeling to determine the import efficiency of synthetic peptides into the cells (and nuclei) (Lin YZ *et al*, 1995).

A target peptide interrupts the interaction between DNA-PKcs and Ku70/Ku80 as well as the binding of Ku complex to DNA. DNA-PKcs and Ku70/Ku80 are abundant proteins approximately 5×10^5 molecules per human cells (Lee and Kim, 2002) and most of Ku70/Ku80 heterodimer exists in cell extracts without forming a complex with DNA-PKcs in the absence of DNA (Hammarsten and Chu, 1998). Therefore, target peptide (HNI-38) was analyzed for its effect on interaction between DNA-PKcs and Ku70/Ku80 in the presence of dsDNA. Varying concentrations of either control (HN-26) or target peptide (HNI-38) was incubated with cell extracts containing DNA-PKcs and Ku complex in the presence of dsDNA cellulose, and examined for its effect on binding of Ku complex and DNA-PKcs to DNA following the dsDNA cellulose pulldown assay (Figure 2A). Although it was marginal, the addition of increasing amount of target peptide (HNI-38) not control peptide (HN-26) led to a decrease in DNA-PKcs associated with dsDNA, suggesting that target peptide binds to DNA-PKcs and inhibits its binding to Ku70/Ku80. It is also noted that the addition of target peptide affected the binding of Ku70/Ku80 to the dsDNA cellulose (Figure 2A).




Figure 2. Effect of the target peptide on interaction of Ku70/Ku80 with DNA-PK or with dsDNA. **Panel A (upper).** The target peptide (HNI-38) interferes with association of DNA-PKcs with dsDNA. Partially purified DNA-PK fraction (100 ng) was incubated with 0 nM (lane 2), 10 nM (lanes 3 & 6), 50 nM (lanes 4 & 7), and 100 nM (lanes 5 & 8) of either control peptide or target peptide prior to dsDNA cellulose pull-down assay. Lane 1 contained partially purified DNA-PK without dsDNA pulldown assay. The protein-dsDNA cellulose complex was analyzed by the procedure described in Methods section. **Panel B (lower).** Effect of HNI-38 on DNA binding activity of Ku70/Ku80 complex. Purified Ku70/Ku80 complex (100 ng) was incubated with 10 nM (lanes 4 & 7), 50 nM (lanes 3 & 6), and 100 nM (lanes 2 & 5) of either control peptide or target peptide prior to dsDNA cellulose pull-down assay. Following the dsDNA cellulose pull-down, Ku70 and Ku80 were analyzed by 10% SDS-PAGE and Western blot.

To further examine the effect of HNI-38 on Ku's DNA binding activity, target peptide was incubated with purified Ku70/Ku80 complex in the presence of dsDNA cellulose, and the reaction mixtures were analyzed for the presence of Ku70 and Ku80 following the dsDNA pulldown assay (Figure 2B). In keeping with Figure 2A, target peptide (HNI-38) significantly interfered with binding of Ku complex to dsDNA under the conditions where control peptide (HN-26) showed virtually no effect (Figure 2B). This result suggests that target peptide not only affects the interaction between DNA-PKcs and Ku, but also interferes with Ku's DNA binding activity.

Effect of target peptide (HNI-38) on DNA-PK kinase activity

Interaction of DNA-PKcs with Ku complex are necessary for activation of its kinase activity (Gottlieb and Jackson, 1993; Hartley et al., 1995), therefore, the efficacy of target peptide was analyzed by measuring DNA-PK kinase activity *in vitro* in the presence of either HI-26 or HNI-38. DNA-PK kinase activity was inhibited up to 50% in the presence of HNI-38 under the conditions where a control peptide (HI-26) showed minimal effect (Figure 3), strongly supporting a notion that target peptide specifically binds to DNA-PKcs and interferes with interaction between DNA-PKcs and Ku complex. Inhibitory effect of target peptide on DNA-PK occurred at low peptide concentration (>20 μ M) and, in the presence of 20 μ M or higher, both target and control peptides inhibited DNA-PK activity (data not shown).




Figure 3. Effect of target peptide on DNA-PK kinase activity *in vitro*. Partially purified DNA-PK fraction was incubated with various concentrations of either control peptide or target peptide prior to the addition of substrate peptide and other components for DNA-PK kinase assay. DNA-PK activity was measured as the relative amounts of ^{32} P transferred to the substrate peptide.

Target peptide interferes with repair of double-stranded DNA breaks induced by IR

IR-induced double-stranded DNA breaks are efficiently repaired by non-homologous end-joining (NHEJ) process. Genetic and biochemical studies strongly indicated that DNA-PK plays an

essential role in NHEJ (Jeggo, 1998; Jin et al., 1997; Blunt et al., 1995). Hence, an alternative way to determine the efficacy of peptide inhibitor is to measure the repair of double-stranded DNA breaks following IR. Breast cancer cells (NCI) grown in the presence of [^{14}C]-thymidine (DiBiase et al, 2000) were treated with either a control or target peptide for 24 hrs. Following irradiation (40 Gy), cells were harvested at various time points and intact chromosomal DNA and DSBs were separated by pulsed field gel electrophoresis (0.5% agarose). Treatment of NCI cells with IR (40 Gy) induced substantial amounts of dsDNA breaks, most of which were repaired within 4 hrs. Cells treated with target or control peptide did not show any difference in generating DSBs following IR (Figure 4A; lane 2 vs. lanes 8 & 14). On the other hand, cells treated with target peptide (Figure 4A; lanes 14-18) compared with those treated with control peptide (Figure 4A; lanes 8-12) showed a noticeable decrease in DSB repair activity. This result suggests that target peptide interfered with dsb repair *in vivo* through the targeted inhibition of DNA-PK activity.




Figure 4. Effect of target peptide on double-stranded DNA breaks (dsb) repair. Breast cancer cells (NCI) grown in ^{14}C -containing media were treated with ionizing radiation (40 Gy) in the presence of no peptide (lanes 1-6), 50 nM of control peptide (lanes 7-12), or 50 nM of target peptide (lanes 13-18). After harvesting the cells at various time points, intact chromosomes and double-stranded DNA breaks (DSBs) were separated by gel electrophoresis (**upper panel**: fluorography) and were quantified by liquid scintillation counter (**lower panel**).

Target peptide inhibits breast cancer cell growth only in the presence of DNA damage

Cells lacking DNA-PK catalytic subunit showed increased sensitivity to DNA damaging drugs or ionizing radiation (IR) (Lees-Miller et al., 1995; Kirchgessner et al., 1995), suggesting that DNA-PK activity is essential for DNA repair and cell survival upon DNA damage. We therefore tested whether a targeted inhibition of DNA-PK by a peptide HNI-38 would sensitize breast cancer cells upon treatment of ionizing radiation or chemotherapeutic drug (cisplatin). Two breast cancer cells (NCI and MDA231) were treated with either control (HI-26) or target peptide (HNI-38) and tested for the efficacy of DNA-PK inhibitory peptide on lowering resistance of cells in response to ionizing radiation using standard colony count cell survival assay. Both control and target peptides did not

show any effect on cell growth in the absence of ionizing radiation. However, cells treated with IR showed significant cell growth inhibition in the presence of target peptide but not with control peptide (Figure 5A), suggesting that cell growth inhibition by target peptide occurs through targeting DNA-PK activity. Cells treated with cisplatin, although not as effective as those treated with IR, also showed inhibitory effect on cell growth in the presence of HNI-38 (Fig 5B).




Figure 5. Effect of control (HN-26) and target (HNI-38) peptides on the growth of breast cancer cells treated with ionizing radiation (**upper panel**) or cisplatin (**lower panel**). Values expressed are means (+ S.E.) of the three replications (*, $p < 0.01$; **, $p < 0.01$). The clonogenic assay was used for the cells treated with ionizing radiation and the cell survival assay (MTT) was employed for those treated with cisplatin (see the Methods section for the detailed procedure).

In summation, DNA-PK activity is essential for DNA repair as well as cell cycle arrest in response to DNA damage, which contributes to cell survival by protecting cells from apoptosis. Cells treated with target peptide not control peptide showed a noticeable decrease in DSB repair following high dose (40 Gy) of IR, suggesting that HNI-38 specifically targets DNA-PK *in vivo* and interferes with dsb repair activity through inhibition of DNA-PK activity. Targeted inhibition of DNA-PK by HNI-38 also caused cell growth inhibition only when cells were treated with IR, suggesting that HNI-38 targeted DNA-PK and lowered resistance of cells in response to ionizing radiation, which eventually causes growth inhibition of both NCI and MDA231. Treatment of cells with HNI-38 also showed additive effect on cell growth inhibition in response to cisplatin treatment. This observation is in keeping with previous findings that DNA-PK is directly involved in NER action in mammals (Muller et al., 1998). It also supports a notion that a targeted inhibition of DNA-PK would sensitize

cancer cells upon treatment of chemotherapeutic drugs such as cisplatin. Together, our study described here not only validates DNA-PK as a useful molecular target for the treatment of drug-resistant cancer cells, but also supports physiologic role for DNA-PK in IR or chemotherapy drug resistance of cancer cells.

KEY RESEARCH ACCOMPLISHMENTS:

1. Demonstration of the relationship between DNA-PK activity and drug resistance of breast cancer cells
2. Validation of a peptide-based inhibitor of DNA-PK on lowering the growth of breast cancer cells following radiation treatment
3. Basic research on the role for DNA-PK in DNA repair and damage-induced cell cycle arrest

REPORTABLE OUTCOMES: (works supported by DAMD 17-00-1-0295)

1. Wang, M, You, J-S, and Lee, S-H. (2001) A structure-function analysis of RPA's zinc finger in redox control. Redox control of zinc finger protein: Mechanisms and role in gene regulation *Antioxidants & Redox Signaling*, 3(4), 657-669.
2. Park, S-J., Oh, E-J., Yoo, M-A, and Lee, S-H. (2001) Involvement of DNA-dependent protein kinase in regulation of stress-induced JNK activation. *DNA & Cell Biol.* 20 (10), 637-645.
3. Lee, S-H. (2001) Recognition of DNA Damage in Mammals. *J. Biochem. & Mol. Biol.* 34 (6), 489-495.
4. Lee, S-H. and Kim, C-H. (2002) DNA-Dependent Protein Kinase Complex: a Multifunctional Protein in DNA Repair and Damage Checkpoint. *Molecules and Cells* 13(2), 159-166.
5. Kim, C-H, Park, S-J, and Lee, S-H (2002) Targeted inhibition of Ku70/Ku80 and DNA-PKcs interaction sensitizes breast cancer cells following ionizing radiation. *J. Pharm. Exp. Ther.* 303 (2), 753-759.
6. You, J-S, Wang, M, and Lee, S-H. (2003) Biochemical analysis of damage recognition process in nucleotide excision repair. *J. Biol. Chem.* 278, 7476-7485.
7. Park, S-J, Ciccone, S., Freie, BA, Kurimasa, A., Chen, DJ, Li, GC, Clapp, DW, and Lee, S-H (2003) A unique role for Ku70/Ku80 complex in chromosomal DNA replication following ionizing radiation. *Mol. Cell. Biol.* (submitted)
8. Park, S-J, Armstrong, S.A., Kim, C-H., Robertson, K., Kelley, M., and Lee, S-H. (2003) Lack of EGF receptor contributes to the drug sensitivity in human germline cells. *Cancer Research* (submitted)
9. Invited presentation at the annual meeting for *The Amelia Project (The Catherine Peachey Fund for Breast Cancer Research)*, Indianapolis, IN, February 2001.
10. Chung-Hui Kim, Su-Jung Park, and Suk-Hee Lee (2002) A targeted inhibition of DNA-dependent protein kinase sensitizes breast cancer cells following ionizing radiation. *DOD Era of Hope-Breast Cancer Research Program*. September 2002, Orlando, FL.
11. Jin-Sam You, Chung-Hui Kim, Mu Wang, and Suk-Hee Lee; Damage recognition occurs in a multi-step process in nucleotide excision repair. *Gordon Conference*, August 2002, Connecticut.
12. Su-Jung Park, Samantha L. M. Ciccone, Brian Freie, Akihiro Kurimasa, David J. Chen, Gloria C. Li, D. Wade Clapp, and Suk-Hee Lee (2003) A novel role for the Ku complex in DNA replication following strand break damage: protection of the integrity of PCNA and replication machinery on chromatin. *The 5th Annual Midwest DNA Repair Symposium*, May 2003, Mayo Clinic, Rochester, MN.

CONCLUSIONS:

The overall goal of this proposal is to explore the role for DNA-PK in the development and progression of breast cancer. Since DNA-PK is a key DNA repair factor as well as involved in damage signaling pathway, levels of DNA-PK activity among breast cancer would contribute to their drug resistance and also provide the basis for selection of patients for treatment with chemotherapy drugs.

From the three-year of study, we concluded that a peptide-based inhibitor preventing DNA-PKs from forming a complex with Ku70/Ku80 significantly lowered DNA-PK activity. Furthermore, treatment of these breast cancer cells with target peptide significantly lowered the cell growth only in the presence of ionizing radiation, indicating that the peptide-based inhibitor exhibited a positive effect on lowering drug resistance by specifically targeting DNA-PK *in vivo*. This result also validates a physiologic role for DNA-PK in chemotherapy drug resistance of breast cancers.

REFERENCES:

- Blunt T, Finnie NJ, Taccioli GE, Smith GC, Demengeot J, Gottlieb TM, Mizuta R, Varghese AJ, Alt FW, Jeggo PA, (1995) Defective DNA-dependent protein kinase activity is linked to V(D)J recombination and DNA repair defects associated with the murine scid mutation. *Cell*. **80**, 813-823.
- Cary RB, Peterson SR, Wang J, Bear DG, Bradbury EM, Chen DJ. (1997) DNA looping by Ku and the DNA-dependent protein kinase. *Proc Natl Acad Sci USA*. **94**, 4267-4272.
- Critchlow, SE and Jackson, SP (1998) DNA end-joining: from yeast to man. *Trends Biochem. Sci.*, **23**, 394-398.
- DiBiase SJ, Zeng ZC, Chen R, Hyslop T, Curran WJ Jr, Iliakis G. (2000) DNA-dependent protein kinase stimulates an independently active, nonhomologous, end-joining apparatus. *Cancer Res*. **60**:1245-1253.
- Featherstone, C, Jackson, SP (1999) Ku, a DNA repair protein with multiple cellular functions? *Mutat Res.*, **434**, 3-15.
- Friedberg, EC (1996) Relationship between DNA repair and transcription. *Ann. Rev. Biochem.*, **65**, 15-42.
- Gell D, Jackson SP. (1999) Mapping of protein-protein interactions within the DNA-dependent protein kinase complex. *Nucleic Acids Res*. **27**, 3494-3502.
- Gottlieb, T.M. and Jackson, S.P. (1993) The DNA-dependent protein kinase: requirement for DNA ends and association with Ku antigen. *Cell*, **72**: 131-142.
- Hammarsten O, Chu G. (1998) DNA-dependent protein kinase: DNA binding and activation in the absence of Ku. *Proc Natl Acad Sci USA*. **95**: 525-530.
- Hanawalt, PC (1994) Transcription-coupled repair and human disease. *Science*, **266**, 1957-1958.

Hartley, K.O., Gel, D., Smith, G.C., Zhang, H., Divecha, N., Connelly, M.A., Admon, A., Lees-Miller, S.P., Anderson, C.W. and Jackson, S.P. (1995) DNA-dependent protein kinase catalytic subunit: a relative of phosphatidylinositol 3-kinase and the ataxia telangiectasia gene product. *Cell*, **82**: 849-856.

Jackson, S.P. (1997) DNA-dependent protein kinase. *Int. J. Biochem. Cell. Biol.*, **29**: 935-938.

Jeggo, PA (1998) DNA breakage and repair. *Adv Genet.* **38**, 185-218.

Jin, S., Inoue, S. and Weaver, D.T. (1997) Functions of the DNA dependent protein kinase. *Cancer Surv.*, **29**: 221-261.

Kim SH, Kim D, Han JS, Jeong CS, Chung BS, Kang CD, Li GC. (1999) Ku autoantigen affects the susceptibility to anticancer drugs. *Cancer Res.* **59**: 4012-4017

Kirchgessner CU, Patil CK, Evans JW, Cuomo CA, Fried LM, Carter T, Oettinger MA, Brown JM. (1995) DNA-dependent kinase (p350) as a candidate gene for the murine SCID defect. *Science*. **267**, 1178-1183.

Kurimasa A, Kumano S, Boubnov NV, Story MD, Tung CS, Peterson SR, Chen DJ. (1999) Requirement for the kinase activity of human DNA-dependent protein kinase catalytic subunit in DNA strand break rejoining. *Mol Cell Biol.* **19**, 3877-3884.

Lee, S-H. and Kim, C-H. (2002) DNA-Dependent Protein Kinase Complex: a Multifunctional Protein in DNA Repair and Damage Checkpoint. *Molecules and Cells* **13**, 159-166.

Lee, S-H. (2001) Recognition of DNA Damage in Mammals. *J. Biochem. & Mol. Biol.* **34**, 489-495.

Lees-Miller, SP., Chen, Y-R. and Anderson, CW (1990) Human cells contain a DNA-activated protein kinase that phosphorylates simian virus 40 T antigen, mouse p53, and the human Ku autoantigen. *Mol. Cell. Biol.*, **10**: 6472-6481.

Lees-Miller SP, Godbout R, Chan DW, Weinfeld M, Day RS 3rd, Barron GM, Allalunis-Turner J. (1995) Absence of p350 subunit of DNA-activated protein kinase from a radiosensitive human cell line. *Science*. **267**, 1183-1185.

Lieber, MR, Grawunder, U, Wu, X, Yaneva, M (1997) Tying loose ends: roles of Ku and DNA-dependent protein kinase in the repair of double-strand breaks. *Curr Opin Genet Dev.*, **7**, 99-104.

Lin Y-Z, Yi SY, Veach R A, Torgerson T R., and Hawiger J (1995) Inhibition of Nuclear Translocation of Transcription Factor NF-B by a Synthetic Peptide Containing a Cell Membrane-permeable Motif and Nuclear Localization Sequence *J. Biol. Chem.* **270**, 14255-14258.

Muller, C, Salles, B (1997) Regulation of DNA-dependent protein kinase activity in leukemic cells. *Oncogene*, **15**, 2343-2348.

Muller, C., Calsou, P., Frit, P., Cayrol, C., Carter, T. and Salles, B. (1998) UV sensitivity and impaired nucleotide excision repair in DNA-dependent protein kinase mutant cells. *Nucleic Acids Res.*, **26**, 1382-1389.

Park, J-S., Park, S-J., Peng, X., Yu, M., Wang, M., and Lee, S-H. (1999) Involvement of DNA-dependent protein kinase in UV-damage induced replication arrest. *J. Biol. Chem.* **274**, 32520-32527.

Shen, H., Kauvar, L. and Tew, K.D. (1997) Importance of glutathione and associated enzymes in drug response. *Oncol. Res.*, **9**: 295-302.

Shen, H, Schultz, M, Kruh, GD, Tew, KD (1998) Increased expression of DNA-dependent protein kinase confers resistance to adriamycin. *Biochim. Biophys. Acta*, **23**, 131-138.

Suwa A, Hirakata M, Takeda Y, Jesch SA, Mimori T, Hardin JA. (1994) DNA-dependent protein kinase (Ku protein-p350 complex) assembles on double-stranded DNA. *Proc Natl Acad Sci USA*. **91**: 6904-6908.

Tew, KD, O'Brien, M, Laing, NM, Shen, H (1998) Coordinate changes in expression of protective genes in drug-resistant cells. *Chem Biol Interact.*, **111**, 199-211.

Weaver, D.T. and Alt, F.W. (1997) V(D)J recombination. From RAGs to stitches. *Nature*, **388**: 428-429.

Woo, R.A., McLure, K.G., Lees-Miller, S.P., Rancourt, D.E., and Lee, P.W.K. (1998) DNA-dependent protein kinase acts upstream of p53 in response to DNA damage. *Nature*, **394**, 700-704.

Wood, RD (1996) DNA repair in eukaryotes. *Ann. Rev. Biochem.*, **65**, 135-167.

Wu, X. And Lieber, MR (1996) Protein-protein and protein-DNA interaction regions within the DNA end-binding protein Ku70-Ku86. *Mol. Cell. Biol.*, **16**, 5186-5193.

APPENDICES:

Six publications supported by DAMD 17-00-1-0295

A Targeted Inhibition of DNA-Dependent Protein Kinase Sensitizes Breast Cancer Cells Following Ionizing Radiation

CHUNG-HUI KIM, SU-JUNG PARK, and SUK-HEE LEE

Department of Biochemistry & Molecular Biology, Indiana University Cancer Center, and Walther Cancer Institute, Indiana University School of Medicine, Indianapolis, Indiana

Received May 3, 2002; accepted June 13, 2002

ABSTRACT

A major mechanism by which cancer cells become resistant to ionizing radiation (IR) and chemotherapy drugs is by enhanced DNA repair of the lesions; therefore, through inhibition of DNA repair pathways that tumor cells rely on to escape chemotherapy, we expect to increase the killing of cancer cells and reduce drug resistance. DNA-dependent protein kinase (DNA-PK) is a nuclear serine/threonine protein kinase essential for DNA repair as well as sensing and transmitting a damage signal to downstream targets leading to cell cycle arrest. We used a peptide cotherapy strategy to see whether a targeted inhibition of DNA-PK activity sensitizes breast cancer cells in response to IR or chemotherapy drug. A synthesized peptide representing the C terminus of Ku80 (HNI-38) selectively targeted and disrupted

interaction between Ku complex and the catalytic subunit of DNA-dependent protein kinase (DNA-PKcs) as well as the DNA binding activity of Ku that led to the inhibition of DNA-PK activity and reduction in double-stranded DNA break (dsb) repair activity. Furthermore, a peptide-based inhibitor with target sequence effectively inhibited the growth of breast cancer cells only in the presence of DNA damage, suggesting that the target peptide sensitizes cancer cells through blocking dsb DNA repair activity. Together, this study not only validates the involvement of the C terminus of Ku80 in Ku's DNA termini binding and interaction with DNA-PKcs, but also supports physiological role for DNA-PK in IR or chemotherapy drug resistance of cancer cells.

DNA-dependent protein kinase (DNA-PK) is a nuclear serine/threonine protein kinase composed of 460-kDa catalytic subunit (DNA-PKcs) and a heterodimer of Ku70 and Ku80, which act as a DNA binding and regulatory component for the complex (Gottlieb and Jackson, 1993; Jin et al., 1997; Lieber et al., 1997). DNA-PK is a key component of the nonhomologous end joining (NHEJ) pathway and V(D)J recombination (Blunt et al., 1995), with the unique property of being activated by DNA ends (Critchlow and Jackson, 1998; Jeggo, 1998; Featherstone and Jackson, 1999). It has long been suspected as a factor involved in sensing and transmitting DNA damage signals to the downstream targets (Jackson, 1997; Weaver and Alt, 1997; Lee and Kim, 2002). Previous studies demonstrated that DNA-PK is necessary for activation of p53 (Woo et al., 1998), nucleotide excision repair (Muller et al., 1998), and damage-induced S-phase arrest (Park et al., 1999) in response to DNA damage, all of which contribute to cell protection from genetic alterations as well

as chemotherapy drug resistance. In vivo observations indicated that DNA-PK mutant cells exhibited sensitivity to ionizing irradiation and chemotherapy drugs and were associated with lower DNA repair activity following DNA damage, suggesting a positive role for DNA-PKcs in DNA repair (Britten et al., 1999; Frit et al., 1999). Also, studies with drug-resistant or drug-sensitive cancer cells suggested that higher levels of DNA-PK expression lead to drug-resistant cells, whereas low DNA-PK activity was associated with cells with drug-sensitive phenotype (Shen et al., 1997) and was linked to cell death via the accumulation of damaged DNA.

The current model of DNA-PK complex activation by DNA is based on the tenet that without DNA, DNA-PKcs is inactive and incapable of binding Ku (Hanawalt, 1994; Suwa et al., 1994; Hartley et al., 1995). When a double-strand break is introduced, Ku complex binds to the DNA because of its high affinity for DNA ends. The binding of Ku induces conformational change that allows it to interact with DNA-PKcs. It is unclear how the Ku/DNA complex activates the kinase activity of DNA-PKcs. One hypothesis is that DNA-PKcs undergoes a conformational change upon association with the Ku/DNA complex, and this conformational change accounts for the activation of kinase activity. The kinase activity associ-

This research was supported by grants from the U.S. Army (DAMD17-00-1-0295) and the National Institutes of Health (CA92111).

Article, publication date, and citation information can be found at <http://jpet.aspetjournals.org>.

DOI: 10.1124/jpet.102.038505.

ABBREVIATIONS: DNA-PK, DNA-dependent protein kinase; DNA-PKcs, catalytic subunit of DNA-dependent protein kinase; NHEJ, nonhomologous end-joining; IR, ionizing radiation; dsb, double-stranded DNA break; dsDNA, double-stranded DNA; MTT, 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazolium bromide; PAGE, polyacrylamide gel electrophoresis.

ated with DNA-PK is needed for DNA repair in vivo, since expression of a kinase-inactive form of DNA-PKcs failed to complement the radiosensitive phenotype of a mammalian cell line lacking the DNA-PKcs protein (Kurimasa et al., 1999). However, the physiological targets of DNA-PK in vivo are still not clear. The DNA-PK complex can physically tether two ends of a dsb in close proximity in vitro, suggesting the hypothesis that the DNA-PK complex acts as a scaffold to assemble the NHEJ pathway proteins at a DSB (Cary et al., 1997).

We hypothesize that 1) DNA-PK plays an important role in conferring cells becoming resistant to ionizing radiation or anticancer DNA-damaging drugs, and 2) targeted inhibition of DNA-PK sensitizes drug resistance of cancer cells and facilitates cell killing. By developing peptides that can directly interfere with DNA-PK activity, one can develop a novel cotherapy that can selectively target and disrupt the IR-induced dsb repair pathway, which will enhance the efficacy of currently available treatments and also broaden the usefulness of chemotherapeutic agents in cancer treatment. We have therefore synthesized a peptide (HNI-38) mimicking the domain of Ku80 essential for interaction with its catalytic subunit (DNA-PKcs) and tested whether it can selectively target and disrupt DNA-PK activity required for dsb repair, which potentiates the effect of chemotherapy drug in cancer treatment. This strategy can be applied to cancer cotherapy, which will broaden the usefulness of chemotherapeutic agents in cancer treatment.

Materials and Methods

Cell Lines, Antibodies, and Chemicals. Two human breast cancer cells, MDA231 and NCI, were obtained from Dr. George Sledge (Indiana University Cancer Center, Indianapolis, IN) and maintained in minimal essential medium supplemented with 10% fetal bovine serum at 37°C in a CO₂ incubator. Antibodies to Ku70/80 and the catalytic subunit of DNA-dependent protein kinase (DNA-PKcs) were obtained from either Upstate Biotechnology (Lake Placid, NY) or BD Pharmingen (San Diego, CA). [γ -³²P]ATP (4500 Ci/mmol) was from ICN Pharmaceuticals (Costa Mesa, CA), and dsDNA cellulose and cisplatin were obtained from Sigma-Aldrich (St. Louis, MO).

dsDNA Cellulose Pull-Down Assay. The dsDNA cellulose fraction (100 μ g) containing DNA-PKcs and Ku70/Ku80 heterodimer was prepared from HeLa cells (Lees-Miller et al., 1990) and incubated with the indicated amount of either control (HN-26) or a target (HNI-38) peptide in the presence of 4 mM ATP and 50 μ l of dsDNA cellulose (3 mg of dsDNA/mg of cellulose; Sigma-Aldrich) for 3 h at 4°C with rocking for the interaction of DNA-PKcs and Ku70/Ku80 heterodimer. Where indicated, purified Ku70/Ku80 complex (100 ng) was used instead of the dsDNA cellulose fraction. After centrifugation at 4000 rpm, the precipitates were collected and washed three times with a buffer (50 mM Tris-HCl, pH 8.0, 200 mM NaCl, 0.5% Nonidet P-40, 1 mM EDTA, 10% glycerol, 1 mM dithiothreitol, and 0.1 mM phenylmethylsulfonyl fluoride) for protein analysis. For Western blot, the precipitates were separated by 8 or 10% SDS-PAGE, transferred to nitrocellulose (Millipore Corp., Bedford, MA), and blotted with primary antibody to Ku70/80 and/or DNA-PKcs followed by a peroxidase-coupled secondary antibody (Amersham Biosciences Inc., Piscataway, NJ) and an enhanced chemiluminescence (ECL kit, Amersham) reaction prior to visualization on Kodak-O-mat film.

DNA-PK Kinase Assay. Reaction mixtures (20 μ l) contained 20 mM HEPES-KOH (pH 7.5), 1 mM EGTA, 1 mM EDTA, 1 mM dithiothreitol, 10 mM MgCl₂, 7 mM MnCl₂, 5 mM NaF, 1 mM

Na₃VO₄, 50 μ l of [³²P]ATP, 150 μ l of substrate peptide, 0.4 μ g of DNase I-activated calf-thymus DNA (Sigma-Aldrich), and 100 ng of partially purified DNA-PK complex. DNA-PK complex was partially purified from HeLa cells according to the procedure described previously (Lees-Miller et al., 1990). Substrate peptide (EP-PLSQEAFADLWKK) representing amino acids 11–24 of p53 was used as a substrate for DNA-PK assay (Lees-Miller et al., 1990). To find out whether the peptide interferes with DNA-PK kinase activity, various amounts of peptide inhibitor were added to the reaction. After incubation at 30°C for 30 min, the reaction mixtures were stopped with 30% acetic acid and a portion of the reaction mixtures (5 μ l) was spotted onto a P81 strip, and after extensive washing, radioactivity was measured. DNA-PK activity was measured as picomoles of ³²P transferred to the substrate peptide.

Cell Survival Assay. Cells (1.0 $\times$ 10⁴ cells/well) were seeded in a 96-well plate in the presence of control or target peptide and incubated for 24 h before the treatment of cells with either ionizing radiation or cisplatin. After further incubation at 37°C, 5% CO₂ for 72 h, cell survival was measured using a colorimetric cell survival assay from Roche Diagnostics (Indianapolis, IN; MTT Cell Proliferation Kit). Alternatively, clonogenic assay was used to measure the ability of cells to form colonies on 100-mm² tissue culture dishes after treatment with ionizing radiation or cisplatin. Controls consisted of cells untreated with peptides, cells untreated with DNA-damaging agent, or cells without either treatment. Cells were continuously exposed for 5 days to the indicated concentrations of the peptide and colonies were stained with crystal violet; then, colonies greater than 50 cells were counted. Each point represents mean values $\pm$ S.E.,




Fig. 1. Panel A, a peptide cotherapy strategy for targeted inhibition of DNA-PK in cancer cell cotherapy. Treatment of cells with ionizing radiation (or chemotherapy drug) induces strand-break DNA damage. To repair DNA damage, DNA-PK heterotrimeric complex (Ku70, Ku80, and DNA-PKcs) needs to be assembled at the ends of DNA. Target peptide representing amino acids 720–732 of Ku80 not only interacts with DNA-PKcs, but may also be involved in Ku heterodimer assembly and DNA-termini binding. As a result, cells treated with target peptide will exhibit poor or no DNA repair and become highly sensitive to ionizing radiation or chemotherapy drug. Panel B, synthetic peptide used for cotherapy study. Sequences of synthetic peptide-based inhibitor and control peptides (single-letter amino acid code). The membrane-translocating hydrophobic signal sequence is indicated in italic letters and the nuclear localization sequence is shown in boldface. Twelve residues of the peptide inhibitor region are indicated at the C terminus of HNI-38.

each conducted with triplicate plates. The *p* values in Fig. 5A (see under *Results*) were obtained from two separate experiments using a one-way analysis of variance method (SigmaStat for Windows, version 2.03; SPSS Science, Chicago, IL).

Double-Stranded DNA Break (dsb) Repair Assay. Kinetics of rejoining of radiation-induced damaged DNA in breast cancer cells following exposure of cells to 40 Gy gamma irradiation (^{137}Cs) were measured by pulsed field gel electrophoresis. Breast cancer cells (NCI) were grown in the presence of 2.5 μM [^{14}C]thymidine (0.1 $\mu\text{Ci/ml}$) (DiBiase et al., 2000) and treated with either a control or target peptide. After irradiation (40 Gy), cells were further incubated at 37°C with prewarmed (42°C) fresh medium to allow DSB repair, and then harvested at various times and resuspended in serum-free medium at a concentration of 2 to 5 $\times 10^6$ cells/ml. Cells were mixed with an equal volume of 1% agarose, and the solidified cell-agarose suspension was lysed with buffer containing 10 mM Tris (pH 8.0), 50 mM NaCl, 0.5 M EDTA, 2% *N*-lauryl sarcosyl, and proteinase E and O (0.1 mg/ml) for 16 to 18 h at 50°C (DiBiase et al., 2000). DNA double-strand breaks were analyzed by asymmetric field inversion gel electrophoresis using 0.5% agarose gel in 0.5 $\times$ Tris borate-EDTA at 10°C for 40 h. After electrophoresis, gels were analyzed by fluorography. For quantification of damaged DNA repair, intact chromosome and damaged DNA were separately removed from the gel and measured for ^{14}C labeling using a liquid scintillation counter.

Results

Targeted Inhibition of DNA-PK. Ku70 and Ku80 form a heterodimeric complex that is important for DNA-termini binding; neither Ku70 nor Ku80 alone is active in DNA

binding activity (Wu and Lieber, 1996; Gell and Jackson, 1999). The C termini of both Ku70 and Ku80 are necessary for heterodimer assembly as well as for DNA-termini binding (Wu and Lieber, 1996; Gell and Jackson, 1999). A recent protein interaction study indicated that the DNA-PKcs interacting domain is localized at the extreme C terminus of Ku80 (amino acids 720–732) (Gell and Jackson, 1999). Since the C terminus of Ku80 is also likely involved in heterodimer assembly and DNA-termini binding, this region (amino acids 720–732 of Ku80) was selected to synthesize a target peptide that would prevent DNA-PKcs from binding to Ku70/Ku80 regulatory subunits (see Fig. 1A). To deliver a peptide to the cancer cells, a cell-permeable peptide import domain and the nuclear localization domain were added to the target peptide to obviate the need for permeabilization or microinjection of individual cells (Lin et al., 1995; Fig. 1B).

A Target Peptide Interrupts the Interaction between DNA-PKcs and Ku70/Ku80 as well as the Binding of Ku Complex to DNA. DNA-PKcs and Ku70/Ku80 are abundant proteins, approximately 5 $\times 10^5$ molecules per human cell (Lee and Kim, 2002, and references therein), and most of the Ku70/Ku80 heterodimer exists in cell extracts without forming a complex with DNA-PKcs in the absence of DNA (Hammarsten and Chu, 1998; Fig. 2A). Therefore, target peptide (HNI-38) was analyzed for its effect on interaction between DNA-PKcs and Ku70/Ku80 in the presence of dsDNA. Varying concentrations of either control (HN-26) or target peptide




Fig. 2. Effect of the target peptide on interaction of Ku70/Ku80 with DNA-PK or with dsDNA. Panel A, Ku70/Ku80 complex is not in complex with DNA-PKcs. Chromatographic separation of DNA-PKcs from Ku70/Ku80 heterodimer. Partially purified DNA-PK fractions (dsDNA cellulose fraction; see Lees-Miller et al., 1990 for details) were subjected to heparin-Sepharose column chromatography and eluted with 100 to 500 mM NaCl gradient. Fractions were analyzed by 6% SDS-PAGE for DNA-PKcs and Ku80 followed by immunoblot using anti-DNA-PKcs and anti-Ku80 antibodies. Panel B, the target peptide (HNI-38) interferes with association of DNA-PKcs with dsDNA. Partially purified DNA-PK fraction (100 ng) was incubated with 0 nM (lane 2), 10 nM (lanes 3 and 6), 50 nM (lanes 4 and 7), and 100 nM (lanes 5 and 8) concentrations of either control peptide or target peptide prior to dsDNA cellulose pull-down assay. Lane 1 contained partially purified DNA-PK without dsDNA. The protein-dsDNA cellulose complex was analyzed by the procedure described under *Materials and Methods*. Panel C, effect of HNI-38 on DNA binding activity of Ku70/Ku80 complex. Purified Ku70/Ku80 complex (100 ng) was incubated with 10 nM (lanes 4 and 7), 50 nM (lanes 3 and 6), and 100 nM (lanes 2 and 5) concentrations of either control peptide or target peptide prior to dsDNA cellulose pull-down assay. After the dsDNA cellulose pull-down, Ku70 and Ku80 were analyzed by 10% SDS-PAGE and Western blot.

(HNI-38) were incubated with cell extracts containing DNA-PKcs and Ku complex in the presence of dsDNA cellulose, and examined for its effect on binding of Ku complex and DNA-PKcs to DNA after the dsDNA cellulose pull-down assay (Fig. 2B). Although it was marginal, the addition of increasing amounts of target peptide (HNI-38), not control peptide (HN-26), led to a decrease in DNA-PKcs associated with dsDNA, suggesting that target peptide binds to DNA-PKcs and inhibits its binding to Ku70/Ku80. It is also noted that the addition of target peptide affected the binding of Ku70/Ku80 to the dsDNA cellulose (Fig. 2B). To further examine the effect of HNI-38 on the DNA binding activity of Ku, target peptide was incubated with purified Ku70/Ku80 complex in the presence of dsDNA cellulose, and the reaction mixtures were analyzed for the presence of Ku70 and Ku80 after the dsDNA pull-down assay (Fig. 2C). In keeping with Fig. 2B, target peptide (HNI-38) significantly interfered with binding of Ku complex to dsDNA under the conditions where control peptide (HN-26) showed virtually no effect (Fig. 2C). This result suggests that target peptide not only affects the interaction between DNA-PKcs and Ku complex, but also interferes with the DNA binding activity of Ku.

Effect of Target Peptide (HNI-38) on DNA-PK Kinase Activity. Interaction of DNA-PKcs with Ku complex is necessary for activation of its kinase activity (Gottlieb and Jackson, 1993; Hartley et al., 1995); therefore, the efficacy of target peptide was analyzed by measuring DNA-PK kinase activity in vitro in the presence of either HN-26 or HNI-38. DNA-PK kinase activity was inhibited up to 50% in the presence of HNI-38 under the conditions where a control peptide (HN-26) showed minimal effect (Fig. 3), strongly supporting the notion that target peptide specifically binds to DNA-PKcs and interferes with interaction between DNA-PKcs and Ku complex. Inhibitory effect of target peptide on DNA-PK occurred at low peptide concentrations (<20 nM) and, in the presence of 20 nM or higher, both target and control peptides inhibited DNA-PK activity (data not shown).




Fig. 3. Effect of target peptide on DNA-PK kinase activity in vitro. Partially purified DNA-PK fraction was incubated with various concentrations of either control peptide or target peptide prior to the addition of substrate peptide and other components for DNA-PK kinase assay (see *Materials and Methods* for details). DNA-PK activity was measured as the relative amounts of ^{32} P transferred to the substrate peptide.

Target Peptide Interferes with Repair of Double-Stranded DNA Breaks Induced by IR. IR-induced double-stranded DNA breaks are efficiently repaired by an NHEJ process. Genetic and biochemical studies strongly indicated that DNA-PK plays an essential role in NHEJ (Blunt et al., 1995; Jin et al., 1997; Jeggo, 1998). Hence, an alternative way to determine the efficacy of peptide inhibitor is to measure the repair of double-stranded DNA breaks following IR. Breast cancer cells (NCI) grown in the presence of [^{14} C]thymidine (DiBiase et al., 2000) were treated with either a control or a target peptide for 24 h. After irradiation (40 Gy), cells were harvested at various time points, and intact chromosomal DNA and DSBs were separated by pulsed field gel electrophoresis (0.5% agarose). Treatment of NCI cells with IR (40 Gy) induced substantial amounts of dsDNA breaks, most of which were repaired within 4 h. Cells treated with target or control peptide did not show any difference in generating DSBs after IR (Fig. 4A; lane 2 versus lanes 8 and 14). On the other hand, cells treated with target peptide (Fig. 4A, lanes 14–18) compared with those treated with control peptide (Fig. 4A, lanes 8–12) showed a noticeable decrease in DSB repair activity. This result suggests that target peptide interfered with dsb repair in vivo through the targeted inhibition of DNA-PK activity.

Target Peptide Inhibits Breast Cancer Cell Growth Only in the Presence of DNA Damage. Cells lacking DNA-PK catalytic subunit showed increased sensitivity to




Fig. 4. Effect of target peptide on double-stranded DNA break repair. Breast cancer cells (NCI) grown in ^{14} C-containing media were treated with ionizing radiation (40 Gy) in the presence of no peptide (lanes 1–6), 50 nM control peptide (lanes 7–12), or 50 nM target peptide (lanes 13–18). After harvesting the cells at various time points, intact chromosomes and double-stranded DNA breaks (DSBs) were separated by gel electrophoresis (panel A, fluorography) and were quantified by liquid scintillation counter (panel B).

DNA-damaging drugs or IR (Kirchgessner et al., 1995; Lees-Miller et al., 1995), suggesting that DNA-PK activity is essential for DNA repair and cell survival upon DNA damage. We therefore tested whether a targeted inhibition of DNA-PK by a peptide, HNI-38, would sensitize breast cancer cells upon treatment of ionizing radiation or chemotherapeutic drug (cisplatin). Two breast cancer cell lines (NCI and MDA231) were treated with either control (HN-26) or target peptide (HNI-38) and tested for the efficacy of DNA-PK inhibitory peptide on lowering resistance of cells in response to ionizing radiation using a standard colony count cell survival assay. Neither control nor target peptides showed any effect on cell growth in the absence of ionizing radiation. However, cells treated with IR showed significant cell growth inhibi-

tion in the presence of target peptide but not with control peptide (Fig. 5A), suggesting that cell growth inhibition by target peptide occurs through targeting DNA-PK activity. Cells treated with cisplatin, although not as effective as those treated with ionizing radiation, also showed inhibitory effect on cell growth in the presence of HNI-38 (Fig. 5B).

Discussion

Many key human DNA repair pathways, such as double-strand break repair or nucleotide excision repair pathway, rely on multimeric polypeptide activities (Friedberg, 1996; Sancar, 1996; Wood, 1996; Lee, 2001). Interactions between damage recognition proteins and those proteins that report




Fig. 5. Effect of control (HN-26) and target (HNI-38) peptides on the growth of breast cancer cells treated with ionizing radiation (panel A) or cisplatin (panel B). Values expressed are means ($\pm$ S.E.) of the three replications (*, $p < 0.01$; **, $p < 0.01$). The clonogenic assay was used for the cells treated with ionizing radiation, and the cell survival assay (MTT) was used for those treated with cisplatin (see *Materials and Methods* for the detailed procedure).

the damage to downstream repair activities are crucial for DNA repair. DNA-PK is a key component of the NHEJ pathway with the unique property of being activated by double-stranded DNA breaks (Blunt et al., 1995). Earlier studies with drug-resistant and -sensitive cancer cells suggested that high level expression of DNA-PK leads to drug-resistant cells, whereas low DNA-PK activity was associated with drug-sensitive phenotype (Muller and Salles, 1997; Shen et al., 1997, 1998; Muller et al., 1998; Tew et al., 1998; Frit et al., 1999; Kim et al., 1999, 2000), implicating a role for DNA-PK in conferring cells becoming drug resistant in response to anticancer DNA-damaging drug. Since the interaction of DNA-PKs to its regulatory subunits, Ku70 and Ku80, is crucial for its function in DNA repair, a targeted inhibition of DNA-PK would sensitize drug resistance of cancer cells and facilitate cell killing. Therefore, we attempted to develop a peptide cotherapy strategy in which a low molecular weight peptide-based inhibitor specifically interferes with interaction between DNA-PKs and Ku complex.

A target peptide (HNI-38) containing the C terminus of Ku80 interfered with the interaction between DNA-PKs and Ku complex. This was a much anticipated result since the C terminus of Ku80 was previously identified as DNA-PKs interacting domain (Gell and Jackson, 1999). Inhibitory effect of HNI-38 on the interaction between DNA-PKs and Ku70/Ku80 directly affected its kinase activity, showing inhibition of DNA-PK activity up to 50% under the conditions where a control peptide (HN-26) showed very little effect (Fig. 3). However, addition of an excess amount of target peptide did not show any further inhibition of DNA-PK kinase activity (data not shown). This is likely due to the fact that DNA-PKs without Ku complex can still function as a kinase, although its activity is low. A target peptide (HNI-38) not only inhibited the interaction of DNA-PKs with Ku complex on dsDNA, but also affected the dsDNA binding activity of Ku (Fig. 2C). It is not clear how HNI-38 interferes with DNA binding activity of the Ku complex; however, the C terminus of both Ku70 and Ku80 has been shown to be important for heterodimer assembly as well as for DNA-termini binding (Wu and Lieber, 1996; Gell and Jackson, 1999). It is possible that HNI-38 may interfere with the Ku70-Ku80 interaction through its binding to Ku70, which would negatively influence the DNA-termini binding activity of Ku.

DNA-PK activity is essential for DNA repair as well as cell cycle arrest in response to DNA damage, which contributes to cell survival by protecting cells from apoptosis. Cells treated with target peptide but not control peptide showed a noticeable decrease in DSB repair after a high dose (40 Gy) of IR, suggesting that HNI-38 specifically targets DNA-PK in vivo and interferes with dsb repair activity through inhibition of DNA-PK activity. Targeted inhibition of DNA-PK by HNI-38 also caused cell growth inhibition only when cells were treated with IR, suggesting that HNI-38 targeted DNA-PK and lowered resistance of cells in response to ionizing radiation, which eventually causes growth inhibition of both NCI and MDA231. Treatment of cells with HNI-38 also showed additive effect on cell growth inhibition in response to cisplatin treatment. This observation is in keeping with previous findings that DNA-PK is directly involved in nucleotide excision repair action in mammals (Muller et al., 1998). It also supports the notion that a targeted inhibition of DNA-PK would sensitize cancer cells upon treatment of chemotherapeutic drugs such as cisplatin. Taken together, our study

results described here not only validate DNA-PK as a useful molecular target for the treatment of drug-resistant cancer cells, but also support a physiological role for DNA-PK in IR or chemotherapy drug resistance of cancer cells.

Acknowledgments

We thank Anita Riemen for technical assistance, Karen Pollock for assisting us in the use of the gamma irradiation facility (^{137}Cs) located in the Wells center, Indiana University School of Medicine, and John Hawes and the Biochemistry Biotechnology Facility for peptide synthesis.

References

- Blunt T, Finnie NJ, Taccioli GE, Smith GC, Demengeot J, Gottlieb TM, Mizuta R, Varghese AJ, Alt FW, Jeggo PA, et al. (1995) Defective DNA-dependent protein kinase activity is linked to V(D)J recombination and DNA repair defects associated with the murine scid mutation. *Cell* 80:813–823.
- Britten RA, Kuny S, and Perdue S (1999) Modification of non-conservative double-strand break (DSB) rejoining activity after the induction of cisplatin resistance in human tumour cells. *Br J Cancer* 79:843–849.
- Cary RB, Peterson SR, Wang J, Bear DG, Bradbury EM, and Chen DJ (1997) DNA looping by Ku and the DNA-dependent protein kinase. *Proc Natl Acad Sci USA* 94:4267–4272.
- Critchlow SE and Jackson SP (1998) DNA end-joining: from yeast to man. *Trends Biochem Sci* 23:394–398.
- DiBiase SJ, Zeng ZC, Chen R, Hyslop T, Curran WJ Jr, and Iliakis G (2000) DNA-dependent protein kinase stimulates an independently active, nonhomologous, end-joining apparatus. *Cancer Res* 60:1245–1253.
- Featherstone C and Jackson SP (1999) Ku, a DNA repair protein with multiple cellular functions? *Mutat Res* 434:3–15.
- Friedberg EC (1996) Relationship between DNA repair and transcription. *Annu Rev Biochem* 65:15–42.
- Frit P, Canitrot Y, Muller C, Foray N, Calsou P, Marangoni E, Bourhis J, and Salles B (1999) Cross-resistance to ionizing radiation in a murine leukemic cell line resistant to cis-dichlorodiammineplatinum(II): role of Ku autoantigen. *Mol Pharmacol* 56:141–146.
- Gell D and Jackson SP (1999) Mapping of protein-protein interactions within the DNA-dependent protein kinase complex. *Nucleic Acids Res* 27:3494–3502.
- Gottlieb TM and Jackson SP (1993) The DNA-dependent protein kinase: requirement for DNA ends and association with Ku antigen. *Cell* 72:131–142.
- Hammarsten O and Chu G (1998) DNA-dependent protein kinase: DNA binding and activation in the absence of Ku. *Proc Natl Acad Sci USA* 95:525–530.
- Hanawalt PC (1994) Transcription-coupled repair and human disease. *Science (Wash DC)* 266:1957–1958.
- Hartley KO, Gell D, Smith GC, Zhang H, Divecha N, Connelly MA, Admon A, Lees-Miller SP, Anderson CW, and Jackson SP (1995) DNA-dependent protein kinase catalytic subunit: a relative of phosphatidylinositol 3-kinase and the ataxia telangiectasia gene product. *Cell* 82:849–856.
- Jackson SP (1997) DNA-dependent protein kinase. *Int J Biochem Cell Biol* 29:935–938.
- Jeggo PA (1998) DNA breakage and repair. *Adv Genet* 38:185–218.
- Jin S, Inoue S, and Weaver DT (1997) Functions of the DNA dependent protein kinase. *Cancer Surv* 29:221–261.
- Kim SH, Um JH, Dong-Won B, Kwon BH, Kim DW, Chung BS, and Kang CD (2000) Potentiation of chemosensitivity in multidrug-resistant human leukemia CEM cells by inhibition of DNA-dependent protein kinase using wortmannin. *Leuk Res* 24:917–925.
- Kim SH, Kim D, Han JS, Jeong CS, Chung BS, Kang CD, and Li GC (1999) Ku autoantigen affects the susceptibility to anticancer drugs. *Cancer Res* 59:4012–4017.
- Kirchgessner CU, Patil CK, Evans JW, Cuomo CA, Fried LM, Carter T, Oettinger MA, and Brown JM (1995) DNA-dependent kinase (p350) as a candidate gene for the murine SCID defect. *Science (Wash DC)* 267:1178–1183.
- Kurimasa A, Kumano S, Boubnov NV, Story MD, Tung CS, Peterson SR, and Chen DJ (1999) Requirement for the kinase activity of human DNA-dependent protein kinase catalytic subunit in DNA strand break rejoining. *Mol Cell Biol* 19:3877–3884.
- Lee S-H (2001) Recognition of DNA damage in mammals. *J Biochem Mol Biol* 34:489–495.
- Lee S-H and Kim C-H (2002) DNA-dependent protein kinase complex: a multifunctional protein in DNA repair and damage checkpoint. *Mol Cells* 13:159–166.
- Lees-Miller SP, Chen Y-R, and Anderson CW (1990) Human cells contain a DNA-activated protein kinase that phosphorylates simian virus 40 T antigen, mouse p53 and the human Ku autoantigen. *Mol Cell Biol* 10:6472–6481.
- Lees-Miller SP, Godbout R, Chan DW, Weinfeld M, Day RS 3rd, Barron GM, and Allalunis-Turner J (1995) Absence of p350 subunit of DNA-activated protein kinase from a radiosensitive human cell line. *Science (Wash DC)* 267:1183–1185.
- Lieber MR, Grawunder U, Wu X, and Yanava M (1997) Tying loose ends: roles of Ku and DNA-dependent protein kinase in the repair of double-strand breaks. *Curr Opin Genet Dev* 7:99–104.
- Lin Y-Z, Yi SY, Veach RA, Torgerson TR, and Hawiger J (1995) Inhibition of nuclear translocation of transcription factor NF- κ B by a synthetic peptide containing a cell membrane-permeable motif and nuclear localization sequence. *J Biol Chem* 270:14255–14258.
- Muller C, Calsou P, Frit P, Cayrol C, Carter T, and Salles B (1998) UV sensitivity

- and impaired nucleotide excision repair in DNA-dependent protein kinase mutant cells. *Nucleic Acids Res* **26**:1382-1389.
- Muller C and Salles B (1997) Regulation of DNA-dependent protein kinase activity in leukemic cells. *Oncogene* **15**:2343-2348.
- Park J-S, Park S-J, Peng X, Yu M, Wang M, and Lee S-H (1999) Involvement of DNA-dependent protein kinase in UV-damage induced replication arrest. *J Biol Chem* **274**:32520-32527.
- Sancar A (1996) DNA excision repair. *Annu Rev Biochem* **65**:43-81.
- Shen H, Kauvar L, and Tew KD (1997) Importance of glutathione and associated enzymes in drug response. *Oncol Res* **9**:295-302.
- Shen H, Schultz M, Kruh GD, and Tew KD (1998) Increased expression of DNA-dependent protein kinase confers resistance to adriamycin. *Biochim Biophys Acta* **23**:131-138.
- Suwa A, Hirakata M, Takeda Y, Jesch SA, Mimori T, and Hardin JA (1994) DNA-dependent protein kinase (Ku protein-p350 complex) assembles on double-stranded DNA. *Proc Natl Acad Sci USA* **91**:6904-6908.
- Tew KD, O'Brien M, Laing NM, and Shen H (1998) Coordinate changes in expression of protective genes in drug-resistant cells. *Chem-Biol Interact* **111**:199-211.
- Weaver DT and Alt FW (1997) V(D)J recombination. From RAGs to stitches. *Nature (Lond)* **388**:428-429.
- Woo RA, McLure KG, Lees-Miller SP, Rancourt DE, and Lee PWK (1998) DNA-dependent protein kinase acts upstream of p53 in response to DNA damage. *Nature (Lond)* **394**:700-704.
- Wood RD (1996) DNA repair in eukaryotes. *Annu Rev Biochem* **65**:135-167.
- Wu X and Lieber MR (1996) Protein-protein and protein-DNA interaction regions within the DNA end-binding protein Ku70-Ku86. *Mol Cell Biol* **16**:5186-5193.

Address correspondence to: Suk-Hee Lee, Department of Biochemistry and Molecular Biology, Indiana University School of Medicine, 635 Barnhill Dr., Indianapolis, IN 46202. E-mail: slee@iupui.edu

Biochemical Analysis of the Damage Recognition Process in Nucleotide Excision Repair*

Received for publication, October 16, 2002, and in revised form, November 20, 2002
Published, JBC Papers in Press, December 13, 2002, DOI 10.1074/jbc.M210603200

Jin-Sam You, Mu Wang†, and Suk-Hee Lee§

From the Department of Biochemistry and Molecular Biology, Indiana University Cancer Center, and Walther Oncology Center, Indiana University School of Medicine, Indianapolis, Indiana 46202

XPA, XPC-hHR23B, RPA, and TFIIH all are the damage recognition proteins essential for the early stage of nucleotide excision repair. Nonetheless, it is not clear how these proteins work together at the damaged DNA site. To get insight into the molecular mechanism of damage recognition, we carried out a comprehensive analysis on the interaction between damage recognition proteins and their assembly on damaged DNA. XPC physically interacted with XPA, but failed to stabilize the XPA-damaged DNA complex. Instead, XPC-hHR23B was effectively displaced from the damaged DNA by the combined action of RPA and XPA. A mutant RPA lacking the XPA interaction domain failed to displace XPC-hHR23B from damaged DNA, suggesting that XPA and RPA cooperate with each other to destabilize the XPC-hHR23B-damaged DNA complex. Interestingly, the presence of hHR23B significantly increased RPA/XPA-mediated displacement of XPC from damaged DNA, suggesting that hHR23B may modulate the binding of XPC to damaged DNA. Together, our results suggest that damage recognition occurs in a multistep process such that XPC-hHR23B initiates damage recognition, which was replaced by combined action of XPA and RPA. XPA and RPA, once forming a complex at the damage site, would likely work with TFIIH, XPG, and ERCC1-XPF for dual incision.

Nucleotide excision repair (NER)¹ is one of the major repair pathways for removal of DNA damage caused by UV irradiation and a wide variety of bulky helix-distorting lesions such as cisplatin (1–4). In mammals, NER requires over 20 polypeptides, including damage recognition and/or structure distortion factors (XPA, XPC-hHR23B, replication protein A (RPA), and a transcription factor, TFIIH), strand separating helicases to create an open preincision complex (TFIIH containing XPB and

XPB DNA helicases), two structure-specific endonucleases (ERCC1-XPF and XPG), and the enzymes needed for gap filling (DNA polymerase δ/ϵ , proliferating cell nuclear antigen, replication factor C, and RPA).

Both RPA and XPA are also known as damage recognition proteins because they preferentially bind to cisplatin- or UV-damaged DNA (5–10). Both proteins may also play a role in subsequent steps in NER through interaction with other repair proteins (8, 11–15). The XPA-DNA interaction is relatively weak and characterized by rapid dissociation, whereas RPA formed a much more stable complex with UV-damaged DNA (16). XPA physically interacts with RPA, which is necessary for efficient NER action (14). Wild-type RPA, but not a mutant lacking the XPA interaction domain, led to stabilization of XPA-damaged DNA complex, implicating a unique role for RPA in stabilizing the XPA-damaged DNA complex. The XPA-damaged DNA interaction is also likely necessary for recruiting other DNA repair proteins such as XPG, ERCC1-XPF, and TFIIH to the damaged site (8, 15, 17). RPA may also be involved in the later stage of NER, gap-filling, that requires proliferating cell nuclear antigen, replication factor C, and DNA polymerase δ (or ϵ) (18).

XPC-hHR23B is a human homolog of yeast Rad4 and Rad23 proteins, respectively, and forms a stable complex in solution. XPC-hHR23B exhibits the strongest affinity for damaged DNA (19–21), as does the yeast counterpart, Rad4-Rad23 (22). Rad23 without Rad4 does not show any DNA binding activity, suggesting that Rad4 is solely responsible for recognition of damaged DNA. Rad23 is essential for XPC function in NER and may also be necessary for the solubility of Rad4 (23). XPC-hHR23B showed a remarkable preference to UV-damaged DNA particularly in the presence of nondamaged competitor DNA and has been suggested as the initiator of global genome NER (19, 24). A recent immunohistochemistry study also strongly supports a role for XPC as a global initiator in repair (25), while suggesting a role for XPA and RPA as repair mediator proteins. XPC-hHR23B is also involved in the recruitment of TFIIH to damaged DNA (26). TFIIH, once recruited, may play a role in distinguishing the damaged strand from the nondamaged one (27) as well as local unwinding of the damaged DNA region. TFIIH with its DNA helicase activity likely generates a junction between single-stranded DNA and duplex DNA that is recognized by two structure-specific endonucleases, XPG and ERCC1-XPF, for dual incision of damaged strand.

An ongoing challenge is to understand how DNA damage is recognized and distinguished from nondamaged sites. In mammalian cells, XPC-hHR23B, XPA, RPA, and TFIIH factors may all have roles in damage recognition during the early stage of NER. In this study we carried out a comprehensive analysis on the interaction between damage recognition proteins and their assembly on damaged DNA. We found that XPC-hHR23B, like

* This work was supported in part by National Institutes of Health Grant CA92111 and United States Army Grant DAMD17-00-1-0295. The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

† Supported by the National Institutes of Health Postdoctoral Fellowship F32 GM20167-01.

§ To whom correspondence should be addressed: 1044 W. Walnut St., Rm. 153, Indiana University Cancer Research Institute Building, Indianapolis, IN 46202. Tel.: 317-278-3464; Fax: 317-274-8046; E-mail: slee@iupui.edu.

¹ The abbreviations used are: NER, nucleotide excision repair; ERCC, excision repair cross-complementing; GST, glutathione *S*-transferase; hHR23, human homolog of the yeast RAD23; ITR-60, intrastrand cross-linked duplex DNA of 60-mer; RPA, replication protein A; ssDNA, single-stranded DNA; dsDNA, double-stranded DNA; TFIIH, transcription factor IIH; XP, xeroderma pigmentosum; TET, tetrachloro-fluorescein.

RPA, physically interacted with XPA. However, the XPA-XPC interaction, unlike the XPA-RPA interaction, failed to stabilize the XPA-damaged DNA complex. Instead, XPA cooperates with RPA to promote the destabilization of the XPC-hHR23B-damaged DNA interaction. This finding supports a notion that the damage recognition process occurs in a stepwise manner such that XPC-hHR23B initiates damage recognition, which was replaced by the combined action of XPA and RPA.

MATERIALS AND METHODS

Preparation of Platinum-induced Damaged DNA—To study the interaction between damaged DNA and damage recognition factors, we constructed a duplex DNA with the cisplatin lesion at a specific site (Fig. 1A). Oligonucleotides containing an intrastrand (ITR-60) cross-link were prepared according to the previously described procedure with some modifications (28). For cross-linking, the top strand was first incubated with cisplatin (in TE, pH 8.0, 2-fold molar excess) at 37 °C in the dark for 48 h, and then ethanol-precipitated. The damaged DNA was purified by 15% denaturing polyacrylamide gel electrophoresis and annealed to the bottom strand (5-fold molar excess). The duplex DNA was purified by 15% native polyacrylamide gel electrophoresis. Purified duplex DNA, 5'-³²P-labeled at top strand, was analyzed for intrastrand cross-link by treatment with 10% dimethyl sulfate at room temperature for 5 min and 1 M piperidine at 90 °C for 30 min, followed by 15% denaturing polyacrylamide gel electrophoresis (Fig. 1B).

Purification of XPA, RPA, and XPC-hHR23B—Both RPA and XPC-hHR23B are multiprotein complexes that can be highly expressed in insect cells by coinfecting recombinant baculoviruses (21, 29). RPA complex was prepared by coinfecting recombinant baculoviruses encoding the individual subunits (29) and the XPC-hHR23B complex was from insect cell lysates infecting a recombinant baculovirus containing both XPC and hHR23B genes (Ref. 21; recombinant baculovirus containing XPC and hHR23B was kindly provided by Dr. A. Sancar, University of North Carolina, Chapel Hill, NC). A high level of histidine-tagged XPA was expressed in *Escherichia coli* under control of the T7 promoter (5). Both RPA and XPA have been isolated with more than 95% purity using a series of conventional column chromatography as described previously (5, 13, 29) (Fig. 1C). XPC-hHR23B complex was purified using a series of column chromatography (UV-damaged dsDNA cellulose, phosphocellulose (P11), and heparin-Sepharose) (Fig. 1C, the details of XPC-hHR23B purification will be described elsewhere). All three proteins (XPA, RPA, and XPC-hHR23B) were functionally active in supporting *in vitro* NER activity with HeLa cell extracts lacking respective protein (data not shown). All GST fusion proteins were purified using glutathione-Sepharose affinity column as described previously (14). TFIIF was isolated from HeLa cell nuclear extracts using an antibody affinity procedure described previously (31). During purification, TFIIF was monitored by Western blot using anti-XPB and -XPD antibodies (Santa Cruz Biotechnology Inc., Santa Cruz, CA).

GST Fusion Pull-down Assay—The indicated amount of GST fusion protein was incubated with glutathione-Sepharose beads (25 μ l) in 0.5% nonfat milk at 4 °C for 30 min, and washed three times with washing buffer (50 mM Tris-HCl, pH 8.0, 1 mM EDTA, 1 mM dithiothreitol, 10% glycerol, 0.5% Nonidet P-40, 150 mM NaCl, 200 μ g/ml bovine serum albumin). After washing the beads 3–5 times, beads were incubated with the indicated protein either in the presence or absence of damaged DNA for an additional 20 min at room temperature on a rocker and then washed 5 times with washing buffer. Beads were then mixed with 25 μ l of SDS-PAGE sample buffer, heated at 95 °C for 5 min, and proteins (or DNA) were resolved on 10% SDS-PAGE. To quantitatively measure the damaged DNA, the gel was directly scanned on a Hitachi FMBIO II Fluorescent Image Scanner.

SDS-PAGE and Western Blot Analysis—Protein samples were separated by 10% SDS-PAGE as described previously (32). Proteins were then transferred to polyvinylidene difluoride membranes, immunoblotted with corresponding antibodies, and detected by ECL plus Western blotting detection reagents (Amersham Biosciences).

Protein-damaged DNA Interaction: Surface Plasmon Resonance Analysis—Interactions of XPA, RPA, and XPC-hHR23B with DNA were monitored using a surface plasmon resonance biosensor instrument, Biacore 3000 (Biacore) as described previously (16, 34). For preparation of the biosensor surface with DNA, 5'-biotinylated 60-mer duplex DNA (prepared in a buffer containing 10 mM sodium acetate, pH 4.8, and 1.0 M NaCl) was manually injected onto a streptavidin-coated surface of a Biacore sensor chip to the desired density in different flow cells. One flow cell was left underivatized to allow for refractive index change

correction. Proteins were diluted in the running buffer containing 10 mM Hepes, pH 7.4, 150 mM NaCl, 2 mM MgCl₂, 0.005% polysorbate-20, and 1 mM dithiothreitol. Each experiment was repeated at least twice to assure reproducibility.

RESULTS

Interaction of Damage Recognition Factors with DNA—To get insight into the molecular mechanism of how damage recognition proteins are assembled and working together at the damaged DNA site, we carried out a comprehensive analysis on the interaction of XPA, RPA, and XPC-hHR23B with damaged DNA. Binding affinity of damage recognition factors to a duplex DNA containing an intrastrand platinum cross-link (ITR-60, Fig. 1) was measured using an electrophoretic mobility shift assay. As previously reported (Refs. 1–4 and references therein), both RPA and XPC-hHR23B showed preferential binding to the cisplatin-damaged DNA over the nondamaged one, whereas the XPA-DNA interaction was observed only in the presence of excess amounts (Fig. 2A). On the other hand, surface plasmon resonance kinetic analysis indicated that XPC-hHR23B compared with RPA exhibited much higher affinity to the damaged DNA (Fig. 2B). This finding is in keeping with the previous observation that XPC has a considerable preference for binding to the damaged DNA over nondamaged competitor DNA (24, 25). XPC-hHR23B also differs from RPA in its preferential binding to dsDNA over ssDNA (35), whereas RPA has a much higher affinity to ssDNA than to dsDNA (Fig. 2, A and B).

XPC Physically Interacts with XPA, but Not with RPA—Because all three proteins (RPA, XPA, and XPC-hHR23B) independently bind to the damaged DNA, we examined a possibility whether all three damage recognition proteins form a complex on the damaged DNA site (11–13). Initially, we examined the interactions among damage recognition factors without DNA. Because we know that XPA and RPA interact with each other in solution as well as on damaged DNA (14, 17), our focus was on the interaction of XPC-hHR23B with RPA or XPA (Fig. 3). GST-XPC or GST-XPC-hHR23B were incubated with an increasing amount of XPA (or RPA), pulled down with glutathione-Sepharose beads for co-precipitation of XPA (or RPA), and analyzed by Western blot (Fig. 3). XPA was co-precipitated with GST-XPC or GST-XPC-hHR23B complex, but not with GST (Fig. 3A), suggesting that XPA physically interacts with XPC-hHR23B. In the XPA-XPC interaction, GST-XPC was almost comparable with GST-XPC-hHR23B (Fig. 3A, lanes 3–4 versus lanes 5 and 6), indicating that XPC not hHR23B physically interacts with XPA. In keeping with this, hHR23B without XPC failed to interact with XPA (data not shown). Unlike XPA, however, RPA had no physical interaction with GST-XPC-hHR23B or GST-XPC in solution (Fig. 3B).

Because both RPA and XPC interact with XPA (Refs. 11–13 and Fig. 3), it is possible that the binding of XPC (or RPA) to XPA may competitively exclude RPA (or XPC) from the complex. If so, we may not be able to see a complex containing all three damage recognition proteins. To examine this possibility, XPA was incubated with GST-XPC-hHR23B in the presence of an increasing amount of RPA and analyzed for the XPA-XPC interaction. Addition of a molar excess of RPA had little or no effect on the XPA-XPC interaction as determined by the GST-XPC-hHR23B pull-down assay (Fig. 3C), suggesting that RPA and XPC-hHR23B may recognize two separate domains of XPA. Nonetheless, we were not able to detect RPA in the GST-XPC pull-down assay in the presence of XPA and RPA (Fig. 3C).

Presence of Damaged DNA Significantly Inhibits the XPA-XPC Interaction—The XPA-damaged DNA interaction is weak, but markedly stimulated in the presence of a high affinity DNA-binding protein, RPA (14). The stimulatory effect of RPA

A**Intra-strand crosslinked duplex DNA (ITR-60)**

5'-TTC CTC CTT CTT CTC TTC CTC CTT CTT GGT TCT CTT CTT CTC CTC TTC CTC CTT CTT CTC-3'
 3'-AAG GAG GAA GAA GAG AAG GAG GAA GAA CCA AGA GAA GAA GAG GAG AAG GAG GAA GAA GAG-5'



B**C**

FIG. 1. *Panel A*, cisplatin-induced intrastrand cross-linked DNA (60-mer; ITR-60). *Underlined* guanine residues (*G*) are the reactive sites for platination. *Panel B*, the duplex DNA with intrastrand cross-link (ITR-60) showed slower mobility on a 15% denatured polyacrylamide gel electrophoresis (*lanes 1 versus 3*). The Maxam-Gilbert G-residue reaction (DMS + piperidine treatment) showed protection of ^{32} P-labeled DNA from cleavage in intrastrand cross-linked DNA (*lanes 2 versus 4*). Cisplatin-damaged duplex DNA was labeled at the 5'-end of the nondamage-containing strand with fluorescence dye (TET, synthesized from the Integrated DNA Technologies, Coralville, IA). *Panel C*, SDS-PAGE and Western blot analysis of purified XPA, RPA, and XPC and/or hHR23B. Individual proteins were electrophoretically separated by 12% SDS-PAGE and visualized by Coomassie Blue staining (*left lane*; see "Materials and Methods" for the details) as well as by immunoblot using the corresponding polyclonal antibody (*right lane*).

on the XPA-damaged DNA interaction occurs through the RPA-XPA interaction (14, 16). Similar to RPA, XPC-hHR23B exhibits a high affinity to damaged DNA (19–23) and physically

interacts with XPA (Fig. 3). We therefore examined whether XPC-hHR23B affects the XPA-damaged DNA interaction. In the GST-XPA pull-down assay, XPA had very little or no bind-




FIG. 2. Binding characteristics of XPA, RPA, and XPC-hHR23B to the cisplatin-damaged DNA (ITR-60). Panel A, increasing amounts of XPC-hHR23B (top panel; lanes 1, 6, and 11, 0 ng; lanes 2, 7, and 12, 20 ng; lanes 3, 8, and 13, 40 ng; lanes 4, 9, and 14, 60 ng; lanes 5, 10, and 15, 100 ng), RPA (middle panel; lanes 1, 6, and 11, 0 ng; lanes 2, 7, and 12, 10 ng; lanes 3, 8, and 13, 20 ng; lanes 4, 9, and 14, 40 ng; lanes 5, 10, and 15, 80 ng), or XPA (bottom panel; lanes 1, 6, and 11, 0 ng; lanes 2, 7, and 12, 35 ng; lanes 3, 8, and 13, 70 ng; lanes 4, 9, and 14, 175 ng; lanes 5, 10, and 15, 350 ng) was incubated with 100 fmol of 5'-³²P-damaged duplex DNA (ITR-60; lanes 1–5), 5'-³²P-nondamaged duplex DNA (lanes 6–10), or 5'-³²P-ssDNA (lanes 11–15) for 15 min at room temperature. The protein-DNA complex was analyzed by 4% polyacrylamide gel in 0.5× TBE (acrylamide:bisacrylamide = 43.2:0.8). For quantification, regions corresponding to protein-DNA complex were excised and measured for radioactivity. Asterisk indicates the ssDNA fragment (60-mer) generated from ITR-60 during gel purification. Panel B, biomolecular interaction (BIAcore) analysis of the interaction of XPC-hHR23B and RPA with damaged DNA (ITR-60). XPC-hHR23B or RPA (25 nM) was injected into ssDNA (SS), nondamaged duplex DNA (ND), or cisplatin-damaged DNA (platinum; Pt) surface (50 response units) using the KINJECT function of BIAcore 3000. The association phase was allowed for 180 s followed by a 120-s buffer injection period for dissociation (see "Materials and Methods" for details).

ing to the damaged DNA (Fig. 4A, lane 2), which was significantly stimulated by addition of RPA (Fig. 4A, lanes 3–5). In contrast, XPC-hHR23B showed no effect on the XPA-damaged DNA interaction (Fig. 4A, lanes 6–8). Interestingly, the stimulatory effect of RPA on the XPA-damaged DNA interaction was markedly reduced in the presence of XPC-hHR23B (Fig. 4A, lane 9), suggesting that XPC-hHR23B, unlike RPA, may not form a stable complex with XPA on damaged DNA. To investigate this further, we examined the effect of damaged DNA on the XPA-XPC interaction (Fig. 4B). XPA was success-

fully co-precipitated with GST-XPC in the pull-down assay in the absence of damaged DNA (Fig. 4B, lanes 4 and 5). In the presence of damaged DNA, however, the amount of XPA co-precipitated with GST-XPC was significantly reduced to an undetectable level (Fig. 4B, lanes 6 and 7), suggesting that the XPA-XPC interaction is destabilized in the presence of damaged DNA.

XPC-hHR23B Is Displaced from Damaged DNA by the Combined Action of RPA and XPA—Our result (Fig. 4) strongly suggests that XPA, RPA, and XPC-hHR23B do not form a




FIG. 3. XPC interacts with XPA, but not with RPA. Panel A, 2 pmol of GST-XPC or GST-XPC-hHR23B was mixed with glutathione-Sepharose beads (25 μ l) and gently rocked in the presence of XPA (2 pmol (lanes 3 and 5) and 5 pmol (lanes 4 and 6)) at room temperature for 20 min. After incubation, beads were washed with buffer (50 mM Tris-HCl, pH 8.0, 1 mM EDTA, 1 mM dithiothreitol, 10% glycerol, 0.5% Nonidet P-40, 150 mM NaCl, 200 μ g/ml bovine serum albumin), and the bound XPA was analyzed by Western blot using an anti-XPA polyclonal antibody. Panel B, GST-XPC or GST-XPC-hHR23B (2 pmol) was mixed with glutathione beads (25 μ l) in the presence of RPA (2 pmol (lanes 3 and 5) or 5 pmol (lanes 4 and 6)) at room temperature for 20 min. After incubation, beads were washed with buffer and bound RPA was analyzed by Western blot using an anti-RPA p34 polyclonal antibody. Lane 1 contained purified RPA as a control.




FIG. 4. Interaction between XPC-hHR23B and XPA on damaged DNA. Panel A, RPA not XPC stabilizes the XPA-damaged DNA interaction. GST-XPA (2 pmol) was incubated with 2 pmol of 5'-TET-labeled cisplatin-damaged DNA (60-mer) in the presence of increasing amounts of RPA (2, 5, and 10 pmol in lanes 3-5, respectively) or XPC-hHR23B (2, 5, and 10 pmol in lanes 6-8, respectively). Following the GST pull-down assay (as described in the legend to Fig. 3A), precipitates were separated by 10% SDS-PAGE and analyzed for TET-labeled damaged DNA using a fluorescent image scanner (Hitachi FMBIO II) (top panel) or GST-XPA using an anti-GST antibody (bottom panel). Panel B, the presence of damaged DNA significantly inhibits the XPA-XPC interaction. Two pmol of GST (lane 2) or GST-XPC-hHR23B (lanes 3-7) was incubated with increasing amount of XPA (2 pmol in lanes 2, 4, and 7; 5 pmol in lanes 5 and 7) for 20 min at room temperature. Where indicated, 2 pmol of cisplatin-damaged DNA (ITR-60) was included. Following the GST pull-down, precipitates were separated (10% SDS-PAGE) and analyzed by Western blot using an anti-XPA (top panel) or -GST (bottom panel) antibody.

stable complex on the damaged DNA. Instead, the damage recognition process may occur in a stepwise manner such that XPC-hHR23B interacts with damaged DNA first and then is replaced by XPA and/or RPA. To test this, biotin-labeled cisplatin-damaged DNA was first incubated with XPC-hHR23B, followed by the addition of RPA (Fig. 5A, lanes 3-5), XPA (lanes 6-8), or XPA + RPA (lanes 9-11) to the reaction mixtures. The streptavidin-Sepharose pull-down assay revealed that XPC-hHR23B formed a stable complex with damaged DNA (Fig. 5A,

top panel, lane 2) and, the addition of RPA (lanes 3-5) or XPA (lanes 6-8) showed very little effect on the XPC-hHR23B-damaged DNA interaction. In the presence of both XPA and RPA, however, the XPC-hHR23B-damaged DNA interaction was significantly inhibited (Fig. 5A, top panel, lanes 9-11). In contrast to XPC, the amount of XPA co-precipitated with damaged DNA was markedly increased in the presence of RPA (Fig. 5A, second panel, lanes 9-11), whereas the RPA-damaged DNA interaction was hardly affected by the presence of XPA (Fig.





FIG. 5. XPC-hHR23B is displaced from the damaged DNA by a combined action of XPA and RPA. *Panel A*, Western blot analysis of individual damage recognition factors associated with biotin-labeled damaged DNA. Biotin-labeled damaged DNA (ITR-60; 2 pmol) was preincubated with streptavidin-Sepharose prior to the addition of GST-XPC-hHR23B (800 ng in lanes 1–11), RPA (2, 5, and 10 pmol in lanes 3–5, respectively), XPA (2, 5, and 10 pmol in lanes 6–8, respectively), and XPA + RPA (2, 5, and 10 pmol each in lanes 9–11, respectively). Following the pull-down of beads, proteins were analyzed by Western blot using an anti-GST (for GST-XPC), -XPA, -RPAp70, or RPAp34 antibody. In lane M, purified protein (GST-XPC, XPA, or RPA for p70 and p34) was included. *Panel B*, glutathione-Sepharose beads (25 μ l) were mixed with GST-XPC-hHR23B (400 ng) and fluorescence (TET)-labeled damaged DNA (2 pmol), and gently rocked for 20 min at room temperature. After challenging the complex with either XPA (2, 5, and 10 pmol in lanes 3–5, respectively) or XPA + RPA (2, 5, and 10 pmol each in lanes 6–8, respectively), beads were pulled down and analyzed by 10% SDS-PAGE for TET-labeled damaged DNA using fluorescence image scanner (*top panel*) or Western blot using an anti-GST antibody (*bottom panel*). Relative amounts of TET-labeled damaged DNA are indicated at the bottom of the figure. *Panel C*, effect of RPA and XPA on the interaction between XPC and damaged DNA in a gel mobility shift assay. GST-XPC-hHR23B (20 ng) was incubated with 100 fmol of ^{32} P-platinum-damaged DNA (ITR-60) and incubated for 20 min at room temperature. Where indicated, XPA (20 ng in lanes 4, 10, 12, and 14; 50 ng in lanes 5, 11, 13, and 15), RPA (20 ng in lanes 7, 10, 12, and 14; 50 ng in lanes 8, 11, 13, and 15), or an anti-GST antibody (5 μ l; 100 μ g/ml) was added to the mixtures and further incubated for 20 min at room temperature prior to gel electrophoresis. The protein-DNA complex was analyzed by 4% polyacrylamide gel in 0.5 $\times$ TBE (acrylamide:bisacrylamide = 43.2:0.8). GST/XPC-damaged DNA complex and its supershifted complexes are indicated by arrows. No protein was included in lane 1.

5A, third and fourth panels, lanes 9–11). Together, this result suggests that XPC-hHR23B can be effectively displaced from damaged DNA by the combined action of RPA and XPA.

We further analyzed the effect of XPA and/or RPA on the interaction between XPC-hHR23B and damaged DNA by measuring damaged DNA co-precipitated with GST-XPC-hHR23B in the glutathione-Sepharose pull-down assay. Fluorescence

(TET)-labeled damaged DNA was first incubated with GST-XPC-hHR23B before the addition of either XPA or XPA + RPA and analyzed quantitatively following GST-XPC pull-down assay (Fig. 5B). Addition of both XPA and RPA drastically reduced the amount of damaged DNA interacting with GST-XPC-hHR23B, whereas XPA alone marginally affected the XPC-damaged DNA interaction (Fig. 5B), suggesting that

A



B




FIG. 6. hHR23B is necessary for the displacement of XPC from damaged DNA. Panel A, XPC-hHR23B not XPC is effectively displaced from damaged DNA. Two pmol of either GST-XPC (lanes 2–4) or GST-XPC-hHR23B (lanes 5–7) was mixed with glutathione-Sepharose beads (25 μ l) and TET-labeled cisplatin-damaged DNA (2 pmol) and rocked for 20 min at room temperature. After adding increasing amounts of RPA + XPA (2 pmol (lanes 3 and 6) or 5 pmol (lanes 4, 5, and 7)), beads were washed and analyzed for TET-labeled cisplatin-damaged DNA by 10% SDS-PAGE. Relative amounts of TET-labeled DNA are indicated at the bottom of the figure. Panel B, hHR23B stimulates XPA/RPA-mediated displacement of XPC from the damaged DNA. Two pmol of GST-XPC (lanes 2–7) or GST-XPC-hHR23B (lanes 8–10) was mixed with glutathione beads (25 μ l) and TET-labeled damaged DNA (2 pmol), and where indicated, 2 pmol of purified hHR23B was included.

both RPA and XPA are necessary to destabilize the interaction between XPC-hHR23B and damaged DNA. Effect of XPA and/or RPA on the GST-XPC-hHR23B-damaged DNA interaction was also analyzed by an antibody supershift assay using an anti-GST antibody (Fig. 5C). On a native gel electrophoresis, GST-XPC-hHR23B-damaged DNA complex was identified as a distinct band (Fig. 5C, lane 2) that was supershifted in the presence of an anti-GST antibody (Fig. 5C, lane 3). The GST-XPC-hHR23B-damaged DNA complex (lanes 12 and 13) or its supershifted complex (lanes 14 and 15) was significantly reduced by addition of an increasing amount of both RPA and XPA, even though the supershift may be in part because of the reaction of GST antibody to XPA and/or RPA (Fig. 5C). This result is in keeping with Fig. 5, A and B, indicating that XPC-hHR23B is displaced from damaged DNA by the combined action of RPA and XPA.

hHR23B Is Necessary for XPA/RPA-mediated Displacement of XPC from Damaged DNA—XPC (Rad4) forms a stable complex with hHR23B (Rad23), which appears to be essential for NER (23). Nonetheless, damaged DNA binding activity of the XPC-hHR23B complex belongs to the XPC subunit (23) and the exact role for hHR23B in repair is unclear. In an effort to explore the role for hHR23B in DNA repair, we examined whether the presence of hHR23B affects RPA/XPA-mediated displacement of XPC from the damaged DNA. GST-XPC-hHR23B (or GST-XPC) was incubated with fluorescence (TET)-labeled damaged DNA in the presence of RPA and XPA, and the amount of damaged DNA precipitated with GST-XPC and measured (Fig. 6A). Both XPC and XPC-hHR23B showed strong interaction with damaged DNA (Fig. 6A, lanes 2 and 5). In the presence of RPA + XPA, however, the damaged DNA-XPC-hHR23B complex, not the damaged DNA-XPC complex, was significantly reduced (Fig. 6A, lanes 3 and 4 versus 6 and 7), suggesting that hHR23B is somehow involved in the displacement of XPC from damaged DNA in the presence of RPA + XPA. To further investigate this, purified hHR23B was added to the reaction mixtures containing GST-XPC for its effect on the XPC-damaged DNA interaction in the presence of RPA + XPA (Fig. 6B). GST-XPC without hHR23B formed a stable complex with damaged DNA (Fig. 6B, lane 2), which was hardly affected by RPA + XPA (Fig. 6B, lanes 3 and 4). In contrast, addition of purified hHR23B to the reaction mixtures significantly decreased the binding of XPC to the damaged DNA in the presence of RPA + XPA (Fig. 6B, lanes 6 and 7). This result strongly suggests that hHR23B is necessary for the RPA/XPA-mediated displacement of XPC from damaged DNA.

The RPA-XPA Interaction Is Essential for the Displacement of XPC-hHR23B from Damaged DNA—Interaction between RPA and XPA is not only required to stabilize the XPA-damaged DNA interaction, but also necessary for NER activity (14, 16). Because both RPA and XPA are essential for the displacement of XPC from damaged DNA, we examined whether the RPA-XPA interaction is necessary for it. For this, wild-type RPA and two RPA mutants (RPAp34C33 lacking the C terminus of p34 subunit (XPA interaction domain); ZFM4, a mutant with cysteine to alanine substitution at the zinc finger domain of the p70 subunit) were compared in the displacement of XPC-hHR23B from damaged DNA. Although both mutants poorly supported NER activity *in vitro* (14, 36), ZFM4 supported the displacement of XPC-hHR23B from damaged DNA, whereas RPAp34C33 did not (Fig. 7). This result not only

After rocking for 20 min at room temperature, equimolar amounts of RPA + XPA (2 pmol (lanes 3, 6, and 9) or 5 pmol (lanes 4, 7, and 10)) were added to the mixtures, and beads were pulled down and analyzed for TET-labeled damaged DNA by 10% SDS-PAGE. Relative amounts of TET-labeled DNA are indicated at the bottom of the figure.

FIG. 7. A mutant RPA lacking the XPA interaction domain did not support the displacement of XPC from damaged DNA. GST-XPC-hHR23B (400 ng) was mixed with glutathione-Sepharose beads and TET-damaged DNA (2 pmol), and rocked for 20 min at room temperature. The mixtures were then challenged with increasing amounts of XPA (2 pmol (lanes 3, 5, and 7) or 5 pmol (lanes 4, 6, and 8)) in the presence of wild-type RPA (lanes 3 and 4), a mutant lacking C terminus of p34 (lanes 5 and 6), or a RPA zinc finger mutant (*ZFM4*) (lanes 7 and 8). Following the GST pull-down, beads were washed and analyzed for TET-labeled damaged DNA by 10% SDS-PAGE. Relative amounts of TET-labeled DNA are indicated at the bottom of the figure.






FIG. 8. Displacement of XPC-hHR23B from damaged DNA is not affected by the order of assembly of damage recognition proteins. The components indicated inside the boxes were incubated together for 20 min at room temperature prior to the addition of the remaining components. Where indicated, 400 ng of GST-XPC-hHR23B, 2 pmol of TET-labeled damaged DNA, and equimolar amounts of XPA + RPA (2 pmol (lanes 3, 6, and 9), 5 pmol (lanes 4, 7, and 10), or 10 pmol (lanes 5, 8, and 11)) were added. Following the GST pull-down, beads were washed and analyzed for TET-labeled damaged DNA by 10% SDS-PAGE. Relative amounts of TET-labeled DNA are indicated at the bottom of the figure.

suggests that RPA and XPA cooperate with each other in the displacement of XPC-hHR23B from damaged DNA, but also supports that the RPA-XPA interaction may be necessary for such cooperation.

Displacement of XPC-hHR23B from Damaged DNA Is Not Affected by the Order of Protein Assembly on Damaged DNA—Our finding that XPA and RPA cooperate with each other to displace XPC-hHR23B from the damaged DNA supports a notion that XPC-hHR23B is the initiator of global genome NER (19, 23, 24). To investigate this further, we examined whether the order of protein assembly at the damaged DNA site affects the XPC-hHR23B-damaged DNA interaction (Fig.

8). The 5'-fluorescence (TET)-labeled damaged DNA (ITR-60) was incubated with GST-XPC-hHR23B (Fig. 8, lanes 3–5), XPA + RPA (Fig. 8, lanes 6–8), or all three proteins (Fig. 8, lanes 9–11), and 20 min later, the remaining factor(s) were added to the reaction mixtures. The mixtures were then analyzed by the GST-XPC pull-down assay for the interaction of GST-XPC-hHR23B with damaged DNA. Regardless of the order of assembly, all assembly groups showed remarkably similar patterns in that the addition of increasing amounts of XPA + RPA proportionally displaced XPC-hHR23B from damaged DNA (Fig. 8). This result not only supports the multistep damage recognition in NER, but is also in keeping

with a notion that XPC-hHR23B is the global initiator in the damage recognition process.

DISCUSSION

Recognition of damaged DNA is a complex process involving a number of proteins (XPA, RPA, XPC-hHR23B, and TFIIH), all of which can independently bind to the damaged DNA (Refs. 1–4 and references therein). Although some biochemical properties of damage recognition proteins are known, the molecular mechanism of how these proteins function at the damaged DNA site is not clear. In this study we carried out a comprehensive biochemical analysis on the interaction of damage recognition factors themselves and with damaged DNA.

XPC as a Global Initiator in NER—Recent *in vitro* studies strongly point to a role for XPC-hHR23B as the initiator of global genomic repair (25). This is primarily based on the findings that: 1) preincubation of UV-damaged plasmid DNA with XPC was preferentially repaired in an *in vitro* kinetic experiment, and 2) XPC shows a considerable preference for binding to UV-damaged DNA in the presence of nondamaged competitor DNA (24, 25). On the other hand, a separate *in vitro* study demonstrated that preincubation of damaged DNA with RPA and XPA, compared with that of XPC-hHR23B, led to a faster repair (10).

From the study described here and the previous studies by others (9, 16, 19–23), it is evident that XPC-hHR23B and RPA share basic properties in damage recognition: 1) preferential binding to the damaged DNA, and 2) the physical interaction with XPA (11–13) (Fig. 3), which makes both XPC-hHR23B and RPA eligible for a global initiator in NER (10, 19). On the other hand, these two proteins exhibit quite different biochemical characteristics: first, XPC has a considerable preference for binding to UV- or cisplatin-damaged DNA in the presence of nondamaged competitor DNA (24, 25) (Fig. 2B), whereas RPA retains only a moderate preference to damaged DNA over non-damaged DNA (16). Second, RPA has a significant preference to ssDNA over dsDNA, whereas XPC (or XPC-hHR23B) shows a higher affinity to dsDNA over ssDNA (35) (Fig. 2). The latter finding suggests that XPC-hHR23B functions at an early stage of duplex DNA damage recognition, whereas RPA is involved in a later stage of damage recognition including a structural distortion of damaged DNA that leads to an unwinding of duplex DNA (or a generation of ssDNA). Third, XPC-hHR23B and RPA differ in their interaction with XPA such that the RPA-XPA interaction stabilizes weak binding of XPA to the damaged DNA (14, 16) (Fig. 4) necessary for NER action (14), whereas the interaction between XPC-hHR23B and XPA did not contribute to a stability of the XPA-damaged DNA complex (Fig. 4). Instead, the presence of XPC-hHR23B interferes with the formation of the RPA-XPA-damaged DNA complex (Fig. 4). Moreover, the XPA-XPC interaction may eventually lead to the displacement of XPC-hHR23B from damaged DNA (Fig. 5). The difference between XPC-hHR23B and RPA in their biochemical characteristics described above not only reflects their unique role(s) in the early stage of NER, but also supports the role for XPC-hHR23B as a global initiator in NER.

Multistep Damage Recognition Process in Early Stage of NER—XPC physically interacted with XPA (Fig. 3), but the interaction was significantly inhibited by the presence of damaged DNA (Fig. 4). This observation suggests that the interaction between XPA and XPC may be necessary to recruit XPA to the damaged DNA site. The XPC-hHR23B, once binding to the damaged DNA, likely recruits XPA (Fig. 3) and other repair factors such as TFIIH and XPG (38) to the damaged DNA site (Fig. 9). XPC-hHR23B is also involved in the recruitment of TFIIH to damaged DNA (26). The XPA-RPA, once introduced to the damaged DNA site, cooperates with each other to destabi-




FIG. 9. A proposed molecular mechanism of damage recognition process in the early stage of nucleotide excision repair. Transient steps are indicated with brackets.

lize the XPC-hHR23B-damaged DNA interaction. A recent *in vitro* study also supported the absence of XPC-hHR23B in the final incision complex (39). Displacement of XPC-hHR23B from the damaged DNA likely requires a physical interaction between XPA and RPA on damaged DNA because a mutant RPA lacking XPA interaction domain poorly functioned in the displacement of XPC from damaged DNA (Fig. 7). The XPA-RPA interaction on the damaged DNA would likely force XPC to dissociate itself from XPA and the damaged DNA (Figs. 5 and 9). These lines of evidence suggest that XPA, RPA, and XPC-hHR23B do not form a stable three-protein complex at the damaged DNA site. Instead, it supports a notion that the damage recognition process occurs in a stepwise manner. However, we cannot rule out a possibility that XPA, XPC-hHR23B, and RPA form a transient complex on the damaged DNA prior to the displacement of XPC-hHR23B (Fig. 9) because RPA and XPC likely recognize two separate domains of XPA (Fig. 3). We should also point out that the *in vitro* study described here did not include TFIIH, a key damage recognition protein involved in distinguishing the damaged strand from the nondamaged one (27) as well as local unwinding of the damaged DNA region. The interaction between XPC and TFIIH appears to be essential for nucleotide excision repair (40). Nonetheless, addition of TFIIH appeared to have no effect on the RPA/XPA-mediated

displacement of XPC-hHR23B from damaged DNA.²

Role for hHR23B in Damage Recognition—The XPC forms a stable complex with hHR23B (41, 42). Although the XPC subunit is solely responsible for the binding of the XPC-hHR23B complex to the damaged DNA (23) and the interaction with XPA (Fig. 3), hHR23B (Rad23) is essential for XPC function in NER (23). Very large amounts of XPC without hHR23B showed some repair activity, but equimolar hHR23B led about 10-fold higher activity (20). Rad23 contains a ubiquitin-associated domain that may play a role in controlling NER through proteasome-mediated degradation of repair factors (33, 43). In addition, hHR23B interacts with the base excision repair protein, *N*-methylpurine-DNA glycosylase, suggesting that it may have a role in mediating various repair pathways (37). In this study we found that hHR23B is necessary for XPA/RPA-mediated displacement of the XPC-hHR23B complex from damaged DNA (Fig. 6). hHR23B does not directly interact with XPA or RPA; we do not know what specific role hHR23B plays in the displacement of XPC from damaged DNA. A recent study showed that the DNA binding domain of XPC overlaps with the hHR23B interaction domain (40), suggesting that hHR23B may facilitate the displacement of XPC through the modulation of its DNA binding activity in the presence of XPA + RPA (Fig. 3).

Physiologic significance of the displacement of XPC from damaged DNA in NER is yet to be determined, however, it at least provides a crucial information that damage recognition occurs in an ordered, multistep process. Our finding that the order of protein assembly had no effect on the displacement of XPC from damaged DNA (Fig. 8) supported a notion that XPC-hHR23B is the initiator of global genomic repair (19, 24). Because XPC, compared with RPA and XPA, exhibited exceptionally strong affinity to damaged DNA (24), it is quite possible that the role for XPC-hHR23B is to effectively identify DNA damage *in vivo*. RPA-XPA-mediated displacement of XPC may be necessary for the formation of the stable XPA-RPA complex at the damaged site, which would allow a proper positioning of the endonucleases (XPG and ERCC1-XPF) for accurate and efficient incisions. It should be pointed out, however, that the RPA-XPA-mediated displacement of XPC from damaged DNA could be a part of the alternative global genomic NER. For example, although XPC-hHR23B does not preferentially bind to CPD DNA (24), repair of *cis-syn* cyclobutane dimer-containing DNA was dependent on XPC-hHR23B, suggesting that there may be alternative pathways for the global genomic NER that requires XPC-hHR23B but can be replaced by XPA and RPA (30). Further functional analysis would be necessary to validate the role for damage recognition complex in NER action.

Acknowledgments—We thank B. Hwang for providing purified XPC-hHR23B protein, A. Riemen for technical assistance, and A. Sancar and J. T. Reardon for XPC-hHR23B expressing recombinant baculoviruses and anti-XPC and hHR23B antibodies. We also thank C-H. Kim for assistance in earlier experiments.

² J-S. You and S-H. Lee, unpublished data.

REFERENCES

- Batty, D. P., and Wood, R. D. (2000) *Gene (Amst.)* **241**, 193–204
- Wood, R. D. (1996) *Annu. Rev. Biochem.* **65**, 135–167
- Sancar, A. (1996) *Annu. Rev. Biochem.* **65**, 43–81
- Lee, S.-H. (2001) *J. Biochem. Mol. Biol.* **34**, 489–495
- Jones, C. J., and Wood, R. D. (1993) *Biochemistry* **32**, 12096–12104
- Asahina, H., Kuraoka, I., Shirakawa, M., Morita, E. H., Miura, N., Miyamoto, I., Ohtsuka, E., Okada, Y., and Tanaka, K. (1994) *Mutat. Res.* **315**, 229–237
- Guzder, S. N., Sung, P., Prakash, L., and Prakash, S. (1993) *Proc. Natl. Acad. Sci. U. S. A.* **90**, 5433–5437
- He, Z., Henricksen, L. A., Wold, M. S., and Ingles, C. J. (1995) *Nature* **374**, 566–569
- Burns, J. L., Guzder, S. N., Sung, P., Prakash, S., and Prakash, L. (1996) *J. Biol. Chem.* **271**, 11607–11610
- Wakasugi, M., and Sancar, A. (1999) *J. Biol. Chem.* **274**, 18759–18768
- Li L., Lu, X., Peterson, C. A., and Legerski, R. J. (1995) *Mol. Cell. Biol.* **15**, 5396–5402
- Saijo, M., Kuraoka, I., Masutani, C., Hanaoka, F., and Tanaka, K. (1996) *Nucleic Acids Res.* **24**, 4719–4724
- Lee, S.-H., Kim, D. K., and Drissi, R. (1995) *J. Biol. Chem.* **270**, 21800–21805
- Stigger, E., Drissi, R., and Lee, S.-H. (1998) *J. Biol. Chem.* **273**, 9337–9343
- Park, C. H., Mu, D., Reardon, J. T., and Sancar, A. (1995) *J. Biol. Chem.* **270**, 4896–4902
- Wang, M., Mahrenholz, A., and Lee, S.-H. (2000) *Biochemistry* **39**, 6433–6439
- Araujo, S. J., and Wood, R. D. (1999) *Mutat. Res.* **435**, 23–33
- Wood, R. D. (1997) *J. Biol. Chem.* **272**, 23465–23468
- Sugasawa, K., Ng, J. M., Masutani, C., Iwai, S., van der Spek, P. J., Eker, A. P., Hanaoka, F., Bootsma, D., and Hoeijmakers, J. H. (1998) *Mol. Cell* **2**, 223–232
- Batty, D., Rapić-Otrin, V., Levine, A. S., and Wood, R. D. (2000) *J. Mol. Biol.* **300**, 275–290
- Reardon, J. T., Mu, D., and Sancar, A. (1996) *J. Biol. Chem.* **271**, 19451–19456
- Jansen, L. E., Verhage, R. A., and Brouwer, J. (1998) *J. Biol. Chem.* **273**, 33111–33114
- Guzder, S. N., Sung, P., Prakash, L., and Prakash, S. (1998) *J. Biol. Chem.* **273**, 31541–31546
- Sugasawa, K., Okamoto, T., Shimizu, Y., Masutani, C., Iwai, S., and Hanaoka, F. (2001) *Gene Dev.* **15**, 507–521
- Volker, M., Mone, M. J., Karmaker, P., van Hoffen, A., Schul, W., Vermeulen, W., Hoeijmakers, J. H., van Driel, R., van Zeeland, A. A., and Mullenders, L. H. (2001) *Mol. Cell* **8**, 213–224
- Yokoi, M., Masutani, C., Maekawa, T., Sugawara, K., Ohkuma, Y., and Hanaoka, F. (2000) *J. Biol. Chem.* **275**, 9870–9875
- Hanawalt, P. C. (2000) *Nature* **405**, 415–416
- Moggs, J. G., Yarema, K. J., Essigmann, J. M., and Wood, R. D. (1996) *J. Biol. Chem.* **271**, 7177–7186
- Stigger, E., Dean, F. B., Hurwitz, J., and Lee, S.-H. (1994) *Proc. Natl. Acad. Sci. U. S. A.* **91**, 579–583
- Kusumoto, R., Masutani, C., Sugawara, K., Iwai, S., Araki, M., Uchida, A., Mizukoshi, T., and Hanaoka, F. (2001) *Mutat. Res.* **485**, 219–227
- LeRoy, G., Drapkin, R., Weis, L., and Reinberg, D. (1998) *J. Biol. Chem.* **273**, 7134–7140
- Park, J. S., Park, S. J., Peng, X., Wang, M., Yu, M. A., and Lee, S.-H. (1999) *J. Biol. Chem.* **274**, 32520–32527
- Gillette, T. G., Huang, W., Russell, S. J., Reed, S. H., Johnston, S. A., and Friedberg, E. C. (2001) *Genes Dev.* **15**, 1528–1539
- You, J.-S., Wang, M., and Lee, S.-H. (2001) *Antioxidants Redox Signaling* **3**, 657–669
- Hey, T., Lipps, G., Sugawara, K., Iwai, S., Hanaoka, F., and Krauss, G. (2002) *Biochemistry* **41**, 6583–6587
- Dong, J., Park, J.-S., and Lee, S.-H. (1999) *Biochem. J.* **337**, 311–317
- Miao, F., Bouziane, M., Dammann, R., Masutani, C., Hanaoka, F., Pfeifer, G., and O'Connor, T. R. (2000) *J. Biol. Chem.* **275**, 28433–28438
- Araujo, S. J., Nigg, E. A., and Wood, R. D. (2001) *Mol. Cell. Biol.* **21**, 2281–2291
- Reardon, J. T., and Sancar, A. (2002) *Mol. Cell. Biol.* **22**, 5938–5945
- Uchida, A., Sugawara, K., Masutani, C., Dohmae, N., Araki, M., Yokoi, O., Ohkuma, Y., and Hanaoka, F. (2002) *DNA Repair*, **1**, 449–461
- Masutani, C., Sugawara, K., Yanagisawa, J., Sonoyama, T., Ui, M., Enomoto, T., Takio, K., Tanaka, K., van der Spek, P. J., and Bootsma, D. (1994) *EMBO J.* **13**, 1831–1843
- Shivji, M. K. K., Eker, A. P. M., and Wood, R. D. (1994) *J. Biol. Chem.* **269**, 22749–22757
- Chen, L., Shinde, U., Ortolan, T. G., and Madura, K. (2001) *EMBO Rep.* **2**, 933–938

Forum Original Research Communication

Role of Zinc-Finger Motif in Redox Regulation of Human Replication Protein A

MU WANG, JIN-SAM YOU, and SUK-HEE LEE

ABSTRACT

Replication protein A (RPA) is a heterotrimeric zinc-finger protein complex involved in DNA replication, repair, and genetic recombination. Unlike other zinc-finger proteins, RPA's zinc-finger motif is not essential for its single-stranded DNA (ssDNA) binding activity, but is involved in redox regulation of its single-stranded DNA (ssDNA) binding activity. To get an insight into the regulation of RPA-ssDNA interaction, wild-type RPA (wt-RPA) and zinc-finger mutant were examined for ssDNA binding activity using surface plasmon resonance technique. Interaction of wt-RPA with ssDNA under nonreducing conditions was very weak ($K_D \times 2.33 \times 10^{-8} M$) compared with that under reducing conditions ($K_D = 7.35 \times 10^{-11} M$), whereas ssDNA binding affinity of the zinc-finger mutant was not affected by redox. The divalent ion chelator, *o*-phenanthroline, significantly reduced wt-RPA-ssDNA interaction, but had no effect on the zinc-finger mutant. The inhibitory effect of *o*-phenanthroline on RPA-ssDNA interaction was reversed by Zn(II), but not by other divalent cations, suggesting that Zn(II) is the unique metal coordinating the zinc-finger cysteines in redox regulation of RPA-ssDNA interaction. In DNA repair, redox affected RPA's interaction with damaged DNA, but not its role in stabilizing the xeroderma pigmentosum group A (XPA)-damaged DNA complex, suggesting that the zinc-finger motif may mediate the transition of RPA-XPA interaction to a stable RPA-XPA-damaged DNA complex in a redox-dependent manner. Antioxid. Redox Signal. 3, 657-669.

INTRODUCTION

REPPLICATION PROTEIN A [RPA; also known as human single-stranded DNA (ssDNA) binding protein, HSSB] is a three-subunit protein complex consisting of 70-, 34-, and 11-kDa subunits involved in DNA replication, repair, and genetic recombination (Fig. 1A) (28). In DNA replication, RPA is involved in unwinding of the replication origin, whereas it interacts with the DNA polymerase α -primase complex at the origin (5, 7, 15, 20). In nucleotide excision repair (NER), RPA interacts with the xeroderma pigmentosum group A complementing protein (XPA) on damaged DNA and

stimulates XPA-DNA interaction and also recruits other repair proteins such as XPG, excision repair cross complementing protein 1 (ERCC1)-XPF, and transcription factor IIH (TFIIH) to the damaged site for the subsequent incision/excision step (11, 19). RPA may also be involved in a later stage of NER, gap-filling, that requires PCNA, RF-C, and DNA polymerase δ (or ϵ) (1).

The large subunit of RPA (p70) contains the ssDNA-binding domain that resides in the middle region of p70 with two homologous subdomains in tandem positions (3, 8, 14, 16). RPA p70 has a 4-cysteine type zinc-finger motif toward the C-terminus (amino acids

Department of Biochemistry and Molecular Biology, Indiana University Cancer Center and Walther Oncology Center, Indiana University School of Medicine, Indianapolis, IN 46202.




FIG. 1. Proteins used for this study. (A) Schematic representation of wild-type RPA (wt-RPA) and a zinc-finger mutant RPA [ZFM4; containing cysteine-to-alanine substitution at all four cysteine sites (amino acids 481, 486, 500, and 503)]. (B) SDS-PAGE of purified proteins. Proteins were separated by 12% SDS-PAGE followed by Coomassie Blue staining (see Materials and Methods).

481–503; Fig. 1A) (21, 30). Unlike other zinc-finger proteins, RPA's zinc-finger domain is not essential for its ssDNA binding activity and has little or no effect on its DNA binding activity (14, 16). Recent studies have shown that RPA's ssDNA binding activity is regulated by redox through the cysteines in a putative zinc-finger domain (21, 30). In fact, all four zinc-finger cysteines are required for redox regulation of RPA's DNA binding activity (30). Under reduced condition, the zinc-finger structure is favorably formed and Zn(II) may protect the zinc-finger cysteines from engaging in disulfide bond formation. Under nonreducing conditions, however, oxidation of the Zn(II)-thiolate bond induces the release of Zn(II) from the pocket (21), which promotes the formation of disulfide bonds (21, 30). The redox status of RPA significantly affected initial interaction with ssDNA, but had no effect after RPA formed a stable complex with DNA (30), suggesting that redox regulation of the zinc finger may be involved in me-

diating initial RPA-ssDNA interaction to form a stable RPA-ssDNA complex.

XPA is also a zinc-finger protein involved in the damage recognition step of NER (10, 13, 24). XPA preferentially binds to (6-4) photoproduct of ultraviolet ray (UV)-damaged DNA and may also play a role in subsequent steps of NER through interaction with other repair proteins (13). XPA exhibits a low-affinity binding to UV-damaged DNA; hence, the interaction between XPA and RPA stimulates the XPA-damaged DNA complex and also recruits other repair proteins such as XPG, ERCC1-XPF, and TFIIH to the damaged site for the subsequent incision/excision step (11, 19, 27). A mutation of RPA at any zinc-finger cysteine abolished its function in repair (6), suggesting a unique role for the zinc-finger domain in the early stage of repair.

To understand the role of the zinc-finger motif in the redox regulation of RPA-DNA interaction, we carried out a real-time analysis of

the interaction between RPA (and XPA) and DNA using the surface plasmon resonance (SPR) technique. We found that interaction of wild-type RPA (wt-RPA) with ssDNA under nonreducing conditions was 1/3,000th of the affinity observed under reducing conditions, whereas the zinc-finger mutant exhibited a strong binding to ssDNA regardless of its redox status. RPA-damaged DNA interaction was also regulated by redox, although RPA's role in stabilization of the XPA-damaged DNA complex was not affected by redox, suggesting that the zinc-finger motif may mediate the transition of RPA-XPA interaction to a stable RPA-XPA-damaged DNA complex in a redox-dependent manner.

MATERIALS AND METHODS

Proteins and antibodies

wt-RPA and a zinc-finger mutant [ZFM4; the mutation with the change of all four cysteines (the sites 481, 486, 500, and 503) into alanines] were prepared according to the procedure described previously with slight modifications (6). In brief, cell lysates were prepared from insect cells (Sf-9), coinfecting with recombinant baculoviruses encoding three subunits (11 kDa, 34 kDa, and either wild-type or zinc-finger mutant 70 kDa). After the salt concentration (0.5 M NaCl) was adjusted, cell lysates were loaded onto a ssDNA cellulose column (1.0 × 8 cm) equilibrated with buffer A [25 mM Tris-HCl (pH 7.5), 10% glycerol, 0.02% Nonidet P-40, 1 mM dithiothreitol (DTT), 0.5 mM EDTA, 0.1 mM phenylmethylsulfonyl fluoride, 0.1 µg/ml leupeptin, and 0.2 µg/ml antipain] containing 0.5 M NaCl. The column was successively washed with 20 column volumes of buffer A containing 0.5 M NaCl and 0.8 M NaCl. The proteins were eluted with buffer A containing 2.0 M NaCl and 40% ethylene glycol. The eluted fractions were diluted fivefold with buffer A, and loaded onto an Affi-Gel Blue (Bio-Rad) column equilibrated with buffer A containing 0.5 M NaCl. After the column was washed with buffer A containing 0.5 M NaCl and 0.8 M NaCl, proteins were eluted with buffer con-

taining 2.5 M NaCl and 40% ethylene glycol. The RPA-containing fractions were pooled and dialyzed against buffer A containing 50 mM NaCl and further purified on a Q-Sepharose column with a linear salt gradient (50 mM to 0.4 M NaCl). During purification, RPA was monitored by immunoblotting using anti-p70 and -p34 antibodies (25). Histidine-tagged XPA was prepared according to the procedure described previously (15).

Biomolecular interaction analysis

Interaction of wt-RPA, a zinc-finger mutant, and/or XPA with ssDNA (or UV-damaged DNA) was monitored using an SPR biosensor instrument, Biacore 3000 (Biacore) as described previously (27). For preparation of the biosensor surface with DNA, 5'-biotinylated 70mer DNA was diluted to 1.5 nM in a buffer containing 10 mM sodium acetate (pH 4.8) and 1.0 M NaCl, and manually injected onto an immobilized streptavidin surface of the Biacore sensor chip to the desired density in different flow cells. One flow cell was left underivatized to allow for refractive index change correction. Proteins were diluted in the running buffer containing 10 mM HEPES (pH 7.4), 150 mM NaCl, 2 mM MgCl₂, 0.005% polysorbate-20, and 1 mM DTT. Each experiment was repeated at least twice to assure reproducibility.

Electrophoretic mobility shift assay of RPA-ssDNA interaction

Oligo(dT)₅₀ was labeled with [γ -³²P]ATP (ICN) and T4 polynucleotide kinase (Roche Molecular Biochemical) based on the manufacturer's instructions. The indicated amount of wt-RPA or zinc-finger mutant was incubated with 100 fmol of 5'-³²P-labeled oligo(dT)₅₀ at room temperature for 25 min in a reaction mixture containing 50 mM HEPES-KOH (pH 7.8), poly(dI-dC) (0.2 µg), bovine serum albumin (0.2 µg/µl), 200 mM NaCl, and the indicated amounts of other chemicals. The RPA-ssDNA complex was analyzed on 5% polyacrylamide gels in 0.5X Tris-borate-EDTA buffer (acrylamide: bisacrylamide = 43.2:0.8). The gels were dried and exposed to x-ray films (Kodak). For quantification, the bands of interest were excised

from the gels and measured for radioactivity using a Beckman Scintillation Counter LS 6500.

Proteolytic analysis of wt-RPA and zinc-finger mutant with trypsin

The reaction mixture (60 μ l) contained 0.1 M Tris-HCl (pH 8.5), 120 ng/ μ l wild-type or mutant RPA, and, where indicated, 200 ng/ μ l oligo(dT)₅₀. After the reaction mixture was incubated at room temperature for 15 min, 50 ng of trypsin (sequencing grade; Boehringer Mannheim) was added into the reaction. The reaction mixture was then immediately incubated at 37°C. Eight or 10 μ l of the sample was removed from the reaction at the indicated time points. The reaction samples were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) gel and were analyzed by the western blot (Fig. 1B).

Western blotting

Western blots were performed as described previously (25). In brief, wild-type or mutant RPA was run in a 12% SDS-PAGE gel and transferred to a nitrocellulose membrane (BA83; Bio-Rad). After the transfer, either anti-p70 or anti-p34 polyclonal antibody was used to detect RPA subunits, which were then revealed by ¹²⁵I-protein A (Amersham Corp.) treatment and visualized by autoradiography.

Circular dichroism (CD) analysis

CD spectra of purified wt-RPA and a zinc-finger mutant were acquired on a Jasco J-720 spectropolarimeter (Japan Spectroscopic Co.). Far-UV CD spectra (200–260 nm) were measured in a 1-mm path length quartz cell and represent averages of four accumulations with a protein concentration of 100 μ g/ml in 25 mM Tris-HCl (pH 7.5), 10% (vol/vol) glycerol, 0.02% (vol/vol) NP40, 0.5 mM EDTA, 0.1 mM phenylmethylsulfonyl fluoride, 0.1 μ g/ml leupeptin, 0.2 μ g/ml antipain, and 200 mM NaCl. All spectra were baseline-corrected by subtraction of an averaged scan derived from the buffer alone to obtain the mean molar residue ellipticity (deg cm² dmol⁻¹). The result from CD spectra was deconvoluted to obtain information on the secondary structures that are mostly affected by the zinc-finger motif during interaction of protein with damaged

DNA. The software program (CDnn) used to deconvolute CD spectra was obtained through the Internet (<http://bioinformatik.biochemtech.uni-halle.de/cdnn/>).

RESULTS

Role of zinc-finger motif in redox regulation: kinetic analysis of RPA's ssDNA binding activity

Our recent study indicates that all four zinc-finger cysteines are essential for redox regulation of RPA's-ssDNA binding activity (30). Redox regulation affects initial RPA-ssDNA interaction, but has no effect after RPA forms a stable complex with DNA (30), suggesting that the zinc-finger motif may mediate the initial interaction of RPA with ssDNA. To get an insight into the RPA-ssDNA interaction, we used the SPR technique to examine the ssDNA binding activity of wt-RPA and zinc-finger mutant (ZFM4) under various redox conditions, which allows macromolecular interactions to be measured in real time (27). wt-RPA or zinc-finger mutant ranging from 0.125 nM to 2 nM was injected onto a biosensor chip containing a low level (33 RU) of 70mer ssDNA for 600 s, followed by 900 s of a buffer injection period for dissociation. The sensorgram indicated that both wt-RPA and zinc-finger mutant bound to ssDNA with high affinity in the presence of 1 mM DTT (Fig. 2B and D). Under nonreducing conditions, however, wt-RPA exhibited very low ssDNA binding activity, whereas ZFM4 maintained its high-affinity binding to ssDNA (Fig. 2A and C, and Table 1). Interaction of wt-RPA with the ssDNA surface under nonreducing conditions was very weak, as indicated by the derived $K_D = 2.33 \times 10^{-8}$ M, 1/3,000th of the affinity observed for RPA under reducing conditions (Table 1). Even though wt-RPA exhibited a low-affinity binding to ssDNA under nonreducing conditions, it maintained a stable interaction with ssDNA once forming a complex.

A unique role for Zn(II) in RPA-ssDNA interaction

To understand the role of Zn(II) in the redox regulation of RPA-ssDNA interaction, we ex-




FIG. 2. Biomolecular interaction analysis of RPA-ssDNA binding under redox conditions. A 5 nM concentration of wt-RPA or a zinc-finger mutant (ZFM4) was injected onto a sensor chip surface containing 33 RU of ssDNA using the KINJECT function of Biacore 3000. The association phase was allowed for 600 s followed by 900 s of a buffer injection period for dissociation. (A) wt-RPA in the absence of DTT. (B) wt-RPA in the presence of 1 mM DTT. (C) ZFM4 in the absence of DTT. (D) ZFM4 in the presence of 1 mM DTT.

amined various divalent ions for RPA's ssDNA binding activity (Fig. 3). The role of Zn(II) in RPA-ssDNA interaction was supported by the fact that the inhibitory effect of the divalent ion chelator, *o*-phenanthroline, was reversed by addition of Zn(II) (30; Fig. 3A, lanes 4–6). In contrast, the interaction of the zinc-finger mutant with ssDNA was not affected by either *o*-phenanthroline or Zn(II) (Fig. 3A, lanes 7–16). Addition of an excess amount of Zn(II) non-specifically inhibited RPA-ssDNA interaction regardless of *o*-phenanthroline's presence, suggesting that Zn(II)-dependent reversal of the inhibitory effect of *o*-phenanthroline functions within stoichiometric concentration (data not shown). To assess the role of Zn(II) in coordination of the zinc finger cysteine, we examined

other divalent ions to see whether they can replace Zn(II) in RPA-ssDNA interaction (Fig. 3B). Mg(II), Ca(II), and Cu(II) were not able to compensate for Zn(II) in overcoming the inhibitory effect of *o*-phenanthroline in RPA-ssDNA interaction, whereas Fe(II) partially replaced Zn(II) (Fig. 3B). This result suggests that Zn(II) is the unique metal element coordinating four cysteine residues at RPA's zinc-finger motif.

Involvement of zinc-finger domain in RPA's structural change following interaction with DNA

To understand further the role of the zinc-finger motif in RPA-ssDNA interaction, we ex-

TABLE 1. EQUILIBRIUM AND KINETIC BINDING CONSTANTS OF WT-RPA AND A ZINC-FINGER MUTANT (ZFM4) TO ssDNA

RPA	$K_D(M)$	$k_a(s^{-1} M^{-1})$	$k_d(s^{-1})$
+ DTT			
Wild-type	$(7.35 \pm 0.01) \times 10^{-11}$	$(2.76 \pm 0.02) \times 10^{-6}$	$(2.03 \pm 0.01) \times 10^{-4}$
ZFM4	$(4.47 \pm 0.04) \times 10^{-11}$	$(6.39 \pm 0.01) \times 10^{-6}$	$(2.86 \pm 0.02) \times 10^{-3}$
- DTT			
Wild-type	$(2.33 \pm 0.20) \times 10^{-8}$	$(2.38 \pm 0.06) \times 10^{-3}$	$(5.53 \pm 0.32) \times 10^{-5}$
ZFM4	$(4.97 \pm 0.03) \times 10^{-11}$	$(4.49 \pm 0.01) \times 10^{-6}$	$(2.23 \pm 0.01) \times 10^{-4}$

The data were globally fit to a 1:1 binding model.

amined far-UV CD spectra of wt-RPA and a zinc-finger mutant to see a change in secondary structure upon binding to DNA. A significant change in CD spectra was observed with both wt-RPA and a zinc-finger mutant upon binding to DNA (Fig. 4), suggesting that RPA undergoes a significant conformational change upon binding to DNA. A noticeable difference between wt-RPA and a zinc-finger mutant in the CD spectra pattern was observed under reducing conditions (Fig. 4A and C, and Table 2). Under nonreducing conditions, however, very little change in CD spectra between wt-RPA and zinc-finger mutant was observed (Fig. 4B and C, and Table 2). This result not only supports a role for the zinc-finger domain in the structural change of RPA, but also establishes a relationship between redox regulation and the structural change of RPA upon binding to DNA.

We also examined p70 and p34 subunits of wt-RPA and zinc-finger mutant for the tryptic digestion patterns in the presence and absence of ssDNA. In the absence of ssDNA, wt-RPA was cleaved into several tryptic products (55-, 47-, 19–21-, and 25-kDa fragments), whereas 19 kDa and 55 kDa were the major cleavage fragments in the presence of ssDNA (Fig. 5A). This result is in keeping with that from far-UV CD spectra analysis (Fig. 4) and suggests that RPA undergoes a conformational change upon binding to DNA (8, 21). The zinc-finger mutant, however showed very little change of its tryptic digestion in the presence of ssDNA (Fig. 5B), suggesting that a mutation at zinc-finger cysteine affects p70's conformational change upon binding to ssDNA. We also examined the tryptic digestion pattern of p34 to see whether the mutation at p70's zinc-finger domain affects

the conformation of the p34 subunit. The p34 subunit of the zinc-finger mutant (ZFM4) was not only protected from tryptic digestion, but also unaffected by the presence of ssDNA (Fig. 5D). On the other hand, wt-RPA showed a change in the p34 tryptic digestion pattern upon binding to ssDNA, such that the p34 subunit was more accessible to tryptic digestion following RPA–ssDNA interaction. This result suggests that the mutation at the zinc-finger domain of p70 may also affect the accessibility of p34 to tryptic digestion.

Redox regulation of RPA's zinc-finger motif in recognition of damaged DNA

RPA not only has a preferential binding to UV- or cisplatin-damaged DNA, but also can stabilize XPA-damaged DNA complex through protein–protein interaction (2, 4, 22, 26, 27). Although zinc-finger mutant RPA failed to support NER activity (6), it also can stabilize the interaction of XPA with damaged DNA (26). We therefore examined whether redox affects RPA's interaction with damaged DNA and/or with XPA. Both XPA and wt-RPA are the 4-cysteine type zinc-finger proteins, and their interactions with UV-damaged DNA were affected by redox (Fig. 6A and B), whereas the interaction of the zinc-finger mutant RPA with damaged DNA was not affected by redox change (Fig. 6B). On the other hand, RPA's role in stabilization of XPA-damaged DNA complex was not affected by redox (Fig. 6A & 6B), suggesting that the failure of zinc-finger mutant in supporting NER may not be due to its role in interaction with damaged DNA or XPA. Redox regulation of the zinc finger significantly affected the initial RPA–ssDNA interaction, but



A

B


FIG. 3. (A) Inhibitory effect of *o*-phenanthroline on RPA ssDNA binding activity is restored by Zn(II). wt-RPA or zinc-finger mutant (20 ng) was treated with 1.0 mM *o*-phenanthroline followed by the addition of increasing amounts of Zn(II) to the reaction mixture. After incubation for 15 min in the presence of 100 fmol of 5'-³²P-labeled oligo(dT)₅₀, the RPA-ssDNA complex was analyzed by gel mobility shift assay (5% PAGE). Lane 1 contained no RPA. (B) Effect of various metal ions on RPA's ssDNA binding activity in the presence of *o*-phenanthroline. wt-RPA (20 ng) was treated with 1.0 mM *o*-phenanthroline and the indicated amounts of divalent metal (100 μ M in lanes 4, 7, 10, 13, and 16; 250 μ M in lanes 5, 8, 11, 14, and 17; 1,000 μ M in lanes 6, 9, 12, 15, and 18) added to the reaction mixture before incubation for 15 min with 100 fmol of 5'-³²P-labeled oligo(dT)₅₀. No RPA was included in lane 1. The RPA-ssDNA complex was analyzed by the procedure as described in A.




FIG. 4. Far-UV CD spectra of wild-type RPA (—) and zinc-finger mutant (ZFM4) (---) under various conditions. Reactions were carried out at 25°C in either the absence (A and B) or presence (C and D) of oligo(dT)₅₀. Reactions contained either 1 mM (A and C) or 0 mM DTT (B and D).

had no effect after RPA formed a stable complex with DNA (30). Similarly, RPA-XPA-damaged DNA complex, once formed, was not affected by redox (Fig. 6C), suggesting that the zinc-finger motif may mediate the transition of RPA-XPA interaction to a stable RPA-XPA-damaged DNA complex in a redox-dependent manner.

DISCUSSION

The zinc finger is one of the key structural motifs found in many DNA binding proteins in eukaryotes and implicates a role in the regulation of DNA binding activity through redox (9, 12, 29). Nonetheless, the role of the zinc finger in redox regulation has not been directly

demonstrated because the zinc finger domain is essential for sequence-specific DNA binding and a mutation at this region would abolish DNA binding activity. RPA's zinc-finger motif is not a DNA binding element, and a mutation at the zinc-finger motif has very little effect on its DNA binding activity (8, 14, 16), which makes RPA an excellent model to study the role of zinc-finger motifs in redox regulation. In this study, we analyzed a role for RPA's zinc-finger motif in the redox regulation of RPA-DNA interaction.

The SPR analysis indicated that interaction of wt-RPA with ssDNA under nonreducing conditions was very weak and was only 1/3,000th of the affinity observed for RPA under reducing conditions (Table 1). Even though wt-RPA showed a low affinity to ssDNA un-

TABLE 2. FAR UV-CD ANALYSIS OF WT-RPA AND ZINC-FINGER MUTANT (ZFM)

	Helix (%)	Antiparallel (%)	Parallel (%)	β -turn (%)	Random coil (%)
wt-RPA (+ DTT)	10.9	29.6	5.5	20.3	36.1
wt-RPA (- DTT)	15.7	28.0	6.3	15.9	32.9
wt-RPA/DNA (+ DTT)	16.3	27.8	6.4	15.5	32.4
wt-RPA/DNA (- DTT)	11.4	31.1	5.8	18.5	35.0
ZFM (+ DTT)	9.9	31.4	5.4	20.3	36.3
ZFM (- DTT)	13.6	30.4	6.2	16.3	33.6
ZFM/DNA (+ DTT)	15.4	30.3	6.5	14.8	32.2
ZFM/DNA (- DTT)	11.7	32.3	6.0	17.4	34.4

der nonreducing conditions, it maintained a stable interaction with ssDNA once forming the RPA-ssDNA complex. This result is in keeping with a recent observation that redox

regulation of the zinc finger significantly affected initial RPA-ssDNA interaction, but had no effect after RPA formed a stable complex with DNA (30). This result also suggests that




FIG. 5. Involvement of zinc-finger motif in the conformational change of RPA. (A and B) Tryptic digestion of RPA p70 subunit upon binding to ssDNA. Reaction mixtures (60 μ l) contained 0.1 M Tris-HCl (pH 8.5) and 120 ng/ μ l of either wt-RPA (A) or a zinc-finger mutant (ZFM4) (B). Where indicated, 40 ng/ μ l oligo(dT) was added. After incubation at room temperature for 15 min, 50 ng of trypsin (sequencing grade; Boehringer Mannheim) was added to the reaction. Reaction mixtures were then incubated at 37°C, and aliquots were removed from the reactions at the indicated time points. Samples were mixed with gel loading buffer, boiled for 5 min, and loaded onto a 14% SDS-PAGE. The gel was then subjected to a western blot analysis using an anti-p70 polyclonal antibody. (C and D). Tryptic digestion pattern of RPA p34 subunit upon binding to ssDNA. Reaction mixtures and the sample preparations were the same as in A and B except that an anti-p34 polyclonal antibody was used to visualize p34 and its cleaved fragments for western analysis.

the zinc-finger motif mediates the transition of RPA-ssDNA interaction to a stable RPA-ssDNA complex in a redox-dependent manner.


The 4-cysteine zinc finger contains Zn(II) that tetrahedrally coordinates four cysteine residues (18). Under reducing conditions, the zinc-finger structure is favorably formed and Zn(II), buried in the interior, stabilizes the module by binding four cysteines (21), whereas the oxidation of the Zn(II)-thiolate bond induces the release of Zn(II) from the zinc finger, which likely promotes disulfide bond formation between the zinc-finger cysteines (21, 30). Other divalent ions were not able to compensate for Zn(II) in overcoming the inhibitory effect of *o*-phenanthroline in RPA-ssDNA interaction (Fig. 3), suggesting that Zn(II) may play a unique role in coordinating the zinc finger cysteines, which in turn controls RPA's ssDNA binding activity. The latter is also supported by the fact that a mutation at zinc-finger cysteines (amino acids 481, 486, 500, or 503) would be sufficient to abolish redox sensitivity of RPA's ssDNA binding activity (21, 30).

A significant change in the tryptic digestion pattern observed with wt-RPA, but not with a zinc-finger mutant, upon binding to ssDNA (Fig. 5A) strongly suggests that the conformational change of p70 occurs in an ssDNA-dependent manner, which mainly protects the large domain of p70 (55-kDa fragment) from proteolytic digestion. One possibility is that the 55-kDa fragment may actually be involved in the conformational change of p70 and protected from tryptic digestion through its interaction with ssDNA. It would be interesting to see whether the protected region of p70 (55-kDa fragment) represents the domain directly in contact with ssDNA. The p34 subunit (wt-RPA) also showed a change in the tryptic digestion pattern, becoming more accessible to

tryptic digestion following its interaction with ssDNA, whereas the tryptic digestion pattern of the p34 subunit from the zinc-finger mutant (ZFM4) was virtually unaffected by the presence of ssDNA (Fig. 5D). This result suggests that the zinc-finger domain of p70 not only is essential for the conformational change of its own subunit, but may also affect the accessibility of p34 to tryptic digestion. The latter possibility was also supported by the previous finding where DNA-dependent protein kinase phosphorylated the p34 subunit from wt-RPA, but not that from the zinc-finger mutant, in the presence of ssDNA (6).

A role for RPA's zinc-finger domain has been demonstrated previously indicating that mutations at the zinc-finger domain differentially affected its function in DNA replication and NER (6, 17). The failure of zinc-finger mutants in supporting NER was not due to its role in the stabilization of the XPA-damaged DNA complex (26, 27). It is possible that the zinc-finger motif may be involved in mediating the generation of ssDNA during RPA-damaged DNA interaction (23). No matter what the role of RPA's zinc-finger domain in DNA repair may be, the initial interaction of XPA with damaged DNA under nonreducing conditions was significantly affected in the presence of wt-RPA, but not in the presence of a zinc-finger mutant (Fig. 6). It is not clear whether the differential effect of wt-RPA and a zinc-finger mutant on the XPA-damaged DNA interaction plays a direct role in NER. Considering the fact that the zinc-finger domain is involved in the conformational change of the RPA p70 subunit, it is conceivable that RPA's zinc-finger domain and its redox regulation may affect the early stage of repair, such as formation of a damage-recognition complex.

FIG. 6. Biacore analysis of redox regulation of RPA stabilization of XPA-(64) photoproduct interaction. wt-RPA or ZFM4 was injected onto a sensor chip surface containing 2,000 RU of double-stranded DNA, which contains a centrally located (6-4) photoproduct, using the KINJECT function of Biacore 3000. The association phase was allowed for 300 s followed by 540 s of a buffer injection period for dissociation. (A) wt-RPA (1 nM), XPA (2.5 nM), and the mixture of XPA (2.5 nM) and wt-RPA (1 nM) in the presence (1 mM) and absence of DTT. (B) ZFM4 RPA (1 nM), XPA (2.5 nM), and the mixture of XPA (2.5 nM) and ZFM4 RPA (1 nM) in the presence (1 mM) and absence of DTT. (C) wt-RPA (1 nM), XPA (2.5 nM), and the mixture of XPA (2.5 nM) and wt-RPA (1 nM) were injected onto the sensor chip surface in the presence of 1 mM DTT for a period of 300 s for association followed by a dissociation of 540 s by washing with a buffer containing either 1 mM or 0 mM DTT.



ACKNOWLEDGMENTS

We would like to thank Dr. John Hawes for help with HPLC purification of oligonucleotides containing DNA damage, and Dr. Alan Mahrenholz for valuable suggestions throughout the course of this study. This research was supported by grants from the U.S. Army (DAMD17-00-1-0295), NIH (CA92111), and the Catherine Peachey Funds. M.W. was supported by a postdoctoral fellowship from NIH (F32 GM20167-01).

ABBREVIATIONS

CD, circular dichroism; DTT, dithiothreitol; ERCC1, excision repair cross complementing protein 1; NER, nucleotide excision repair; RPA, replication protein A; SDS-PAGE, sodium dodecyl sulfate-polyacrylamide gel electrophoresis; SPR, surface plasmon resonance; ssDNA, single-stranded DNA; TFIIF, transcription factor IIF; UV, ultraviolet ray; wt-RPA, wild-type RPA; XPA, xeroderma pigmentosum group A protein; ZFM, zinc-finger mutant RPA.

REFERENCES

1. Aboussekhra A, Biggerstaff M, Shivji MK, Vilpo JA, Moncollin V, Podust VN, Protic M, Hubscher U, Egly JM, and Wood RD. Mammalian DNA nucleotide excision repair reconstituted with purified protein components. *Cell* 80: 859-868, 1995.
2. Batty DP and Wood RD. Damage recognition in nucleotide excision repair of DNA. *Gene* 241: 193-204, 2000.
3. Bochkarev A, Pfuetzner RA, Edwards AM, and Frappier L. Structure of the single-stranded DNA-binding domain of replication protein A bound to DNA. *Nature* 385: 176-181, 1997.
4. Burns JL, Guzder SN, Sung P, Prakash S, and Prakash L. An affinity of human replication protein A for ultraviolet-damaged DNA. Implications for damage recognition in nucleotide excision repair. *J Biol Chem* 271: 11607-11610, 1996.
5. Collins KL and Kelly TJ. Effects of T antigen and replication protein A on the initiation of DNA synthesis by DNA polymerase α -primase. *Mol Cell Biol* 11: 2108-2115, 1991.
6. Dong J, Park JS, and Lee SH. In vitro analysis of the zinc-finger motif in human replication protein A. *Biochem J* 337: 311-317, 1999.
7. Dornreiter I, Erdile LF, Gilbert IU, von Winkler D, Kelly TJ, and Fanning E. Interaction of DNA polymerase α -primase with cellular replication protein A and SV40 T antigen. *EMBO J* 11: 769-776, 1992.
8. Gomes XV and Wold MS. Functional domains of the 70-kilodalton subunit of human replication protein A. *Biochemistry* 35: 10558-10568, 1996.
9. Gowen LC, Avrutskaya AV, Latour AM, Koller BH, and Leadon SA. BRCA1 required for transcription-coupled repair of oxidative DNA damage. *Science* 281: 1009-1012, 1998.
10. Guzder SN, Sung P, Prakash L, and Prakash S. Yeast DNA-repair gene RAD14 encodes a zinc metalloprotein with affinity for ultraviolet-damaged DNA. *Proc Natl Acad Sci U S A* 90: 5433-5437, 1993.
11. He Z, Henricksen LA, Wold MS, and Ingles CJ. RPA involvement in the damage-recognition and incision steps of nucleotide excision repair. *Nature* 374: 566-569, 1995.
12. Hutchison KA, Matic G, Meshinchi S, Bresnick EH, and Pratt WB. Redox manipulation of DNA binding activity and BuGR epitope reactivity of the glucocorticoid receptor. *J Biol Chem* 266: 10505-10509, 1991.
13. Jones CJ and Wood RD. Preferential binding of the xeroderma pigmentosum group A complementing protein to damaged DNA. *Biochemistry* 32: 12096-12104, 1993.
14. Kim DK, Stigger E, and Lee SH. Role of the 70-kDa subunit of human replication protein A (I). Single-stranded DNA binding activity, but not polymerase stimulatory activity, is required for DNA replication. *J Biol Chem* 271: 15124-15129, 1996.
15. Lee SH and Kim DK. The role of the 34-kDa subunit of human replication protein A in simian virus 40 DNA replication in vitro. *J Biol Chem* 270: 12801-12807, 1995.
16. Lin YL, Chen C, Keshav KF, Winchester E, and Dutta A. Dissection of functional domains of the human DNA replication protein complex replication protein A. *J Biol Chem* 271: 17190-17198, 1996.
17. Lin YL, Shivji MKK, Chen C, Kolodner R, Wood RD, and Dutta A. The evolutionarily conserved zinc finger motif in the largest subunit of human replication protein A is required for DNA replication and mismatch repair but not for nucleotide excision repair. *J Biol Chem* 273: 1453-1461, 1998.
18. Luisi BF, Xu WX, Otwinowski Z, Freedman LP, Yamamoto KR, and Sigler PB. Crystallographic analysis of the interaction of the glucocorticoid receptor with DNA. *Nature* 352: 497-505, 1991.
19. Matsunaga T, Park CH, Bessho T, Mu D, and Sancar A. Replication protein A confers structure-specific endonuclease activities to the XPF-ERCC1 and XPG subunits of human DNA repair excision nuclease. *J Biol Chem* 271: 11047-11050, 1996.
20. Melendy T and Stillman B. An interaction between replication protein A and SV40 T antigen appears es-

- essential for primosome assembly during SV40 DNA replication. *J Biol Chem* 268: 3389–3395, 1993.
21. Park JS, Wang M, Park SJ, and Lee SH. Zinc finger of replication protein A, a non-DNA binding element, regulates its DNA binding activity through redox. *J Biol Chem* 274: 29075–29080, 1999.
22. Patrick SM and Turchi JJ. Human replication protein A preferentially binds cisplatin-damaged duplex DNA in vitro. *Biochemistry* 37: 8808–8815, 1998.
23. Patrick SM and Turchi JJ. Replication protein A (RPA) binding to duplex cisplatin damaged DNA is mediated through the generation of single-stranded DNA. *J Biol Chem* 274: 14972–14978, 1999.
24. Robins P, Jones CJ, Biggerstaff M, Lindahl T, and Wood RD. Complementation of DNA repair in xeroderma pigmentosum group A cell extracts by a protein with affinity for damaged DNA. *EMBO J* 10: 3913–3921, 1991.
25. Stigger E, Dean FB, Hurwitz J, and Lee SH. Reconstitution of functional human single-stranded DNA-binding protein from individual subunits expressed by recombinant baculoviruses. *Proc Natl Acad Sci U S A* 91: 579–583, 1994.
26. Stigger E, Drissi R, and Lee SH. Functional analysis of human replication protein A in nucleotide excision repair. *J Biol Chem* 273: 9337–9343, 1998.
27. Wang M, Mahrenholz A, and Lee SH. RPA stabilizes the XPA-damaged DNA complex through protein-protein interaction. *Biochemistry* 39: 6433–6439, 2000.
28. Wold MS. Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Annu Rev Biochem* 66: 61–92, 1997.
29. Wu X, Bishopric NH, Discher DJ, Murphy BJ, and Webster KA. Physical and functional sensitivity of zinc finger transcription factors to redox change. *Mol Cell Biol* 16: 1035–1046, 1996.
30. You JS, Wang M, and Lee SH. Functional characterization of zinc-finger motif in redox regulation of RPA–ssDNA interaction. *Biochemistry* 39: 12953–12958, 2000.

Address reprint requests to:

Dr. S.-H. Lee

Department of Biochemistry and

Molecular Biology

Indiana University School of Medicine

635 Barnhill Dr.

Indianapolis, IN 46202

E-mail: slee@iupui.edu

Received for publication November 2, 2000; accepted January 28, 2001.

Journal of | **Biochemistry &
Molecular Biology**

Volume 34, 2001

<http://www.biochem.or.kr/jbmb.htm>

Recognition of DNA Damage in Mammals

Suk-Hee Lee*

Department of Biochemistry and Molecular Biology, and the Walther Oncology Center,
Indiana University School of Medicine, and the Walther Cancer Institute, Indianapolis, Indiana 46202

Received 11 November 2001

DNA damage by UV and environmental agents are the major cause of genomic instability that needs to be repaired, otherwise it give rise to cancer. Accordingly, mammalian cells operate several DNA repair pathways that are not only responsible for identifying various types of DNA damage but also involved in removing DNA damage. In mammals, nucleotide excision repair (NER) machinery is responsible for most, if not all, of the bulky adducts caused by UV and chemical agents. Although most of the proteins involved in NER pathway have been identified, only recently have we begun to gain some insight into the mechanism by which proteins recognize damaged DNA. Binding of *Xeroderma pigmentosum* group C protein (XPC)-hHR23B complex to damaged DNA is the initial damage recognition step in NER, which leads to the recruitment of XPA and RPA to form a damage recognition complex. Formation of damage recognition complex not only stabilizes low affinity binding of XPA to the damaged DNA, but also induces structural distortion, both of which are likely necessary for the recruitment of TFIIH and two structure-specific endonucleases for dual incision.

Keywords: DNA damage, Nucleotide excision repair, Replication protein A, XPA, and XPC-hHR23B

DNA damage recognition in nucleotide excision repair

Nucleotide excision repair (NER) action involves interaction of multiple proteins locating the damaged DNA sites, removal of short oligonucleotides containing DNA adducts, and synthesizing a replacement patch. In mammals, NER requires over 20 polypeptides including damage recognition/structure distortion factors [*Xeroderma pigmentosum* group A protein (XPA), XPC-hHR23B, replication protein A (RPA), also

known as human single-stranded DNA binding protein, hSSB), and XPE], TFIIH containing two DNA helicases (XPB and XPD) that separate the strands to create an open preincision complex, two structure-specific endonucleases, ERCC1-XPF and XPG, and the enzymes necessary for filling-in the gap [DNA polymerase δ/ϵ , proliferating cell nuclear antigen (PCNA), replication factor C (RF-C), and RPA] (see Fig. 1).

An ongoing challenge is to understand how DNA damage is recognized and distinguished from non-damaged sites during NER. Recognition of DNA damage is the initial, but complicated event for NER. It involves multiple proteins: XPA, RPA, XPC-hHR23B, XPE and TFIIH, all of which can independently bind to damaged DNA and may have a role in damage recognition (Sancar, 1996; Dawn and Wood, 2000). Although some characteristics of the damage recognition proteins are known, the molecular mechanism of how these proteins function at the damaged DNA site is not clear. A recent *in vitro* study suggested that XPC-hHR23B is an initiator of global genomic repair, but not of transcription-coupled repair (Sugasawa *et al.*, 1998; Sugawara *et al.*, 2001). This is based on the findings that preincubation of UV-damaged plasmid DNA with XPC was preferentially repaired in an *in vitro* kinetic experiment and also that XPC has considerable preference for binding to UV-damaged DNA in the presence of non-damaged competitor DNA (Sugasawa *et al.*, 1998). It should be noted however that the binding of XPC-hHR23B to UV-damaged DNA in the absence of non-damaged DNA showed only moderate preference for damaged DNA (Guzder *et al.*, 1998; Jansen *et al.*, 1998; Wakasugi and Sancar, 1999).

There is current discussion as to whether XPA or XPC functions first in the recognition of damage or helix distortion. Both RPA and XPA preferentially bind to cisplatin- or UV-damaged DNA (Jones and Wood, 1993; He *et al.*, 1995; Lee *et al.*, 1995; Li *et al.*, 1995; Burns *et al.*, 1996; Patrick and Turchi, 1998) and form a stable complex on damaged DNA (Wang *et al.*, 2000). Interaction between XPA and RPA is not only crucial for damage-recognition but also for recruiting other repair factors such as XPG, ERCC1-XPF, and TFIIH to

*To whom correspondence should be addressed.

Tel: 1-317-278-3464; Fax: 1-317-274-4686

E-mail: slee@iupui.edu




Fig. 1. A model for the nucleotide excision repair of UV-damaged DNA.

the damaged site (He *et al.*, 1995; Park *et al.*, 1995; Nocentini *et al.*, 1997; Bessho *et al.*, 1997; de Laat *et al.*, 1998). The XPA-DNA interaction is relatively weak and characterized by rapid dissociation, whereas RPA formed a 100-fold more stable complex with UV-damaged DNA (Wang *et al.*, 2000). A mutant RPA lacking XPA interaction domain leads to destabilization of XPA-damaged DNA complex, implicating a role for RPA in the early stage of DNA repair.

XPE (also known as a UV damage-specific DNA binding protein (UV-DDB) is another UV-damage recognition protein found in human and monkey cells (Payne and Chu, 1994; Nichols *et al.*, 2000). XPE binds to various damaged DNA induced by UV, cisplatin, depurination, and nitrogen mustard (Payne and Chu, 1994). Recent *in vivo* and *in vitro* studies showed that UV-DDB stimulates the excision of cyclobutane pyrimidine dimer, but not (6-4) photoproducts (Wakasugi *et al.*, 2001). However, XP-E mutant cells show the mildest NER defect among the XP groups and an *in vitro* NER reaction can be reconstituted without XPE (Aboussekhra *et al.*, 1995; Kazantsev *et al.*, 1998; Nichols *et al.*, 2000), suggesting that XPE may not be an essential factor for NER. In addition to the repair factors described above, high mobility group (HMG) domain protein and human Ku autoantigen (Ku70-Ku80 complex, a regulatory subunit of DNA-dependent protein kinase) bind to cisplatin-DNA adducts (Hughes *et al.*, 1992; Turchi and Henkels, 1996), although they are not essential for NER activity. The binding of HMG protein to cisplatin-damaged DNA inhibited an *in vitro* excision repair activity, suggesting that HMG domain protein may influence DNA repair *in vivo* (Huang *et al.*, 1994).

DNA structure requirements

NER machinery is responsible for recognizing and removing

most of the chemically modified bulky adducts, whereas non-bulky base adducts such as those caused by hydrolysis, oxygen free radicals, and simple alkylating agents are recognized and repaired by the base excision repair (BER). In NER, presence of non-complementary bases alone does not cause oligonucleotides excision. The loss of hydrogen bonding between two helical strands is an essential signal for the recruitment of NER enzymes. Oligonucleotides excision also requires chemical modification of DNA. For example, cisplatin covalently binds to the N7 positions of guanine and adenine bases and mainly forms 1, 2-intrastrand d(GpG) or d(ApG) adducts. The 1,3-intrastrand d(GpTpG) cisplatin is more structurally distorted than 1,2-intrastrand adducts and is therefore recognized and repaired more efficiently (Bellon *et al.*, 1991; Patrick and Turchi, 2001). Inclusion of non-complementary residue opposite the 1, 2-intrastrand d(GpG) adducts also increases the structural distortion of DNA and also increases repair efficiency significantly (Moggs *et al.*, 1997; Mu *et al.*, 1997). These findings suggest that local disruption of Watson-Crick base pairing as well as chemical modification of bases are indispensable components of the molecular signal that attracts human NER enzymes to the damaged DNA sites (Buschta-Hedayat *et al.*, 1999).

The cisplatin-DNA adduct containing 1, 2-intrastrand d(GpG) cross-link not only generates structural distortion of DNA, but also induces DNA unwinding and DNA bending ($>40^\circ$) toward the major groove (Rice *et al.*, 1988). All these changes in DNA structure are likely important for damage recognition in mammals since XPA binds most efficiently to rigidly bent duplexes but not to single-stranded DNA (Missura *et al.*, 2001), while RPA preferentially recognizes single-stranded DNA site rather than bent DNA. The association of XPA with RPA may generate dual sensorship that detects both backbone and base pair distortion of DNA. Since the affinity of XPA for bent duplexes is not compatible with its function in recognition of nucleotide lesions, XPA in collaboration with RPA may be involved in monitoring the integrity of the Watson-Crick base-pairing to verify the damage-specific localization of repair complexes (Buschta-Hedayat *et al.*, 1999; Missura *et al.*, 2001).

Damage recognition proteins

Requirement of multiple damage recognition factors in NER indicates that XPC-hHR23B, XPA, RPA, and TFIIH may all have unique role in damage recognition and the early stage of repair. It is still in debate however how these damage recognition proteins function at the damaged DNA site. XPA is a multifunctional zinc-finger protein involved in the damage recognition step of NER (Robins *et al.*, 1991; Guzder *et al.*, 1993; Jones *et al.*, 1993). Its zinc-finger motif is essential for its DNA binding activity as well as its function in NER (Matsuda *et al.*, 1995). XPA preferentially binds to (6-4) photoproduct of UV-irradiated DNA, although it exhibits a low affinity binding to UV-damaged DNA (Wang *et al.*,

Table 1. Mammalian proteins involved in early stage of nucleotide excision repair

Repair factor	(yeast homolog)	Size (kDa)	Role in repair
XPA	(RAD14)	31-kDa	Damage recognition, Interaction with RPA
RPA	(RFA1, 2, 3)	70, 34, 11-kDa	Damage recognition, DNA binding Interaction with XPA
XPC-hHR23B	(RAD4)	125-kDa (XPC)	Damage recognition, DNA binding
	(RAD23)	58-kDa (HR23B)	Stabilization of XPC, control of DNA repair?
TFIIH	(RAD25)	89-kDa (XPB)	DNA helicase
	(RAD3)	80-kDa (XPD)	DNA helicase
	(TFB-1)	62-kDa	Core TFIIH subunit
	(SSL1)	44-kDa	Core TFIIH subunit
	(KIN28)	41-kDa (Cdk7)	cyclin-dependent protein kinase
	(CCL1)	38-kDa	cyclin H
		34-kDa (CAK)	Cdk activating kinase
XPF	(RAD1)	112-kDa (XPF)	5'-incision
	(RAD10)	33-kDa (ERCC1)	5'-incision
XPG	(RAD2)	135-kDa (ERCC5)	3'-incision

2000). XPA's binding to damaged DNA is stabilized through interaction with RPA on damaged DNA (Stigge *et al.*, 1998; Wang *et al.*, 2000). XPA also plays a role in recruiting TFIIH to the damaged site for subsequent incision/excision step (Park *et al.*, 1995; Nocentini *et al.*, 1997).

RPA is a heterotrimeric protein complex consisting of 70-, 34-, and 11-kDa subunits involved in DNA replication, repair, and recombination (Wold, 1997). RPA is a 4-cysteine type zinc-finger protein and a mutation at any zinc-finger cysteine abolishes its function in repair (Stigge *et al.*, 1998; Dong *et al.*, 1999), suggesting a unique role for the zinc-finger domain in early stage of repair. In contrast to XPA, RPA's zinc-finger domain is not essential for DNA binding activity, but is involved in regulation of RPA's DNA binding activity through redox change [Lin *et al.*, 1998; Park *et al.*, 1999]. RPA preferentially binds to UV-damaged DNA and its interaction with UV-irradiated DNA was not affected by prior enzymatic photoreactivation of DNA, suggesting a preferred binding of RPA to the (6-4) photoproduct (Burns *et al.*, 1996). It is interesting to note that RPA, unlike XPA, may bind to duplex cisplatin-damaged DNA via generating single-stranded DNA at the lesion (Patrick *et al.*, 1999). The findings that RPA stimulates the XPA-DNA interaction through interaction with XPA on damaged DNA and also interacts with two endonucleases, XPG and ERCC1-XPF, suggest that the XPA-RPA complex, once formed on damaged DNA, recruits together XPG, ERCC1-XPF, and TFIIH to the damaged site for subsequent incision/excision step (He *et al.*, 1995; Matsunaga *et al.*, 1996). RPA may also be involved in a later stage of NER, gap-filling, that requires PCNA, RF-C, and DNA polymerase δ (or ϵ) (Aboussekhr *et al.*, 1995).

XPC-hHR23B is a human homolog of yeast Rad4 and Rad23 proteins, respectively, and has been shown to exhibit strong affinity for damaged DNA (Sugasawa *et al.*, 1998;

Batty *et al.*, 2000), as does the yeast counterpart, Rad4-Rad23 complex (Jansen *et al.*, 1998). The 106-kDa XPC protein forms a stable complex with 43-kDa HR23B (Masutani *et al.*, 1994; Shivji *et al.*, 1994). In yeast, Rad23 without Rad4 does not show any DNA binding under conditions where Rad4-Rad23 complex does (Guzder *et al.*, 1998), suggesting that Rad4 may be responsible for recognition of UV-damaged DNA. Rad23 (HR23B) is essential for XPC function in NER and may be necessary for the solubility of Rad4 (Guzder *et al.*, 1998). Recent studies also indicate that HR23B may be involved in regulation of repair activity. For example, Rad23 contains ubiquitin-associated (UBA) domain that may play a role in controlling NER through proteasome-mediated ubiquitin-dependent degradation of repair factors (Gillett *et al.*, 2001; Chen *et al.*, 2001). HR23B also interacts with base excision repair factor, MPG, suggesting that HR23B may have a unique role in DNA repair accommodating various repair pathways (Miao *et al.*, 2000).

Evidence suggests that XPC-hHR23B is involved in the recruitment of TFIIH to the damaged site through physical interaction with TFIIH (Yokoi *et al.*, 2000). XPC-hHR23B may collaborate with XPA in recruiting TFIIH since XPA also interacts with TFIIH on damaged DNA (Park *et al.*, 1995; Nocentini *et al.*, 1997). TFIIH, once loaded on the damaged DNA, may play a role in distinguishing the damaged strand from the non-damaged one since the translocation along a DNA strand by TFIIH DNA helicases is hampered by encountering a lesion (Naegeli *et al.*, 1992; Naegeli *et al.*, 1993). Local unwinding of the damaged DNA region by TFIIH likely generates junction between single-stranded DNA and duplex DNA that is recognized by two structure-specific endonucleases, XPG and ERCC1-XPF, for dual incision on the damaged strand.

Interaction of damage recognition proteins with DNA lesions

Both RPA and XPA preferentially bind to DNA lesions induced by cisplatin or UV-irradiation (Jones *et al.*, 1993; Guzder *et al.*, 1993; Asahina *et al.*, 1994; He *et al.*, 1995; Burns *et al.*, 1996; Wagasuki *et al.*, 1999; Wang *et al.*, 2000). RPA binds to (6-4) photoproduct-containing DNA with nearly 2-fold higher affinity relative to *cis-syn* cyclobutane dimer and nearly 4-fold greater than non-damaged DNA (Wang *et al.*, 2000). Although XPA showed preferential binding to (6-4) photoproduct-containing DNA, its interaction with damaged DNA was significantly weaker ($K_D = 2.13 \times 10^{-8}$ M) than RPA (2.02×10^{-10} M). Real-time analysis showed that XPA quickly dissociated from damaged DNA, while RPA remained associated with damaged DNA (Wang *et al.*, 2000; Wang *et al.*, 2001). Low affinity binding of XPA to damaged DNA is stabilized by RPA. A mutant RPA lacking XPA interaction domain failed to stabilize XPA-damaged DNA interaction, indicating that a role for RPA in NER may be to stabilize the interaction between XPA and damaged DNA through protein-protein interaction. Interaction of XPA with RPA on damaged DNA is not only crucial for damage-recognition and but may also play a role in subsequent steps in recruiting other DNA repair proteins such as XPG, ERCC1-XPF, and TFIIH to the damaged site (He *et al.*, 1995; Park *et al.*, 1995; Lee *et al.*, 1995; Li *et al.*, 1995; Saijo *et al.*, 1996; Stigger *et al.*, 1998).

The fact that both XPA and RPA showed only moderate preference to damaged DNA over non-damaged DNA raises a question as to whether XPA and/or RPA are responsible for recognition of damaged DNA *in vivo*. XPC-hHR23B showed a remarkable preference to UV-damaged circular DNA particularly in the presence of non-damaged competitor DNA (Sugasawa *et al.*, 1998; Sugasawa *et al.*, 2001). Preferential binding of XPC-hHR23B to damaged circular DNA was likely due to its higher binding affinity to UV-damaged DNA than to a non-damaged counterpart, but is also possible that high affinity binding may be due to structural alteration caused by UV-damage on circular DNA, but not on duplex DNA fragment. It is interesting that XPC-hHR23B, like RPA, physically interacts with XPA and stabilizes the XPA-damaged DNA interaction (Wang *et al.*, 2001). In fact, XPC-hHR23B and RPA exhibited remarkably similar biochemical properties: their binding to damaged DNA, physical interaction with XPA, and stabilization of XPA on damaged DNA. Nonetheless, preincubation of damaged DNA with XPC-hHR23B but not with RPA significantly enhanced XPA-damaged DNA interaction, suggesting that damage recognition complex may be formed in an ordered process, such that XPC-hHR23B interacts with damaged DNA before addition of XPA and RPA to form a stable damage recognition complex (Sugasawa *et al.*, 2001). A recent immunohistochemistry study also strongly supports a role for XPC as a global initiator in repair (Volker *et al.*, 2001) while suggesting a role for XPA and RPA as repair mediator

proteins. Biochemical basis for the interaction of XPC and damaged DNA needs to be addressed to clarify the discrepancy.

It is highly speculated that three damage recognition proteins (XPA, RPA, and XPC-hHR23B) form a complex at the damaged DNA site, although there is no direct evidence on that. A potential implication of forming damage recognition complex on damaged DNA is to position the preincision machinery for dual incision reaction at damaged DNA strand. In DNA replication, origin binding protein such as SV40 T-antigen form a multiprotein complex with RPA and DNA polymerase (pol) α -primase complex for the proper positioning of pol α -primase to initiate DNA synthesis at the specific site (Hurwitz *et al.*, 1990; Bullock *et al.*, 1991). Likewise, the formation of multiprotein complex at damaged site may be necessary for proper positioning of two endonucleases XPG and ERCC1-XPF for accurate incision at the 3' and 5' sites of the damaged strand (de Laat *et al.*, 1998).

Protein-induced structural distortion of damaged DNA

In DNA replication, binding of origin recognition factor(s) to the replication origin leads to a significant structural distortion at AT-rich region, which is necessary for loading of DNA helicase(s) (Borowiec *et al.*, 1990 and the references therein). During NER, distortion of damaged DNA may also be essential for loading of two DNA helicases (XPB and XPD of the TFIIH complex) to the damaged site. Structural distortion at damaged DNA is not only crucial for the formation of preincision complex (PIC), but also necessary for dual incisions. Since the formation of PIC at damaged site requires RPA, XPA, XPC-hHR23B, and TFIIH in the expense of ATP hydrolysis (Evans *et al.*, 1997a; Evans *et al.*, 1997b; Mu *et al.*, 1997a), it is conceivable that multiple repair factors contribute to a structural distortion of damaged DNA. RPA is a leading candidate for structural distortion of damaged DNA. RPA's zinc-finger motif is essential for its function in NER (Stigger *et al.*, 1998). The failure of a zinc-finger mutant in supporting NER was not due to its role in stabilization of XPA-damaged DNA complex (Park *et al.*, 1999; Wang *et al.*, 2000), suggesting that RPA may have a unique role in early stage of repair. Tryptic digestion pattern and far-UV CD spectra analysis indicated that RPA undergoes a significant conformational change upon binding to DNA in its zinc-finger motif-dependent manner (Wang *et al.*, 2001). RPA's zinc-finger motif may mediate the transition of RPA-XPA interaction into a stable RPA-XPA-damaged DNA complex through its conformational change (Park *et al.*, 1999; Wang *et al.*, 2001). Alternatively, both RPA and XPA, two zinc-finger proteins, undergo conformational changes upon binding to the damaged DNA (Wang *et al.*, 2001) to further distort the damaged DNA. It is also interesting to note that requirement for XPC-hHR23B in dual incisions can be bypassed by the use of a bubble-like DNA substrate containing a thymidine

dimer (Mu *et al.*, 1997b), suggesting that XPC-hHR23B may also play a role in inducing a structural distortion and/or local opening of the damaged DNA site.

Concluding Remarks

The NER machinery is primarily responsible for the repair of various intrastrand cross-linked DNA (Sancar, 1996; Wood, 1996), although some NER factors such as ERCC1-XPF may also participate in the repair of interstrand cross-linked damage (Kuraoka *et al.*, 2000; Mu *et al.*, 2000). Although the details of damage recognition process is still not clear, all three damage recognition factors, XPA, RPA, and XPC-hHR23B, play an essential role in damage recognition and recruiting other repair factors to the damaged site for dual incision. Protein-induced structural distortion at the damaged DNA is likely necessary for the loading of TFIIH complex (XPB and XPD helicases) to initiate unwinding of both 3'- and 5'-sides of DNA damage. Therefore, a new challenge in early stage of repair is to unravel the role of damage recognition factors in further helix destabilization at damaged DNA site.

Another future challenge is to understand how damage recognition factors effectively recognize DNA damage on chromatin as compared to that on naked DNA as the chromosomal DNA *in vivo* is assembled into a compact nucleosome structure. We do not know the answer, however, chromatin accessibility factors such as GADD45 may play a role in mediating the recognition of damaged DNA on chromatin (Carrier *et al.*, 1999). Also, various DNA damages induce chromatin modifications such as phosphorylation of histone H2AX and acetylation of histone H3 (Paull *et al.*, 2000; Brand *et al.*, 2001). These modifications would likely facilitate the access of damage recognition proteins and/or DNA repair machinery to lesions.

Acknowledgments This work was supported by grants from the National Institute of Health (R01 CA92111) and the U.S. Army (DAMD17-00-1-0295).

References

- Aboussekhra, A., Biggerstaff, M., Shivji, M. K., Vilpo, J. A., Moncollin, V., Podust, V. N., Protic, M., Hubscher, U., Egly, J. M. and Wood, R. D. (1995) Mammalian DNA nucleotide excision repair reconstituted with purified protein components. *Cell* **80**, 859-868.
- Asahina, H., Kuraoka, I., Shirakawa, M., Morita, E. H., Miura, N., Miyamoto, I., Ohtsuka, E., Okada, Y. and Tanaka, K. (1994) The XPA protein is a zinc metalloprotein with an ability to recognize various kinds of DNA damage. *Mutat. Res.* **315**, 229-237.
- Batty, D., Rapic'Otrin, V., Levine, A. S. and Wood, R. D. (2000) Stable Binding of Human XPC Complex to Irradiated DNA Confers Strong Discrimination for Damaged Sites. *J. Mol. Biol.* **300**, 275-290.
- Batty, D. P. and Wood, R. D. (2000) Damage recognition in nucleotide excision repair of DNA. *Gene* **241**, 193-204.
- Bellon, S. F., Coleman, J. H. and Lippard, S. J. (1991) DNA unwinding produced by site-specific intrastrand cross-links of the antitumor drug cis-diamminedichloroplatinum (II). *Biochemistry* **30**, 8026-8035.
- Bessho, T., Sancar, A., Thompson, L. H. and Telen, M. P. (1997) Reconstitution of human excision nuclease with recombinant XPF-ERCC1 complex. *J. Biol. Chem.* **272**, 3833-3837.
- Borowiec, J. A., Dean, F. D., Bullock, P. A. and Hurwitz, J. (1990) Binding and unwinding-how T antigen engages the SV40 origin of replication. *Cell* **60**, 181-184.
- Brand, M., Moggs, J. G., Oulad-Abdelghani, M., Lejeune, F., Dilworth, F. J., Stevenin, J., Almouzni, G. and Tora, L. (2001) UV-damaged DNA-binding protein in the TFIIH complex links DNA damage recognition to nucleosome acetylation. *EMBO J.* **20**, 3187-3196.
- Bullock, P. A., Seo, Y. S. and Hurwitz, J. (1991) Initiation of simian virus 40 DNA synthesis *in vitro*. *Mol. Cell Biol.* **11**, 2350-2361.
- Burns, J. L., Guzder, S. N., Sung, P., Prakash, S. and Prakash, L. (1996) An affinity of human replication protein A for ultraviolet-damaged DNA. *J. Biol. Chem.* **271**, 11607-11610.
- Buschta-Hedayat, N., Buterin, T., Hess, M. T., Missura, M. and Naegeli, H. (1999) Recognition of nonhybridizing base pairs during nucleotide excision repair of DNA. *Proc. Natl. Acad. Sci. USA* **96**, 6090-6095.
- Carrier, F., Georgel, P. T., Pourquier, P., Blake, M., Kontny, H. U., Antinore, M. J., Gariboldi, M., Myers, T. G., Weinstein, J. N., Pommier, Y. and Fornace, A. J. Jr. (1999) Gadd45, a p53-responsive stress protein, modifies DNA accessibility on damaged chromatin. *Mol. Cell Biol.* **19**, 1673-1685.
- Chen, L., Shinde, U., Ortolan, T. G. and Madura, K. (2001) Ubiquitin-associated (UBA) domains in Rad23 bind ubiquitin and promote inhibition of multi-ubiquitin chain assembly. *EMBO Rep.* **2**, 933-938.
- Dawn, P. B. and Wood, R. D. (2000) Damage recognition in nucleotide excision repair of DNA. *Gene* **241**, 193-204.
- de Laat, W. L., Appeldoorn, E., Sugawara, K., Weterings, E., Jaspers, N. G. and Hoeijmakers, J. H. (1998) DNA-binding polarity of human replication protein A positions nucleases in nucleotide excision repair. *Genes Dev.* **12**, 2598-2609.
- Dong, J., Park, J.-S. and Lee, S.-H. (1999) In vitro analysis of zinc-finger motif in human replication protein A. *Biochem. J.* **337**, 311-317.
- Evans, E., Fellows, J., Coffey, A. and Wood, R. D. (1997a) Open complex formation around a lesion during nucleotide excision repair provides a structure for cleavage by human XPG protein. *EMBO J.* **16**, 625-638.
- Evans, E., Moggs, J. G., Hwang, J. R., Egly, J. M. and Wood, R. D. (1997b) Mechanism of open complex and dual incision formation by human nucleotide excision repair factors. *EMBO J.* **16**, 6559-6573.
- Gillette, T. G., Huang, W., Russell, S. J., Reed, S. H., Johnston, S. A. and Friedberg, E. C. (2001) The 19S complex of the proteasome regulates nucleotide excision repair in yeast. *Genes Dev.* **15**, 1528-1539.
- Guzder, S. N., Sung, P., Prakash, L. and Prakash, S. (1993) Yeast DNA-repair gene RAD14 encodes a zinc metalloprotein with affinity for ultraviolet-damaged DNA. *Proc. Natl. Acad. Sci.*

- USA **90**, 5433-5437.
- Guzder, S. N., Sung, P., Prakash, L. and Prakash, S. (1998) Affinity of yeast nucleotide excision repair factor 2, consisting of the Rad4 and Rad23 proteins, for ultraviolet damaged DNA. *J. Biol. Chem.* **273**, 31541-31546.
- He, Z., Henriksen, L. A., Wold, M. S. and Ingles, C. J. (1995) RPA involvement in the damage-recognition and incision steps of nucleotide excision repair. *Nature* **374**, 566-569.
- Huang, J. C., Zamble, D. B., Reardon, J. T., Lippard, S. J. and Sancar, A. (1994) HMG-domain proteins specifically inhibit the repair of the major DNA adduct of the anticancer drug cisplatin by human excision nuclease. *Proc. Natl. Acad. Sci. USA* **91**, 10394-10398.
- Hughes, E. N., Engelsberg, B. N. and Billings, P. C. (1992) Purification of nuclear proteins that bind to cisplatin-damaged DNA. Identity with high mobility group proteins 1 and 2. *J. Biol. Chem.* **267**, 13520-13527.
- Hurwitz, J., Dean, F. B., Kwong, A. D. and Lee, S.-H. (1990) The *in vitro* Replication of DNA containing the SV40 origin. *J. Biol. Chem.* **266**, 594-602.
- Jansen, L. E., Verhage, R. A. and Brouwer, J. (1998) Preferential binding of yeast Rad4-Rad23 complex to damaged DNA. *J. Biol. Chem.* **273**, 33111-33114.
- Jones, C. J. and Wood, R. D. (1993) Preferential binding of the Xeroderma pigmentosum group A complementing protein to damaged DNA. *Biochemistry* **32**, 12096-12104.
- Kazantsev, A., Bullock, S. K., Sancar, A. and Cordeiro-Stone, M. (1998) Generation and characterization of an immortal cell line of xeroderma pigmentosum group E. *Mutat. Res.* **407**, 55-65.
- Kuraoka, I., Kobertz, W. R., Ariza, R. R., Biggerstaff, M., Essigmann, J. M. and Wood, R. D. (2000) Repair of an interstrand DNA cross-link initiated by ERCC1-XPF Repair/Recombination nuclease. *J. Biol. Chem.* **275**, 26632-26636.
- Lee, S. H., Kim, D. K. and Drissi, R. (1995) Human xeroderma pigmentosum group A protein interacts with human replication protein A and inhibits DNA replication. *J. Biol. Chem.* **270**, 21800-21805.
- Li, L., Lu, X., Peterson, C. A. and Legerski, R. J. (1995) An interaction between the DNA repair factor XPA and replication protein A appears essential for nucleotide excision repair. *Mol. Cell Biol.* **15**, 5396-5402.
- Lin, Y. L., Shivji, M. K., Chen, C., Kolodner, R., Wood, R. D. and Dutta, A. (1998) The evolutionarily conserved zinc finger motif in the largest subunit of human replication protein A is required for DNA replication and mismatch repair but not for nucleotide excision repair. *J. Biol. Chem.* **273**, 1453-1461.
- Masutani, C., Sugawara, K., Yanagisawa, J., Sonoyama, T., Ui, M., Enomoto, T., Takio, K., Tanaka, K., van der Spek, P. J. and Bootsma, D. (1994) Purification and cloning of a nucleotide excision repair complex involving the xeroderma pigmentosum group C protein and a human homologue of yeast RAD23. *EMBO J.* **13**, 1831-1843.
- Matsuda, T., Saijo, M., Kuraoka, I., Kobayashi, T., Nakatsu, Y., Nagai, A., Enjoji, T., Masutani, C., Sugawara, K. and Hanaoka, F. (1995) DNA repair protein XPA binds replication protein A (RPA). *J. Biol. Chem.* **270**, 4151-4157.
- Matsunaga, T., Park, C. H., Bessho, T., Mu, D. and Sancar, A. (1996) Replication protein A confers structure-specific endonuclease activities to the XPF-ERCC1 and XPG subunits of human DNA repair excision nuclease. *J. Biol. Chem.* **271**, 11047-11050.
- Miao, F., Bouziane, M., Dammann, R., Masutani, C., Hanaoka, F., Pfeifer, G. and O'Connor, T. R. (2000) 3-Methyladenine-DNA glycosylase (MPG protein) interacts with human RAD23 proteins. *J. Biol. Chem.* **275**, 28433-28438.
- Missura, M., Buterin, T., Hindges, R., Hübscher, U., Kaspárková, J., Brabec, V. and Naegeli, H. (2001) Double-check probing of DNA bending and unwinding by XPARPA: an architectural function in DNA repair. *EMBO J.* **20**, 3554-3564.
- Moggs, J. G., Yarema, K. J., Essigmann, J. M. and Wood, R. D. (1997) Analysis of incision sites produced by human cell extracts and purified proteins during nucleotide excision repair of a 1,3-intrastrand d(GpTpG)-cisplatin adduct. *J. Biol. Chem.* **271**, 7177-7186.
- Mu, D., Wakasugi, M., Hsu, D. S. and Sancar, A. (1997) Characterization of reaction intermediates of human excision repair nuclease. *J. Biol. Chem.* **272**, 28971-28979.
- Mu, D. and Sancar, A. (1997) Model for XPC-independent transcription-coupled repair of pyrimidine dimers in humans. *J. Biol. Chem.* **272**, 7570-7573.
- Mu, D., Bessho, T., Nechev, L. V., Chen, D. J., Harris, T. M., Hearst, J. E. and Sancar, A. (2000) DNA interstrand cross-links induce futile repair synthesis in mammalian cell extracts. *Mol. Cell Biol.* **20**, 2446-2454.
- Naegeli, H., Bardwell, L. and Friedberg, E. C. (1992) The DNA helicase and adenosine triphosphatase activities of yeast Rad3 protein are inhibited by DNA damage. A potential mechanism for damage-specific recognition. *J. Biol. Chem.* **267**, 392-398.
- Naegeli, H., Bardwell, L. and Friedberg, E. C. (1993) Inhibition of Rad3 DNA helicase activity by DNA adducts and abasic sites: implications for the role of a DNA helicase in damage-specific incision of DNA. *Biochemistry* **32**, 613-621.
- Nichols, A. F., Itoh, T., Graham, J. A., Liu, W., Yamaizumi, M. and Linn, S. (2000) Human damage-specific DNA-binding protein p48. Characterization of XPE mutations and regulation following UV irradiation. *J. Biol. Chem.* **275**, 21422-21428.
- Nocentini, S., Coin, F., Saijo, M., Tanaka, K. and Egly, J. M. (1997) DNA damage recognition by XPA protein promotes efficient recruitment of transcription factor II H. *J. Biol. Chem.* **272**, 22991-22994.
- Park, C. H., Mu, D., Reardon, J. T. and Sancar, A. (1995) The general transcription-repair factor TFIIH is recruited to the excision repair complex by the XPA protein independent of the TFIIIE transcription factor. *J. Biol. Chem.* **270**, 4896-4902.
- Park, J. S., Park, J. S., Wang, M., Park, S. J. and Lee, S. H. (1999) Zinc-finger of replication protein A, a non-DNA binding element, regulates its DNA binding activity through redox. *J. Biol. Chem.* **274**, 29075-29080.
- Patrick, S. M. and Turchi, J. J. (1998) Human replication protein A preferentially binds cisplatin-damaged duplex DNA *in vitro*. *Biochemistry* **37**, 8808-8815.
- Patrick, S. M. and Turchi, J. J. (1999) Replication protein A (RPA) binding to duplex cisplatin-damaged DNA is mediated through the generation of single-stranded DNA. *J. Biol. Chem.* **274**, 14972-14978.
- Patrick, S. M. and Turchi, J. J. (2001) Stopped-flow kinetic analysis of replication protein A-binding DNA: damage recognition and affinity for single-stranded DNA reveal differential contributions of k_{on} and k_{off} rate constants. *J. Biol. Chem.* **276**, 22630-22637.

- Paull, T. T., Rogakou, E. P., Yamazaki, V., Kirchgessner, C. U., Gellert, M. and Bonner, W. M. (2000) A critical role for histone H2AX in recruitment of repair factors to nuclear foci after DNA damage. *Curr. Biol.* **10**, 886-895.
- Payne, A. and Chu, G. (1994) Xeroderma pigmentosum group E binding factor recognizes a broad spectrum of DNA damage. *Mutat. Res.* **310**, 89-102.
- Rice, J. A., Crothers, D. M., Pinto, A. L. and Lippard, S. J. (1988) The major adduct of the antitumor drug cis-diamminedichloroplatinum(II) with DNA bends the duplex by approximately equal to 40 degrees toward the major groove. *Proc Natl Acad Sci USA* **85**, 4158-4161.
- Robins, P., Jones, C. J., Biggerstaff, M., Lindahl, T. and Wood, R. D. (1991) Complementation of DNA repair in xeroderma pigmentosum group A cell extracts by a protein with affinity for damaged DNA. *EMBO J.* **10**, 3913-3921.
- Saijo, M., Kuraoka, I., Masutani, C., Hanaoka, F. and Tanaka, K. (1996) Sequential binding of DNA repair proteins RPA and ERCC1 to XPA in vitro. *Nucleic Acids Res.* **24**, 4719-4724.
- Sancar, A. (1996) DNA excision repair. *Ann. Rev. Biochem.* **65**, 43-81.
- Shivji, M. K. K., Eker, A. P. M. and Wood, R. D. (1994) DNA repair defect in xeroderma pigmentosum group C and complementing factor from HeLa cells. *J. Biol. Chem.* **269**, 22749-22757.
- Stigger, E., Drissi, R. and Lee, S-H. (1998) Functional Analysis of human replication protein A in Nucleotide Excision Repair. *J. Biol. Chem.* **273**, 9337-9343.
- Sugasawa, K., Ng, J. M., Masutani, C., Iwai, S., van der Spek, P. J., Eker, A. P., Hanaoka, F., Bootsma, D. and Hoeijmakers, J. H. (1998) Xeroderma pigmentosum group C protein complex is the initiator of global genome nucleotide excision repair. *Mol. Cell.* **2**, 223-232.
- Sugasawa, K., Okamoto, T., Shimizu, Y., Masutani, C., Iwai, S. and Hanaoka, F. (2001) A multistep damage recognition mechanism for global genomic nucleotide excision repair. *Genes Dev.* **15**, 507-521.
- Turchi, J. J. and Henkels, K. (1996) Human Ku autoantigen binds cisplatin-damaged DNA but fails to stimulate human DNA-activated protein kinase. *J. Biol. Chem.* **271**, 13861-13867.
- Volker, M., Mone, M. J., Karmakar, P., van Hoffen, A., Schul, W., Vermeulen, W., Hoeijmakers, J. H., van Driel, R., van Zeeland, A. A. and Mullenders, L. H. (2001) Sequential assembly of the nucleotide excision repair factors *in vivo*. *Mol. Cell* **8**, 213-224.
- Wakasugi, M. and Sancar, A. (1999) Order of assembly of human DNA repair excision nuclease. *J. Biol. Chem.* **274**, 18759-18768.
- Wagasuki, M., Shimizu, M., Morioka, H., Linn, S., Nikaido, O. and Matsunaga, T. (2001) Damaged DNA binding protein DDB stimulates the excision of cyclobutane pyrimidine dimers in vitro in concert with XPA and replication protein A. *J. Biol. Chem.* **276**, 15434-15440.
- Wang, M., Mahrenholz, A. and Lee, S. H. (2000) RPA stabilizes the XPA-damaged DNA complex through protein-protein interaction. *Biochemistry* **39**, 6433-6439.
- Wang, M., You, J-S. and Lee, S-H. (2001) Functional analysis of XPC-hHR23B in early stage of nucleotide excision repair. (submitted)
- Wood, R. D. (1996) DNA repair in eukaryotes. *Ann. Rev. Biochem.* **65**, 135-167.
- Wold, M. S. (1997) Replication protein A: a heterotrimeric, single-stranded DNA-binding protein required for eukaryotic DNA metabolism. *Ann. Rev. Biochem.* **66**, 61-92.
- Yokoi, M., Masutani, C., Maekawa, T., Sugasawa, K., Ohkuma, Y. and Hanaoka, F. (2000) The xeroderma pigmentosum group C protein complex XPC-hHR23B plays an important role in the recruitment of transcription factor IIH to damaged DNA. *J. Biol. Chem.* **275**, 9870-9875.
- Wang, M., You, J-S. and Lee, S-H. (2001) Role of zinc finger motif in redox regulation of human replication protein A. *Antioxidants & Redox Signaling*, **3**, 657-669.

Involvement of DNA-Dependent Protein Kinase in Regulation of Stress-Induced JNK Activation

SU-JUNG PARK, EUN-JIN OH,¹ MI-AE YOO,¹ and SUK-HEE LEE

ABSTRACT

DNA-dependent protein kinase (DNA-PK) is composed of a 460-kDa catalytic subunit and the regulatory subunits Ku70 and Ku80. The complex is activated on DNA damage and plays an essential role in double-strand-break repair and V(D)J recombination. In addition, DNA-PK is involved in S-phase checkpoint arrest following irradiation, although its role in damage-induced checkpoint arrest is not clear. In an effort to understand the role of DNA-PK in damage signaling, human and mouse cells containing the DNA-PK catalytic subunit (DNA-PKcs proficient) were compared with those lacking DNA-PKcs for c-Jun N-terminal kinase (JNK) activity that mediates physiologic responses to DNA damage. The DNA-PKcs-proficient cells showed much tighter regulation of JNK activity after DNA damage, while the level of JNK protein in both cell lines remained unchanged. The JNK proteins physically associated with DNA-PKcs and Ku70/Ku80 heterodimer, and the interaction was significantly stimulated after DNA damage. Various JNK isoforms not only contained a DNA-PK phosphorylation consensus site (serine followed by glutamine) but also were phosphorylated by DNA-PK *in vitro*. Together, our results suggest that DNA damage induces physical interaction between DNA-PK and JNK, which may in turn negatively affect JNK activity through JNK phosphorylation by DNA-PK.

INTRODUCTION

CELLULAR RESPONSES to various genotoxic stresses include cell-cycle arrest, DNA repair, and apoptosis, all of which occur through damage signaling pathways initiated from either DNA damage or the plasma membrane (Angel and Karin, 1991; Liu *et al.*, 1996). DNA-dependent protein kinase (DNA-PK), ataxia telangiectasia-mutated gene (*ATM*), and the AT-related gene (*ATR*) are upstream components of damage signaling pathways and are linked to cell-cycle checkpoint arrest (Canman *et al.*, 1994; Lees-Miller *et al.*, 1995; Park *et al.*, 1999; Rathmell *et al.*, 1997). The c-Jun N-terminal kinase (JNK) pathway is most notably initiated from the plasma membrane; however, it is strongly induced on treatment with various DNA-damaging drugs, suggesting that the JNK signaling pathway mediates a physiologic response to DNA damage (Coso *et al.*, 1995; Devary *et al.*, 1991; Liu *et al.*, 1996; Minden *et al.*, 1995).

DNA-PK is a heterotrimeric enzyme composed of a 460-kDa

catalytic subunit (DNA-PKcs) and the two regulatory subunits, Ku80 and Ku70. DNA-PKcs is a serine/threonine protein kinase and is recruited to DNA strand-break sites necessary to activate its kinase activity (Casciola-Rosen *et al.*, 1995; Finnie *et al.*, 1995; Gupta *et al.*, 1995; Hendrickson *et al.*, 1991; Lieber *et al.*, 1997; Peterson *et al.*, 1997; Ramsden and Gellert, 1998; Savitsky *et al.*, 1995). Studies with DNA-PK-defective mutant cells indicated that DNA-PK is essential in the repair of double-strand breaks (DSBs) and for V(D)J recombination (Blunt *et al.*, 1995; Hendrickson *et al.*, 1991; Jackson, 1997). Compared with their wildtype counterparts, DNA-PKcs-deficient cell lines were additionally sensitive to heat-induced apoptosis (Hendrickson *et al.*, 1991), suggesting a role for DNA-PK in anti-apoptotic responses (Gupta *et al.*, 1995; Hendrickson *et al.*, 1991; Muller *et al.*, 1998; Peterson *et al.*, 1995). *In vitro*, DNA-PK phosphorylates heat shock protein 90 (hsp90) and a number of DNA-binding proteins, including the Ku autoantigen, replication protein A (RPA), simian virus 40 large tumor anti-

Department of Biochemistry and Molecular Biology and the Indiana University Cancer Center and Walther Oncology Center, Indiana University School of Medicine, Indianapolis, Indiana.

¹Permanent address: Department of Molecular Biology, Pusan National University, Pusan, Korea.

gen (SV40 Tag), the tumor suppressor protein p53, and several transcription factors (Sp-1, Oct-1, Fos, and serum response factor) (Lees-Miller *et al.*, 1990). The physiologic role of DNA-PK in phosphorylation of these proteins, however, is not clear.

The JNK/SAPK signaling pathway in response to genotoxic stress is thought to be largely a compensatory and protective response that positively regulates genes involved in cellular protective functions, such as glutathione-S-transferase (GST), quinone reductase, and MTIIA (Bergelson *et al.*, 1994; Litz-Jackson *et al.*, 1992; Pinkus *et al.*, 1996). On the other hand, persistent JNK activation may promote, rather than protect against, apoptosis in certain cells, although JNK/SAPK alone may not be sufficient to induce apoptosis (Sanchez-Perez *et al.*, 1998; Widmann *et al.*, 1997). Apoptosis signal-regulating kinase (ASK1) has been identified as a mitogen-activated protein kinase (MAPK) kinase kinase (MAPKKK) that activates the JNK and p38 MAP kinase pathways (Ichijo *et al.*, 1997). It is also required for TNF- α -induced apoptosis; however, whether JNK activation is directly involved in this apoptotic pathway remains controversial (Faris *et al.*, 1998; Guo *et al.*, 1998; Roulston *et al.*, 1998). In addition, the possible involvement of the JNK/SAPK signaling pathway, such as the MAPK cascades, in cell proliferation and control of DNA metabolism has been suggested (Auer *et al.*, 1998; Potapova *et al.*, 1997). At present, a role for the JNK signaling pathway in cellular proliferation, DNA replication, or apoptosis has yet to be defined.

The JNK protein was initially characterized as a MAPK family member that binds and phosphorylates c-Jun at serines 63 and 73 located within its transactivation domain (Angel *et al.*, 1988; Kyriakis *et al.*, 1994; Pulverer *et al.*, 1991; Smeal *et al.*, 1991). Phosphorylation of these serine sites resulted in increased transactivation potential of c-Jun (Derijard *et al.*, 1994; Smeal *et al.*, 1991). The substrates of JNK have now been extended to include other transcription factors, such as ATF-2 and ELK-1 (Gupta *et al.*, 1995, 1996; Livingstone *et al.*, 1995). Ten JNK isoforms have been identified from human brain, and two main forms, of 46 kDa and 55 kDa, correspond to alternatively spliced variants of JNK1, JNK2, and JNK3 (Davis, 1994; Gupta *et al.*, 1995). The activity of JNK is regulated by dual phosphorylation of specific threonine and tyrosine residues by MKK4 (otherwise known as SEK1) (Duyster *et al.*, 1995; Jackson and Jeggo, 1995; Johnson *et al.*, 1996). Activation of JNK has been shown to occur in response to various types of external stress such as heat shock, inflammatory cytokines (Foltz *et al.*, 1998; Moriguchi *et al.*, 1997; Raingeaud *et al.*, 1995) and DNA-damaging agents: ultraviolet light, ionizing radiation, cisplatin (Liu *et al.*, 1996; Moriguchi *et al.*, 1997; Schreiber *et al.*, 1995) and mitomycin C (MMC; Liu *et al.*, 1996). Unlike p53 induction, JNK activation is not a universal response to DNA damage. Different genotoxic agents vary greatly in their ability to activate JNK yet are effective in inducing p53 accumulation, growth arrest, and eventual cell death.

Here, we report our investigation into the involvement of DNA-PK in the regulation of the JNK signaling pathway after genotoxic stress. We found that JNK activity in DNA-PKs-proficient cells, in contrast to the activity in DNA-PKs-deficient cells, was tightly regulated. In addition, our *in vitro* studies revealed that JNK protein associated with DNA-PK and was efficiently phosphorylated by DNA-PK, suggesting that DNA-

PK plays an important role in controlling JNK activity in response to DNA damage.

MATERIALS AND METHODS

Proteins, chemicals, and antibodies

Polyclonal antibodies to JNK1 and JNK1/2 were obtained from Santa Cruz Biotechnology (Santa Cruz, CA) or Pharmin-gen (San Diego, CA). The GST fusion form of c-Jun protein (GST-cJun; a kind gift from Dr. S. Boswell, Indiana University School of Medicine, Indianapolis, IN) containing residues 1 through 79 of human c-Jun was overexpressed in *Escherichia coli* and purified using glutathione-agarose affinity column chromatography. Purified GST-JNK2 (77-kDa) protein and GST-JNK1 expression plasmid were obtained from Santa Cruz Biotechnology. DNA-PK holoenzyme was purified from HeLa cells according to the procedure described previously (Lees-Miller *et al.*, 1990). The [γ - ^{32} P]-ATP (4500 Ci/mmol) was obtained from ICN. Epidermal growth factor (EGF), Adriamycin (doxorubicin), MMC, and cisplatin were purchased from Sigma Chemical Co. (St. Louis, MO), and methylmethane sulfate (MMS) was obtained from Aldrich (Milwaukee, WI).

Cell culture and treatment conditions

Malignant glioblastoma cells, M059K (DNA-PKcs proficient) and M059J (DNA-PKcs deficient) (obtained from Dr. M. J. Allalunis-Turner, Cross Cancer Institute, Edmonton, Alb., Canada), were grown in tissue culture dishes in DMEM/F12 supplemented with 10% FBS at 37°C in a CO₂ incubator. Mouse cell lines CB-17 (DNA-PKcs proficient) and SCID (DNA-PKcs deficient) (obtained from Drs. J.M. Brown and C. Kirchgessner, Department of Radiation Oncology, Stanford University, Stanford, CA) were maintained in DMEM supplemented with 10% FBS. To see the effect of the EGF (20 ng/ml) and genotoxic agents, cells were treated with MMS (0.1 or 1.0 mM), MMC (0.1 mM), or Adriamycin (1.0 μ g/ml) and incubated until harvest.

Cell preparation and JNK immune complex assay

Cells were grown in culture medium containing 0.5% FBS for 16 h prior to treatment with EGF or genotoxic agents. Cells were washed in ice-cold PBS, with addition of 0.5 ml of JNK lysis buffer (25 mM HEPES, pH 7.5; 0.3 M NaCl, 1.5 mM MgCl₂, 0.2 mM EDTA, 0.5 mM dithiothreitol, 0.5% Triton X-100, 20 mM β -glycerophosphate, 1 mM sodium vanadate, 0.1 μ M okadaic acid, 1 mM phenylmethylsulfonyl fluoride, aprotinin 20 μ g/ml, leupeptin 50 μ g/ml, and 10 μ M pepstatin to the dishes (150 $\times$ 25 mm) before harvest. After 30 min on ice, insoluble material was removed by sedimentation for 30 min at 12,000 $\times$ g. The JNK activity was determined by an immunocomplex assay essentially as described (Duyster *et al.*, 1995; Litz-Jackson *et al.*, 1992). Briefly, cell extracts (200 μ g) were mixed with 1.5 μ l of anti-JNK1/JNK2 polyclonal antibody, and 15 μ l of protein A-Sepharose beads was added for further incubation at 4°C for 3 h. The immune complexes were washed three times with JNK lysis buffer (0.5 ml each time) and once with JNK reaction buffer (20 mM HEPES, pH 7.5; 10 mM

MgCl₂, 7 mM MnCl₂, 1 mM EGTA, 1 mM sodium fluoride, 1 mM sodium vanadate, and 1 mM DTT). The immunoprecipitate was then resuspended in 30 μ l of JNK reaction buffer containing 2 μ g of GST-c-Jun (Litz-Jackson *et al.*, 1992), 50 μ M ATP; and the reaction was initiated by the addition of 1.0 μ l of [γ -³²P]-ATP (45,000 Ci/mmol). After incubation for 20 min at 30°C, the reaction was terminated by the addition of 8 μ l of 5 $\times$ SDS sample buffer (Laemmli, 1970) and heating to 95°C for 5 min. Samples were analyzed by 12% SDS-PAGE.

Coimmunoprecipitation and Western blot analysis

Cell extracts (500 μ g) were incubated with the respective antibodies and protein A-Sepharose for 3 h at 4°C. Precipitates were washed three times with JNK lysis buffer, suspended in SDS sample buffer, boiled for 3 min, and subjected to SDS-PAGE on 6% or 10% gels. Following gel electrophoresis, proteins were transferred to a nitrocellulose membrane and immunoblotted with primary antibody followed by a peroxidase-coupled secondary antibody (Amersham) and an enhanced chemiluminescence reaction mixture (Amersham) prior to visualization on Kodak X-O-mat film.

In vitro phosphorylation of JNK

Purified DNA-PK holoenzyme or cell extracts were mixed with 20 μ l of buffer containing 20 mM HEPES, pH 7.5; 10 mM MgCl₂, 0.1 mM EDTA, 2 mM EGTA, 0.125 mM ATP, 1.0 μ g

of substrate peptide (EPPLSQEAFADLWKK) representing amino acids 11 through 24 of p53 (Lees-Miller *et al.*, 1990), and 1.0 μ l of [γ -³²P]-ATP (45,000 Ci/mmol) for 20 min at 30°C. Where indicated, 1.0 μ g of activated calf thymus DNA (Sigma) was included in the reaction mixtures. Phosphorylation of protein was analyzed by 10% SDS-PAGE followed by autoradiography.

RESULTS

Stress-induced JNK activation is tightly regulated in DNA-PKcs-proficient cells

Although the JNK signaling pathway is largely responsible for the activity of genes involved in cellular protective functions, such as GST, quinone reductase, and MTIIA (Bergelson *et al.*, 1994; Litz-Jackson *et al.*, 1992; Pinkus *et al.*, 1996), continuous JNK activation may induce apoptosis in certain cells (Sanchez-Perez *et al.*, 1998; Widmann *et al.*, 1997), suggesting that regulation of JNK is important for cell protection against apoptosis. In contrast, the physiologic role of DNA-PK may be to protect cells from DNA damage because DNA-PK is involved in immediate S-phase arrest and may be necessary for the efficient repair of DNA damage (Anderson and Carter, 1996; Park *et al.*, 1999). DNA-PK may be directly involved in S-phase arrest, targeting replication




FIG. 1. Damage-induced JNK1 activation in DNA-PKcs-deficient and DNA-PKcs-proficient cells. Two human malignant glioblastoma cells (M059K [DNA-PKcs proficient] and M059J [DNA-PKcs deficient]) (A) or two mouse cell lines (CB-17 [DNA-PKcs proficient] and SCID-st [DNA-PKcs deficient]) (B) were treated with 0.1 mM MMS and harvested at times indicated. Cell lysates were examined for JNK activity using GST-c-JUN as a substrate (upper panel). The levels of JNK1 protein were also measured by immunoblot analysis using an anti-JNK1 antibody (lower panel). For JNK assay, 200 μ g of cell lysates was mixed with an anti-JNK-1 polyclonal antibody and protein A-Sepharose beads and incubated with rocking for 3 h at 4°C. After the beads had been washed three times, the JNK immunocomplex was precipitated and used for an *in vitro* kinase assay in the presence of GST-cJun and [γ -³²P]-ATP. Phosphorylation of GST-JNK1 was analyzed by 10% SDS-PAGE and autoradiography.

factors after DNA damage (Park *et al.*, 1999), or it may participate in the damage signaling pathway, targeting protein kinases such as JNK.

In an effort to understand the role of DNA-PK in the damage signaling pathway, we examined the JNK activity of two human glioblastoma cells, M059K (DNA-PKcs proficient) and M059J (DNA-PKcs deficient), in response to genotoxic stress. Cells were treated with a DNA alkylating agent, MMS (0.1 mM), and examined for *in vitro* JNK activity using GST-cJun as a substrate (Fig. 1; upper panel) and JNK protein level (lower panel) following cell harvest at various time points (Fig. 1; panel A). Although JNK activity was induced by MMS treatment in both DNA-PKcs-proficient and DNA-PKcs-deficient cells, cells lacking DNA-PKcs showed remarkably greater activation than the DNA-PKcs-proficient cells (Fig. 1A). Amounts of JNK protein remained unchanged following genotoxic stress in both DNA-PKcs-deficient and DNA-PKcs-proficient cells, suggesting that the difference in JNK activity in the two cell lines is not at the level of gene expression.

Glioblastoma cells may carry multiple mutations; hence, the difference in JNK activity between DNA-PKcs-proficient

(M059K) and DNA-PKcs-deficient (M059J) cells could also be attributable to mutations of factor(s) other than DNA-PK. Therefore, we also examined two mouse cell lines, SCID (DNA-PKcs deficient) and its parental line, CB-17 (DNA-PKcs proficient) (Blunt *et al.*, 1995; Hendrickson *et al.*, 1991), to see whether DNA-PKcs is responsible for downregulation of JNK activity. Similar to the result with human glioblastoma cells (Fig. 1), SCID cells showed significantly higher JNK activation after MMS treatment than did CB-17 cells (Fig. 2). Together, our results suggest that DNA-PKcs (or its holoenzyme) may be involved in controlling JNK activation in response to genotoxic stress.

Tight regulation of JNK activity in DNA-PKcs-proficient cells occurs only in the presence of DNA damage

DNK-PK gets activated only in the presence of DNA damage, whereas JNK activation can be induced either by DNA damage or by other agents such as EGF (Adler *et al.*, 1992; Foltz *et al.*, 1998; Roulston *et al.*, 1998). We therefore exam-




FIG. 2. Activation by JNK of DNA-PKcs-proficient and DNA-PKcs-deficient cells following the treatment with MMC, Adriamycin, or EGF. Human and mouse cells (DNA-PKcs proficient and DNA-PKcs deficient) were treated with 0.1 mM MMS (A), Adriamycin 1.0 μ g/ml (B, D), or EGF 20 ng/ml (C) and harvested at times indicated. Cell extracts were examined for JNK activity using GST-c-Jun as a substrate (upper panel) and for JNK1 protein by Western blot analysis using an anti-JNK1 antibody (lower panel). The JNK immunocomplex assay was carried out as described in the legend to Figure 1.

defined whether tight regulation of JNK activity in DNA-PKcs-proficient cells occurs in a DNA damage-dependent manner. DNA-PKcs-proficient and DNA-PKcs-deficient cells were examined for JNK activation after treatment with DNA-damaging drugs or EGF. Similar to MMS treatment (see Fig. 1), DNA-PKcs-proficient cells (M059K) compared with the DNA-PKcs-deficient cells (M059J) showed significantly tighter regulation of JNK activity in response to treatment with the DNA-damaging drugs MMC or Adriamycin (Fig. 2A, B). In contrast, DNA-PKcs-proficient and DNA-PKcs-deficient cells showed similar JNK activation after EGF treatment (Fig. 2C), suggesting that damage-induced activation of DNA-PK may be necessary for regulation of JNK activity after treatment with DNA-damaging agents.

The JNK protein associates with DNA-PK in response to DNA damage

The results shown in Figures 1 and 2 strongly suggested that DNA damage-induced activation of DNA-PK is involved in

regulation of JNK activity. Hence, we examined whether DNA-PK physically interacts with JNK in response to DNA damage. DNA-PKcs-proficient cells were treated with 0.1 mM MMS and harvested at various times. Cell extracts were then incubated with anti-JNK1/2 polyclonal antibody for immunoprecipitation of JNK1/2 protein and its associated proteins. Western blot analysis indicated that cells treated with MMS showed a marked increase in interaction of both DNA-PKcs and Ku70 with JNK1/2 (Fig. 3A, B). Levels of DNA-PKcs and Ku70 proteins remained unchanged, suggesting that the interaction between JNK1/2 and DNA-PK is altered by DNA damage.

Phosphorylation of JNK by DNA-PK in vitro

DNA-PK phosphorylates a number of proteins involved in transcription, replication, and DNA repair after DNA damage (Jackson and Jeggo, 1995; Lees-Miller *et al.*, 1990). Downregulation of JNK activity in DNA-PKcs-proficient cells may be secondary to the phosphorylation of JNK protein by DNA-PK




FIG. 3. *In vivo* association of JNK protein(s) with DNA-PKcs or Ku70/80. Two human malignant glioblastoma cells (M059K [DNA-PKcs proficient] and M059J [DNA-PKcs deficient]) were treated with 0.1 mM MMS and harvested at the indicated time points. (A) Cell lysates were subjected to immunoprecipitation with anti-JNK1/2 (upper panel) and anti-DNA-PKcs (lower panel). Immunoprecipitates were analyzed by immunoblotting with anti-DNA-PKcs. Lysates from DNA-PKcs-proficient glioblastoma cells were precipitated without antibody and used as a negative control (lane 9). (B) Cell lysates were used for immunoprecipitation of JNK1/2 with an anti-JNK1/2 antibody, and the precipitates were analyzed for coimmunoprecipitation of Ku70 by immunoblot using an anti-Ku70 antibody (upper panel). As a control experiment (lower panel), an anti-Ku70 antibody was also used for immunoprecipitation and Western blotting. No antibody was included in lane 9 during immunoprecipitation.

in response to DNA damage. At least 10 JNK isoforms have been identified (Gupta *et al.*, 1996), and all isoforms contain the conserved TPY consensus phosphorylation site for SEK1/MEKK4 that is required for JNK activation (Moriguchi *et al.*, 1997). All members of JNK kinase family also contain at least one consensus phosphorylation site (-SQ-) for DNA-PK (Anderson and Miller, 1992; Lees-Miller *et al.*, 1990; Fig. 4A), suggesting that JNK may actually be a substrate for DNA-PK. We therefore examined whether DNA-PK phosphorylates JNK protein *in vitro*. Purified DNA-PK was incubated with a GST fusion form of either JNK1 or JNK2 and examined for

JNK phosphorylation. DNA-PK phosphorylated the GST fusion form of JNK1 (71 kDa) and JNK2 (77 kDa) in a DNA-dependent manner but did not phosphorylate GST alone (Fig. 4). This result suggests that the members of JNK family may be the actual substrates for DNA-PK *in vivo*.

We further examined whether DNA-PK is the major protein kinase responsible for JNK phosphorylation. A nonphosphorylated form of GST-JNK2 from *E. coli* was incubated with extracts of DNA-PKcs-proficient (M059K) or DNA-PKcs-deficient (M059J) cells as a source for protein kinase (Fig. 5). Phosphorylation of JNK2 (77-kDa GST-JNK2) was observed




FIG. 4. *In vitro* phosphorylation of JNK proteins by DNA-PK. (A) DNA-PK phosphorylation consensus sites among members of the JNK kinase family. TPY represents the phosphorylation sites for MEK4/SEK1 (or MKK7), and SQ represents the phosphorylation sites for DNA-PK. (B) *In vitro* phosphorylation of JNK 1 and JNK 2 by DNA-PK. Reaction mixtures contained either 0.05 μ g (lanes 1 and 4) or 0.1 μ g (lanes 2, 3, and 5–8) of purified DNA-PK holoenzyme. Where indicated, 1.0 μ g of activated calf thymus DNA and 1.0 μ g of substrate (GST-JNK1 [lanes 1–3], GST-JNK2 [lanes 4–6], or 1.0 μ g of GST [lane 8]) were included. * and ** = Positions of phosphorylated Ku80 and Ku70 proteins, respectively. Detailed procedure for protein kinase assay is described in Materials and Methods.




FIG. 5. *In vitro* phosphorylation of JNK2 in the presence of cell extracts from DNA-PKcs-proficient cells or DNA-PKcs-deficient cells. Cell extracts (5.0 μ g) from DNA-PKcs-deficient (MO59J; lanes 1–3) or DNA-PKcs-proficient (MO59K; lanes 4–6) cells were used as the source of protein kinase. Reaction mixtures contained 20 mM HEPES, pH 7.5; 1 mM EGTA, 1 mM EDTA, 1 mM DTT, 10 mM $MgCl_2$, 7 mM $MnCl_2$, 5 mM NaF, 1 mM Na_3VO_4 , and 50 μ M of [γ - ^{32}P]-ATP (sp. act. 30,000 cpm/pmol). Where indicated, 1.0 μ g of purified GST-JNK2 (77 kDa) or 1.0 μ g of activated calf thymus DNA was included. After incubation for 30 min at 30°C, reaction mixtures were subjected to 10% SDS-PAGE; and, after drying, the gel was exposed to X-ray film. The position of GST-JNK2 is indicated at left.

only when it was incubated with extracts of DNA-PKcs-proficient cells, not with extracts of DNA-PKcs-deficient cells (Fig. 5), suggesting that DNA-PK is a major protein kinase responsible for JNK phosphorylation. Because DNA-PK is activated in response to DNA damage, we also examined JNK phosphorylation with extracts of nonirradiated and irradiated DNA-PKcs-proficient cells. Extracts of irradiated cells showed higher levels of JNK phosphorylation *in vitro* than did those of non-irradiated cells, suggesting that JNK phosphorylation by DNA-PK is induced in response to DNA damage.

DISCUSSION

The stress-induced signaling pathway involves a cascade of protein phosphorylation, including JNK/SAPK and c-jun (AP-1) phosphorylation, that enhances the transcription of factors involved in cellular protection, such as DNA polymerase, thiol

synthesis, and DNA repair enzymes (Auer *et al.*, 1998; Bergelson *et al.*, 1994; Litz-Jackson *et al.*, 1992; Nishina *et al.*, 1997; Pinkus *et al.*, 1996; Potapova *et al.*, 1997). However, prolonged activation of JNK can lead to apoptosis (Auer *et al.*, 1998; Johnson *et al.*, 1995; Sanchez-Perez *et al.*, 1998; Widmann *et al.*, 1997), suggesting that the JNK/SAPK pathway may mediate either cell protection or programmed cell death after DNA damage. Hence, regulation of the JNK signaling pathway appears to be crucial for cell survival. The JNK/SAPK protein is activated by SEK1/MKK4 through the phosphorylation of the consensus site TPY that can also be dephosphorylated by MAP kinase phosphatases (MKPs). This implies a role for MKPs in blocking activation of the physiologic target of JNK/SAPK activation, the transcription factor c-Jun (Franklin *et al.*, 1998; Guo *et al.*, 1998b; Hirsh and Stork, 1997).

In this study, we found that DNA-PKcs-proficient cells in contrast to those lacking DNA-PKcs exhibited tight regulation of JNK/SAPK activation in response to genotoxic stress. This finding is in keeping with the previous observation that Ku70- or Ku80-deficient cells exhibit a significantly higher level of JNK than wildtype cells (Kim *et al.*, 1999). It is highly unlikely that the enhanced JNK activation in response to DNA damage in DNA-PKcs-deficient cells is reflective of the DNA repair deficiency of these cells, as there is no noticeable difference between DNA-PKcs-proficient and DNA-PKcs-deficient cells in nucleotide excision repair activity responsible for the removal of cisplatin or UV adducts (Park *et al.*, 1999). Regulation of JNK activity may occur *in vivo* through phosphorylation by DNA-PK, as both JNK1 and JNK2 were good substrates for DNA-PK *in vitro*. Our findings described here not only support a role for DNA-PK in the stress-induced signaling pathway but also imply a role for DNA-PK in protecting cells from apoptosis (Canman *et al.*, 1994; Lees-Miller *et al.*, 1995; Park *et al.*, 1999). The latter idea is further supported by previous observations that DNA-PK is the primary target for the CPP-32 (ICE-like) proteasome (Song *et al.*, 1996).

Activation of JNK occurs through phosphorylation by SEK1/MKK4 (or MKK7) in a dose (damage)-dependent manner (Derijard *et al.*, 1994; Devary *et al.*, 1991) and is critical for induction of cell-cycle arrest and DNA repair because high doses of genotoxic stress may lead to the apoptotic response. Stress-induced JNK activation is tightly regulated in DNA-PKcs-proficient cells compared with that observed in DNA-PKcs-deficient cells but only when cells were treated with low levels of irradiation or DNA-damaging agents (see Figs. 1 and 2). In contrast, significantly higher JNK activity was observed in both DNA-PKcs-proficient and DNA-PKcs-deficient cells treated with high doses of UV irradiation (50 J/m²), MMS (1 mM), or cisplatin (1 mM) (data not shown). This result suggests that regulation of JNK by DNA-PK may be related to cell protection in response to physiologic doses of irradiation/DNA damaging agents.

Phosphorylation of JNK by SEK-1/MKK4 leads to the activation of JNK kinase, which then activates transcription factors (c-Jun, ATF-2, and ELK-1) through phosphorylation. The SEK-1/MKK4 phosphorylates TPY sites of JNK and activates its kinase. Following JNK activation, JNK kinase activity needs to be inactivated in a timely manner. Downregulation of JNK may occur through dephosphorylation of TPY sites by MKPs (Guo *et al.*, 1998a, b; Sanchez-Perez *et al.*, 1998); or, alternatively, direct phosphorylation of JNK by DNA-PK may also down-

regulate JNK activity. DNA-PK may selectively interact with JNKs/SAPK that is already phosphorylated at threonine and tyrosine residues (TPY sites) by MEKK4/7, as the interaction between JNK and DNA-PK is induced by DNA damage. However, it is also possible that DNA-PK phosphorylates JNK regardless of its phosphorylation status, which in turn alters JNK activity. A detailed *in vivo* analysis would be necessary to clarify the role of DNA-PK in regulation of JNK activity.

ACKNOWLEDGMENTS

We thank Dr. Scott Boswell for providing us with the GST-c-Jun plasmid and stimulating discussion in early stage of the project. This work was supported by grants from the U.S. National Institutes of Health (IR01 CA92111), the U.S. Army (DAMD17-00-1-0295), and the Indiana University Center for Aging Research.

REFERENCES

- ADLER, V., FRANKLIN, C., and KRAFT, A. (1992). Phorbol ester stimulates the phosphorylation of c-Jun but not v-Jun: Regulation by the N-terminal delta domain. *Proc. Natl. Acad. Sci. USA* **89**, 5341-5345.
- ANDERSON, C.W., and CARTER, T.H. (1996). The DNA-activated protein kinase DNA-PK. *Curr. Top. Microbiol. Immunol.* **217**, 91-111.
- ANDERSON, C.W., and MILLER, S.P. (1992). The nuclear serine:threonine protein kinase DNA-PK. *Crit. Rev. Eukaryot. Gene Expr.* **2**, 283-314.
- ANGEL, P., and KARIN, M. (1991). The role of Jun, Fos and the AP-1 complex in cell proliferation and transformation. *Biochem. Biophys. Acta* **1072**, 129-157.
- ANGEL, P., HATTORI, K., SMEAL, T., and KARIN, M. (1988). The c-jun proto-oncogene is positively autoregulated by its product. *Jun/AP-1*. *Cell* **55**, 875-885.
- AUER, K.L., CONTESSA, J., BRENN-VERCA, S., PIROLA, L., RUSCONI, S., COOPER, G., ABO, A., WYMAN, M.P., DAVIS, R.J., BIRRE, M., and DENT, P. (1998). The RAS/Rac1/Cdc42/SEK/JNK/c-Jun cascade is a key pathway by which agonists stimulate DNA synthesis in primary cultures of rat hepatocytes. *Mol. Biol. Cell* **9**, 561-573.
- BERGELSON, S., PINKUS, R., and DANIEL, V. (1994). Induction of Ap-1 (Fos/Jun) by chemical agents mediates activation of glutathione S-transferase and quione reductase gene expression. *Oncogene* **9**, 565-571.
- BLUNT, T., FINNIE, N.J., TACCIOLI, G.E., SMITH, G.C., DEMENGEOT, J., GOTTLIEB, T.M., MIZUTA, R., VARGHESE, A.J., ALT, F.W., and JEGGO, P.A. (1995). Defective DNA-dependent protein kinase activity is linked to V(D)J recombination and DNA repair defects associated with the murine SCID mutation. *Cell* **80**, 813-823.
- CANMAN, C.E., WOLFF, A.C., CHEN, C.Y., FOMACE, A.J., JR., and KASTAN, M.B. (1994). The p53-dependent G1 checkpoint pathway and ataxia-telangiectasia. *Cancer Res.* **54**, 5054-5058.
- CASCIOLA-ROSEN, L.A., ANHALT, G.J., and ROSEN, A. (1995). DNA-dependent protein kinase is one of a subset of autoantigens specifically cleaved early during apoptosis. *J. Exp. Med.* **182**, 1625-1634.
- COSO, O.A., CHIARIELLO, M., YU, J.-C., TERAMOTO, H., CRESCO, P., XU, N., MIKI, T., and GUTKIND, J.S. (1995). The small GTP-binding proteins Rac-1 and Cdc42 regulate the activity of the JNK/SAPK signaling pathway. *Cell* **81**, 1137-1146.
- DAVIS, R.J. (1994). MAPKs: New JNK expands the group. *Trends Biochem. Sci.* **19**, 470-473.
- DERIJARD, B., HIBI, M., WU, I.-H., BARRETT, T., SU, B., DENO, T., KARIN, M., and DAVIS, R.J. (1994). JNK1: A protein kinase stimulated by UV light and Ha-Ras that binds and phosphorylates the c-Jun activation domain. *Cell* **76**, 1025-1037.
- DEVARY, Y., GOTTLIEB, R.A., LAU, L.F., and KARIN, M. (1991). Rapid and preferential activation of the c-jun gene during the mammalian UV response. *Mol. Cell. Biol.* **11**, 2804-2811.
- DUYSTER, J., BASKARAN, R., and WANG, J.Y. (1995). Src homology 2 domain as a specificity determinant in the c-Abl-mediated tyrosine phosphorylation of RNA polymerase II carboxyl-terminal repeated domain. *Proc. Natl. Acad. Sci. USA* **92**, 1555-1559.
- FARIS, M., KOKOT, N., LATINIS, K., KASIBHATLA, S., GREEN, D.R., KORETZKY, G.A., and NEL, A. (1998). The c-Jun N-terminal kinase cascade plays a role in stress-induced apoptosis in Jurkat cells by upregulating Fas ligand expression. *J. Immunol.* **160**, 134-144.
- FINNIE, N.J., GOTTLIEB, T.M., BLUNT, T., JEGGO, P.A., and JACKSON, S.P. (1995). DNA-dependent protein kinase activity is absent in xrs-6 cells: Implications for site-specific recombination and DNA double-strand break repair. *Proc. Natl. Acad. Sci. USA* **92**, 320-324.
- FOLTZ, I.N., GERL, R.E., WIELER, J.S., LUCKACH, M., SALMON, R.A., and SCHRADER, J.W. (1998). Human mitogen-activated protein kinase kinase 7 (MKK7) is a highly conserved c-Jun N-terminal kinase/stress-activated protein kinase (JNK/SAPK) activated by environmental stresses and physiological stimuli. *J. Biol. Chem.* **273**, 9344-9351.
- FRANKLIN, C.C., SRIKANTH, S., and KRAFT, A.S. (1998). Conditional expression of mitogen-activated protein kinase phosphatase-1, MKP-1, is cytoprotective against UV-induced apoptosis. *Proc. Natl. Acad. Sci. USA* **95**, 3014-3019.
- GUO, Y.L., BAYSAL, K., KANG, B., YANG, L.J., and WILLIAMSON, J.R. (1998a). Correlation between sustained c-Jun N-terminal protein kinase activation and apoptosis induced by tumor necrosis factor-alpha in rat mesangial cells. *J. Biol. Chem.* **273**, 4027-4034.
- GUO, Y.L., KANG, B., and WILLIAMSON, J.R. (1998b). Inhibition of the expression of mitogen-activated protein phosphatase-1 potentiates apoptosis induced by tumor necrosis factor-alpha in rat mesangial cells. *J. Biol. Chem.* **273**, 10362-10366.
- GUPTA, S., BARRETT, T., WHITMARSH, A.J., CAVANAGH, J., SLUSS, H.K., DERIJARD, B., and DAVIS, R.J. (1996). Selective interaction of JNK protein kinase isoforms with transcription factors. *EMBO J.* **15**, 2760-2770.
- GUPTA, S., CAMPBELL, D., DERIJARD, B., and DAVIS, R.J. (1995). Transcription factor ATF-2 regulation by the JNK signal transduction pathway. *Science* **267**, 389-393.
- HENDRICKSON, E.A., QIN, X.Q., BUMP, E.A., SCHATZ, D.G., OETTINGER, M., and WEAVER, O.T. (1991). A link between double-strand break-related repair and V(D)J recombination: The SCID mutation. *Proc. Natl. Acad. Sci. USA* **88**, 4061-4065.
- HIRSH, D.D., and STORK, P.J. (1997). Mitogen-activated protein kinase phosphatase inactivates stress-activated protein kinase pathways *in vivo*. *J. Biol. Chem.* **272**, 4568-4575.
- ICHIJO, H., NISHIDA, E., IRIE, K., DIJKE, P., SAITOH, M., MORIGUCHI, T., TAKAGI, M., MATSUMOTO, K., MIYAZONO, K., and GOTOH, Y. (1997). Induction of the apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways. *Science* **275**, 90-94.
- JACKSON, S.P. (1997). DNA-dependent protein kinase. *Int. J. Biochem. Cell. Biol.* **29**, 935-938.
- JACKSON, S.P., and JEGGO, P.A. (1995). DNA double-strand break repair and V(D)J recombination: Involvement of DNA-PK. *Trends Biochem. Sci.* **20**, 412-415.
- JOHNSON, N.L., GARDNER, A.M., DIENER, K.M., LANGE-CARTER, C.A., GLEAVY, J., JARPE, M.B., MINDEN, A., KARIN, M., ZON, L.I., and JOHNSON, G.L. (1996). Signal transduction pathways regulated by mitogen activated/extracellular response kinase kinase induced cell death. *J. Biol. Chem.* **271**, 3229-3237.

- KIM, S.H., KIM, D., HAN, J.S., JEONG, C.S., CHUNG, B.S., KANG, C.D., and LI, G.D. (1999). Ku autoantigen affects the susceptibility to anticancer drugs. *Cancer Res.* **59**, 4012–4017.
- KYRIAKIS, J.M., BANERJEE, P., NIKOLAKAKI, E., DAI, T., RUBIE, E.A., AHMAD, M.F., AVRUCH, J., and WOODGETT, J.R. (1994). The stress-activated protein kinase subfamily of c-jun kinase. *Nature* **369**, 156–160.
- LAEMMLI, U.K. (1970). Cleavage of structural proteins during assembly of the head of bacteriophage T4. *Nature* **227**, 680–685.
- LEES-MILLER, S.P., GODBOUT, R., CHAN, D.W., WEINFELD, M., DAY, R.S. III, BARRON, G.M., and ALLALUNIS-TURNER, J. (1995). Absence of p350 subunit of DNA-activated protein kinase from a radiosensitive human cell line. *Science* **267**, 1183–1185.
- LEES-MILLER, S.P., CHEN, Y.R., and ANDERSON, C.W. (1990). Human cells contain a DNA-activated protein kinase that phosphorylates simian virus 40 T antigen, mouse p53, and the human Ku autoantigen. *Mol. Cell. Biol.* **10**, 6472–6481.
- LIEBER, M.R., GRAWUNDER, U., WU, X., and YANEVA, M. (1997). Tying loose ends: Roles of Ku and DNA-dependent protein kinase in the repair of double-strand breaks. *Curr. Opin. Genet. Dev.* **7**, 99–104.
- LIU, Z.G., BASKARAN, R., LEA-CHOU, E., WOOD, L.D., CHEN, Y., KARIN, M., and WANG, J.Y. (1996). Three distinct signalling responses by murine fibroblasts to genotoxic stress. *Nature* **384**, 273–276.
- LIVINGSTONE, C., PATEL, G., and JONES, N. (1995). ATF-2 contains a phosphorylation-dependent transcriptional activation domain. *EMBO J.* **14**, 1785–1797.
- LITZ-JACKSON, S., MILLER, A.H., BURGESS, G.S., and BOSWELL, H.S. (1992). Dissociation of nuclear events on p21 Ras transformation of FDC-P1 myeloid cells: C-jun/AP-1 expression versus c-myc transcription. *Blood* **79**, 2404–2414.
- MINDEN, A., LIN, A., CLARET, F.X., ABO, A., and KARIN, M. (1995). Selective activation of the JNK signaling cascade and c-JUN transcriptional activity by the small GTPases Rac and Cdc42Hs. *Cell* **81**, 1147–1157.
- MORIGUCHI, T., TOYOSHIMA, F., MASUYAMA, M., HANAFUSA, H., GOTOH, Y., and NISHIDA, E. (1997). A novel SAPK/JNK kinase, MKK7, stimulated by TNF α and cellular stresses. *EMBO J.* **16**, 7045–7053.
- MULLER, C., CALSOU, P., FRIT, P., CAYROL, C., CARTER, T., and SALLES, B. (1998). UV sensitivity and impaired nucleotide excision repair in DNA-dependent protein kinase mutant cells. *Nucleic Acids Res.* **26**, 1382–1389.
- NISHINA, H., FISCHER, K.D., RADVANYI, L., SHAHINIAN, A., HAKEM, R., RUBIE, E.A., BERNSTEIN, A., MAK, T.W., WOODGETT, J.R., and PENNINGER, J.M. (1997). Stress-signaling kinase Sek1 protects thymocytes from apoptosis mediated by CD95 and CD3. *Nature* **385**, 350–353.
- PARK, J.S., PARK, S.J., PENG, X., WANG, M., YU, M.A., and LEE, S.H. (1999). Involvement of DNA-dependent protein kinase in UV-induced replication arrest. *J. Biol. Chem.* **274**, 32520–32527.
- PETERSON, S.R., KURIMASA, A., OSHIMURA, M., DYNAN, W.S., BRADBURY, E.M., and CHEN, D.J. (1995). Loss of the catalytic subunit of the DNA-dependent protein kinase in DNA double-strand-break-repair mutant. *Proc. Natl. Acad. Sci. USA* **92**, 3171–3174.
- PETERSON, S.R., STACKHOUSE, M., WALTMAN, M.J., CHEN, F., SATO, K., and CHEN, D.J. (1997). Characterization of two DNA double-stranded break repair-deficient cell lines that express inactive DNA-dependent protein kinase catalytic subunits. *J. Biol. Chem.* **272**, 10227–10231.
- PINKUS, R., WEINER, L.M., and DANIEL, V. (1996). Role of oxidants and antioxidants in the induction of AP-1, NF-kappaB and glutathione S-transferase gene expression. *J. Biol. Chem.* **271**, 13422–13429.
- POTAPOVA, O., HAGHIGHI, A., BOST, F., LIU, C., BIRRER, M.J., GJERSET, R., and MERCOLA, D. (1997). The Jun kinase/stress-activated protein kinase pathway functions to regulate DNA repair and inhibition of the pathway sensitizes tumor cells to cisplatin. *J. Biol. Chem.* **272**, 14041–14044.
- PULVERER, B., KYRIAKIS, J.M., AVRUCH, J., NIKOLAKAKIE, E., and WOODGETT, J.R. (1991). Phosphorylation of c-jun mediated by MAP kinases. *Nature* **353**, 670–674.
- RAINGEAUD, J., GUPTA, S., ROGER, J.S., DICKENS, M., HAN, J., ULEVITCH, R.J., and DAVIS, R.J. (1995). Pro-inflammatory cytokines and environmental stress cause p38 mitogen-activated protein kinase activation by dual phosphorylation on tyrosine and threonine. *J. Biol. Chem.* **270**, 7420–7426.
- RAMSDEN, D.A., and GELLERT, M. (1998). Ku protein stimulates DNA end joining by mammalian DNA ligases: A direct role for Ku in repair of DNA double-strand breaks. *EMBO J.* **17**, 609–614.
- RATHMELL, W.K., KAUFMANN, W.K., HURT, J.C., BYRD, L.L., and CHU, G. (1997). DNA-dependent protein kinase is not required for accumulation of p53 or cell cycle arrest after DNA damage. *Cancer Res.* **57**, 68–74.
- ROULSTON, A., REINHARD, C., AMIRI, P., and WILLIAMS, L.T. (1998). Early activation of c-Jun N-terminal kinase and p38 kinase regulate cell survival in response to tumor necrosis factor alpha. *J. Biol. Chem.* **273**, 10232–10239.
- SANCHEZ, I., HUGHES, R.T., MAYER, B.J., YEE, K., WOODGETT, J.R., AVRUCH, J., KYRIAKIS, J.M., and ZON, L.I. (1994). Role of SAPK/ERK kinase-1 in the stress-activated pathway regulating transcription factor c-Jun. *Nature* **372**, 794–798.
- SANCHEZ-PEREZ, I., MURGUIA, J.R., and PERONA, R. (1998). Cisplatin induces a persistent activation of JNK that is related to cell death. *Oncogene* **16**, 533–540.
- SAVITSKY, K., BAR-SHIRA, A., GILAD, S., ROTMAN, G., ZIV, Y., VANAGAITE, L., TAGLE, D.A., SMITH, S., UZIEL, T., SFEZ, S., ASHKENAZI, M., ET AL. (1995). A single ataxia telangiectasia gene with a product similar to PI-3 kinase. *Science* **268**, 1749–1753.
- SCHREIBER, M., BAUMANN, B., COTTON, M., ANGEL, P., and WAGNER, E.F. (1995). Fos is an essential component of the mammalian UV response. *EMBO J.* **14**, 5338–5349.
- SMEAL, T., BINETRUY, B., MERCOLA, D.A., BIRRER, M., and KARIN, M. (1991). Oncogenic and transcriptional cooperation with Ha-Ras requires phosphorylation of c-Jun on serines 63 and 73. *Nature* **354**, 494–496.
- SONG, Q., LEES-MILLER, S.P., KUMAR, S., ZHANG, N., CHAN, D.W., SMITH, G.C.M., JACKSON, S.P., ALNEMRI, E.S., LITWACK, G., KHANNA, K.K., and LAVIN, M.F. (1996). DNA-dependent protein kinase catalytic subunit: A target for an ICE-like protease in apoptosis. *EMBO J.* **15**, 3238–3246.
- WIDMANN, C., JOHNSON, N.L., GARDNER, A.M., SMITH, R.J., and JOHNSON, G.L. (1997). Potentiation of apoptosis by low dose stress stimuli in cells expressing activated MEK kinase 1. *Oncogene* **15**, 2439–2447.

Address reprint requests to:

Dr. Suk-Hee Lee

Department of Biochemistry and Molecular Biology
Indiana University School of Medicine
Indianapolis, IN 46202

E-mail: slee@iupui.edu

Received for publication June 15, 2001; received in revised form August 2, 2001; accepted August 3, 2001.

Minireview

DNA-dependent Protein Kinase Complex: a Multifunctional Protein in DNA Repair and Damage Checkpoint

Suk-Hee Lee* and Chung-Hui Kim

Department of Biochemistry and Molecular Biology, and the Walther Oncology Center, Indiana University School of Medicine, and the Walther Cancer Institute, Indianapolis, Indiana 46202, USA.

(Received February 26, 2002; Accepted March 5, 2002)

DNA-dependent protein kinase (DNA-PK) is a nuclear serine/threonine protein kinase that is activated upon DNA damage generated by ionizing radiation or UV-irradiation. It is a three-protein complex consisting of a 470-kDa catalytic subunit (DNA-PKcs) and the regulatory DNA binding subunits, Ku heterodimer (Ku70 and Ku80). Mouse and human cells deficient in DNA-PKcs are hypersensitive to ionizing radiation and defective in V(D)J recombination, suggesting a role for the kinase in double-strand break repair and recombination. The Ku heterodimer binds to double-strand DNA breaks produced by either DNA damage or recombination, protects DNA ends from degradation, orients DNA ends for re-ligation, and recruits its catalytic subunit and additional factors necessary for successful end-joining. DNA-PK is also involved in an early stage of damage-induced cell cycle arrest, however, it remains unclear how the enzyme senses DNA damage and transmits signals to downstream gene(s) and proteins.

Keywords: Damage Checkpoint; DNA Damage; DNA-PK; DSB Repair; Ku Complex; Nucleotide Excision Repair.

Introduction

DNA damage is constantly generated by radiation and chemotherapy drugs, which must be repaired to prevent genomic alterations that could otherwise contribute to cancer progression and/or generation of cancer. For this reason, cells invoke various mechanisms to repair DNA damage while operating damage checkpoint pathways that

are responsible for sensing DNA damage and causing arrest of cell cycle progression until the damage is repaired. It is not clear, however, which genes are involved in this pathway or how DNA damage induces cell cycle arrest while permitting DNA repair. The DNA-dependent protein kinase (DNA-PK) is a member of the phosphatidylinositol-3 kinase (PI-3 kinase) family and shares amino acid sequence homology in its carboxy-terminal kinase domain with other family members, including ataxia telangiectasia mutated gene (ATM), ATM-related (ATR), and p110 PI-3 kinase (Hartley *et al.*, 1995; Poltoratsky *et al.*, 1995). All members of the PI 3-kinase family are activated by stress; PI-3 kinase is regulated by heat shock and DNA-damage, and ATM, ATR, and DNA-PK are activated by DNA damage (Gottlieb and Jackson, 1993; Lin *et al.*, 1997; Rotman and Shiloh, 1997; Yuan *et al.*, 1997). The DNA-PK is a unique damage-response element not only because it functions as a DNA damage sensor, but also it is a key player in various DNA repair pathways. In this review, we focus on recent advances on the DNA-PK in DNA repair and damage checkpoint pathways.

The DNA-PK complex and its regulation

Association of the DNA end-binding Ku70/Ku80 heterodimer with the 470 kDa serine/threonine kinase catalytic subunit forms the DNA-PK holoenzyme that is essential for double-strand break (DSB) repair by non-homologous recombination in mammalian cells. The DNA-PK catalytic subunit (DNA-PKcs) is a very abun-

Abbreviations: ATM, ataxia telangiectasia mutated; DNA-PK, DNA-dependent protein kinase; DSB, double-strand break; JNK, c-Jun N-terminus protein kinase; NHEJ, non-homologous end joining; PCNA, proliferating cell nuclear antigen; PI-3 kinase, phosphatidylinositol-3 kinase; RPA, replication protein A.

* To whom correspondence should be addressed.

Tel: 1-317-278-3464; Fax: 1-317-274-4686

E-mail: slee@iupui.edu

dant nuclear protein in human cells and a very large polypeptide of 4127 amino acids giving a mass of 470 kDa (Gottlieb and Jackson, 1993). Its imaging structure indicates that DNA-PKcs has an open, pseudo 2-fold symmetric structure with a gap separating a crown-shaped top from a rounded base (Chiu *et al.*, 1998). With the exception of the 400 amino acids of the C-terminus, the amino acid sequence of DNA-PKcs shows little amino acid similarity to any protein in the databases. The C-terminus of DNA-PKcs are similar to proteins of the phosphatidylinositol (PI) 3-kinase superfamily (Hartley *et al.*, 1995; Poltoratsky *et al.*, 1995), both of which possess the DXXXXN and the DXG amino acid motifs that are found in protein kinase and are required for catalysis (Fig. 1). Despite the amino acid similarity with lipid kinases, DNA-PKcs acts as protein kinase, and no lipid kinase activity has been detected with purified DNA-PKcs or DNA-PK complex *in vitro* (Hartley *et al.*, 1995).

Ku complex, consisting of Ku70 and Ku80 subunits, serves as regulatory subunits for DNA-PK kinase activity essential for dsb rejoining (Kurimasa *et al.*, 1999). It is a very abundant nuclear protein, estimated at 4×10^5 molecules per cell and is present in vertebrates, insects, yeast, and worms (Mimori *et al.*, 1986). The Ku complex binds to DNA ends, nicks, gaps, and regions of transition between single and double-stranded structure (Blier *et al.*, 1993; Falzon *et al.*, 1993). *In vitro* photocross-linking of Ku protein bound to single DNA-strand ends showed that the preferred orientation is when Ku70 contacts the major groove and Ku80 contacts the minor groove (Yoo *et al.*, 1999). The crystal structure of the human Ku heterodimer revealed that Ku70 and Ku80 share a common topology and form a dyad-symmetrical molecule with a preformed ring that encircles duplex DNA (Walker *et al.*, 2001). The binding site can cradle two full turns of DNA while encircling only the central 3–4 base pairs. Surprisingly, Ku makes minimal contacts with DNA bases and with the sugar-phosphate backbone, however its ring structure nicely fits into major and minor groove contours in order to support its interaction with broken DNA ends and to recruit DNA-PKcs to the DNA (Walker *et al.*, 2001).

Regulation of DNA-PK activity While the activation of DNA-PK kinase requires DNA ends, its activity can be regulated by several mechanisms. Heterodimerization is not only essential for Ku-dependent DNA repair *in vivo*, but may be also a way to regulate DNA-PK activity. A central region within Ku80 is required for heterodimerization with Ku70. Evidences suggest that heterodimerization is required for the stabilization of Ku70 and Ku80. Each Ku subunit can translocate to the nucleus not only through its own nuclear localization signal but also through heterodimerization with each other, suggesting that the heterodimerization of these Ku subunits is important for their nuclear entry (Gell and Jackson, 1999; Ko-

ike *et al.*, 1999; 2001; Osipovich *et al.*, 1999). *In vitro*, work showed that DNA-PK kinase activity involving DNA-PKcs controls Ku entry into DNA which may regulate DNA transactions including transcription in the vicinity of double-strand breaks (Frit *et al.*, 2000).

Phosphorylation of both DNA-PKcs and Ku subunits (Ku70 and Ku80) was observed *in vivo* and *in vitro*. Phosphorylation of either DNA-PKcs or Ku subunit resulted in inactivation of the serine/threonine protein kinase activity, suggesting that protein phosphorylation is a way to regulate DNA-PK protein kinase activity (Chan and Lees-Miller, 1996; Douglas *et al.*, 2001) and that the protein phosphatase responsible for reactivation *in vivo* is a PP2A-like enzyme (Douglas *et al.*, 2001). Post-translational regulation of DNA-PK kinase activity may also occur through modulation of Ku70/80 DNA-binding activity observed in human neoplastic tissues, suggesting a possible role for heterodimer activity in tumor development (Pucci *et al.*, 2001). Also, irradiation induces the expression of Ku70, but not Ku80, in both p53 and ATM-dependent manner, providing cells with means of assuring proper DNA repair and/or cellular response to DNA damage (Brown *et al.*, 2000).

Role in DSB repair

DSBs are deleterious and potentially lethal form of DNA damage because they physically disrupt the continuity of the genome. Accordingly, eukaryotic cells have evolved two distinct DSB repair pathways. One pathway is homologous recombination repair (HRR) that predominantly occurred in lower eukaryotes such as *S. cerevisiae* and is mediated by Rad51 and Rad52 family proteins. The other pathway is non-homologous end joining (NHEJ) that is a major DSB repair pathway in mammals, at least in G0 or G1-phase of the cell cycle (Lee *et al.*, 1997). The NHEJ machinery consists of the DNA-PK complex and a complex of XRCC4 and DNA ligase IV. Although both HRR and NHEJ require the Rad50/Mre11/NBS1 complex in yeast and presumably also in mammalian cells (Critchlow and Jackson, 1998), it is unclear how DSB repair proteins are coordinated at the damage site. The current model of DNA-PK complex activation by dsDNA break is based on the tenet that without DNA, DNA-PKcs is inactive and incapable of binding Ku complex (Suwa *et al.*, 1994). When a DSB is introduced, Ku binds to the DNA because of its high affinity for DNA ends. The binding of Ku elicits conformational changes that allow it to bind DNA-PKcs. Thus the presumed roles of Ku complex are to first bind to DNA and then recruit DNA-PKcs to the DNA. Ku may also serve as an alignment factor that not only increases NHEJ efficiency but also the accuracy. Furthermore, a secondary NHEJ activity is present in the absence of Ku, which is error-prone and possibly assisted by base

pairing interaction (Feldmann *et al.*, 2000). Upon the assembly of DNA-PK holoenzyme on DNA breaks, this DNA repair complex activates its serine/threonine protein kinase activity and phosphorylates target substrates that colocalize with it on the ends of broken DNA.

The assembly and activation of the DNA-PK complex at a DNA strand break is central to NHEJ, although the *in vivo* study suggests that Ku complex but not DNA-PKcs may be essential for DSB repair (Gu *et al.*, 2000). The Ku complex and DNA-PKcs bind to DNA ends and are capable of physically tethering two ends of DNA molecules (Cary *et al.*, 1997; Yaneva *et al.*, 1998). Low resolution structures of DNA-PKcs revealed that DNA-PKcs has a cage-like structure with channels and cavities within the interior of the structure (Chiu *et al.*, 1998; Leuther *et al.*, 1999). Nonetheless, it is still unclear how the Ku/DNA complex activates the kinase activity of DNA-PKcs during DSB repair. One hypothesis is that DNA-PKcs undergoes a conformational change upon association with the Ku/DNA complex and this conformational change accounts for the activation of kinase activity. The kinase activity associated with DNA-PK is needed for DSB repair *in vivo*, since expression of a kinase-dead form of DNA-PKcs fails to complement the radiosensitive phenotype of a mammalian cell line that lacks the DNA-PKcs protein (Kurimasa *et al.*, 1999). However, the physiological targets of DNA-PK *in vivo* are unknown. The DNA-PK complex can physically tether two ends of a DSB in close proximity *in vitro* suggesting the hypothesis that the DNA-PK complex acts as a scaffold to assemble the NHEJ pathway proteins at a DSB (Cary *et al.*, 1997). In support of this, Ku recruits XRCC4/LigaseIV to DNA termini (McElhinny *et al.*, 2000), and Ku binds to MRE11/Rad50/NBS complex (Goeddecke *et al.*, 1999). In *S. cerevisiae*, NHEJ of plasmid DSBs requires Ku, Xrcc4, and DNA ligase IV, as well as Mre11, Rad50, Xrs2, and DNA damage checkpoint proteins. In *S. pombe*, however, DSB repair is dependent on Ku complex and DNA ligase IV, but does not require Rad32, Rad50 (the *S. pombe* homologues of Mre11 and Rad50, respectively) and checkpoint proteins for NHEJ, suggesting that NHEJ pathway in eukaryotes may occur in more than one mechanism (Manolis *et al.*, 2001).

Other repair pathways DNA-PKcs-deficient cells exhibited sensitivity not only to ionizing irradiation but also in response to chemotherapy drugs and were associated with lower nucleotide excision repair activity, suggesting that DNA-PK may be involved in the repair of various types of DNA damage (Britten *et al.*, 1999; Frit *et al.*, 1999). Also, studies with drug-resistant or drug-sensitive cancer cells suggested that higher levels of DNA-PK expression lead to a drug-resistant cells, whereas the low DNA-PK activity was associated with cells with drug-sensitive phenotype (Shen *et al.*, 1997) and was linked to

cell death via the accumulation of damaged DNA. More than 20-fold overexpression of the catalytic subunit of DNA-PK (DNA-PKcs) was observed with a drug-resistant cancer cells (Muller and Salles, 1997; Shen *et al.*, 1997). The Ku complex also interacts with a key base excision repair enzyme, AP-endonuclease (Chung *et al.*, 1996), suggesting a possible involvement of DNA-PK complex in various DNA repair pathways. It is not clear however whether the Ku or DNA-PK complex has a direct role in these repair pathways.

Role in damage checkpoint

Upon DNA damage, DNA-PK phosphorylates a number of proteins, including p53, RNA-polymerase II, RPA, topoisomerases, hsp90, SV-40 large T antigen, and many transcription factors such as c-Jun, c-Fos, oct-1, sp-1, c-Myc, TFIID, and many more. Among them, p53 is one of the legitimate targets for damage-induced cell cycle arrest. The p53 deficiency prolongs the survival of DNA-PKcs-deficient cells harboring DNA damage by allowing the accumulation of aneuploid cells, suggesting that the DNA-PKcs mutation links to a p53-mediated DNA damage checkpoint (Guidos *et al.*, 1996). The *in vivo* role of DNA-PK in the transduction of the DNA damage signal to p53 remains unclear (Jimenez *et al.*, 1999; Woo *et al.*, 1998), however, DNA-PK along with ATM and ATR collectively detect DSBs and transmit this signal to p53 by phosphorylation. This phosphorylation dissociates p53 from its negative regulator, mdm2, which allows p53 undergo further modification and activate transcription of the genes responsible for cell cycle arrest such as p21waf1/cip1. Both control and DNA-PKcs-null murine embryonic fibroblast cells showed phosphorylation and accumulation of p53 in response to irradiation. However, the rise in p21cip1/waf1 and mdm2 was found to be delayed and attenuated in DNA-PK-deficient cells, which correlated in time with delayed onset of G1/S arrest by flow cytometric analysis, suggesting that loss of DNA-PK activity appears to attenuate the kinetics of p53 to activate downstream genes, implying that DNA-PK plays a role in post-translational modification of p53, without affecting the increase in levels of p53 in response to DNA damage (Kachnic *et al.*, 1999).

Another potentially important role for DNA-PK in damage signaling stems from a functional interaction with c-Abl in response to DNA damage (Kharbanda *et al.*, 1997; Kumaravel *et al.*, 1998). Ionizing radiation stimulates the binding of c-Abl to DNA-PK and induces the association of c-Abl with Ku antigen (Jin *et al.*, 1997a; Kharbanda *et al.*, 1997). DNA-PK phosphorylates c-Abl *in vitro* and activates its tyrosine kinase activity. Moreover, via a potential feedback mechanism, c-Abl phosphorylates DNA-PK, which can then no longer form a complex with DNA

(Kharbanda *et al.*, 1997). Furthermore Ku associates with c-Abl and p21cip1/waf1 after irradiation (Kumaravel *et al.*, 1998), and interestingly, the p21cip1/waf1 only transiently associates with Ku complex at the low dose of irradiation, suggesting a possible role for DNA-PK in the damage-checkpoint pathway to control downstream DNA metabolism. The DNA-PKcs and Ku70/Ku80 heterodimer also can associate with another stress-response protein, the c-Jun N-terminus protein kinase (JNK), and the interaction was significantly stimulated following DNA damage. DNA-PKcs-proficient cells compared to those lacking DNA-PKcs exhibited a tight regulation of JNK activation in response to genotoxic stress. Based on the observation that the DNA-PK phosphorylates JNK *in vitro*, it is possible that DNA-PK is involved in regulation of JNK signaling pathway in response to stress/DNA damage (Park *et al.*, 2001).

G1/S-phase arrest Mounting evidences point to the role of DNA-PK in damage-induced S-phase arrest. Both *in vivo* and *in vitro* studies suggested that DNA-PK kinase activity is necessary for damage-induced replication arrest and its reversal (Park *et al.*, 1999; Wang *et al.*, 1999; 2001). Damage-induced S-phase arrest was reversed *in vitro* by the addition of a DNA-PK inhibitor or by immunodepletion of DNA-PKcs, suggesting that DNA-PKcs may be directly involved in damage-induced S-phase arrest through modulation of replication protein(s) (Park *et al.*, 1999; Wang *et al.*, 1999). Replication protein A (RPA) is a trimeric, multifunctional protein complex involved in DNA replication, DNA repair, and recombination and is a likely target for DNA-PK in damage-induced S-phase arrest because phosphorylation of the RPA2 subunit is observed after exposure of cells to ionizing radiation. DNA-PKcs interacted directly with RPA, and causes RPA2 phosphorylation in response to DNA damage. Phosphorylated RPA has a higher affinity for nuclear structures than unphosphorylated RPA suggesting the modified protein may be involved in the regulation of DNA replication after DNA damage or in DNA repair. RPA phosphorylation is delayed in cells deficient in ATM or DNA-PK, suggesting that DNA-PK and ATM may cooperate to phosphorylate RPA after DNA damage to redirect the functions of the protein from DNA replication to DNA repair (Wang *et al.*, 2001). DNA-PK-mediated RPA phosphorylation requires ongoing DNA replication but is prevented by the cell-cycle checkpoint abrogator, suggesting that the replication fork-associated RPA, when encounters DNA double-strand end-associated DNA-PK, leads to RPA phosphorylation which may signal the presence of DNA damage to an S-phase checkpoint mechanism (Shao *et al.*, 1999). Phosphorylation of RPA in yeast is also dependent on the central checkpoint regulator Mec1p. Because RPA's ssDNA binding activity of the RPA heterotrimer and is required for DNA replication,

repair and recombination, it is possible that phosphorylation of this subunit is directly involved in modulating RPA activity during the checkpoint response (Brush and Kelly, 2000).

DNA damage induces the change in other repair/replication proteins that also affects DNA repair and/or G1/S-phase cell cycle arrest. The proliferating cell nuclear antigen (PCNA) forms specific foci in response to DNA damage and an increased interaction of PCNA with the Ku heterodimer after DNA damage, suggesting a role for PCNA in the NHEJ repair pathway of DNA strand breaks (Balajee and Geard, 2001). Similarly, Ku80-deficient cells showed a significant decrease in the amount of PCNA associated with chromatin following ionizing radiation, suggesting that the Ku-complex by binding at the ends of damaged DNA may also protect the key replication factors from dissociating from chromatin (Park *et al.*, submitted). The Ku (or DNA-PK complex) also associates with ors8 and ors12 and this association was significantly enhanced in cells synchronized at the G1/S border. In addition, *in vitro* DNA replication activity with the use of extracts from Ku80-deficient cells was significantly lower compared to that with wild-type cell extracts, suggesting a novel function for Ku that may act at the initiation step of DNA replication and dissociates after origin firing (Novac *et al.*, 2001).

G2 arrest In addition to its role in damage-induced G1/S-phase arrest, DNA-PK may also have a distinct role in G2 checkpoint traversal in response to DNA damage since its activity is required for exit from a DNA damage-induced G2 checkpoint arrest (Lee *et al.*, 1997). This finding is supported by the observation that the activity of the DNA-PK complex is regulated in a cell cycle-dependent manner peaking its activity at the G1/S boundary and at G2, which is suggestive of the involvement of DNA-PK at multiple checkpoints (Lee *et al.*, 1997). Ku80-deficient cells compared to wild-type cells were highly cytotoxic to various stress agents, which led to their long-term accumulation in the G2 phase. This differential response was not due to differences in DNA repair, since DNA damage was repaired with comparable efficiency in both wild-type and Ku80-deficient cells, but was associated with differences in the expression of important cell cycle regulatory genes, supporting the notion that Ku80-mediated cytoprotection and G2-progression are not only dependent on DNA repair but may also reflect its influence on other cellular processes (Arrington *et al.*, 2000).

Other DNA metabolism

Besides their role in DNA repair, Ku complex also binds to chromosome ends (telomeres) protecting them from




Fig. 2. Involvement of DNA-PK in damage-induced cell cycle arrest and DNA repair.

Concluding Remarks

JNK activation and its signaling pathway are linked to the induction of cell cycle arrest and DNA repair. Prolonged activation of JNK, however, can lead to apoptosis (Fig. 2; Auer *et al.*, 1998; Johnson *et al.*, 1996; Sanchez-perez *et al.*, 1998; Widmann *et al.*, 1997), suggesting that the JNK pathway may mediate either cell protection or programmed cell death following DNA damage. These lines of evidences implicate a role for DNA-PK in protecting cells from apoptosis (Canman *et al.*, 1994; Lees-Miller *et al.*, 1995; Park *et al.*, 1999). In this regard, it is obvious why DNA-PK is the primary target for the CPP-32 proteasome (Song *et al.*, 1996).

Acknowledgments This work was supported by grants from the National Institute of Health (CA92111) and the U.S. Army (DAMD17-00-1-0295 and DOD-OC000113).

References

- Anderson, C. W. (1993) DNA damage and the DNA-activated protein kinase. *Trends Biochem. Sci.* **18**, 433–437.
- Arrington, E. D., Caldwell, M. C., Kumaravel, T. S., Lohani, A., Joshi, A., Evans, M. K., Chen, H. T., Nussenzweig, A., Holbrook, N. J., and Gorospe, M. (2000) Enhanced sensitivity and long-term G2 arrest in hydrogen peroxide-treated Ku80-null cells are unrelated to DNA repair defects. *Free Radical Biol. Med.* **29**, 1166–1176.
- Auer, K. I., Contessa, J., Brenz-Verca, S., Pirola, L., Rusconi, S., Cooper, G., Abo, A., Wymann, M. P., Davis, R. J., Birrer, M., and Dent, P. (1998) The RAS/Rac1/Cdc42/SEK/JNK/c-Jun cascade is a key pathway by which agonists stimulate DNA synthesis in primary cultures of rat hepatocytes. *Mol. Biol. Cell.* **9**, 561–573.
- Bailey, S. M., Meyne, J., Chen, D. J., Kurimasa, A., Li, G. C., Lehnert, B. E., and Goodwin, E. H. (1999) DNA double-strand break repair proteins are required to cap the ends of mammalian chromosomes. *Proc. Natl. Acad. Sci. USA* **96**, 14899–14904.
- Balajee, A. S. and Geard, C. R. (2001) Chromatin-bound PCNA complex formation triggered by DNA damage occurs independent of the ATM gene product in human cells. *Nucleic Acids Res.* **29**, 1341–1351.
- Blier, P. R., Griffith, A. J., Craft, J., and Hardin, J. A. (1993) Binding of Ku protein to DNA. Measurement of affinity for ends and demonstration of binding to nicks. *J. Biol. Chem.* **268**, 7594–7601.
- Britten, R. A., Kuny, S., and Perdue, S. (1999) Modification of non-conservative double-strand break (DSB) rejoining activity after the induction of cisplatin resistance in human tumour cells. *Br. J. Cancer* **79**, 843–849.
- Brown, K. D., Lataxes, T. A., Shangary, S., Mannino, J. L., Giardina, J. F., Chen, J., and Baskaran, R. (2000) Ionizing radiation exposure results in up-regulation of Ku70 via a p53/ataxia-telangiectasia mutated protein-dependent mechanism. *J. Biol. Chem.* **275**, 6651–6656.
- Brush, G. S. and Kelly, T. J. (2000) Phosphorylation of the replication protein A large subunit in the *Saccharomyces cerevisiae* checkpoint response. *Nucleic Acids Res.* **28**, 3725–3732.
- Bryntesson, F., Regan, J. C., Jeggo, P. A., Taccioli, G. E., and Hubank, M. (2001) Analysis of gene transcription in cells lacking DNA-PK activity. *Radiat. Res.* **156**, 167–176.
- Burma, S., Kurimasa, A., Xie, G., Taya, Y., Araki, R., Abe, M., Crissman, H. A., Ouyang, H., Li, G. C., and Chen, D. J. (1999) DNA-dependent protein kinase-independent activation of p53 in response to DNA damage. *J. Biol. Chem.* **274**, 17139–17143.
- Canman, C. E., Wolff, A. C., Chen, C. Y., Fomace, Jr. A. J., and Kastan, M. B. (1994) The p53-dependent G1 checkpoint pathway and ataxia-telangiectasia. *Cancer Res.* **54**, 5054–5058.
- Cary, R. B., Chen, F., Shen, Z., and Chen, D. J. (1997) DNA looping by Ku and the DNA-dependent protein kinase. *Proc. Natl. Acad. Sci. USA* **94**, 4267–4272.
- Chan, D. W. and Lees-Miller, S. P. (1996) The DNA-dependent protein kinase is inactivated by autophosphorylation of the catalytic subunit. *J. Biol. Chem.* **271**, 8936–8941.
- Chibazakura, T., Watanabe, F., Kitajima, S., Tsukada, K., Yasukochi, Y., and Teraoka, H. (1997) Phosphorylation of human general transcription factors TATA-binding protein and transcription factor IIB by DNA-dependent protein kinase-synergistic stimulation of RNA polymerase II basal transcription *in vitro*. *Eur. J. Biochem.* **247**, 1166–1173.
- Chiu, C. Y., Cary, R. B., Chen, D. J., Peterson, S. R., and Stewart, P. L. (1998) Cryo-EM imaging of the catalytic subunit of the DNA-dependent protein kinase. *J. Mol. Biol.* **284**, 1075–1081.
- Chung, U., Igarashi, T., Nishishita, T., Iwanari, H., Iwamatsu, A., Suwa, A., Mimori, T., Hata, K., Ebisu, S., Ogata, E., Fujita, T., and Okazaki, T. (1996) The interaction between Ku antigen and REF1 protein mediates negative gene regulation by extracellular calcium. *J. Biol. Chem.* **271**, 8593–8598.
- Critchlow, S. E. and Jackson, S. P. (1998) DNA end-joining: from yeast to man. *Trends Biochem. Sci.* **23**, 394–398.
- d'Adda di Fagagna, F., Hande, M. P., Tong, W., Roth, D., Lansdorp, P. M., Wang, Z., and Jackson, S. P. (2001) Effects of DNA nonhomologous end-joining factors on telomere length and chromosomal stability in mammalian cells. *Curr. Biol.* **11**, 1192–1196.
- Douglas, P., Moorhead, G. B., Ye, R., and Lees-Miller, S. P. (2001) Protein phosphatases regulate DNA-dependent protein kinase activity. *J. Biol. Chem.* **276**, 18992–18998.
- Falzon, M., Fewell, J. W., and Kuff, E. L. (1993) EBP-80, a transcription factor closely resembling the human autoantigen Ku, recognizes single- to double-strand transitions in DNA. *J. Biol. Chem.* **268**, 10546–10552.
- Featherstone, C. and Jackson, S. P. (1999) Ku, a DNA repair protein with multiple cellular functions? *Mutat. Res.* **434**, 3–15.
- Feldmann, E., Schmiemann, V., Goedecke, W., Reichenberger, S., and Pfeiffer, P. (2000) DNA double-strand break repair in cell-free extracts from Ku80-deficient cells: implications for Ku serving as an alignment factor in non-homologous DNA end joining. *Nucleic Acids Res.* **28**, 2585–2596.
- Frit, P., Canitrot, Y., Muller, C., Foray, N., Calsou, P., Marangoni, E., Bourhis, J., and Salles, B. (1999) Cross-resistance to ionizing radiation in a murine leukemic cell line resistant to cis-dichlorodiammine platinum (II): role of Ku autoantigen. *Mol. Pharmacol.* **56**, 141–146.
- Frit, P., Li, R. Y., Arzel, D., Salles, B., and Calsou, P. (2000) Ku entry into DNA inhibits inward DNA transactions *in vitro*. *J. Biol. Chem.* **275**, 35684–35691.
- Gell, D. and Jackson, S. P. (1999) Mapping of protein-protein interactions within the DNA-dependent protein kinase complex. *Nucleic Acids Res.* **27**, 3494–3502.
- Goedecke, W., Eijpe, M., Offenberg, H. H., van Aalderen, M., and Heyting, C. (1999) Mre11 and Ku70 interact in somatic cells, but are differentially expressed in early meiosis. *Nat. Genet.* **23**, 194–198.
- Gottlieb, T. M. and Jackson, S. P. (1993) The DNA-dependent protein kinase: requirement for DNA ends and association with Ku antigen. *Cell* **72**, 131–142.
- Goytisolo, F. A., Samper, E., Edmonson, S., Taccioli, G. E., and Blasco, M. A. (2001) The absence of the dna-dependent protein kinase catalytic subunit in mice results in anaphase bridges and in increased telomeric fusions with normal telomere length and G-strand overhang. *Mol. Cell. Biol.* **21**,

- 3642-3651.
- Gu, Y., Sekiguchi, J., Gao, Y., Dikkes, P., Frank, K., Ferguson, D., Hasty, P., Chun, J., and Alt, F. W. (2000) Defective embryonic neurogenesis in Ku-deficient but not DNA-dependent protein kinase catalytic subunit-deficient mice. *Proc. Natl. Acad. Sci. USA* **97**, 2668-2673.
- Guidos, C. J., Williams, C. J., Grandal, I., Knowles, G., Huang, M. T., and Danska, J. S. (1996) V(D)J recombination activates a p53-dependent DNA damage checkpoint in scid lymphocyte precursors. *Genes Dev.* **10**, 2038-2054.
- Hartley, K. O., Gel, D., Smith, G. C., Zhang, H., Divecha, N., Connelly, M. A., Admon, A., Lees-Miller, S. P., Anderson, C. W., and Jackson, S. P. (1995) DNA-dependent protein kinase catalytic subunit: a relative of phosphatidylinositol 3-kinase and the ataxia telangiectasia gene product. *Cell* **82**, 849-856.
- Hsu, H. L., Gilley, D., Blackburn, E. H., and Chen, D. J. (1999) Ku is associated with the telomere in mammals. *Proc. Natl. Acad. Sci. USA* **96**, 12454-12458.
- Jackson, S. P. (1996) DNA damage detection by DNA dependent protein kinase and related enzymes. *Cancer Surv.* **28**, 261-279.
- Jhappan, C., Yusufzai, T. M., Anderson, S., Anver, M. R., and Merlino, G. (2000) The p53 response to DNA damage *in vivo* is independent of DNA-dependent protein kinase. *Mol. Cell. Biol.* **20**, 4075-4083.
- Jimenez, G. S., Bryntesson, F., Torres-Arzayus, M. I., Priestley, A., Beeche, M., Saito, S., Sakaguchi, K., Appella, E., Jeggo, P. A., Taccioli, G. E., Wahl, G. M., and Hubank, M. (1999) DNA-dependent protein kinase is not required for the p53-dependent response to DNA damage. *Nature* **400**, 81-83.
- Jin, S., Kharbanda, S., Mayer, B., Kufe, D., and Weaver, D. T. (1997a) Binding of Ku and c-Abl at the kinase homology region of DNA-dependent protein kinase catalytic subunit. *J. Biol. Chem.* **272**, 24763-24766.
- Jin, S., Inoue, S., and Weaver, D. T. (1997b) Functions of the DNA dependent protein kinase. *Cancer Surv.* **29**, 221-261.
- Johnson, N. L., Gardner, A. M., Diener, K. M., Lange-Carter, C. A., Gleavy, J., Jarpe, M. B., Minden, A., Karin, M., Zon, L. I., and Johnson, G. L. (1996) Signal transduction pathways regulated by mitogen activated/extracellular response kinase kinase induced cell death. *J. Biol. Chem.* **271**, 3229-3237.
- Kachnic, L. A., Wu, B., Wunsch, H., Mekeel, K. L., DeFrank, J. S., Tang, W., and Powell, S. N. (1999) The ability of p53 to activate downstream genes p21(WAF1/cip1) and MDM2, and cell cycle arrest following DNA damage is delayed and attenuated in scid cells deficient in the DNA-dependent protein kinase. *J. Biol. Chem.* **274**, 13111-13117.
- Kharbanda, S., Pandey, P., Jin, S., Inoue, S., Bharti, A., Yuan, Z. M., Weichselbaum, R., Weaver, D., and Kufe, D. (1997) Functional interaction between DNA-PK and c-Abl in response to DNA damage. *Nature* **386**, 732-735.
- Koike, M., Ikuta, T., Miyasaka, T., and Shiomi, T. (1999) Ku80 can translocate to the nucleus independent of the translocation of Ku70 using its own nuclear localization signal. *Oncogene* **18**, 7495-7505.
- Koike, M., Shiomi, T., and Koike, A. (2001) Dimerization and nuclear localization of ku proteins. *J. Biol. Chem.* **276**, 11167-11173.
- Kuhn, A., Gottlieb, T. M., Jackson, S. P., and Grummt, I. (1995) DNA-dependent protein kinase: a potent inhibitor of transcription by RNA polymerase I. *Genes Dev.* **9**, 193-203.
- Kumaravel, T. S., Bharathy, K., Kudoh, S., Tanaka, K., and Kamada, N. (1998) Expression, localization, and functional interactions of Ku70 subunit of DNA-PK in peripheral lymphocytes after irradiation. *Int. J. Radiat. Biol.* **74**, 481-489.
- Kurimasa, A., Kumano, S., Boubnov, N. V., Story, M. D., Tung, C.-S., Peterson, S. R., and Chen, D. J. (1999) Requirement for the kinase activity of human DNA-dependent protein kinase catalytic subunit in DNA strand break rejoining. *Mol. Cell. Biol.* **19**, 3877-3884.
- Labhart, P. (1995) DNA-dependent protein kinase specifically represses promoter-directed transcription initiation by RNA polymerase I. *Proc. Natl. Acad. Sci. USA* **92**, 2934-2938.
- Lee, S. E., Mitchell, R. A., Cheng, A., and Hendrickson, E. A. (1997) Evidence for DNA-PK-dependent and -independent DNA double-strand break repair pathways in mammalian cells as a function of the cell cycle. *Mol. Cell. Biol.* **17**, 1425-1433.
- Lees-Miller, S. P., Godbout, R., Chan, D. W., Weinfeld, M., Day III, R. S., Barron, G. M., and Allalunis-Turner, J. (1995) Absence of p350 subunit of DNA-activated protein kinase from a radiosensitive human cell line. *Science* **267**, 1183-1185.
- Leuther, K. K., Hammarsten, O., Kornberg, R. D., and Chu, G. (1999) Structure of DNA-dependent protein kinase: implication for its regulation by DNA. *EMBO J.* **18**, 1114-1123.
- Li, B. and Comai, L. (2001) Requirements for the nucleolytic processing of DNA ends by the Werner syndrome protein-Ku70/80 complex. *J. Biol. Chem.* **276**, 9896-9902.
- Lin, R. Z., Hu, Z. W., Chin, J. H., and Hoffman, B. B. (1997) Heat shock activates c-Src tyrosine kinases and phosphatidylinositol 3-kinase in NIH3T3 fibroblasts. *J. Biol. Chem.* **272**, 31196-31202.
- Manolis, K. G., Nimmo, E. R., Hartsuiker, E., Carr, A. M., Jeggo, P. A., and Allshire, R. C. (2001) Novel functional requirements for non-homologous DNA end joining in *Schizosaccharomyces pombe*. *EMBO J.* **20**, 210-221.
- McConnell, K. and Dynan, W. S. (1996) The DNA-dependent protein kinase: a matter of life and (cell) death. *Curr. Opin. Cell. Biol.* **8**, 325-330.
- McElhinny, S. A., Snowden, C. M., McCarville, J., and Ramsden, D. A. (2000) Ku recruits the XRCC4-ligase IV complex to DNA ends. *Mol. Cell. Biol.* **20**, 2996-3003.
- Mimori, T., Hardin, J. A., and Steitz, J. A. (1986) Characterization of the DNA-binding protein antigen Ku recognized by autoantibodies from patients with rheumatic disorders. *J. Biol. Chem.* **261**, 2274-2278.
- Muller, C. and Salles, B. (1997) Regulation of DNA-dependent protein kinase activity in leukemic cells. *Oncogene* **15**, 2343-2348.
- Novac, O., Matheos, D., Araujo, F. D., Price, G. B., and Zannis-Hadjopoulos, M. (2001) *In vivo* association of Ku with Mammalian origins of DNA replication. *Mol. Biol. Cell.* **12**, 3386-3401.
- Osipovich, O., Duhe, R. J., Hasty, P., Durum, S. K., and Muegge, K. (1999) Defining functional domains of Ku80: DNA end binding and survival after radiation. *Biochem. Biophys. Res. Commun.* **261**, 802-807.
- Park, J. S., Park, S. J., Peng, X., Wang, M., Yu, M. A., and Lee, S. H. (1999) Involvement of DNA-dependent protein kinase

- in UV-induced replication arrest. *J. Biol. Chem.* **274**, 32520–32527.
- Park, S. J., Oh, E. J., Yoo, M. A., and Lee, S. H. (2001) Involvement of DNA-dependent protein kinase in regulation of stress-induced JNK activation. *DNA Cell. Biol.* **20**, 637–645.
- Peterson, S. R., Jesch, S. A., Chamberlin, T. N., Dvir, A., Rabindran, S. K., Wu, C., and Dynan, W. S. (1995) Stimulation of the DNA-dependent protein kinase by RNA polymerase II transcriptional activator proteins. *J. Biol. Chem.* **270**, 1449–1454.
- Poltoratsky, V. P., Shi, X., York, J. D., Lieber, M. R., and Carter, T. H. (1995) Human DNA-activated protein kinase (DNA-PK) is homologous to phosphatidylinositol kinases. *J. Immunol.* **155**, 4529–4533.
- Pucci, S., Mazzearelli, P., Rabitti, C., Giai, M., Gallucci, M., Flammia, G., Alcini, A., Altomare, V., and Fazio, V. M. (2001) Tumor specific modulation of KU70/80 DNA binding activity in breast and bladder human tumor biopsies. *Oncogene* **20**, 739–747.
- Rotman, G. and Shiloh, Y. (1997) Ataxia-telangiectasia: is ATM a sensor of oxidative damage and stress? *Bioessays* **19**, 911–917.
- Sanchez-perez, I., Murguia, J. R., and Perona, R. (1998) Cisplatin induces a persistent activation of JNK that is related to cell death. *Oncogene* **16**, 533–540.
- Shao, R. G., Cao, C. X., Zhang, H., Kohn, K. W., Wold, M. S., and Pommier, Y. (1999) Replication-mediated DNA damage by camptothecin induces phosphorylation of RPA by DNA-dependent protein kinase and dissociates RPA: DNA-PK complexes. *EMBO J.* **18**, 1397–1406.
- Shen, H., Kauvar, L., and Tew, K. D. (1997) Importance of glutathione and associated enzymes in drug response. *Oncol. Res.* **9**, 295–302.
- Song, K., Jung, D., Jung, Y., Lee, S. G., and Lee, I. (2000) Interaction of human Ku70 with TRF2. *FEBS Lett.* **481**, 81–85.
- Song, Q., Lees-Miller, S. P., Kumar, S., Zhang, N., Chan, D. W., Smith, G. C. M., Jackson, S. P., Alnemri, E. S., Litwack, G., Khanna, K. K., and Lavin, M. F. (1996) DNA-dependent protein kinase catalytic subunit: a target for an ICE-like protease in apoptosis. *EMBO J.* **15**, 3238–3246.
- Suwa, A., Hirakata, M., Takeda, Y., Jesch, S. A., Mimori, T., and Hardin, J. A. (1994) DNA-dependent protein kinase (Ku protein-p350 complex) assembles on double-stranded DNA. *Proc. Natl. Acad. Sci. USA* **91**, 6094–6098.
- Walker, J. R., Corpina, R. A., and Goldberg, J. (2001) Structure of the Ku heterodimer bound to DNA and its implications for double-strand break repair. *Nature* **412**, 607–614.
- Wang, H., Guan, J., Wang, H., Perrault, A. R., Wang, Y., and Iliakis, G. (2001) Replication protein A2 phosphorylation after DNA damage by the coordinated action of ataxia telangiectasia-mutated and DNA-dependent protein kinase. *Cancer Res.* **61**, 8554–8563.
- Wang, S., Guo, M., Ouyang, H., Li, X., Cordon-Cardo, C., Kurimasa, A., Chen, D. J., Fuks, Z., Ling, C. C., and Li, G. C. (2000) The catalytic subunit of DNA-dependent protein kinase selectively regulates p53-dependent apoptosis but not cell-cycle arrest. *Proc. Natl. Acad. Sci. USA* **97**, 1584–1588.
- Wang, Y., Zhou, X. Y., Wang, H., Huq, M. S., and Iliakis, G. (1999) Roles of replication protein A and DNA-dependent protein kinase in the regulation of DNA replication following DNA damage. *J. Biol. Chem.* **274**, 22060–22064.
- Widmann, C., Johnson, N. L., Gardner, A. M., Smith, R. J., and Johnson, G. L. (1997) Potentiation of apoptosis by low dose stress stimuli in cells expressing activated MEK kinase 1. *Oncogene* **15**, 2439–2447.
- Woo, R. A., McLure, K. G., Lees-Miller, S. P., Rancourt, D. E., and Lee, P. W. (1998) DNA-dependent protein kinase acts upstream of p53 in response to DNA damage. *Nature* **394**, 700–704.
- Yaneva, M., Kowalewski, T., and Lieber, M. R. (1998) Interaction of DNA-dependent protein kinase with DNA and with Ku: biochemical and atomic-force microscopy studies. *EMBO J.* **16**, 5098–5112.
- Yang, C. R., Leskov, K., Hosley-Eberlein, K., Criswell, T., Pink, J. J., Kinsella, T. J., and Boothman, D. A. (2000) Nuclear clusterin/XIP8, an x-ray-induced Ku70-binding protein that signals cell death. *Proc. Natl. Acad. Sci. USA* **97**, 5907–5912.
- Yoo, S., Kimzey, A., and Dynan, W. S. (1999) Photocross-linking of an oriented DNA repair complex. Ku bound at a single DNA end. *J. Biol. Chem.* **274**, 20034–20039.
- Yuan, Z. M., Utsugisawa, T., Huang, Y., Ishiko, T., Nakada, S., Kharbanda, S., Weichselbaum, R., and Kufe, D. (1997) Inhibition of phosphatidylinositol 3-kinase by c-Abl in the genotoxic stress response. *J. Biol. Chem.* **272**, 23485–23488.