

AD \_\_\_\_\_

Award Number: DAMD17-01-1-0676

TITLE: Preventing the Consequences of Alcohol Abuse:  
Identification of Soldiers at High Risk for Fatal  
and Serious Injuries

PRINCIPAL INVESTIGATOR: Nicole S. Bell, SC.D., MPH

CONTRACTING ORGANIZATION: Social Sectors Development  
Strategies, Inc  
Boston, MA 02118-2025

REPORT DATE: July 2003

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20040220 043

# REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

<b>1. AGENCY USE ONLY</b> (Leave blank)	<b>2. REPORT DATE</b> July 2003	<b>3. REPORT TYPE AND DATES COVERED</b> Annual (1 Jul 2002 - 30 Jun 2003)	
<b>4. TITLE AND SUBTITLE</b> Preventing the Consequences of Alcohol Abuse: Identification of Soldiers at High Risk for Fatal and Serious Injures		<b>5. FUNDING NUMBERS</b> DAMD17-01-1-0676	
<b>6. AUTHOR(S)</b>  Nicole S. Bell, SC.D., MPH		<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>	
<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b> Social Sectors Development Strategies, Inc Boston, MA 02118-2025  E-Mail: nbell@ssds.net			
<b>9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b> U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012		<b>10. SPONSORING / MONITORING AGENCY REPORT NUMBER</b>	
<b>11. SUPPLEMENTARY NOTES</b>			
<b>12a. DISTRIBUTION / AVAILABILITY STATEMENT</b> Approved for Public Release; Distribution Unlimited			<b>12b. DISTRIBUTION CODE</b>
<b>13. ABSTRACT (Maximum 200 Words)</b>  The main goal of this study is to improve our understanding of the relationship between alcohol problems and risk of serious injury by documenting the prevalence of alcohol-related diagnoses among soldiers admitted to the hospital with and without injuries. In this study's first year we have formulated diagnostic criteria for identifying alcohol- and injury-related hospitalizations and verified the continuity of diagnostic codes between the ICD-9 and the ICD-9-CM. Additionally, we have developed a method to identify unique alcohol-related hospitalizations using episodes of care in order to improve the accuracy of the frequency and prevalence estimates for alcohol-related diagnoses within the Army. Descriptive analyses of the prevalence of hospitalizations and characteristics of soldiers hospitalized for alcohol abuse have begun. Analyses comparing Army and civilian alcohol-related hospitalizations are also underway. The work to date highlights the importance of an expanded list of alcohol-related diagnostic criteria to ensure complete case ascertainment, and the necessity of constructing a valid episode of care methodology in order to obtain accurate estimates of alcohol-related hospitalizations.			
<b>14. SUBJECT TERMS</b> Alcohol, Army, injury, hospitalization, recurrent injuries, death, epidemiology, prevention, TAIHOD, behavior, risk factors			<b>15. NUMBER OF PAGES</b> 24
<b>17. SECURITY CLASSIFICATION OF REPORT</b> Unclassified			<b>16. PRICE CODE</b>
			<b>20. LIMITATION OF ABSTRACT</b> Unlimited
<b>18. SECURITY CLASSIFICATION OF THIS PAGE</b> Unclassified	<b>19. SECURITY CLASSIFICATION OF ABSTRACT</b> Unclassified		

## Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Background.....	4
First Year Progress.....	6
SOW Task 1.....	6
SOW Task 2.....	7
SOW Task 3.....	18
SOW Task 4a.....	18
Other Progress.....	18
Key Research Accomplishments.....	20
Reportable Outcomes.....	21
Conclusions.....	22
References.....	23
Appendices.....	none

## INTRODUCTION

The main goal of this study is to improve our understanding of the relationship between alcohol problems and risk of serious injury. Specifically, this study will document the prevalence of alcohol-related diagnoses among soldiers admitted to the hospital with and without injuries, and ascertain the relationship between alcohol-related diagnoses and injury. We will also document patterns of injury (e.g., by major cause or mechanism) associated with different types of alcohol problems (e.g., acute intoxication, chronic abuse/dependence). Additionally, we will compare injured patients with and without alcohol comorbidities, in order to evaluate whether or not alcohol involvement increases the risk of recurrent injury hospitalization or subsequent injury death.


## BACKGROUND

Injuries have a greater impact on the health and readiness of the U.S. armed forces than any other category of medical complaint. Acute injuries and musculoskeletal conditions (which are often the result of old injuries) account for more than 20% of outpatient visits (1), approximately 26% of hospitalizations (1), almost 60% of permanent disabilities (2), and almost 80% of active-duty deaths (3-5). The impact of injuries is felt in many ways, including lost productivity, decreased mission effectiveness, human suffering, and in the huge economic expenses associated with the care and rehabilitation of the injured (2, 6). Alcohol abuse and related problems are also an issue of great concern for the Army, and it has been estimated that the prevalence of alcohol problems is similar, if not greater, in the Army than for comparable age groups of the U.S. population (7). Alcohol abuse is an important potentially modifiable risk factor for injury, particularly off-duty injuries such as those sustained in motor vehicle crashes, falls, and assaults. For these reasons, the Department of Defense (DoD) has identified injuries and alcohol abuse as two of the top three targeted areas for health promotion and preventive care research (8). Despite these observations, many aspects of the quantitative relationship between alcohol abuse and injuries are not well understood. Improving our understanding of these associations is a necessary first step in the development of targeted interventions that can identify individuals at high risk of injury-related morbidity, mortality or repeat injury.

This research draws upon data from the Total Army Injury Health Outcomes Database (TAIHOD) (9, 10). Established at the U.S. Army Research Institute of Environmental Medicine (USARIEM) in 1994 to specifically examine the impact of injury and disability outcomes among U.S. Army soldiers, the TAIHOD now contains electronic records for all soldiers who have been on active duty since 1971 (nearly 5 million individuals). These data sources, which are linked at the level of the individual soldier, contain information on demographic and occupational characteristics, inpatient and outpatient health care utilization, and a variety of adverse health outcomes such as casualties, disabilities, accidents, and early separation from the Army (see Figure 1).

Figure 1. Components of the Total Army Injury Health Outcomes Database (TAIHOD)

## Outcomes Data



## Exposure Data

## FIRST YEAR PROGRESS

This section of the report outlines our research progress to date on each of our approved Statement of Work (SOW) objectives. Note that this grant was originally awarded to the Johns Hopkins University School of Hygiene and Public Health, where the original Principal Investigator, Dr. Gordon Smith, was employed. Upon Dr. Smith's departure from Johns Hopkins, the grant was transferred to Dr. Nicole Bell, of SSDS, Inc. However, due to administrative difficulties, funding was not officially transferred to SSDS until January 20, 2003. Therefore, although this annual report covers the time period from August 1, 2002, through July 31, 2002, it in fact addresses only the progress made by SSDS toward the approved SOW objectives between January 20, 2003, and July 31, 2003.

### SOW TASK 1

***Hire staff for project. Establish formal contract and budget arrangements. Establish access to all necessary databases; complete human subjects review and establish confidentiality safeguards for this project.***

### Progress

In January 2003, Dr. Katy Benjamin joined SSDS, Inc., as a Senior Scientist. Dr. Benjamin serves as project manager for this study, coordinating all analyses and overseeing the daily operations of the research. She is also responsible for several products associated with the study, including the descriptive epidemiology of alcohol hospitalizations and the paper describing the relationship of alcohol to injury and other comorbidities among hospitalized Army personnel.

We have sought and received approval from our IRB, the ARIEM Human Use Review Committee (HURC) for this project. The local approval from our HURC was forwarded to the Human Subjects Research Review Board (HSRRB), who completed their review of the protocol and approved the project in early January 2003. Both committees have approved the measures we have taken to safeguard human subject confidentiality. All team members and consultants have complied with training requirements pertaining to human subject confidentiality.

We have begun extracting information from the TAIHOD and constructing the analytic datasets that are needed to address our Year 1 SOW objectives. The data sources that will be instrumental in addressing our Year 1 Specific Aims (namely, personnel and hospitalization files) are up to date through 1999 and have been cleaned and examined for completeness. We have recently acquired Army hospitalization data through December 2002, and will file an amended study protocol with our IRB, the ARIEM HURC, seeking permission to extend the period of follow up for these analyses. We will also seek permission from our Grants Officer to amend our SOW accordingly.

We have obtained the public access data files from the National Hospital Discharge Survey (NHDS) and have begun exploring these files in order to perform comparative analyses between military and civilian alcohol-related hospitalizations.

## **SOW TASK 2**

***Identify cases with alcohol diagnoses in the hospital discharge data. Calculate incidence and prevalence rates of alcohol diagnoses as both principle and secondary diagnoses.***

### **Progress: Establishing Diagnostic Criteria for Alcohol-Related Hospitalizations**

A major focus of our work to date has been on the formulation of the diagnostic criteria to identify alcohol- and injury-related hospitalizations in the Army databases. Although the grant proposal contained a preliminary list of alcohol-related diagnoses, we began with a thorough review of the literature and consultation with our consultants to be sure that we had carefully enumerated the study population. For example, ingestion of nonbeverage alcohol such as mouthwash, aftershave lotion, and alcohol-based fuels is not uncommon among alcoholics (11, 12). However, diagnoses associated with the toxic effects of these substances are rarely included in studies of the prevalence of alcohol abuse in hospitalized patients. Similarly, discharge diagnoses of adverse effects of alcohol deterrents (ICD-9-CM codes 977.3 and E947.3) or history of mental disorders due to alcohol (V11.3) are infrequently used in alcohol epidemiology research. The diagnosis of fetal alcohol syndrome (760.71) was specifically excluded from the recent National Institute on Alcohol Abuse and Alcoholism (NIAAA) surveillance report because "childbirth is not an illness," and so all patients hospitalized for this purpose were eliminated from the denominator population (13). The omission of these codes from many civilian studies has potentially resulted in an underassessment of the morbidity burden associated with alcohol abuse. Based upon these considerations and further discussions with our consultants, we formulated a revised list of diagnostic criteria for identifying alcohol-related hospitalizations (see Table 1).

**Table 1. Diagnostic codes for identification of alcohol-related hospitalizations**

<b>Category</b>	<b>Classification in ICD-9-CM</b>
<b>Alcoholic psychoses</b>	291.0 Alcohol withdrawal delirium
	291.1 Alcohol amnestic syndrome
	291.2 Other alcoholic dementia
	291.3 Alcohol withdrawal hallucinosis
	291.4 Idiosyncratic alcohol intoxication
	291.5 Alcoholic jealousy
	291.8 Other specified alcoholic psychosis
	291.81 Alcohol withdrawal
	291.89 Alcohol psychosis—Other
	291.9 Unspecified alcoholic psychosis
<b>Alcohol dependence syndrome</b>	303.0 Acute alcoholic intoxication
	303.00 Acute alcoholic intoxication—Unspecified
	303.01 Acute alcoholic intoxication—Continuous
	303.02 Acute alcoholic intoxication—Episodic
	303.03 Acute alcoholic intoxication—In remission
	303.9 Other and unspecified alcohol dependence
	303.90 Other and unspecified alcohol dependence—Unspecified
	303.91 Other and unspecified alcohol dependence—Continuous
	303.92 Other and unspecified alcohol dependence—Episodic
	303.93 Other and unspecified alcohol dependence—In remission
	357.5 Alcoholic polyneuropathy
	425.5 Alcoholic cardiomyopathy
	535.3 Alcoholic gastritis
	535.30 Alcohol gastritis without mention of hemorrhage
535.31 Alcohol gastritis with hemorrhage	
<b>Nondependent abuse of alcohol</b>	305.00 Alcohol abuse—Unspecified
	305.01 Alcohol abuse—Continuous
	305.02 Alcohol abuse—Episodic
	305.03 Alcohol abuse—In remission
<b>Chronic liver disease and cirrhosis</b>	571.0 Alcoholic fatty liver
	571.1 Acute alcoholic hepatitis
	571.2 Alcoholic cirrhosis of liver
	571.3 Alcoholic liver damage, unspecified
	571.5 Cirrhosis of liver without mention of alcohol
	571.6 Biliary cirrhosis
	571.8 Other chronic nonalcoholic liver disease
	571.9 Unspecified chronic liver disease without mention of alcohol
	<b>Other conditions</b>
790.3 Excessive blood level of alcohol	
977.3 Poisoning by alcohol deterrents	
980.0 Toxic effect of ethyl alcohol	
980.1 Toxic effect of methyl alcohol	
980.2 Toxic effect of isopropyl alcohol	
980.8 Toxic effect of other specified alcohols	
980.9 Toxic effect of unspecified alcohol	
V11.3 Personal history of alcoholism	
V79.1 Screening for alcoholism	
E947.3 Adverse effects of alcohol deterrents	

The study period spans the years from 1980–1999. Beginning in approximately 1986 (the roll-out date was not uniform), many military medical facilities switched from using the ICD-9 to the ICD-9-CM when assigning diagnoses. In order to ensure that we were capturing all cases with alcohol involvement in the same way, we explored differences between the two versions of diagnostic codes. We compared all ICD-9-CM diagnoses included in the case criterion to those in the ICD-9 (Table 2). This analysis revealed that no substantive diagnostic coding differences existed to the level of two decimal places. We concluded that the shift from ICD-9 to ICD-9-CM would not affect our analyses.

**Table 2. Comparison of codes for alcohol-related diagnoses in ICD-9-CM and ICD-9**

Category	ICD-9-CM Description	ICD-9 Description	
Alcoholic psychoses	291.0 Alcohol withdrawal delirium	291.0 Delirium tremens	
	291.1 Alcohol amnesic syndrome	Korsakoff's psychosis, alcoholic	
	291.2 Other alcoholic dementia	Other alcoholic dementia	
	291.3 Alcohol withdrawal hallucinosis	Other alcoholic hallucinosis	
	291.4 Idiosyncratic alcohol intoxication	Pathological drunkenness	
	291.5 Alcoholic jealousy	Alcoholic jealousy	
	291.8 Other specified alcoholic psychosis	Other specified alcoholic psychosis	
	291.81 Alcohol withdrawal	Alcohol withdrawal	
	291.89 Alcohol psychosis—Other	Other specified alcohol psychosis	
	291.9 Unspecified alcoholic psychosis	Unspecified alcoholic psychosis	
	Alcohol dependence syndrome	303.0 Acute alcoholic intoxication	303.0 Alcohol dependence with acute intoxication
		303.00 Acute alcoholic intoxication—Unspecified	Alcohol dependence, unspecified
		303.01 Acute alcoholic intoxication—Continuous	Alcohol dependence, Continuous
		303.02 Acute alcoholic intoxication—Episodic	Alcohol dependence, Episodic
		303.03 Acute alcoholic intoxication—In remission	Alcohol dependence, In remission
		303.9 Other and unspecified alcohol dependence	Alcohol dependence, unspecified
		303.90 Other and unspecified alcohol dependence—Unspecified	Unspecified dependence
		303.91 Other and unspecified alcohol dependence—Continuous	Unspecified dependence, Continuous
		303.92 Other and unspecified alcohol dependence—Episodic	Unspecified dependence, Episodic
303.93 Other and unspecified alcohol dependence—In remission		Unspecified dependence, In remission	
357.5 Alcoholic polyneuropathy		Alcoholic polyneuropathy	
425.5 Alcoholic cardiomyopathy		Alcoholic cardiomyopathy	
535.3 Alcoholic gastritis		Alcoholic gastritis	
535.30 Alcohol gastritis without mention of hemorrhage		Alcohol gastritis without hemorrhage	
535.31 Alcohol gastritis with hemorrhage		Alcohol gastritis with hemorrhage	
Nondependent use of alcohol		305.0 Alcohol Abuse—Unspecified	305.0 Alcohol Abuse—Unspecified
		305.00 Alcohol Abuse—Unspecified	Alcohol Abuse—Unspecified
		305.01 Alcohol Abuse—Continuous	Alcohol Abuse—Continuous
		305.02 Alcohol Abuse—Episodic	Alcohol Abuse—Episodic
		305.03 Alcohol Abuse—In remission	Alcohol Abuse—In remission
	Chronic liver disease and cirrhosis	571.0 Alcoholic fatty liver	571.0 Alcoholic fatty liver
571.1 Acute alcoholic hepatitis		Acute alcoholic hepatitis	
571.2 Alcoholic cirrhosis of liver		Alcoholic cirrhosis of liver	
571.3 Alcoholic liver damage, unspecified		Alcoholic liver damage, unspecified	
571.5 Cirrhosis of liver without mention of liver		Cirrhosis of liver without mention of liver	
571.6 Biliary cirrhosis		Biliary cirrhosis	
571.8 Other chronic nonalcoholic liver disease		Other nonalcoholic chronic liver disease	
571.9 Unspecified chronic liver disease without mention of alcohol		Unspecified nonalcoholic chronic liver disease	
Other conditions		760.71 Alcohol affecting fetus via placenta or breast milk	760.71 Fetal alcohol syndrome
	790.3 Elevated blood level of alcohol	790.3 Elevated blood alcohol level	

Category	ICD-9-CM Description	ICD-9 Description
	977.3 Poisoning by alcohol deterrents 980.0 Toxic effect of ethyl alcohol 980.1 Toxic effect of methyl alcohol 980.2 Toxic effect of isopropyl alcohol 980.8 Toxic effect of other specified alcohols 980.9 Toxic effect of unspecified alcohol V11.3 Personal history of alcoholism V79.1 Screening for alcoholism E947.3 Adverse effects of alcohol deterrents	977.3 Poisoning by alcohol deterrents 980.0 Toxic effect of ethyl alcohol 980.1 Toxic effect of methyl alcohol 980.2 Toxic effect of isopropyl alcohol 980.8 Toxic effect of other specified alcohols 980.9 Toxic effect of unspecified alcohol V11.3 Personal history of alcoholism V79.1 Screening for alcoholism E947.3 Alcohol deterrents causing adverse effects in therapeutic use

## **Progress: Establishing Diagnostic Criteria for Injury-Related Hospitalizations**

In conjunction with this work, we have developed the diagnostic criteria for identifying injury-related hospitalizations in Army databases, using ICD-9-CM diagnostic codes. Once again we began with a review of the literature and met with consultants to review and consider all options. Some conditions listed in the 800-999 ICD-9-CM injury category, such as complications from medical procedures, are not considered by most injury epidemiologists to be true "injuries" and not likely to be causally associated with alcohol abuse. We have excluded these conditions from our case definition. Moreover, several codes within the range of injury codes pertain to alcohol-related poisoning. We have excluded these from our injury case definition because they are included in the case definition criteria for alcohol-related hospitalizations. We are also evaluating the inclusion of several conditions that lay outside the traditional range of injury and poisoning codes (800-999). For example, the codes in ICD-9-CM between 710-739 are typically used to code chronic musculoskeletal conditions, but several of these codes may also apply to acute musculoskeletal injuries (e.g., internal derangement of the knee). Based upon our literature review and input from our consultants, we will examine these codes and decide which of them to include in our injury case definition. The list of all injury diagnoses to be used in this study will potentially include:

- All Injury and Poisoning diagnoses (800 – 999) EXCEPT:
  - ICD-9-CM 905–909, Late effects of injuries, poisonings, toxic effects, and other external causes. These were excluded on the grounds that they do not represent acute injuries that could be associated with current alcohol abuse.
  - ICD-9-CM 980.0–980.2; 980.8, 980.9, Toxic effects of alcohol (various)
  - ICD-9-CM 977.3, Poisoning by alcohol deterrents
- ICD-9-CM 310.2, Post-concussion syndrome
- Selected conditions from the ICD-9-CM Musculoskeletal System and Connective Tissue category (codes 710 – 739).

In addition to refining our injury case definition we have also spent time considering the ways in which cause of injury is to be coded for some or our analyses. Military hospitals do not use the ICD-9-CM system of E-codes for indicating external cause of injury. They instead use the North Atlantic Treaty Organization Standardization Agreement (STANAG) 2050 coding system (14, 15). The STANAG system was established by treaty among the NATO member nations to facilitate sharing information about battle casualties. It codes all injuries (whether combat related or not) according to intent, duty status, and specific cause. Coding compliance for injuries in military healthcare facilities is excellent, as the computer system automatically prompts for the selection of a STANAG code if the record contains an ICD-9-CM code for injury or poisoning (i.e., in the range 800-999) anywhere in the record. The STANAG codes comprise two elements, the Trauma code and the Injury code. The Trauma code identifies duty status at the time of the injury (see Table 3).

**Table 3. STANAG Trauma codes**

General Trauma Class	Code	Definition
Battle wound or injury	0	Direct result of action by or against an organized enemy
	1	Other battle casualties
Intentionally inflicted nonbattle injuries	2	Result of intervention of legal authority
	3	Assault, or intentionally inflicted by another person
	4	Intentionally self-inflicted
Accidental injury	5	Occurring while off-duty (includes leave, pass, AWOL, and other off-duty)
	6	Schemes and exercises
	7	All other scheduled training (including basic training), assault courses, etc.
	8	Occurring while on duty
	9	Unknown whether on or off duty

The STANAG codes describing specific external causes of injury appear in Table 4.

**Table 4. STANAG Injury codes**

Category	Major Group	Minor Group	Category Description
I	000-059		Accidents in air transport, as specifically defined, spacecraft accidents and escape system injuries
		000-029	Air transport involving military aircraft
		030-039	Air transport involving nonmilitary and unspecified aircraft
		040-049	Accidents involving spacecraft
		051-057	Escape system injuries
II	100-149		Accidents in land transport, as specifically defined
		100-109	Private vehicle
		110-119	Military vehicle
		120-129	Nontraffic private
		130-139	Nontraffic military
140-149	Rail and other land transport		
III	150-199		Accidents in water transport, as specifically defined
IV	200-249		Athletics and sports, including physical training
V	250-299		Reactions, complications, and misadventures in medical or surgical procedures and late effects
VI	300-479		Instrumentalities of war, when employed by the enemy in wartime
		300-319	Agents of nuclear warfare
		320-339	Agents of chemical warfare, excluding incendiaries
		340-359	Agents of biological warfare
		360-399	Other unconventional instrumentalities of war
		400-419	Conventional weapons injury to occupant of aircraft
		420-439	Conventional weapons injury to person on board ship
440-459	Conventional weapons injury to person on and or in unspecified location		
460-479	Indirect or secondary effects of instrumentalities of war, when employed as such in wartime		
VII	480-499		Accidents in connection with own instrumentalities of war, when employed as such in wartime
VII	50*-59*		Guns, explosives, and related agents, except when used as instrumentalities of war in wartime
IX	60*-69*		Machinery, tools, and selected agents
X	70*-79*		Poisons, fire, hot or corrosive substances
XI	80*-89*		Specified environmental factors
XII	90*-99*		Falls and miscellaneous other or unspecified agents

Note that Injury codes in Categories VII-XII also require place-of-occurrence codes ranging from 0 to 9 (see Table 5).

**Table 5. STANAG Place of occurrence codes**

Code	Place of Occurrence of Injury
0	On board aircraft or spacecraft or in the air, or in space
1	On board ship or other water transport, or in water (sea, rivers, lakes, etc).
2	On land and at an airfield
3	On land and at a dock
4	On land and at an industrial plant (e.g. ordnance factory, supply warehouse, repair shop)
5	On land and on a firing range or drill field
6	On land and on obstacle course
7	On land and in kitchen (other than home), mess hall, or bakery
8	On land and in the home, quarters or barracks.
9	On land other or unspecified

In addition to cause of injury codes, cases will also be grouped into similar categories of injury (e.g., traumatic, poisoning, environmental exposure). These injury groupings will be determined in later analytic planning meetings with input from consultants.

### **Progress: Identifying Unique Episodes of Care**

Most population-based epidemiologic studies of the prevalence of alcohol-related hospitalizations rely on datasets without unique identifiers and are therefore unable to link records belonging to the same individual. Thus, individuals admitted to the hospital and then transferred to another care facility would likely be counted twice for the same episode of care. The problem of accurately counting hospitalizations without over-counting due to transfer cases has been well documented in the medical literature (16, 17) (18) although our recent review of the literature did not identify any publications that discussed it specifically with respect to hospitalizations for alcohol-related problems.

One advantage to hospitalization data contained in the TAIHOD is the presence of identifiers and the relative completeness of data on all hospitalizations, which allows us to identify unique hospitalization events as distinct from continuations of care for the same event. This represents an important advantage over many studies in that it gives us the ability to arrive at more accurate estimates of the incidence and prevalence of alcohol conditions among hospitalized soldiers. Identification of transfers in the TAIHOD hospital database is possible through the use of specific variables in the hospital database that indicate whether the hospitalization is a transfer into or out of the hospital from or /to another facility. Our investigation of these variables indicated that the transfer codes were not always complete. Thus we decided to not rely solely upon the transfer codes to determine whether or not consecutive hospitalizations represented a single episode of care or were two unique events. We developed a method that defined continuations of care (transfer cases) based upon the presence of transfer codes in both the initial hospitalization record and the following transfer record, or cases where there are no transfer codes present but where soldiers have consecutive, contiguous hospitalizations for alcohol-related conditions, and there are no more than 3

days between admissions. Given our findings, this should serve as a conservative measure of transfer cases. Using this method, we have identified 65,681 hospitalizations representing 53,730 individual patients. Approximately 8% of all hospitalizations over the entire study interval appear to be transfers for the same episode of care. Table 6 provides descriptive information on the frequency of alcohol-related hospitalizations for the entire study period.

**Table 6. Frequency of alcohol-related hospitalizations among active duty Army soldiers, 1980-1999**

	N
Individual Alcohol-related hospitalizations, 1980-1999	71,009
Transfer hospitalizations	5,328
Alcohol-related episodes of hospital care, 1980-1999	
Total	65,681
1980-1984	15,066
1985-1989	18,240
1990-1994	23,853
1995-1999	8,522

We have begun descriptive analyses of demographic characteristics in the case population (Table 7), as well as the incidence rates of alcohol-related hospitalizations over time, by diagnosis type and position (primary diagnosis and any other position in the discharge diagnosis), and in relationship to comorbid conditions also listed in the hospital record. These findings will be presented in our first manuscript.

**Table 7. Demographic characteristics of active duty Army soldiers hospitalized for alcohol-related conditions, 1980–1999**

Demographic characteristic	Frequency	Percent
Gender		
Male	60,363	92%
Female	5,294	8%
Age		
< 21	10,083	15%
21 – 25	21,561	33%
26 – 30	12,731	19%
31 – 35	9,986	15%
36 – 40	7,658	12%
> 40	3,559	5%
Ethnicity		
White	45,695	70%
African-American	15,060	23%
Hispanic	2,261	3%
Indian/Alaskan	893	1%
Asian/Pacific Island	492	0.75%
Other	1,252	2%
Unknown	28	0.04%
Education		
< High school	3,738	6%
High School/GED	54,683	83%
Some college	4,117	6%
College	1,946	3%
> College	740	0.01%
Rank		
E1 – E4	38,008	58%
E5 – E9	25,446	39%
Warrant Officer	519	0.8%
O1 – O3	869	1%
O4 – O5	699	1%
O6 – O11	129	0.2%
Time in service		
< 2 years	18,619	28%
2 – 5 years	16,804	26%
> 5 years	19,224	29%

In addition, we propose producing an additional manuscript under this SOW objective that describes our methodology for constructing episodes of care, using examples from the alcohol- and injury-related diagnoses examined in this study.

### **Progress: Comparisons to National Hospital Discharge Survey Data**

As stated in our specific aims, the first paper will compare incidence rates of alcohol-related hospitalizations in the military to those for the same time period within the general civilian population. Towards this end, we have procured data from the National Hospital Discharge Survey (NHDS) for the years 1979-1997. NHDS data for the years 1998-1999 are not presently available, and we will adjust the military data used for the comparative analyses to take this into account.

Using the same case diagnosis definition for alcohol-related hospitalizations that we are employing in our Army data, we plan to calculate annual rates of alcohol-related hospitalizations for the civilian population, adjusting for differences in age, gender, and

race between military and nonmilitary populations. It should be noted that the NHDS data cannot be linked at the patient level and thus cannot be used to construct episodes of care, as we intend to do with the military data. Thus, we expect some discrepancies between military and civilian rates of alcohol-related hospitalizations, because we will not be able to parse the civilian data for unique episodes of care. We will discuss this in the limitations section of the paper.

### **SOW TASK 3**

***Develop linkage between cases in hospitalization databases; obtain, analyze, and link data from mortality databases with hospitalization records.***

#### **Progress: Link Army Data Sources on Nonfatal and Fatal Injuries**

All linkages necessary for the analysis of case characteristics and alcohol-related hospitalizations have been formulated, using data from the TAIHOD personnel and hospitalization databases. Some of the SOW objectives for later years of this study will employ Army data on fatalities. While the original proposal indicated that mortality data would be linked at the beginning of the study, we find it more efficient to link data closer to the time when the analyses will be performed. Thus, mortality data will be examined in subsequent years of this study, when the outcomes of alcohol-related hospitalizations are analyzed. We therefore defer this portion of the task until Year 03.

### **SOW TASK 4A**

***Begin preparation of paper on the prevalence of alcohol diagnoses in the army.***

#### **Progress: Calculating Incidence of Alcohol-Related Hospitalizations**

As indicated in the discussion of SOW Task 2 progress, we are calculating rates of alcohol-related hospitalizations among active-duty Army personnel and have begun the construction of rates for various subgroups of interest (e.g., gender, age, rank, race) and by year. We will also calculate rates according to various categories of alcohol-related conditions, as displayed in Table 1. Prevalence is being calculated for hospitalizations with alcohol diagnoses in the primary position and other diagnostic positions. Where alcohol-related conditions are not the primary diagnosis at discharge, we are constructing rates by type of comorbidity, using the major ICD-9-CM diagnostic categories to describe these other conditions. Additionally, we are constructing the dataset of all alcohol-related hospital discharges to be used in the analyses comparing military and nonmilitary personnel.

### **OTHER PROGRESS**

We were recently contacted by Dr. Les McFarling of the Army Center for Substance Abuse Programs (ACSAP). One of the many programs the ACSAP oversees is the Drug and Alcohol Management Information System (DAMIS), the current and historical repository of data submitted by Army Substance Abuse Programs (ASAP) worldwide along with U.S. Army and contract forensic and toxicology drug testing laboratories (FTDTLs). It provides information on ASAP screenings and/or enrollments, urinalysis testing, prevention and education efforts and ASAP resource and

performance milestones. The goals and objectives of DAMIS are to provide policy makers and managers at all levels with timely, accurate data on the extent, nature and trends of substance abuse in the Army; a centralized means for tracking an individual's current status or previous history through the ASAP from identification to final disposition; and information on the effectiveness of the ASAP. Although the TAIHOD hospitalization files contain information on some soldiers who are treated for alcohol abuse on an inpatient basis, the ACSAP files are more comprehensive. We are exploring ways to incorporate these data into our study. Towards that end, we are corresponding with Dr. McFarling about obtaining DAMIS data to link to the TAIHOD, and will seek approval from the ARIEM HURC to use these new data in the analyses for this project. In accordance with the terms of our grant, we will also, of course, apply to our Grants Officer for permission to revise our approved SOW objectives to use these data.

## KEY RESEARCH ACCOMPLISHMENTS

- Formulated diagnostic criteria for both alcohol-related and injury-related hospitalizations to be examined in this study.
- Verified the continuity of diagnostic codes between the ICD-9 and the ICD-9-CM.
- Constructed a dataset of civilian alcohol-related hospitalizations and designed comparative analyses to explore prevalence rates of alcohol abuse in civilian versus Army populations. This will aid in the interpretation and generalization of our results to the community at large.
- Began preparation of a manuscript describing the epidemiology of alcohol-related hospitalizations among active-duty Army personnel.
- Developed a method to measure hospitalizations using episodes of care as opposed to counting each hospitalization as a discrete event. This will improve the accuracy of the frequency and prevalence estimates for alcohol-related and injury diagnoses within the Army.
- Began preparation of a manuscript discussing the need for an episode of care methodology specific to alcohol epidemiologic research and describing the method we have designed, using examples from this study.

## REPORTABLE OUTCOMES

Manuscript: Alcohol abuse among U.S. Army soldiers, 1980-1999. *In preparation*. This paper will describe the basic epidemiology of hospitalizations for alcohol abuse. It will provide frequencies and rates of alcohol-related conditions at both the hospital and individual patient level, by year and by type of alcohol diagnosis. Demographic information will also be provided. Comparison of military to civilian populations will be made using data from the National Hospital Discharge Survey.

Manuscript: The use of episodes of care methods in epidemiology research (working title). *In preparation*. This paper will discuss the limitations of epidemiological estimates based upon hospitalizations or other health care encounters where transfers or continuation of care are not taken into account. A method of estimating episodes of care by evaluating the lag time between hospitalizations will be presented, and examples of method application will be provided using data from the present study.

## CONCLUSIONS

Although we are still in the very early stages of this study, our preliminary work has revealed some important considerations regarding alcohol epidemiology research and interpretation of these studies. The modifications we have made in diagnostic inclusion criteria and in developing a methodology for identifying unique episodes of care should result in more complete and accurate ascertainment of cases and rates of alcohol abuse.

- Few published studies typically include patients who have ingested nonbeverage alcohols, or who are diagnosed with an adverse reaction to an alcohol deterrent. For example, recent studies have specifically excluded births, thus eliminating those cases associated with Fetal Alcohol Syndrome. Thus, rates of alcohol abuse based upon the criteria used in these studies may underestimate the problem due to incomplete case ascertainment. We are taking a more comprehensive approach to defining alcohol-related diagnoses and expect a more complete accounting of alcohol-related morbidity.
- While hospital admissions and discharges are commonly used to estimate the prevalence of alcohol abuse within a community, most studies do not or cannot take into account transfers for continuation of care, treating each hospitalization as a separate event. This can lead to an overestimate of the problem by inflating the total number of such hospitalizations. Transfer cases should be counted as a single episode of care. We have designed a methodology to address this issue.

## REFERENCES

1. Frequencies, rates, and trends of hospitalizations and associated lost duty time among active soldiers, 1998. *Med Surv Monthly Rep* 1999;5(3):3-17.
2. Jones BH, Hansen BC. Injuries in the military: a hidden epidemic. A report for the Armed Forces Epidemiological Board. Aberdeen Proving Ground, MD: USACHPPM; 1996 November 1996. Report No.: 29-HA-4844-97.
3. U.S. Department of Health and Human Services. National mortality profile of active duty personnel in the U.S. Armed Forces: 1980-1993. Cincinnati, OH: U.S. Department of Health and Human Services Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health September 1996. Report No.: Publication No. 96-103.
4. Powell KE, Fingerhut LA, Branche CM, Perrotta DM. Deaths due to injury in the military. *Am J Prev Med* 2000;18(3 Suppl):26-32.
5. Helmkamp J, Gardner JW, Amoroso PJ. Chapter 2. Deaths due to injuries: casualty office data. *Mil Med* 1999;164(8 Suppl):1-72.
6. Jones BH, Hansen BC. An armed forces epidemiological board evaluation of injuries in the military. *Am J Prev Med* 2000;18(3 Suppl):14-25.
7. Bray RM, Sanches RP, Ornstein ML, Lentine D, Vincus AA, Baird TU, et al. 1998 Department of Defense Survey of Health Related Behaviors Among Military Personnel. Research Triangle Park, NC: Research Triangle Institute; 1998 1998.
8. DefenseLINK News. DoD Looks to Prevention as Cure for Costly Care. In: Defense LINK News; 1998.
9. Amoroso PJ, Swartz WG, Hoin FA, Yore MM. Total Army Injury and Health Outcomes Database: Description and capabilities. Technical Note. Natick, MA: U.S. Army Research Institute of Environmental Medicine; 1997 February 21. Report No.: TN97-2.
10. Amoroso PJ, Yore MM, Weyandt B, Jones BH. Chapter 8. Total Army injury and health outcomes database: a model comprehensive research database. *Mil Med* 1999;164(8 Suppl):1-36.
11. Egbert AM, Reed JS, Powell BJ, Liskow BI, Liese BS. Alcoholics who drink mouthwash: the spectrum of nonbeverage alcohol use. *J Stud Alcohol* 1985;46(6):473-81.
12. Burkhart KK, Kulig KW. The other alcohols. Methanol, ethylene glycol, and isopropanol. *Emerg Med Clin North Am* 1990;8(4):913-28.
13. Whitmore CC, Yi H, Chen CM, Dufour M. Surveillance Report #58 (Revised): Trends in Alcohol-Related Morbidity Among Short-Stay Community Hospital Discharges, United States, 1979-99. Rockville, MD: National Institute on Alcohol Abuse and Alcoholism, Division of Biometry and Epidemiology; 2002 April 2002.
14. Amoroso PJ, Smith GS, Bell NS. Qualitative assessment of cause-of-injury coding in U.S. military hospitals: NATO standardization agreement (STANAG) 2050. *Am J Prev Med* 2000;18(3 Suppl):174-87.
15. Amoroso PJ, Bell NS, Smith GS, Senior L, Pickett D. Viewpoint: a comparison of cause-of-injury coding in U.S. military and civilian hospitals. *Am J Prev Med* 2000;18(3 Suppl):164-73.

16. Goldacre MJ, Ferguson JA, Welch A, Kettlewell MG. Profiles of workload in general surgery from linked hospital statistics. *Br J Surg* 1993;80(8):1073-7.
17. Ferguson JA, Goldacre MJ, Bulstrode CJ. Workload in trauma and orthopaedic surgery: use of linked statistics to profile a specialty. *Health Serv Manage Res* 1995;8(1):55-63.
18. Westfall JM, McGloin J. Impact of double counting and transfer bias on estimated rates and outcomes of acute myocardial infarction. *Med Care* 2001;39(5):459-68.
19. Wingert TD, Kralewski JE, Lindquist TJ, Knutson DJ. Constructing episodes of care from encounter and claims data: Some methodological issues. *Inquiry* 1995;32(Winter 1995/96):430-443.