

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)

17-05-2004

REPRINT

4. TITLE AND SUBTITLE

Coronal Shocks and Solar Energetic Proton Events

5a. CONTRACT NUMBER

5b. GRANT NUMBER

5c. PROGRAM ELEMENT NUMBER

61102F

6. AUTHOR(S)

E.W. Cliver, S.W. Kahler, and D.V. Reames*

5d. PROJECT NUMBER

2311

5e. TASK NUMBER

RD

5f. WORK UNIT NUMBER

A1

7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)

Air Force Research Laboratory/VSBXS
29 Randolph Road
Hanscom AFB MA 01731-3010

8. PERFORMING ORGANIZATION REPORT NUMBER

20040526 037

9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)

AFRL/VSBXS

11. SPONSOR/MONITOR'S REPORT NUMBER(S)

AFRL-VS-HA-TR-2004-1087

12. DISTRIBUTION / AVAILABILITY STATEMENT

Approved for Public Release; Distribution Unlimited.

*NASA Goddard Space Flight Center, Greenbelt, MD

13. SUPPLEMENTARY NOTES

REPRINTED FROM: PROCEEDINGS OF THE 28TH INTERNATIONAL COSMIC RAY CONFERENCE, TSUKUBA, JAPAN, pp 3331-3334, 2003.

14. ABSTRACT

From July 1996 - June 2001, < 50% of favorably-located metric type II radio bursts had associated solar energetic protons (SEPs). When western hemisphere metric IIs were accompanied by decametric-hectometric (DH; 1-14 MHz) type II emission, their association with ~20 MeV SEP events was 90% vs. only 25% for metric IIs without a DH counterpart. Overall, 82% (63%) of all SEP events with visible disk origins were associated with metric (DH) type IIs, with percentage associations increasing with SEP event size to 88% (96%), respectively, for ~20 MeV SEP events with peak intensities $\geq 10^{-1}$ pr cm⁻² s⁻¹ sr⁻¹ MeV⁻¹. Our results are consistent with the following (not mutually exclusive) possibilities: (1) large ~20 MeV SEP events result from strong shocks that can persist well beyond ~3 R_☉; (2) shock acceleration is most efficient above ~3 R_☉; (3) shocks that survive to ~3 R_☉ are more likely to have broad longitudinal extents.

15. SUBJECT TERMS

Interplanetary shock waves
Solar radio emission

Solar energetic particles
Solar coronal mass ejection

16. SECURITY CLASSIFICATION OF:

a. REPORT
UNCLAS

UNCLAS

c. THIS PAGE
UNCLAS

17. LIMITATION OF ABSTRACT

SAR

18. NUMBER OF PAGES

4

19a. NAME OF RESPONSIBLE PERSON

E.W. Cliver

19b. TELEPHONE NUMBER (include area code)
781-377-3975

Coronal Shocks And Solar Energetic Proton Events

E.W. Cliver,¹ S.W. Kahler,¹ and D.V. Reames²

(1) *Space Vehicles Directorate, Hanscom AFB MA, 01731, USA*

(2) *NASA/Goddard Space Flight Center, Code 661, Greenbelt, MD 20771, USA*

Abstract

From July 1996 - June 2001, < 50% of favorably-located metric type II radio bursts had associated solar energetic protons (SEPs). When western hemisphere metric IIs were accompanied by decametric-hectometric (DH; 1-14 MHz) type II emission, their association with ~ 20 MeV SEP events was 90% vs. only 25% for metric IIs without a DH counterpart. Overall, 82% (63%) of all SEP events with visible disk origins were associated with metric (DH) type IIs, with percentage associations increasing with SEP event size to 88% (96%), respectively, for ~ 20 MeV SEP events with peak intensities $\geq 10^{-1}$ pr cm⁻² s⁻¹ sr⁻¹ MeV⁻¹. Our results are consistent with the following (not mutually exclusive) possibilities: (1) large ~ 20 MeV SEP events result from strong shocks that can persist well beyond $\sim 3 R_{\odot}$; (2) shock acceleration is most efficient above $\sim 3 R_{\odot}$; (3) shocks that survive to $\sim 3 R_{\odot}$ are more likely to have broad longitudinal extents.

1. Introduction

It is generally accepted that metric type II solar radio bursts are manifestations of shock waves caused by disturbances moving outward through the solar corona with speeds of ~ 500 - 1000 km s⁻¹. Metric type II bursts have typical starting frequencies ~ 100 MHz ($\sim 1.5 R_{\odot}$) and drift toward the lowest frequencies commonly observable from Earth (~ 20 MHz; $\sim 2.5 R_{\odot}$) with a drift rate of ~ 0.1 - 1 MHz s⁻¹. Coronal shock waves are thought to be the principal acceleration mechanism for the largest ("gradual") SEP events observed at 1 AU [7].

While coronal shocks are held to be a necessary condition for large SEP events, metric type II bursts are not a sufficient condition for a SEP event to occur. Kahler [4] found that only about half (31/58) of favorably connected (W10-W85) metric IIs observed from June 1973 through June 1980 were associated with >20 MeV SEP events. We investigate the possibility that the presence or absence of a DH type II burst observed in the 1-14 MHz range ($\sim 10 R_{\odot}$ - $3 R_{\odot}$) by the WAVES experiment on the Wind spacecraft may distinguish between SEP-associated and non-SEP-associated metric type II bursts (see [2]).




Fig. 1. Percentage association of ~ 20 MeV SEP events with metric type II (dashed line) and DH type II (solid line) bursts as a function of SEP event peak intensity.

2. Analysis

2.1. Event Associations

Metric type IIs and H α flares: We began with the list of metric type II bursts reported in Solar-Geophysical Data (SGD) between 01 July 1996 and 30 June 2001. During this period, 447 separate metric type II bursts were reported, of which 57% (253/447) were associated with H α disk flares.

DH type IIs and H α flares: For a list of DH type IIs, we used the compilation on the WAVES website, considering only those events with starting frequencies ≥ 1 MHz. During the period of interest, 172 independent DH IIs were reported, of which 54% (93/172) were associated with H α disk flares.

SEP events and H α flares: To compile a SEP event list, we used 19-22 MeV data from the EPACT investigation on the Wind spacecraft. This channel has a background of $\sim 2-3 \times 10^{-4}$ pr cm⁻² s⁻¹ sr⁻¹ MeV⁻¹. We identified 134 increases of ~ 20 MeV protons $\geq 10^{-3}$ pr cm⁻² s⁻¹ sr⁻¹ MeV⁻¹ during the five year interval and found disk flare associations for 88 of these events.

SEP events with metric and DH type IIs: We find that 82% (72/88) of our SEP events are associated with metric IIs, and 63% (55/88) are associated with DH IIs. 91% (80/88) of SEP events have metric and/or DH type II emission.

In Figure 1 it can be seen that smaller SEP events are more likely to be associated with metric IIs than with DH IIs and that the association of SEP events with DH type IIs increases rapidly with SEP event size. 80% (51/64) of SEP events with ~ 20 MeV intensities < 0.1 pr had metric IIs, vs. 88% (21/24) of SEP events with peak intensities \geq this threshold. Corresponding figures for DH type II associations with SEP events are 50% and 96%, while the rates for SEP events with associated metric and/or DH IIs are 88% and 100%.




Fig. 2. Percentages of metric type II only (dashed line) and metric II + DH II (solid line) bursts associated with ~ 20 MeV SEP events as a function of flare longitude.

Table 1.

WH metric II	SEP Event?		Total
	Yes	No	
w/ DH II	26	3	29
w/o DH II	17	52	69
Total	43	55	98

2.2. Factors affecting metric type II - SEP association

Flare longitude: "SEP visibility functions" for metric only type IIs (dashed line) and metric + DH type IIs (solid line) are shown in Figure 2. To construct these curves, we began with our samples of these two categories of type II bursts and determined their percentage SEP event associations as a function of flare longitude. In the figure it can be seen that SEP visibility for metric + DH type IIs is a factor of three or more times that for metric only IIs at all longitudes.

DH association: The matrix in Table 1 includes only the metric type IIs from western hemisphere sources in Figure 2. Comparing the columns of the matrix confirms Kahler's [4] finding that approximately half of favorably located type II bursts ($55/98 = 56\%$) are not accompanied by SEP events at Earth. Comparing the rows of the matrix indicates that only 30% ($29/98$) of metric type II bursts have DH counterparts.

The key result of our study is the marked difference in the degree of SEP association between metric IIs with and without DH type IIs. Only 25% ($17/69$) of western hemisphere metric type IIs without a DH II counterpart are followed by SEPs at Earth vs. 90% ($26/29$) of metric IIs with associated DH type II bursts.

3. Discussion

Our principal finding is that favorably located metric type II bursts that have a DH counterpart are much more likely to be followed by SEPs than metric IIs that lack such low frequency emission. How do we interpret this result? There are several possibilities, which are not mutually exclusive.

First, the strong/fast shocks that survive into the DH range [6] are also those that are most likely to be efficient proton accelerators (at all heights in the corona) since CME speed is correlated with peak ~ 20 MeV SEP intensity [5]. An alternative (or contributing) explanation for the high association between DH type IIs and SEP events is suggested by studies (e.g., [1]) indicating that the Alfvén speed in the corona typically has a peak value around $\sim 3 R_{\odot}$, corresponding to the ~ 14 MHz plasma level in the DH range. Above this height, shock acceleration will become more efficient as the Alfvén speed decreases. Support for the viewpoint that shock acceleration of SEPs is most efficient above $\sim 3 R_{\odot}$ comes from persistent evidence for delayed injection onsets of high-energy protons and mildly relativistic electrons [e.g., 3]. A third possible explanation for the link between DH type IIs and SEP events indicated by Table 1 might be that those shocks that survive to greater coronal heights will also span a broader range of longitudes and thus be more likely to reach the magnetic fieldline connected to Earth [8]. This effect is best illustrated by considering eastern hemisphere solar activity: eastern hemisphere metric IIs without accompanying DH type IIs are much less likely to have SEP association (8%; 4/50) than is the case for such events with associated DH type II emission (52%; 12/23).

In sum, the statistics presented here support the view that the solar energetic protons (SEPs) observed near Earth are accelerated at coronal shocks. But not all shocks are "SEP-effective"; only about half of favorably-located metric type IIs are linked to SEPs. The shocks that are strong enough to produce type IIs in the DH range ($> 3 R_{\odot}$) are the ones most likely to have SEP association.

4. References

1. Gopalswamy, N., et al. 2001a, *J. Geophys. Res.*, 106, 25,261
2. Gopalswamy, N., et al. 2002, *ApJ*, 572, L103
3. Haggerty, D.K., and Roelof, E.C. 2002, *ApJ*, 579, 841
4. Kahler, S.W. 1982, *ApJ*, 261, 710
5. Kahler, S.W. 2001, *J. Geophys. Res.*, 106, 20,947
6. Lara, A., et al. 2003, *Geophys. Res. Lett.* (in press)
7. Reames, D.V. 1999, *Space Sci. Rev.*, 90, 413
8. Torsti, J. 1999, *ApJ*, 510, 460