

## Dissertation Summary

# High-Precision Optical Interferometry and Application to Be Stars

CHRISTOPHER TYCNER

Current address: US Naval Observatory, Flagstaff Station, P.O. Box 1149, Flagstaff, AZ 86002; tycner@nofs.navy.mil

Thesis work conducted at University of Toronto

Ph.D. thesis directed by John B. Lester; Ph.D. degree awarded 2004 November

Received 2004 September 20; accepted 2004 September 20; published 2004 October 22

This dissertation presents a new technique for calibrating optical long-baseline interferometric observations that are obtained simultaneously in a number of spectral channels. For certain classes of objects, such as emission-line sources or binary systems, the new technique allows the calibration corrections and source characteristics to be obtained from the observations of a program star alone. This is because for such sources, the parameters describing the characteristics of the source have a different functional dependence than the calibration parameters.

The new calibration method is applied to long-baseline interferometric observations of four non-supergiant B-type stars with H $\alpha$  emission lines, which are also known as classical Be stars. The H $\alpha$  emission associated with these stars is also known to be the strongest line emission, and therefore it is an excellent probe of the shape and extent of the H $\alpha$ -emitting circumstellar region. The observations presented in this dissertation were obtained with the Navy Prototype Optical Interferometer,<sup>1</sup> which utilizes measurements obtained simultaneously in many spectral channels covering a wide spectral range. The alignment of the spectral channels was such that the H $\alpha$  emission was located in one or two spectral channels, and therefore it was possible to use the measurements from the continuum channels to derive the calibration corrections, which in turn were applied to the interferometric observables associated with the H $\alpha$  emission.

Circularly symmetric and elliptical Gaussian models were fitted to the calibrated observations in H $\alpha$ . Models with uniform brightness distributions were also successfully fitted. Because of the limited  $(u, v)$ -plane coverage, it was not possible to preferentially choose one set of models over the other. For consistency (and compatibility) with the already published interferometric results in the literature, the elliptical Gaussian models were chosen as the representative models describing the angular size and the apparent ellipticity of the circumstellar regions. In at least one of the stars analyzed, a signature indicative of a circumstellar disk that is not concentric with the central star was detected. Possible causes are discussed, and these range from the presence of one-armed density waves in the disk to a simple rotation signature.

To expand the data set, thereby making correlations more

apparent, the model parameters for the four Be stars investigated in this dissertation were combined with three Be stars from the study of Quirrenbach et al. (1997, ApJ, 479, 477). After accounting for the different distances to the sources and the different stellar continuum flux levels (with respect to which the H $\alpha$  emission is typically measured), it was possible to study the relationship between the net H $\alpha$  emission (in ergs s<sup>-1</sup>) and the physical extent (in m) of the H $\alpha$ -emitting circumstellar region. For the first time, a clear dependence of the net H $\alpha$  emission on the extent of the circumstellar region is demonstrated (see Fig. 1). These results are consistent with an optically thick line emission that is directly proportional to the effective area of the emitting disk. Within the small sample of stars considered in this analysis, no clear dependence on the spectral type or stellar rotation is established, although the results do suggest that hotter stars might have more extended H $\alpha$ -emitting regions.




FIG. 1.—log-log plot of the H $\alpha$  luminosity (in ergs s<sup>-1</sup>), corrected for the projection effect (through the division by the axial ratio  $r$ ), as a function of the size (in m) of the major axis of the disk. *Filled circles*: stars observed with the NPOI; *open circles*: stars observed with the Mark III interferometer (Quirrenbach et al. 1997, ApJ, 479, 477). A linear fit in the log-log plot produces a best-fit slope of  $2.24 \pm 0.26$  (*dashed line*) consistent with an optically thick H $\alpha$  emission from a geometrically thin disk of a relatively constant surface brightness.

<sup>1</sup> The NPOI is a joint project of the Naval Research Laboratory and the US Naval Observatory, in cooperation with Lowell Observatory, and is funded by the Office of Naval Research and the Oceanographer of the Navy.

# Report Documentation Page

Form Approved  
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE <b>22 OCT 2004</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>High-Precision Optical Interferometry and Application to Be Stars</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>U.S. Naval Observatory Library 3450 Massachusetts Avenue, N.W. Washington, DC 20392-5420</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>1</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			