



Simulation Studies of Cyanide-Caused Cardiac Toxicity

by C. K. Zoltani, COL G. E. Platoff, and S. I. Baskin

ARL-TR-3443

March 2005

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

Destroy this report when it is no longer needed. Do not return it to the originator.

Army Research Laboratory

Aberdeen Proving Ground, MD 21005-5067

ARL-TR-3443

March 2005

Simulation Studies of Cyanide-Caused Cardiac Toxicity

C. K. Zoltani

Computational and Information Sciences Directorate, ARL

COL G. E. Platoff and S. I. Baskin

**U.S. Army Medical Research Institute of Chemical Defense
Pharmacology Division**

REPORT DOCUMENTATION PAGE			<i>Form Approved</i> <i>OMB No. 0704-0188</i>		
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) March 2005		2. REPORT TYPE Final		3. DATES COVERED (From - To) October 2004–June 2004	
4. TITLE AND SUBTITLE Simulation Studies of Cyanide-Caused Cardiac Toxicity			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S) C. K. Zoltani, COL G. E. Platoff, and S. I. Baskin			5d. PROJECT NUMBER 4U74CC		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Research Laboratory AMSRD-ARL-CI-HC Aberdeen Proving Ground, MD 21005-5067			8. PERFORMING ORGANIZATION REPORT NUMBER ARL-TR-3443		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT A series of computer studies showing the effect of cyanide (CN) on the electrophysiology of cardiac tissue is presented. A mathematical model of the electrophysiology of cardiac tissue, with initial and boundary conditions based on experimental data from studies using CN as metabolic blockers from the literature, was used to simulate changes in the electrical activity of the heart. Emphasis was on the modulation of ion concentrations in the cells, changes in current magnitudes, and the activation of currents that are dormant under normal circumstances. These calculations showed for the first time: (1) disturbance of the energy homeostasis and ion concentrations in cardiac tissue due to CN results in the reversal of the direction from the normal and change in magnitudes of cellular membrane currents. These, in turn, change the morphology of the action potential and the electrocardiogram (ECG). This is the initial step leading to ventricular fibrillation, the usual endpoint in the effect of CN on the heart. (2) CN causes cell swelling and hemorrhaging in cardiac tissue. Cell swelling activates chloride membrane currents affecting homeostasis of the tissue. These effects were shown to be important for the electrical state of the CN-affected tissue and were included for the first time in a model of CN-affected cardiac tissue. (3) The calculations reproduced aspects of the changes in an ECG of a subject under the effect of a lethal dose of CN. (4) The obtained results suggest and define the characteristics required of a pharmacological intervention needed to overcome or reverse CN poisoning, of vital importance for development of therapeutics for force protection. Primarily, such intervention needs to reduce the calcium overload of the cardiac cell, restore the depolarizing sodium current, and alleviate the accumulation of potassium ions exterior to the cell.					
15. SUBJECT TERMS computer simulation, cardiac toxicity, cyanide					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UL	18. NUMBER OF PAGES 20	19a. NAME OF RESPONSIBLE PERSON C. K. Zoltani
a. REPORT UNCLASSIFIED	b. ABSTRACT UNCLASSIFIED	c. THIS PAGE UNCLASSIFIED			19b. TELEPHONE NUMBER (Include area code) 410-278-6650

Contents

List of Figures	iv
List of Tables	iv
Acknowledgments	v
1. Introduction	1
2. Effect of Cyanide on Cardiac Tissue	1
3. Computer Model	2
4. Results	4
5. Discussion	8
6. References	9
Distribution List	11

List of Figures

Figure 1. The baseline mathematically-generated ECG of the cardiac tissue with AP shown in the insert.....	4
Figure 2. CN notably shortens the CL of affected tissue (blue trace) in comparison with the baseline (red trace).....	5
Figure 3. Simulated ECG (blue trace) of severely CN-affected cardiac tissue. $[Ca^{2+}] = 0.0009$, $[ATP] = 1.0$, and $K^+_o = 12.0$ mM/L. The baseline is shown in red.	5
Figure 4. When the normoxic dormant currents are activated, further morphological changes in the ECG are evident.	5
Figure 5. ECG from a CN-intoxicated individual from Wexler et al. (1947).....	6
Figure 6. The effect of calcium-overload, reduced availability of ATP, and high external potassium concentration.....	6
Figure 7. The sodium concentrations returned to normal, and, with internal calcium at 0.0004 mM/L, the CL is improved, but tachycardia still is in evidence.....	7
Figure 8. Elimination of the exterior potassium overload helps (green curve), but baseline behavior is not achieved.....	7
Figure 9. Simulated pharmacological intervention: the red trace shows the baseline. The blue trace shows the case with ion concentrations restored except for sodium, showing a 50% cell overload.....	8

List of Tables

Table 1. Initial values.....	3
------------------------------	---

Acknowledgments

It is a pleasure to thank Dr. John Pormann of Duke University for his advice in the use of Cardiowave. Special thanks go to the U.S. Army Research Laboratory Major Shared Resource Center at Aberdeen Proving Ground, MD for the use of its high-performance computer assets, including the SP-4.

INTENTIONALLY LEFT BLANK.

1. Introduction

Cyanide (CN) is one of the most potent poisons used in warfare since ancient times. Development of effective antidotes for the warfighter is still a work in progress because many of the CN-affected cellular processes are not completely understood. These processes can now be mathematically modeled, based on experimental measurements, and the effects of changes in cellular parameters due to the presence of threat agents predicted. Only recently, with scalable computers coming online, has computational capability improved to the point where such simulations could be attempted. This report demonstrates the power of *in silico* (computational) experiments as a cost-effective, ancillary means of advancing the development of threat agent antidotes that is an essential part of force protection. We believe that this new approach will have a major impact on future therapeutic developments (see also Zoltani et al., 2003, 2004a, 2004b, and 2004c).

The strategy adopted was to use tissue parameter values in the mathematical simulations available from experimental investigations where CN was used as a metabolic blocking agent. The deviations from the baseline morphology in the calculated action potential (AP), expressing the voltage in the tissue as a function of time, and the mathematically-generated electrocardiogram (ECG), was related to and correlated with the measured tissue parameters. This gives a direct indication of the effects that have the strongest influence on the deviation of electrical state of the tissue from baseline and are the best candidates for pharmacological intervention in case of CN intoxication.

2. Effect of Cyanide on Cardiac Tissue

Dose-dependent heterogeneity marks the deposition of CN in cardiac tissue. Histological changes include cell swelling and hemorrhaging. CN has a strong attraction for iron ions, and its deposition prevents the transfer of electrons to molecular oxygen. Though there is oxygen available, it cannot perform the normoxic function in the adenosine triphosphate (ATP) generation needed by the tissue. The tissue becomes oxygen “starved,” resulting in the modulation of the energy homeostasis. Initially, glycolysis attempts to replenish the ATP, the energy source, but the replenishment is short lived. The whole tissue substrate changes; lactic acid and catecholamines are formed. The stage is set for changes and deviation from the norm in the electrical activity of the tissue.

One of the first manifestations of the changed electrophysiology is bradycardia that may soon change to Torsade de Pointes and possible culmination in ventricular fibrillation. On the ECG, the P-wave, the atrial depolarization, is eliminated. ST-segment deviation, usually a rise in the

slope, becomes noticeable, followed by modulation of the T-wave. The changed morphology is expressed in steepening and coalescing of the QRS and the T-waves. A J-wave becomes noticeable. Additional details are given in Katzman and Penny (1993) and Wexler et al. (1947).

On the cellular level, changes in the ion concentrations become important, especially calcium overload of the cell and increase in the extra cellular potassium concentration, $[K^+]_o$. The cell's energy homeostasis (Balaban, 2002) is profoundly disturbed, and several compensatory membrane currents are activated and others diminished. Three of the most important ones are the ATP-dependent I_{KATP} (Elliott et al., 1989), the osmotic swelling-activated $I_{Cl,swell}$, and the calcium-dependent $I_{Ca(L)}$. The disequilibrium in the membrane currents caused by the CN has grave implications for the cell's electrophysiology. CN-caused cardiac toxicity shares some commonality with ischemia but is different in the level of acidity of the tissue and the nature of some of the activated currents. A number of ancillary effects, including enhanced catecholamine (CA) secretion, the effect of the increase in free Mg^{2+} , and pH changes, are not addressed in this report, but we note that CA binds to α and β receptors that affect membrane currents. Additional details are given in Baskin and Brewer (1997), Baskin et al. (2004), Leimdorfer (1950), and Van der Heyden et al. (1985).

3. Computer Model

The effect of the presence of CN in the tissue was modeled by changing the tissue parameters to those measured in CN-caused metabolic blockade of cardiac tissue available in the open literature and including the currents activated under these conditions. Two of the more important ones for this model are the activation of I_{KATP} due to decline in the ATP stores and $I_{Cl,swell}$ when the cell volume is modulated. Change in cellular ion concentrations is also an important aspect of CN toxicity. Calcium overload causes the activation of K^+ channels (Inoue, 1998). Rise in $[Na^+]_i$ and $[Ca^{2+}]_i$ enhances the i_{Ks} current that is activated at voltage values much higher than for i_{Kr} . i_{Kr} is reduced by acidification and the presence of external divalent cations, noticeable in CN-affected tissue.

For these numerical simulations, cardiac tissue cells of three kinds were considered: epicardial, midmyocardial, and endocardial (representing the ventricular wall). The important distinction among these cells for these calculations is in the value of the maximum cell conductance. The model of Vandenberg et al. (1997) for osmotic swelling-activated chloride current was modified and incorporated into the simulations. It accounts for the expression of some of the changes in the membrane currents caused by CN-caused lesions (Suzuki, 1968). Cell swelling contributes to the rise of the resting membrane potential and the shortening of the AP.

For the calculations reported here, a monodomain approach was adopted with fiber orientation (one of the diffusion matrix entries) assumed to be uniform. The propagation of the AP was based on the following cable equation:

$$\frac{\partial V}{\partial t} = -I_{ion}/C_m + D \left(\frac{\partial^2 V}{\partial x^2} + \frac{\partial^2 V}{\partial y^2} \right). \quad (1)$$

In this equation, V is the membrane voltage and I_{ion} is the transmembrane ionic current, mainly made up from the sodium, potassium, calcium, and chloride currents, and pumps and exchangers. D is the diffusion constant and C_m the membrane capacitance. Appended are the gating variables of the ionic channels and equations for the change of the ion concentrations. No flux boundary conditions were used. The computational domain consisted of a fiber of 165 cardiac cells incorporating the regional differences of the tissue parameters in the ventricular wall. The signal propagation was described by a reaction diffusion equation. Explicit Euler method is adequate for the solution algorithm, although Crank-Nicholson can give more accuracy.

After determining the baseline in these simulations, the approach was to vary the ion channel conductivities thought to be affected in CN-affected tissue. In addition, a series of simulations were performed increasing the extra cellular potassium concentration, the calcium concentration in the cell, and the internal sodium level. Two additional currents, the ATP dependent potassium current and the cell volume dependent chlorine current, were activated in the simulation of severely-affected tissue. Using the conditions of table 1, first a baseline AP and then a pseudo-ECG were calculated. The former used the Luo-Rudy model of the cellular processes described in Ferrero et al. (1996) and Shaw and Rudy (1997). A pulse of 200 mA/cm² of 0.5-ms duration initiated the AP.

Table 1. Initial values.

Model Quantity	Baseline (mM)	CN-Affected Cell (mM)	Ischemia (mM)
[Na ⁺] _i	10.0	Incr. ~2.5 x	10 - 20
[Na ⁺] _o	145.0	134.0	140.0
[K ⁺] _i	150.0	125.0	125.0 (acidic)
[K ⁺] _o	4.0	10.0	4 -16
[Ca ²⁺] _i	0.0003	Incr. >3 x	0.0003 – 0.0009
[Ca ²⁺] _o	1.8	2.0	~2.0
[ATP]	5.0	3.0	3.0 <

Using the formulation of Plonsey and Barr (2000) with the pseudo-ECG electrode at 2 cm from epicardium, the extra cellular unipolar potential generated by the fiber in the surrounding field was calculated from

$$\Phi_e = \frac{a^2 \sigma_i}{4\sigma_e} \int \left(-\nabla V_m \right) \left[\nabla \frac{1}{r} \right] dx, \quad (2)$$

where a is the radius of the fiber, r is the distance from the source to the field point, V_m is the

transmembrane potential, and σ_i/σ_e is the ratio of the intracellular to the extra cellular conductivities.

The mathematically-generated ECG was calculated on a strip of cardiac tissue made up of three types of cells: endocardial, midmyocardial, and epicardial. A distinctive difference among these cells is the maximum value of the channel conductance for the i_{Kr} and the i_{Ks} currents. The conductance ratios for these cells were set at 11:1, 4:1, and 35:1, respectively, following experimentally-obtained values (Viswanathan and Rudy, 2000).

4. Results

The ECG and the AP morphologies are markers of the electrophysiological state of cardiac tissue. The baseline, figure 1 with the insert showing the epicardial AP, was the starting point in these simulations. The parametric values of the substrate were modified within the guidelines of table 1, using data from Carmeliet (1999), Antzelevitch and Zygmunt (2003), Lukas and Antzelevitch (1993), Mejia-Alvarez and Marban (1992), and Ju and Allen (2003), in a series of calculations.




Figure 1. The baseline mathematically-generated ECG of the cardiac tissue with AP shown in the insert.

Figure 2 shows the effect of CN intoxication on the AP of ventricular tissue. Notable are the shortened cycle length (CL), a sign of tachycardia, the lowering of the wave amplitude, and the rise of the resting potential. Figure 3 shows the effect on the ECG, whose morphology has been drastically altered. A T-wave as a separate entity is no longer in evidence, the shape of the QRS portion has been altered, and the peak has been halved. With the chloride current activated, a further deviation of the shape of the wave from the norm occurs (figure 4).




Figure 2. CN notably shortens the CL of affected tissue (blue trace) in comparison with the baseline (red trace).




Figure 3. Simulated ECG (blue trace) of severely CN-affected cardiac tissue. $[Ca^{2+}] = 0.0009$, $[ATP] = 1.0$, and $K^+_o = 12.0$ mM/L. The baseline is shown in red.




Figure 4. When the normoxic dormant currents are activated, further morphological changes in the ECG are evident.

The calculations reproduced aspects of the changes in an ECG of a subject under the effect of a lethal dose of CN (figure 5 [Wexler et al., 1947]). The change in the morphology is reproduced in figure 6.




Figure 5. ECG from a CN-intoxicated individual from Wexler et al. (1947).




Figure 6. The effect of calcium-overload, reduced availability of ATP, and high external potassium concentration.

A series of simulations to determine the effect of the individual parametric changes are summarized in figure 6, concentrating on the effect of intracellular calcium overload, reduced availability of ATP, and high external potassium concentration. The red trace shows the effect of a 50% overload of internal sodium. The blue curve represents potassium and calcium overload. The green ECG trace shows the effect of the activation of the chloride channel under ion concentration overload conditions. The reversal from normal direction of the membrane currents was noted in the data.

Disturbance of the energy homeostasis and ion concentrations in cardiac tissue due to CN results in the reversal of the direction from the normal and change in magnitudes of cellular membrane currents. These, in turn, change the morphology of the AP and the ECG. This is the initial step leading to ventricular fibrillation, the usual endpoint in the effect of CN on the heart. Several of the membrane currents reverse direction under these tissue conditions. The negative trending T-wave in the ECG indicates pathological behavior and abnormal repolarization of the ventricle. CN causes cell swelling and hemorrhaging in cardiac tissue. Cell swelling activates chloride

membrane currents affecting homeostasis of the tissue. These effects were shown to be important for the electrical state of the CN-affected tissue and were included for the first time in a model of CN-affected cardiac tissue.

Pharmacological intervention can reverse the effect of CN intoxication, as shown in figures 7–9. Restoration of the cell's energy source, the ATP concentration, and blocking calcium accumulation within the cell are primary objectives of therapeutic strategies.




Figure 7. The sodium concentrations returned to normal, and, with internal calcium at 0.0004 mM/L, the CL is improved, but tachycardia still is in evidence.




Figure 8. Elimination of the exterior potassium overload helps (green curve), but baseline behavior is not achieved.




Figure 9. Simulated pharmacological intervention: the red trace shows the baseline. The blue trace shows the case with ion concentrations restored except for sodium, showing a 50% cell overload.

5. Discussion

Exposure to CN has immediate consequences on the electrophysiology of the heart. This computational study focused on several of the determining factors characterizing the morphology of the AP and the ECG.

CN changes the energy homeostasis and ion concentrations, and causes the activation of several otherwise dormant membrane currents. These include I_{KATP} , $I_{CL,swell}$, and the throttling of $I_{Ca(L)}$. Their inclusion into the model presented is vital for the simulation of the CN-caused ECG modulation. Considerable refinements in the results were possible by inclusion into the model of the swelling-activated chloride current. The simulation was able to approximate changes observed in the ECG of a subject under CN toxicity. The calculations showed the rise in the resting voltage and demonstrated how it could be restored to baseline. Means of recapturing the width of the AP post-CN-modulation, important for preventing arrhythmia, was also illustrated. These approaches also allow modification of the ST segment and J-wave modulations caused by CN. The ST denivelation seen in the experimental data was clearly reproduced in the model figure 3. The QRS widening and modulation of the wave is also seen in the calculational results (e.g., figure 4). Parametric changes of the substrate, as shown in figure 7, can change the morphology of the AP and the electrophysiology of the tissue.

In silico techniques are powerful ancillary tools for the understanding of the modulation of cellular processes by threat agents. The results of this study narrow the search on the requirements on the means of pharmacological intervention to counter the effect of CN-caused cardiac toxicity. Of special importance may be the restoration of ATP, potassium, and calcium concentrations in the ventricular cells, increase of the level of $I_{Ca(L)}$, and modulating of $I_{CL,swell}$.

6. References

- Antzelevitch, C.; Zygmunt, A. C.; Dumaine, R. Electrophysiology and Pharmacology of Ventricular Repolarization. In *Contemporary Cardiology: Cardiac Repolarization: Bridging Basic and Clinical Science*; Gussak, I. et al., Eds.; Humana Press Inc.: Totowa, NJ, 2003.
- Balaban, R. S. Cardiac Energy Metabolism Homeostasis: Role of Cytosolic Calcium. *J. Mol. Cell. Cardiol.* **2002**, *34*, 1259–71.
- Baskin, S. I.; Brewer, T. G. Cyanide Poisoning. *Medical Aspects of Chemical and Biological Warfare*; Sidell, F. R., Takafuji, E. T., Franz, D. R., Eds.; Borden Institute: Walter Reed Army Medical Center, 1997; pp 271–286.
- Baskin, S. I.; Petrikovics, I.; Kurche, J. S.; Nicholson, J. D.; Logue, B. A.; Maliner, B. I.; Rockwood, G. A. Insights on Cyanide Toxicity and Methods of Treatment. In *Pharmacological Perspectives of Toxic Chemicals and Their Antidotes*; Flora, E. S. J. et al. Eds.; Narosa Publishing House: New Delhi, India, 2004.
- Carmeliet, E. Cardiac Ionic Currents and Acute Ischemia: From Channels to Arrhythmias. *Physiol. Rev.* **1999**, *79*, 917–1017.
- Elliott, A. C.; Smith, G. L.; Allen, D. G. Simultaneous Measurement of Action Potential Duration and Intracellular Atp in Isolated Ferret Hearts Exposed to Cyanide. *Circ. Res.* **1989**, *64*, 583–591.
- Ferrero, J. M., Jr.; Saiz, J.; Ferrero, J. M.; Thakor, N. V. Simulation of Action Potentials From Metabolically Impaired Cardiac Myocytes. *Circ. Res.* **1996**, *79*, 208–221.
- Inoue, M.; Imanaga, I. Activation of Ca^{2+} -Dependent K^{+} Channels by Cyanide in Guinea Pig Adrenal Chromaffin Cells. *Am. J. Physiol.* **1998**, *274* (*Cell Physiol.* *43*), C105–C111.
- Ju, Y. K.; Allen, D. G. Early Effects of Metabolic Inhibition on Intracellular Ca^{2+} in Toad Pacemaker Cells: Involvement of Ca^{2+} Stores. *Am. J. Physiol. Heart Circ. Physiol.* **2003**, *284*, H1087–1094.
- Katzman, G. M.; Penney, D. G. Electrocardiographic Responses to Carbon Monoxide and Cyanide in the Conscious Rat. *Toxicology Letters* **1993**, *69*, 139–153.
- Leimdorfer, A. About Anoxia of the Heart Produced by Intravenous Sodium Cyanide Injections. *Arch. Int. Pharmacodyn.* **1950**, *84*, (2–3), 181–188.
- Lukas, A.; Antzelevitch, C. Differences in the Electrophysiological Response of Canine Ventricular Epicardium and Endocardium to Ischemia. *Circulation* **1993**, *88*, 2903–2915.

- Mejia-Alvarez, R.; Marban, E. Mechanism of the Increase in Intracellular Sodium During Metabolic Inhibition: Direct Evidence Against Mediation By Voltage-Dependent Sodium Channels. *J. Mol. Cell Cardiol.* **1992**, *24*, 1307–1320.
- Plonsey, R.; Barr, R. C. *Bioelectricity*, 2nd ed.; Kluwer: New York, 2000.
- Shaw, R. M.; Rudy, Y. Electrophysiologic Effects of Acute Myocardial Ischemia. *Circ. Res.* **1997**, *80*, 124–138.
- Suzuki, T. Ultrastructural Changes of Heart Muscle in Cyanide Poisoning. *Tohoku J. Exp. Med.* **1968**, *95*, 271–87.
- Vandenberg, J. I.; Bett, G. C. L.; Powell, T. Contribution of a Swelling-Activated Chloride Current to Changes in the Cardiac Action Potential. *Am. J. Physiol.* **1997**, *273*, C541–C547.
- Van der Heyden, G.; Vereecke, J.; Carmeliet, E. The Effect of Cyanide on the K-Current in Guinea-Pig Ventricular Myocytes. *Basic Res. Cardiol.* **1985**, *80* Suppl. 1, 93–96.
- Viswanathan, P. C.; Rudy, Y. Cellular Arrhythmogenic Effects of Congenital and Acquired Long-QT Syndrome in the Heterogeneous Myocardium. *Cir. Res.* **2000**, *101*, 1192–1198.
- Wexler, J.; Whittenberger, J. L.; Dumke, P. R. The Effect of Cyanide on the Electrocardiogram of Man. *Am. Heart J.* **1947**, *34*, 163–173.
- Zoltani, C. K.; Baskin, S. I. *In Silico* Cardiac Toxicity: Increasing the Discovery of Therapeutics Through High-Performance Computing. In *Alternate Toxicological Methods*; Salem, H., Katz, S. A., Eds.; CRC Press: Boca Raton, 2003.
- Zoltani, C. K.; Baskin, S. I.; Platoff, G. E. ECGs and Metabolic Networks: An *In Silico* Exploration of Cyanide-Caused Cardiac Toxicity. In *Pharmacological Perspectives of Some Toxic Chemicals and Their Antidotes*; Flora, S. J. S., Romano, J. A., Baskin, S. I., Eds.; Narosa Publishing House: New Delhi, India, 2004a.
- Zoltani, C. K.; Platoff, G. E.; Baskin, S. I. Acquired Brugada-Like Symptoms of Cyanide-Caused Cardiac Toxicity: A Computational Study. Accepted by *Computers in Cardiology*, 2004b.
- Zoltani, C. K.; Platoff, G. E.; Baskin, S. I. *In Silico* Techniques Aid the Decryption of Cellular Mechanisms of Cyanide-Caused Cardiac Toxicity. *Proc. Bioscience 2004c Medical Defense Review*, submitted for publication, 2004c.

NO. OF
COPIES ORGANIZATION

1 DEFENSE TECHNICAL
(PDF INFORMATION CTR
ONLY) DTIC OCA
8725 JOHN J KINGMAN RD
STE 0944
FORT BELVOIR VA 22060-6218

1 US ARMY RSRCH DEV &
ENGRG CMD
SYSTEMS OF SYSTEMS
INTEGRATION
AMSRD SS T
6000 6TH ST STE 100
FORT BELVOIR VA 22060-5608

1 INST FOR ADVNCD TCHNLGY
THE UNIV OF TEXAS
AT AUSTIN
3925 W BRAKER LN STE 400
AUSTIN TX 78759-5316

1 US MILITARY ACADEMY
MATH SCI CTR EXCELLENCE
MADN MATH
THAYER HALL
WEST POINT NY 10996-1786

1 DIRECTOR
US ARMY RESEARCH LAB
IMNE ALC IMS
2800 POWDER MILL RD
ADELPHI MD 20783-1197

3 DIRECTOR
US ARMY RESEARCH LAB
AMSRD ARL CI OK TL
2800 POWDER MILL RD
ADELPHI MD 20783-1197

3 DIRECTOR
US ARMY RESEARCH LAB
AMSRD ARL CS IS T
2800 POWDER MILL RD
ADELPHI MD 20783-1197

NO. OF
COPIES ORGANIZATION

ABERDEEN PROVING GROUND

1 DIR USARL
AMSRD ARL CI OK TP (BLDG 4600)

NO. OF
COPIES ORGANIZATION

ABERDEEN PROVING GROUND

- | | |
|---|--|
| 8 | DIR USARL
AMSRD ARL CI H
C NIETUBICZ
AMSRD ARL CI HC
R NAMBURU
C ZOLTANI (6 CPS) |
| 4 | CMDR
USAMRICD
PHARMACOLOGY DIV
G E PLATOFF (2 CPS)
S I BASKIN (2 CPS)
3100 RICKETTS POINT RD
APG MD 21020-5400 |