

AD _____

Award Number: DAMD17-01-1-0076

TITLE: Role of the Human Polyomavirus, BKV, in Prostate Cancer

PRINCIPAL INVESTIGATOR: Michael Imperiale, Ph.D.

CONTRACTING ORGANIZATION: The University of Michigan
Ann Arbor, Michigan 48109-1274

REPORT DATE: August 2004

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

20050715 041

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 074-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE August 2004	3. REPORT TYPE AND DATES COVERED Annual (1 Aug 2003 - 31 Jul 2004)	
4. TITLE AND SUBTITLE Role of the Human Polyomavirus, BKV, in Prostate Cancer			5. FUNDING NUMBERS DAMD17-01-1-0076	
6. AUTHOR(S) Michael Imperiale, Ph.D.				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) The University of Michigan Ann Arbor, Michigan 48109-1274 <i>E-Mail:</i> imperial@umich.edu			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012			10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES Original contains color plates: All DTIC reproductions will be in black and white.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited				12b. DISTRIBUTION CODE
13. ABSTRACT (Maximum 200 Words) BKV is a human polyomavirus that establishes a lifelong, persistent infection of the urinary tract. The virus encodes oncoproteins that induce tumors in animal models, and BKV DNA has been detected in human urinary tract tumors, including prostate. Prostate tumors have a relatively low frequency of mutations in the <i>p53</i> and <i>Rb1</i> genes, indicating that an agent such as a virus may be inactivating their functions. The aims of this proposal are to determine if BKV is present in prostate tumors and, if so, whether viral oncogenes are expressed. To accomplish this, normal and tumor tissue from individual patients will be analyzed. PCR, in situ PCR, and in situ hybridization will be performed to determine the presence of viral sequences, and RT-PCR and immunohistochemistry will be used to examine gene expression. Viral sequences from patients will be cloned and their function compared to wild type virus. This past year, we optimized most of these assays and began to analyze samples. We detect the presence and expression of the virus in a subset of normal and cancer cells. At this time, firm conclusions cannot be drawn as more samples need to be analyzed. If BKV is associated with some prostate cancers, our knowledge of the virus will be useful in designing drugs and vaccines for treatment.				
14. SUBJECT TERMS DNA tumor virus, T antigen, polymerase chain reaction, prostate				15. NUMBER OF PAGES 33
				16. PRICE CODE
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

Table of Contents

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	6
Reportable Outcomes.....	6
Conclusions.....	6
References.....	7
Appendices.....	7

INTRODUCTION

BK virus (BKV) is a human polyomavirus that establishes a lifelong, persistent infection of the urinary tract. The virus encodes two oncoproteins, the large T and small t antigens, that have been shown to cause cancer in animal models, and recently BKV DNA has been detected in various urinary tract tumors including tumors of the prostate. Among the spectrum of human cancers, prostate tumors have a relatively low frequency of mutations in the *p53* and *RB1* genes, indicating that an outside agent such as a virus may be inactivating the function of their gene products. Both large T and small t antigens are capable of doing so. The aims of this proposal are to determine if BKV is indeed present in these tumors and, if so, whether these oncogenic proteins are expressed and function to deregulate cell growth control.

BODY

During the third year of this project we have made significant progress on our analysis of tumor samples. Our progress is described below after the appropriate items from the approved Statement of Work. Please note that rather than reproducing figures from our manuscript in press in *Oncogene* in the body of this report, I have referred to them instead. The manuscript is attached as an Appendix.

Task 1. To examine prostate tumors and matched normal tissue for the presence of BKV sequences (months 1-30).

- isolate DNA and RNA from microdissected normal and tumor tissue (months 1-6)
- perform PCR analysis of DNA (months 3-18)
- perform IS-PCR and ISH analysis of DNA (months 3-24)

This Task is almost complete in that we are now assaying tissue microarrays to boost the number of samples which we have analyzed. In the tissues that we have examined, however, we now have clear evidence for the presence of BKV in both normal ducts and in atrophic epithelium. This has been shown both by solution PCR and by in situ hybridization, both using probes that are highly specific and sensitive for BKV. Thus, the extra time it took us to develop these assays at the beginning of this project has paid off in that our results are very clean and unambiguous. These results are shown in Figures 2 and 4 of the *Oncogene* manuscript, and are also included in the summary (Table 1).

- perform sequence analysis of DNA (months 6-24)

The DNA sequence analysis is shown in Figure 3 of the manuscript. The analysis of the T antigen coding region (early region) proves that the virus we are detecting is BKV and not SV40 or JCV, which are highly homologous to BKV. In addition, the sequence analysis of the regulatory regions indicates that at least three different strains of virus are present in the samples. Indeed, since submitting this manuscript we have performed additional analyses which indicate the presence of at least one other strain in prostate tumors. Of interest is that we do not detect the archetype strain, which is the strain that is thought to remain latent in the urinary tract. Rather, we find strains that are associated with acute disease. Moreover, the fact that we find strains that

we do not study in the laboratory argues that we have not inadvertently contaminated our samples with laboratory virus.

- perform RT-PCR and IS-RT-PCR analysis of RNA (months 12-30)

Having obtained a good handle on the status and sequence of the viral DNA, during the past few months we have turned our attention to the RNA. Again, we have taken our time to carefully determine the best conditions for isolation of the RNA and its amplification for sequencing. We initially have concentrated on the *p53* gene since we would predict that it would not be mutant in the BKV-positive cells. This is because the presence of T antigen is equivalent to mutations due to its inactivation of the *p53* protein. We began by trying to amplify the entire *p53* cDNA in a single reaction. However, this was not possible, probably due to the nature of the RNA extracted from paraffin-embedded tissue. Therefore, we are now amplifying subregions of the mRNA, concentrating on exons 5-9, in which most *p53* mutations are found. Rather than sequence the products from RNA isolated from bulk sections, however, we have decided to use laser capture microdissection to purify the cells that are virus-positive, as judged by in situ hybridization and immunohistochemistry (see below). This is because we desire to only analyze the *p53* from the affected cells (i.e., cells in the normal and abnormal ducts) and not the total *p53* mRNA, which would include undoubtedly wildtype, no matter what the pathology, derived from the stroma. We believe that the sensitivity of our RT-PCR reaction has reached the point where we can now perform it on the small number of cells we will obtain by laser capture.

As we discussed in the past two years' reports, we have added immunohistochemical assays for T antigen to this task. All samples have been analyzed by this method and a fraction of the samples are positive for T antigen expression. Interestingly, while both normal and atrophic epithelia are positive for viral DNA, only the atrophic lesions express T antigen (Figure 5 of the manuscript). Also of interest is the finding that the T antigen is cytoplasmic. This indicates that we are not detecting viral replication, which would require T antigen to be present in the nucleus. We have also stained for *p53* protein and in the T antigen positive cells, the *p53* co-localizes to the cytoplasm (Figure 6 and 7), indicating that its function has been compromised: as a transcription factor, it must be in the nucleus to function. In those samples that are T antigen negative, the *p53* is nuclear. As discussed above, we are very interested in examining the sequence of the *p53* gene in these different samples.

Task 2. To determine how alterations in viral sequences in tumors affect the replication and transformation properties of the virus (months 9-36)

- construct genomic viral clones containing patient-derived sequences (months 9-24)
- determine the ability of these clones to replicate in permissive host cells (months 12-36)
- construct early region expression vectors containing patient-derived sequences (months 9-24)
- determine the ability of these clones to interfere with normal growth control pathways involving pRb and *p53* (months 18-36)

We have begun to clone the regulatory regions from the patient derived materials. We have also developed an in vitro cell culture system, using primary renal proximal tubule epithelial cells, to study BKV replication in its natural host cell (*Virology* manuscript by Low, et al., see Appendix). While this was not originally proposed in the application, we found it necessary to develop this system as we found other cell lines suboptimal for studying the viral life cycle. We will begin examining the function of the patient isolates very shortly, which is one of the reasons we have requested and been granted a no-cost time extension for the project.

KEY RESEARCH ACCOMPLISHMENTS

- detection of BKV in normal and atrophic prostatic epithelia
- detection of BKV T antigen in atrophic cells
- co-localization of p53 with T antigen in cells
- sequence analysis of BKV isolates from prostate
- development of in vitro system for assaying BKV replication in its natural host cell

REPORTABLE OUTCOMES

Two abstracts were presented at international meetings this year:

1. Oral presentation by Dr. Das:

Das, D., Shah, R., and Imperiale, M.J. (2004). Detection and expression of human BK virus sequences in neoplastic prostate tissues. *Molecular Biology of DNA Tumor Viruses Conference*, Madison, WI.

2. Poster presentation by Ms. Abend:

Abend, J., Low, J., and Imperiale, M.J. (2004). BKV and SV40 interactions with human kidney proximal tubule epithelial cells. *Molecular Biology of DNA Tumor Viruses Conference*, Madison, WI.

Two manuscripts resulted from this work:

Low, J., Humes, H.D., Szczypka, M., and Imperiale, M.J. (2004). BKV and SV40 infection of human kidney tubular epithelial cells in vitro. *Virology* 323:182-188.

Das, D., Shah, R.B., and Imperiale, M.J. (2004). Detection and expression of human BK virus sequences in neoplastic prostate tissues. *Oncogene*, in press.

CONCLUSIONS

BKV is associated with atrophic lesions of the prostate, which are thought to be the first step in the progression from normal to cancerous epithelium. Viral sequences are present in both normal and atrophic epithelial cells, and viral T antigen, the transforming protein of the virus, is expressed in the latter cells. Moreover, the T antigen co-localizes in the cytoplasm with p53, indicating that the virus is not replicating and the p53 cannot function as a tumor suppressor. The

future goals of this work are to further analyze the BKV-positive cells in an attempt to collect additional evidence that the virus is indeed contributing to neoplastic progression.

REFERENCES

none

APPENDICES

Copies of abstracts and manuscripts reported above in the Outcomes section.

BKV and SV40 Interactions with Human Kidney Proximal Tubule Epithelial Cells

Johanna Abend, Jonathan Low, and Michael Imperiale

University of Michigan Medical School

We have previously described human proximal tubule epithelial (HPTE) cells as a model system to study BKV in an environment similar to that which the virus encounters in an immunocompromised host. HPTE cells regulate blood pressure and volume, return soluble molecules to the bloodstream, secrete vitamin precursors, and mediate immune responses to the kidney through the secretion of cytokines and chemokines. The HPTE cells used in our assays are able to maintain their differentiated structure and function for up to six passages *in vitro*. BKV undergoes a lytic infection in these cells, while SV40 cannot enter them. SV40 is capable of lytic infection after the addition of the ganglioside GM1, recently shown to be an SV40 receptor. We have now elucidated additional interactions between the virus and the cell. Upon infection, BKV downregulates the expression of a set of inflammatory molecules expressed by the HPTE cells. We also find that infecting HPTE cells in the presence of IFN- γ downmodulates the expression of T antigen. These studies suggest that inflammatory molecules produced by HPTE cells may play a role in keeping BKV in a subclinical state either through direct suppression of viral genes, or through the recruitment of immune cells to the site of BKV infection. Upon infecting HPTE cells with SV40 in the presence of GM1, we observe that while the majority of cells are destroyed due to a lytic infection, a subset of cells continues to grow past the point at which uninfected HPTE cells senesce. These cells undergo changes in morphology and produce infectious SV40.

**BKV and SV40 Infection of Human Kidney Tubular Epithelial Cells
*In Vitro***

**Jonathan Low, David Humes, Mark Szczypka, and Michael
Imperiale, Department of Microbiology and Immunology University
of Michigan Medical School, Michigan, USA, 48108**

Infection with BK virus (BKV) is nearly ubiquitous in the human population. Primary infection with BKV most often occurs as an asymptomatic infection during childhood followed by a persistent or latent infection of the kidney that can be reactivated upon immunosuppression, often leading to hemorrhagic cystitis or nephritis. Previous studies have found BKV to be tumorigenic in rodents, and there have been reports of the presence of BKV sequences in various human tumors. Little is known about the interaction of BKV with its natural target cells, human kidney epithelial cells. *In vitro* infections of human primary kidney epithelial cells were performed to simulate a BKV infection in its natural host cell. BKV goes through a lytic replication cycle in this system: high levels of T antigen expression were first detected at 36 hours post-infection, while viral DNA replication and capsid protein expression were observed at 48 hours post-infection. A corresponding production of infectious BKV was also seen at 48 hours post-infection. Levels of p53 in the infected cells increased, and hyperphosphorylated forms of pRb appeared. SV40 causes a similar disease in monkeys and is believed to be able to infect humans. It was observed that SV40 is incapable of infecting human kidney epithelium except in the presence of the GM1 ganglioside, recently reported to be an SV40 receptor.

This work was supported by Department of Defense CDMRP grant PC001595.

Jonathan Low, (734) 936-0876, X5

Detection and Expression of Human BK Virus Sequences in Neoplastic Prostate Tissues

Dweeapanita Das¹, Rajal B. Shah^{2,3}, and Michael J. Imperiale*,^{1,3}

¹Department of Microbiology and Immunology, ²Departments of Pathology and Urology, and ³Comprehensive Cancer Center, University of Michigan Medical School, Ann Arbor, Michigan 48109-0942

BK virus is ubiquitous in the human population and establishes a lifelong, subclinical persistent infection in the urinary tract. When the immune system is compromised, it can cause severe disease in the kidney and bladder. Detection of BKV sequences in urinary tract neoplasms has led to the postulation that this virus may induce human oncogenesis through the function of its large T antigen (TAg). In this study, examination of prostate tumor tissue sections using in situ hybridization shows the presence of BKV sequences in benign and atrophic epithelium. Solution PCR on DNA extracted from the tissues and sequence analysis of the products reveal the presence of BKV regulatory and early region sequences. Sequence analysis of the regulatory region reveals the presence of the TU strain in most of the samples, along with the wt-501 and Dunlop strains to a lesser extent. Some of the regulatory regions showed patchy homology to the Gardner, renal and urinary tract isolates of BKV. Additionally, immunohistochemical analysis using monoclonal antibodies specific to TAg or p53 shows the expression of TAg in some of the samples and p53 staining which can be correlated to TAg expression. Though the normal cellular localization of TAg and p53 is nuclear, double immunofluorescence labeling with p53 and TAg antibodies indicates co-localization of p53 and TAg to the cytoplasm in the glandular epithelial cells of the sections. Although BKV DNA was found in benign and atrophic lesions, TAg and p53 co-expression was observed only in atrophic lesions. The implications of these results as they relate to a possible role for BKV in prostate cancer will be discussed.



ORIGINAL PAPER

Detection and expression of human BK virus sequences in neoplastic prostate tissues

Dweepanita Das¹, Rajal B Shah^{2,3} and Michael J Imperiale^{*1,3}¹Department of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor, MI 48109-0942, USA;²Departments of Pathology and Urology, University of Michigan Medical School, Ann Arbor, MI 48109-0942, USA; ³Comprehensive Cancer Center, University of Michigan Medical School, Ann Arbor, MI 48109-0942, USA

BK virus (BKV) is ubiquitous in the human population and establishes a lifelong, subclinical persistent infection in the urinary tract. When the immune system is compromised, it can cause severe disease in the kidney and bladder. Detection of BKV sequences in urinary tract neoplasms has led to the postulate that this virus may induce human oncogenesis through the function of its large tumor antigen (TAG). In this study, examination of prostate tumor tissue sections using *in situ* hybridization shows the presence of BKV sequences in atrophic epithelium. Solution polymerase chain reaction on DNA extracted from the tissues and sequence analysis of the products reveal the presence of BKV regulatory and early region sequences. In addition, immunohistochemical analysis using monoclonal antibodies specific to TAG or p53 shows the expression of TAG in some of the samples and p53 staining that can be correlated to TAG expression. Although the normal cellular localization of TAG and p53 is nuclear, double immunofluorescence labeling with anti-p53 and TAG antibodies indicates colocalization of p53 and TAG to the cytoplasm in the glandular epithelial cells of the sections. Although BKV DNA was found in benign and atrophic lesions, TAG and p53 coexpression was observed only in atrophic lesions.

Oncogene advance online publication, 19 July 2004; doi:10.1038/sj.onc.1207920

Keywords: polyomavirus BK; large T antigen; p53; postatrophic hyperplasia; proliferative inflammatory atrophy; *in situ* hybridization; immunohistochemistry; immunofluorescence; high-grade prostate intraepithelial neoplasia

Introduction

Prostate cancer (PCA) is the second leading cause of cancer deaths in older men in the US (Gonzalzo and Isaacs, 2003; Jemal *et al.*, 2003). In spite of the

availability of biomarkers for PCA, the basic molecular mechanisms regulating its development and progression are still very poorly understood (Nupponen and Visakorpi, 1999; Jiang *et al.*, 2001; Kuefer *et al.*, 2002; Rubin *et al.*, 2002; Varambally *et al.*, 2002; Yang *et al.*, 2002). Recognition of the precursor lesions of PCA has become a primary focus in recent years. High-grade prostate intraepithelial neoplasm (HGPIN), characterized by proliferation of epithelial cells in ducts and acini (Bostwick and Brawer, 1987), has been reported to be the most likely precursor lesion for PCA in the US (Myers *et al.*, 1994; Bostwick, 1995; reviewed in Gonzalzo and Isaacs, 2003). Postatrophic hyperplasia (PAH), also referred to as proliferative inflammatory atrophy (PIA), is a proliferative atrophic lesion characterized by a decrease of cytoplasmic mass as compared to benign glands. Atrophic epithelium in PAH has recently been proposed to be a precursor to PIN and PCA (De Marzo *et al.*, 1999; Putzi and De Marzo, 2000; Shah *et al.*, 2001; reviewed in De Marzo *et al.*, 2003; Nelson *et al.*, 2003). Several studies have identified focal areas of epithelial atrophy and have suggested their importance in prostatic carcinogenesis (Franks, 1954; Gardner and Bennett, 1992; De Marzo *et al.*, 2003; Nelson *et al.*, 2003). It has also been reported that atrophy of prostatic glands is common in middle-aged and elderly men (Gardner and Culbertson, 1987; McNeal, 1988).

The role of tumor suppressor genes (*p53* and *Rb1*) in PCA is not very clear. Mutations in the *Rb1* gene are infrequent in PCA (Bookstein *et al.*, 1990; Sarkar *et al.*, 1992; Melamed *et al.*, 1997; reviewed in Abate-Shen and Shen, 2000). Although the mutation rate of the *p53* gene in PCA is also low, *p53* inactivation has been implicated in PCA progression (Bookstein *et al.*, 1993; Bauer *et al.*, 1995; Brewster *et al.*, 1999; Osman *et al.*, 1999; reviewed in Abate-Shen and Shen, 2000). These observations have led researchers to speculate that an infectious agent might play a role in PCA (McNicol and Dodd, 1991; Ibrahim *et al.*, 1992; Rotola *et al.*, 1992; Cuzik, 1995; Noda *et al.*, 1998). Polyomaviruses are candidates for this agent, although the etiological role of polyomaviruses in human tumors remains debated despite several reports demonstrating the presence and expression of sequences from these viruses (Imperiale, 2000, 2001; Replogue *et al.*, 2001; Tognon *et al.*, 2003).

*Correspondence: MJ Imperiale, Department of Microbiology and Immunology, University of Michigan Medical School, 1500 E. Medical Center Dr. 6304 Cancer Center, Ann Arbor, MI 48109-0942, USA; E-mail: imperial@umich.edu

Received 28 February 2004; revised 25 May 2004; accepted 9 June 2004

BK virus (BKV), a polyomavirus, was first isolated from the urine of a renal transplant patient (Gardner *et al.*, 1971). It persistently infects almost 100% of the human population by early childhood (Gardner, 1973; Padgett, 1981; Dörries *et al.*, 1994). It resides in the kidneys in a latent or persistent state, but can be reactivated upon immunosuppression of the host and is associated with hemorrhagic cystitis and polyomavirus nephropathy (Arthur *et al.*, 1986; Hiraoka *et al.*, 1991; Pappo *et al.*, 1996; Randhawa *et al.*, 1999; Li *et al.*, 2002). BKV oncogenically transforms rodent cells in culture, causes kidney tumors in transgenic mice, and transforms primary human cells in culture when coexpressed with an activated *ras* oncogene (Portolani *et al.*, 1975; Pater and Pater, 1986; Small *et al.*, 1986;

Dalrymple and Beemon, 1990). BKV has been reported to be detected in a number of human tumors, including those in the urinary tract, brain, pancreatic islet, lung, eye, liver, Kaposi's sarcoma, and bone (reviewed in Imperiale, 2000, 2001; Reploeg *et al.*, 2001; Tognon *et al.*, 2003).

The BKV genome is divided into regulatory, early, and late regions, and encodes five major proteins (Figure 1; Seif *et al.*, 1979; Frisque *et al.*, 1984; Pipas, 1992). The early proteins, encoded by alternatively spliced mRNAs, are the large tumor antigen (TAg), the main regulatory protein, and the supporting small tumor antigen. TAg aids in the oncogenic transformation of cells through its ability to bind and inactivate the products of tumor suppressor genes, including p53 and




Figure 1 Schematic of BKV genome and oligonucleotides and primers used in the study. (a) Schematic of BKV genome depicting the early and late regions, with the regulatory region shown as a gray rectangle. ORI: Origin of replication. The schematic to the right depicts major viral genes. The angled lines represent introns. The hatched sequences are common to TAg and tAg. Unique sequences for tAg and TAg are represented by bricked and lined sequences, respectively. The dotted sequence represents VP1. Identical sequences for VP2 and VP3 are denoted by the checkered box, whereas VP2 unique sequence is represented by the black box. (b) Schematic of the early region. Boxes, protein coding sequences; V-shape, intron. Numbered arrows refer to the indicated primers and oligonucleotide probe specific to BKV. (c) Schematic of the regulatory region. Numbered arrows refer to the indicated primers

the pRb family (Nakashatri *et al.*, 1988; Dyson *et al.*, 1989, 1990; Kang and Folk, 1992; Simmons, 1995; Staib *et al.*, 1996; Shivakumar and Das, 1996; Harris *et al.*, 1996, 1998a, b). TAg also regulates viral DNA replication and gene expression, and coregulates activated and basal cellular transcription factors (Moens and Rekvig, 2001). Small tumor antigen is also involved in transformation and viral and cellular DNA replication (Moens and Rekvig, 2001). The late proteins, the capsid proteins VP1, VP2, and VP3, are also expressed from alternatively spliced mRNAs, after viral DNA replication. The regulatory region contains promoter/enhancer elements of the early and late genes and the origin of DNA replication (Imperiale, 2001; Reploeg *et al.*, 2001; Hirsch and Steiger, 2003). Different BKV strains show sequence variability in their regulatory regions. These strains derive from a common archetype strain, which infects the general population, due to deletions, duplications, and rearrangements in the regulatory region (Rubinstein *et al.*, 1987; Shah, 1996).

The role of BKV in human neoplasia is not clear, due to conflicting reports. In the urinary tract, Monini *et al.* (1995) demonstrated the presence of BKV sequences by polymerase chain reaction (PCR) in more than 50% of both normal and tumor tissues, including prostate. Viral DNA could be detected in some tumors by Southern blotting, whereas PCR amplification was always required to find BKV in normal tissues, and further analysis revealed that viral sequences were integrated into the host chromosome in the six tumors that were positive by Southern blotting. Sequence analysis of the viral origin of replication discovered the presence of point mutations in one of the samples, defining it as a new strain of BKV, designated as URO1. It has been shown in cell culture that similar mutations impair the replication ability of the virus, leading to an increased transformation potential (Shinohara *et al.*, 1993). Hence, it was speculated that mutant genomes impaired for replication might integrate into the host genome and eventually lead to cancer. Another report found high levels of BKV TAg expression in primary and metastatic bladder carcinoma but not in non-neoplastic urothelium (Geetha *et al.*, 2002). In these studies, BKV TAg and p53 were colocalized to the nuclei of tumor cells. Given the results obtained with prostate and bladder carcinoma, along with kidney being the main site for BKV persistence, tumors of the urinary tract are the most logical target sites for an etiological association with BKV. Since p53 and Rb1 mutations are not ubiquitous in PCA, we reasoned that TAg might inactivate their function in a subset of tumors, similar to the role of the human papillomavirus E6 and E7 gene products in cervical carcinomas (Scheffner *et al.*, 1991; Ibrahim *et al.*, 1992; Iwasaka *et al.*, 1993; Cuzik, 1995). Hence, we were interested in screening prostatic carcinomas for BKV sequences and, if present, in determining if DNA could be correlated to protein expression.

In this report, we demonstrate the presence of BKV sequences in the epithelium of benign ducts and atrophic lesions of neoplastic prostate tissues using *in situ* and solution analysis. The primers and probe used in our

studies are BKV specific and do not crossreact with JCV or SV40. Using immunohistochemistry, we also detected the expression of TAg in the atrophic lesions. Moreover, TAg expression could be correlated to p53 expression in this set of samples. Interestingly, double immunofluorescence labeling studies demonstrate that TAg and p53 colocalize to the cytoplasm, but in the TAg-negative tumors, p53 is nuclear.

Results

Before beginning to analyse resection specimens (patient samples) for the presence of BKV, we developed PCR and *in situ* hybridization (ISH) conditions and reagents that would have a high degree of specificity and sensitivity for BKV and not the related viruses, SV40 and JCV, which share about 69–75% sequence identity (Seif *et al.*, 1979; Frisque *et al.*, 1984; Pipas, 1992). To demonstrate this specificity, PCR amplification of plasmids containing SV40, JCV, and BKV sequences was performed (Figure 2a). Using the early region primers, BKV_{for}(4935–4907) and BKV_{rev}(4358–4382), a product of 577 base pair (bp) was observed with the BKV plasmid and no product was detected with SV40 or JCV. A Southern blot of the products using the BKV2_(4434–4478) probe, which was also used for ISH studies, confirmed the sequence-specific amplification of the BKV early region. In these reactions, we could detect as little as 0.1 fg of BKV DNA (data not shown). Similarly, both the nested primers to the early region and the regulatory region were determined to be sequence specific, although the regulatory region assay is less sensitive (data not shown). A direct Southern blot of BKV, JCV, and SV40 plasmids with the BKV2 probe confirms its specificity (Figure 2b).

Having developed specific assays, we began analysing patient samples. In all, 21 neoplastic adenocarcinoma prostate tissue samples were selected for evaluation (Table 1). To test for the presence of BKV sequences, DNA extracted from formalin-fixed paraffin-embedded samples was analysed by solution PCR using regulatory and early region primers. Extreme care was taken to avoid sample contamination: plasmid controls were not analysed with the tissue samples, different sets of pipettors were used for the tissues and plasmids, and high-quality DNA grade water was used. The integrity of the DNA was evaluated by amplification of the β -globin gene. Nested amplification was employed for the regulatory and early regions since it was possible that the copy number of the viral genome would be low. First-round amplification was performed for 30 cycles for the early region, after which a product was detectable only by Southern blot (data not shown); after a second round (30 cycles) of amplification with nested primers, enough product resulted for direct detection by ethidium bromide staining. For the regulatory region, two rounds of 40 cycles were required to detect the PCR product due to the lower sensitivity of this assay. Representative EtBr-stained agarose gels of




Figure 2 Detection of BKV DNA in human PCA. (a) Top panel is an EtBr-stained gel showing the PCR amplification products of SV40, JCV, or BKV plasmids (left to right; 10 ng, 1 ng, 100 pg, 10 pg, and 1 pg of each plasmid) with BKV_{for}(4935-4907) and BKV_{rev}(4358-4382) primers. Control lane (c) represents PCR reaction using water as template. Bottom panel is a Southern blot of the same gel probed with the BKV2₍₄₄₃₄₋₄₄₇₈₎ probe. Asterisk represents the hybridization of the probe with the BKV PCR product. (b) Direct Southern blot of SV40, JCV, or BKV plasmid (200, 100, 50, and 25 ng, left to right) using BKV2 probe. (c) EtBr-stained gels showing the PCR products of prostate DNA amplified with β -globin primers (top) and early region primers (bottom). The control lane is from a reaction using water as the template. The sizes of the PCR products are indicated. (d) EtBr-stained gels showing the PCR products of prostate DNA amplified with β -globin primers (top) and regulatory region primers (bottom). The control lane is from a reaction using water as the template

the PCR amplification products of DNA are shown in Figures 2c and d and the results from an average of at least three independent amplifications are summarized in Table 1. A correct size early region product of 452 bp was obtained in 12/16 of the samples every time they were tested. Some samples, such as 4, 5, and 13, were

positive for the early region all but one time. Of the samples, 13 yielded PCR products of approximately 300 bp with the regulatory region primers, of which 11 were positive each time they were assayed. The product size for the regulatory region is variable due to possible rearrangements. Seven of the regulatory regions were

Table 1 Data summary

Sample	Gleason score	PCR		ISH		IHC
		Early	Regulatory ^a	BKV	TAg	p53 ^b
11.	7	+	+TU	+	+	+
12	7	+	+TU	+	+	+
1	7	+	-	+	+	+
2	7	+	+ND	+	+	+
16	7	+	+ND	+	+	+
13	7	+ ^c	+ ^a ND	+	+	+
17	7	ND	ND	+	+	+
14	9	+	-	+	-	-
15	7	+	-	+	-	-
7	7	+	+TU	+	-	+(N)
8	7	+	+TU	+	-	-
3	9	+	+ND	+	-	-
4	9	+ ^c	+ND	-	-	-
5	7	+ ^c	+TU, wt-501	-	-	+(N)
6	7	+	+TU	-	-	-
10	7	+	+ND	-	-	-
9	7	-	+ ^a Du	-	-	+(N)
18	7	ND	ND	-	-	-
19	7	ND	ND	+	+	+
20	7	ND	ND	+	+	ND
21	6	ND	ND	+	-	-

The samples have been listed in groups based on staining patterns. ^aStrain homology is indicated after + symbol. ND, sequence not determined. ^b-, negative; +, positive and cytoplasmic; + (N), positive and nuclear. ^cSample was PCR positive with early region primers four out of five times. ^dSample was PCR positive three out of four times. ^eSample was PCR positive with early region primers and β -globin primers in the subsequent extraction from that shown in Figure 2. ND, DNA unable to be recovered

confirmed to be BKV by sequence analysis (see below). Sample 9 was positive for the early region in three out of six amplifications and was positive for the regulatory region.

To confirm the identity of the PCR products, they were sequenced (Figure 3). Identical results were obtained from all the early region samples and the sequences were identical to the published BKV genomic sequence (Figure 3a). The majority of the regulatory regions showed identity or similarity to the TU strain (Sundsfjord *et al.*, 1990), while sample 9 appears to be most similar to Dunlop (Figure 3b). Interestingly, one sample (#5) appeared to have two different regulatory regions, one identical to TU and the other most similar to wt-501 (Watanabe and Yoshiike, 1989). We were unable to obtain clean sequencing data from some of the regulatory region products. Products that were not able to be sequenced directly were cloned into a plasmid vector and sequenced, but these sequences did not appear to be homologous to BKV. Thus, 44% of the samples contained regulatory regions that could be confirmed to be BKV.

We wished to use an independent means of detecting BKV DNA in these samples. We chose ISH using the BKV2 probe. Control experiments on BSC-1 (TAG-negative) and BSC-BKT (TAG-positive) cells demonstrate the specificity of the ISH conditions (Figure 4). Panels C-G show ISH on a representative prostate sample. This section, which was positive for BKV by

PCR, also showed significant hybridization to the BKV2 probe as compared to scrambled probe, which has the same G + C content as the BKV2 probe. The signal was localized specifically to the nuclei of epithelial cells in the benign ducts (panel G) and atrophic lesions (panels D and E). We did not detect any signal in stromal cells or in invasive tumor cells. Breast carcinoma tissue sections processed in the same manner as the prostate tissues did not react with this probe (panel H). The ISH studies revealed BKV sequences in 15/21 of the prostate samples.

We next examined the expression of TAg in these samples using immunohistochemical (IHC) labeling. The absence of TAg would be a strong evidence against a role for BKV in oncogenesis. We first examined BSC-1 and BSC-BKT cells as controls (Figure 5). TAg was localized to the nucleus in BSC-BKT cells, whereas BSC-1 cells were negative for TAg. IHC labeling of prostate samples revealed the expression of TAg in the epithelial cells of atrophic lesions but not in tumor or normal epithelium, or stromal cells (panels D and E). Surprisingly, the TAg appeared to be localized to the cytoplasm/perinuclear compartment. A serial section probed with an isotype-matched control antibody gave no signal (panel C). To ensure that the cytoplasmic staining was not an artifact, sections were also probed with anti-androgen receptor antibody, which gave nuclear staining (panel F). In addition, breast carcinoma sections showed no signal with anti-TAg (panel G), while a monoclonal antibody to a 90 kDa tumor-associated glycoprotein expressed in mammary carcinomas gave nuclear staining in the breast samples (data not shown) (Colcher *et al.*, 1981). The IHC labeling results reveal that 9/21 of the prostate tissue sections are positive for TAg expression, that the TAg is mostly cytoplasmic, and that it is expressed in atrophic lesions.

Next, we were interested in examining the prostate samples for the expression of p53, which becomes stabilized and inactivated on association with viral oncoproteins including TAg (Lane and Crawford, 1979; reviewed in Pipas and Levine, 2001). Anti-p53 staining results fall into three categories (Figure 6): TAg and BKV DNA negative (nuclear p53); p53 and TAg positive (cytoplasmic/perinuclear localization of both proteins in atrophic epithelium, panels D and E); and TAg negative but BKV DNA positive (nuclear p53 in atrophic prostate gland; panels F and G). TAg-negative BSC-1 cells stain weakly for p53, whereas the TAg-positive BSC-BKT show nuclear staining (panels A and B). Breast carcinoma tissue sections had robust p53 staining (panel H) (Cattoretti *et al.*, 1988; Moll *et al.*, 1992), and isotype-matched control antibodies did not stain the prostate (panel C). We did not detect any p53 staining in stromal cells or normal epithelium. Since p53 and TAg expression could be correlated, with what appeared to be identical staining patterns, it was of interest to examine p53 and TAg colocalization using double immunofluorescence labeling. The results show staining in the epithelial cells of the atrophic lesions, with cytoplasmic colocalization of both proteins (Figure 7).

a									
Sample 12	-----GCTGTTG	CTTCTTCATCACTGG	CAAACATATCTTCAT	GGCAAAATAAATCTT	CATCCCATTTTTTCAT	TAAAGGAACCTCCACC			
Sample 7	-----GCTGTTG	CTTCTTCATCACTGG	CAAACATATCTTCAT	GGCAAAATAAATCTT	CATCCCATTTTTTCAT	TAAAGGAACCTCCACC			
Sample 6	-----GCTGTTG	CTTCTTCATCACTGG	CAAACATATCTTCAT	GGCAAAATAAATCTT	CATCCCATTTTTTCAT	TAAAGGAACCTCCACC			
BKV	GAGAAATCTGCTGTTG	CTTCTTCATCACTGG	CAAACATATCTTCAT	GGCAAAATAAATCTT	CATCCCATTTTTTCAT	TAAAGGAACCTCCACC			
JCV	---GGGATCTGCTGT	TTTCATCATCACTGG	CAAACATTTCTTCAT	GGCAAAACAGGTCTT	CATCCCACTTCTCAT	TAAATGTATTCCACC			
SV40	GAGAGTCAGCAGTAG	CCTCATCATCACTAG	ATGGCATTTCTTCG	AGCAAAACAGGTITT	CCT-----CAT	TAAAGGCAITCCACC			
Sample 12	AGGACTCCCACTCTT	CTGTTCCATAGGTTG	GCACCTATAAAAAAA	ATAATTACTTAGGGC	CTTTTAATATTTTAT	TATTTATCTAAATAT			
Sample 7	AGGACTCCCACTCTT	CTGTTCCATAGGTTG	GCACCTATAAAAAAA	ATAATTACTTAGGGC	CTTTTAATATTTTAT	TATTTATCTAAATAT			
Sample 6	AGGACTCCCACTCTT	CTGTTCCATAGGTTG	GCACCTATAAAAAAA	ATAATTACTTAGGGC	CTTTTAATATTTTAT	TATTTATCTAAATAT			
BKV	AGGACTCCCACTCTT	CTGTTCCATAGGTTG	GCACCTATAAAAAAA	ATAATTACTTAGGGC	CTTTTAATATTTTAT	TATTTATCTAAATAT			
JCV	AGGATTCOCATTCAT	CTGTTCCATAGGTTG	GCACCTATAAAAAAA	CAATTAAGTTTATTG	TAAAAACAAAATGC	CCTGCAAAAGAAAAA			
SV40	ACTGCTOCCATTCAT	CAGTTCCATAGGTTG	GAATCTA-AAATACA	CAACAATTAGATATC	AGTAGITTAAACACAT	TATACACTTAAAAAT			
Sample 12	AAGTTAGTTAOCCTTA	AAGCTTTAGATCTCT	GAAGGGAGTTTCTCC	AATTATTGGACCCA	CCATTGCAGAGTTTC	TTCACTTAGGTCCTAA			
Sample 7	AAGTTAGTTAOCCTTA	AAGCTTTAGATCTCT	GAAGGGAGTTTCTCC	AATTATTGGACCCA	CCATTGCAGAGTTTC	TTCACTTAGGTCCTAA			
Sample 6	AAGTTAGTTAOCCTTA	AAGCTTTAGATCTCT	GAAGGGAGTTTCTCC	AATTATTGGACCCA	CCATTGCAGAGTTTC	TTCACTTAGGTCCTAA			
BKV	AAGTTAGTTAOCCTTA	AAGCTTTAGATCTCT	GAAGGGAGTTTCTCC	AATTATTGGACCCA	CCATTGCAGAGTTTC	TTCACTTAGGTCCTAA			
JCV	TAGTGGTTTAAOCCTTA	AAGCTTTAGATCCCT	GTAGGGGGTGTCTCC	AAGAATTTCTCCCA	GCAATGAAGAGCTTC	TTGGGTAAAGTCACA			
SV40	TTTATATTAAOCCTTA	GAGCTTTAAATCTCT	GTAGTAGTTTGTCTCC	AATTATGTCACCCA	CAGAAGTAAGGTTCC	TTCACTTAGGTCCTAA			
Sample 12	GCACAAACCACTGTGT	GAAGCAGTCAATGCA	GTAGCAATCTATCCA	AACCAAGGGCTCTTT	TCTTAAAAATTTTCT	ATTTAAATGCCTTAA			
Sample 7	GCACAAACCACTGTGT	GAAGCAGTCAATGCA	GTAGCAATCTATCCA	AACCAAGGGCTCTTT	TCTTAAAAATTTTCT	ATTTAAATGCCTTAA			
Sample 6	GCACAAACCACTGTGT	GAAGCAGTCAATGCA	GTAGCAATCTATCCA	AACCAAGGGCTCTTT	TCTTAAAAATTTTCT	ATTTAAATGCCTTAA			
BKV	GCACAAACCACTGTGT	GAAGCAGTCAATGCA	GTAGCAATCTATCCA	AACCAAGGGCTCTTT	TCTTAAAAATTTTCT	ATTTAAATGCCTTAA			
JCV	CCCAAAACCATTTGCT	GAAGCAATCAAGCA	ATAGCAATCTATCCA	CACAAGTGGGCTGCT	TCTTAAAAATTTTCT	GTTCCTATGCCTTAA			
SV40	TCCAAACCACTTCT	AAAGCAATCGAAGCA	GTAGCAATCAAGCCA	CACAAGTGGATCTTT	CCTGTATAATTTTCT	ATTTTCATGCTTCAT			
Sample 12	TCTAAGCTGACATAG	CATGCAAGGGCAGTG	CACAGAAGGCTTTTT	GGAACAAATAGGCOA	TTCCCTGCACTACAG	GGTATCTGGCCAAA-			
Sample 7	TCTAAGCTGACATAG	CATGCAAGGGCAGTG	CACAGAAGGCTTTTT	GGAACAAATAGGCOA	TTCCCTGCACTACAG	GGTATCTGGCCAAA-			
Sample 6	TCTAAGCTGACATAG	CATGCAAGGGCAGTG	CACAGAAGGCTTTTT	GGAACAAATAGGCOA	TTCCCTGCACTACAG	GGTATCTGGCCAAA-			
BKV	TCTAAGCTGACATAG	CATGCAAGGGCAGTG	CACAGAAGGCTTTTT	GGAACAAATAGGCOA	TTCCCTGCACTACAG	GGTATCTGGCCAAA-			
JCV	TTTATGATGACATAG	TAAACAGGGGCAATG	CACTGAAGGATTAGT	GGCAGAGTTAGGCOA	TTCCCTGCAATAAAG	GGTATCAGAATTAGG			
SV40	CCTCAGTAAGCAGAG	CAAGCATATGCAATT	AGCAGACATTTTCTT	TGCACACTCAGGCOA	TTGTTGCACTACAT	TGCATCAACACAGG			

Figure 3 Sequence alignment of BKV isolates from prostate samples. (a) Alignment of three patient samples with early regions of BKV, JCV, and SV40. Identical sequences are in bold. (b) Alignment of four patient samples with BKV regulatory regions of archetype, MM, Dun, TU, UR01, wt-501, and Gardner strains (Seif *et al.*, 1979; Yang and Wu, 1979; Berg *et al.*, 1988; Watanabe and Yoshiike, 1989; Sundsfjord *et al.*, 1990; Negri *et al.*, 1991; Monini *et al.*, 1995). Sample 5 yielded two PCR products shown as 5 and 5(L), the latter being the lower molecular weight product. Identical sequences are shown in bold

Discussion

The experiments in this study were designed to investigate if BKV sequences are present in neoplastic prostatic tissues. In addition, we wanted to examine viral oncoprotein expression and cellular p53 levels, and determine if there was a correlation. Since BKV is ubiquitous in the human population, there is speculation concerning the oncogenic potential of the virus (Imperiale, 2000, 2001; Tognon *et al.*, 2003). The presence of BKV sequences in tumors has been reported in several studies (De Mattei *et al.*, 1995; Monini *et al.*, 1995, 1996; Flaegstad *et al.*, 1999; Jorgensen *et al.*, 2000; Geetha *et al.*, 2002). There are contradictory reports on the presence of BKV DNA in urinary tract tumors: Monini *et al.* (1995) detected BKV DNA using PCR in 31/52 samples, whereas Völter *et al.* (1997) were unable to find it (0/15). Zambrano *et al.* (2002) detected BKV in 4/30 fresh tissue prostate samples and were unsuccessful with archival specimens. These investigators did not examine TAg expression, however. The presence of BKV TAg in neuroblastomas has been reported

(Flaegstad *et al.*, 1999), but these studies have been called into question by a report which demonstrated that the antibody used by these authors is not specific for TAg (Zambrano and Villarreal, 2002). The antibody we used is specific for TAg and does not cross react with any cellular proteins (Harlow *et al.*, 1981). A report on bladder carcinoma found high expression of BKV TAg in primary or metastatic tumors but not in normal epithelium (Geetha *et al.*, 2002). In these studies, p53 was colocalized with TAg to the nuclei of tumor cells, suggesting stabilization and inhibition of p53 function by TAg. This report by Geetha *et al.* (2002) is the first in which BKV localization to tumor cells was demonstrated, suggesting a causal role for BKV in oncogenesis.

We analysed PCA specimens for BKV because of these previous reports along with the facts that the urinary tract is the main site for BKV persistence and that prostate carcinomas have a low frequency of p53 and Rb1 mutations. Using ISH and solution PCR, we detect BKV sequences in neoplastic prostate specimens. Detection of viral sequences in tumors using solution PCR has been viewed with skepticism by some, due to

b

Sample 5	-----TTGAGAGAA AGGGTGGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
Sample 12	-----GGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
Sample 8	-----TTGAGAGAA AGGGTGGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
TU	TACTACTTGAGAGAA AGGGTGGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
MM	TACTACTTGAGAGAA AGGGTGGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
Gardner	TACTACTTGAGAGAA AGGGTGGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
Dunlop	TACTACTTGAGAGAA AGGGTGGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
Sample 9	-----TTGAGAGAA AGGGTGGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
URO1	-----CTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
Archetype	-----GCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
wt-501	TACTACTTGAGAGAA AGGGTGGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
Sample 5(L)	-----TTGAGAGAA AGGGTGGAGGCAGAG GCGGCCTCGGCCTCT TATATATTATAAAAA AAAAGGCCACA--GG GAGGAGCTGCTTACC
Sample 5	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
Sample 12	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
Sample 8	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
TU	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
MM	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
Gardner	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
Dunlop	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
Sample 9	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
URO1	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
Archetype	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
wt-501	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
Sample 5(L)	CATGGAATGCAGCCA AACCATGACCTCAGG AAGGAAAGTGCATGA CT-----GGGCAGCC AGCCAG--TGGC--AGTTA--ATAGTG
Sample 5	AAACC-----CGC C-CCTAAA-ATTCTC TCTTACCATGGAAT GCAGCCAAACCATGA CCTCAGGAA--GG A--AAGTGCATGACT
Sample 12	AAACC-----CGC C-CCTAAA-ATTCTC TCTTACCATGGAAT GCAGCCAAACCATGA CCTCAGGAA--GG A--AAGTGCATGACT
Sample 8	AAACC-----CTGC C-CCTAAA-ATTCTC TCTTACCATGGAAT GCAGCCAAACCATGA CCTCAGGAA--GG A--AAGTGCATGACT
TU	AAACC-----CGC C-CCTAAA-ATTCTC TCTTACCATGGAAT GCAGCCAAACCATGA CCTCAGGAA--GG A--AAGTGCATGACT
MM	AAACC-----CGC C-CCTAAA-ATC-TC TCTTACCATGGAAT GCAGCCAAACCATGA CCTCAGGAA--GG A--AAGTGCATGACT
Gardner	A--GC-----CAGA C--AGA-CAT-TT GCTTACCATGGAAT GCAGCCAAACCATGA CCTCAGGAA--GG A--AAGTGCATGACT
Dunlop	CATGA-----CTCA C-AGGGAG-GAGCT- GCTTACCATGGAAT GCAGCCAAACCATGA CCTCAGGAA--GG A--AAGTGCATGACT
Sample 9	CATGA-----CTCA C-AGGGAG-GAGCT- GCTTACCATGGAAT GCAGCCAAACCATGA CCTCAGGAA--GG A--AAGTGCATGACT
URO1	-----AGC-AAACCATGA CCTCAGGAA--GG A--AAGTGCATGACT
Archetype	AAACC-----CGC C-CCTAAA-ATTCTC -T-----CAA--ATAA ACACAAGAG--G--AAGTGCATGACT
wt-501	AAACC-----CGC C-GACAGACATGTTT TGGGAGCCTAGGAAT CTTG-----GCCTTGT COCCAGTTAACTGG ACACAAGGOCATGTT
Sample 5(L)	AAACC-----CGC C-GACAGACATGTTT TGGGAGCCTAGGAAT CTTG-----GCCTTGT COCCAGTTAACTGG ACACAAGGOCATGTT
Sample 5	G-----GGCAG CC-AGCCAG--TGGC--AGTTAATAGTGAA ACCGGAAAGC--AG-CCA-----GACAGA-CATGTTT
Sample 12	G-----GGCAG CC-AGCCAG--TGGC--AGTTAATAGTGAA ACCGGAAAGC--AG-CCA-----GACAGA-CATGTTT
Sample 8	G-----GGCAG CC-AGCCAG--TGGC--AGTTAATAGTGAA ACCGGAAAGC--AG-CCA-----GACAGA-CATGTTT
TU	G-----GGCAG CC-AGCCAG--TGGC--AGTTAATAGTGAA ACCGGAAAGC--AG-CCA-----GACAGA-CATGTTT
MM	G-----GGCAG CC-AGCCAG--TGGC--AGTTAATAGTGAA ACCGGAAAGC--AG-CCA-----GACAGA-CATGTTT
Gardner	G-----GGCAG CC-AGCCAG--TGGC--AGTTAATAGTGAA ACCGGAAAGC--AG-CCA-----GACAGA-CATGTTT
Dunlop	GACATGTTTTCGAG CCTAGGAATCTTGGC CTGTGCCCCAGTTAA ACTGGA--CAAGG CCAATGGTTC--TG CCGCAGC--TGTC
Sample 9	GACATGTTTTCGAG CCTAGGAATCTTGGC --
URO1	G-----GGCAG CC-AGCCAG--TGGC--AGTTAATAGTGAA ACCGGAAAGC--AG-CCA-----GACAGA-CATGTTT
Archetype	G-----G-----CCAA--AGGC--AGTTAATAGTGAA ACCGGAAAGC--AG-CCA-----GACAGA-CATGTTT
wt-501	C-----TGC-G CC-AGC-----TGTC--ACGACAAAGCT--
Sample 5	TG-CGAGC-CTAGG AATCTTGGCA----
Sample 12	TG-CGAGC-CTAGG AATCTTGGCA----
Sample 8	TG-CGAGC-CTAGG AATCTTGGC----
TU	TG-CGAGC-CTAGG AATCTTGGCCTTGT
MM	GG-GCAGC-CAGCC AGTGGCAGTTAATT
Gardner	TG-CGAGC-C-----
Dunlop	CGACAGCTTCAGTG AAGTTGGTAAAC
URO1	TG-CGAGC-CTAG- -
Archetype	-----G AATCTTGGCCTTGT

Figure 3 Continued

the possibility of false positives caused by contamination of reagents with laboratory viral sequences. Our controls without any DNA gave no PCR product even after both rounds of PCR amplification, indicating that the positive signals were due to authentic amplification

of DNA from the tissues. Moreover, our ISH results confirm the presence of viral DNA in the samples. We amplified the BKV early region in 94% and regulatory region in 81% of the archival specimens, although we could only confirm the sequence of the regulatory region




Figure 4 Detection of BKV DNA in human PCA using ISH. (a and b) BSC-1 and BSC-BKT cells, respectively, stained with BKV specific probe. (c-e) Atrophic regions of the prostate tissue section stained with scrambled probe (c) and BKV-specific probe (d and e). (f and g) Benign regions of the prostate tissue section stained with scrambled and BKV specific probe, respectively. (h) Breast carcinoma stained with BKV probe. Magnification: $\times 400$




Figure 5 Expression of TAg in prostate samples. (a and b) BSC-1 and BSC-BKT cells, respectively, immunostained with anti-TAg antibody. (c-e) Atrophic regions of prostate tissue sections stained with IgG_{2a} isotype control (c) or anti-TAg (d and e). (f) Prostate tissue section stained with anti-AR. (G) Breast carcinoma stained with anti-TAg. Magnification: $\times 400$ (a-c, g, and f); $\times 100$ (d); and $\times 1000$ (e)




Figure 6 Expression of p53 in prostate samples. (a and b) BSC-1 and BSC-BKT cells, respectively, immunostained with anti-p53 antibody. (c-e) Atrophic regions of prostate sections stained with IgG_{2a} isotype control (c) or anti-p53 (d and e). (f and g) TAG-negative prostate tissue samples stained with anti-p53 (f) or anti-TAG (g). (h) Breast carcinoma stained with anti-p53. Magnification: × 400 (a-c, f, and g); × 100 (d); and × 1000 (e)




Figure 7 Colocalization of TAG and p53 expression in prostate samples. (a–d) BSC-1 cells. (a) DAPI staining; (b) anti-p53- and FITC-conjugated secondary; (c) anti-TAG- and TR-conjugated secondary; (d) merged image of (b) and (c). (e–h) BSC-BKT cells. (e) DAPI staining; (f) anti-p53- and FITC-conjugated secondary; (g) anti-TAG- and TR-conjugated secondary; (h) merged image of (f) and (g). (i–l) Atrophic regions of the prostate. (i) Anti-p53- and FITC-conjugated secondary; (j) anti-TAG- and TR-conjugated secondary; (k) merged image of (i) and (j); (l) image (k) merged with DAPI stain to show nuclei

in 44% of the samples. Some samples were positive for the early region but not the regulatory region. Since the regulatory region has been shown to be prone to rearrangement, it is possible that rearrangements led to lack of recognition by the primers. In addition, the regulatory region PCR assay is less sensitive than that used for the early region. Our DNA sequencing data indicate that the early regions are identical to the early region of BKV. We found a number of regulatory region sequences, which were homologous to the TU, Dunlop, and wt-501 strains. There is no obvious correlation between the regulatory region sequences and the histopathology of the samples. Owing to the heterogeneity of the cells in each sample, drawing such a correlation will require sequence analysis of DNA isolated from specific cells in each section rather than from bulk DNA isolated from an entire section. We have not determined if BKV is present episomally or is integrated into the genome. Our ISH studies detected BKV sequences in 71% of the samples, and in all cases, the signal localized to the benign and proliferative atrophic lesions.

IHC analysis detected TAG expression in 43% of the samples. TAG was expressed in focal atrophic lesions (simple atrophy) and in hyper-PIA, but was undetectable in either benign or tumor areas of the tissues. Since our ISH and PCR analyses specifically detect BKV

DNA that can be correlated to TAG expression in the same location, we are confident that we are detecting BKV TAG and not that of JCV or SV40, which are also recognized by this antibody. We speculate that BKV infects normal cells, induces the cells to PAH or PIA, and these eventually transform to tumor cells, since atrophic lesions are proposed to be precursors to HGPIN. It is unclear why BKV might be lost upon progression to frank carcinoma, but it is possible that the reported proapoptotic effects of TAG, mediated through interactions with pRB family proteins, or chromosomal damaging effects of TAG, might be incompatible with growth at that stage of carcinogenesis (Theile and Grabowski, 1990; Tognon *et al.*, 1996; TrabANELLI *et al.*, 1998).

TAG expression could be correlated to elevated p53 expression in the atrophic lesions and both proteins were localized to the cytoplasm of the secretory cells. In the absence of TAG, p53, where detectable, was nuclear. It is unclear why these two proteins, particularly TAG, are not entering the nucleus, although this might be related to the function of the cellular protein, Parc, which has previously been shown to sequester p53 in the cytoplasm (Nikolaev *et al.*, 2003). In the cytoplasm, TAG would not be able to support replication of the virus. We propose that TAG drives proliferation of these cells through its interactions with pRB and, by sequestering p53 in the

cytoplasm, inhibits p53-mediated cell cycle arrest or apoptosis (Knippschild *et al.*, 1996; Moll *et al.*, 1996). Two groups have observed cytoplasmic BKV and SV40 TAG in neuroblastoma and thyroid tumors, respectively (Jorgensen *et al.*, 2000; Vivaldi *et al.*, 2003). There is also precedence for the occurrence of cytoplasmic p53 in human tumors, including inflammatory breast cancer, neuroblastoma, colon carcinoma, and malignant melanoma (Moll *et al.*, 1992; Castresana *et al.*, 1993; Bosari *et al.*, 1994, 1995; Moll *et al.*, 1995; Weiss *et al.*, 1995). To our knowledge, there is no report of cytoplasmic p53 in PCA. Since the cytoplasmic staining of p53 is mostly in atrophic lesions, it is possible that others might not have classified these lesions as cancer, however. Together, these results suggest that a defect in p53 nuclear import could contribute to tumorigenesis.

Our present findings support the hypothesis that PIA lesions are precursors to HGPIN or PCA, as suggested previously. Distinct histopathological relationships between cells found in PIA, HGPIN, and PCA have been reported, suggesting that PIA develops into HGPIN or PCA (De Marzo *et al.*, 1999). PIA is frequently associated with acute and chronic inflammatory processes and is highly proliferative when compared with normal epithelium (Ruska *et al.*, 1998). Atrophic regions containing proliferative epithelial cells, which are unable to differentiate into columnar secretory cells, are usually located in the periphery of the prostate, where PCAs most often occur (Nelson *et al.*, 2003). Although there is evidence to support the association between inflammation and PCA, the exact mechanism still remains elusive. We speculate that BKV infects benign glands, transforms them to atrophic lesions through TAG expression, and this ultimately leads to HGPIN and cancer.

The results in this report are in agreement with an earlier report of the presence of BKV DNA in prostate tumors (Monini *et al.*, 1995). Our data extend this finding by demonstrating expression of TAG to atrophic lesions in the tumors, where it colocalizes with p53. The presence of the BKV genome and, more importantly, expression of its early protein TAG, is suggestive of a role in the development of PCA. These observations emphasize the need for rigorous and more detailed investigations of the association of this virus with PCA and the elucidation of pathways that may be regulated by BKV.

Materials and methods

Human tissue specimens

Paraffin-embedded adenocarcinoma prostate resection specimens from radical prostatectomies, staged using standard TNM (primary tumor, regional nodes, metastasis staging) criteria, were obtained from the Tissue Procurement Core at the University of Michigan Comprehensive Cancer Center (UMCCC). One section from each of the tissue specimens was stained with hematoxylin and eosin to confirm diagnosis. The samples were a mixture of three different histological types, benign, atrophic, and tumor. Each case was evaluated for

benign, atrophic, or tumors by the pathologist. Paraffin-embedded breast tissue blocks were obtained from the breast tissue procurement core at the UMCCC. The Gleason scores of the tissue specimens are listed in Table 1.

DNA extraction

Eight serial paraffin-embedded 5- μ m tissue sections were used for DNA extraction. The sections were treated with 100% xylene for 2 $\times$ 10 min at room temperature to remove the paraffin, washed with 100% ethanol, and dried in a Savant SpeedVac using medium heat. The EX-WAX™ extraction kit (Serological Corp.) was used to extract DNA as indicated by the manufacturer, with minor modification. DNA was precipitated from the tissues by incubation at -80°C for 2 h and dissolved in 50 μ l of resuspension solution from the manufacturer. The DNA was subsequently stored at -20°C. A 2- μ l aliquot was used in the PCR assays. The extraction process was performed in an area that was BKV free and care was taken to avoid crosscontamination of the samples by frequent changing of gloves between samples.

Oligonucleotide primers and probes

The sequence of oligonucleotide primers and probes used for these studies are shown in Figure 1. The human β -hemoglobin gene primers, KM38 and PCO3, amplify a 167 bp fragment (Vilchez *et al.*, 2002).

The first-round BKV early region PCR was performed using primers BKV_{for}(4935-4907) and BKV_{rev}(4358-4382) to generate a 577 bp PCR product. The second, nested round was performed using BKV_{nesfor}(4895-4871) and BKV_{nesrev}(4443-4467), which amplify a 452 bp product. The first round BKV regulatory region PCR was performed using primers BKRRLT1₍₄₀₀₋₃₈₁₎ and BKRRLT6₍₂₁₋₄₀₎ (Monini *et al.*, 1995), yielding a 379 bp product. The nested, second round was performed using primers BRP1₍₈₂₋₁₀₁₎ and BRP2₍₃₅₈₋₃₃₉₎, resulting in a 276 bp product if there is no rearrangement of the regulatory region.

The BKV early region oligonucleotide probe for ISH and Southern blotting, BKV2₍₄₄₃₄₋₄₄₇₈₎, was 45 bases, and is modified from a previous report (Bersagel *et al.*, 1992). A scrambled probe with the same length and G + C content was used as a control in the ISH studies. The sequence of the scrambled probe is: 5'TGTTGATGTAGGATATGCGTCTGTCTTCTTACACCGTGACACAA3'.

Plasmids

The plasmids used to determine the specificity of the PCR and probe were pBK-GEM, containing the BKV early region (Harris *et al.*, 1996); pJCV, containing the complete JCV genome, strain Mad1 (ATCC #45027); and pJYM, containing the complete SV40 genome, strain 776 (Lusky and Botchan, 1981).

PCR amplification

Extracted DNA was amplified using Titanium™ Taq DNA polymerase (Clontech) in a ThermoHybaid thermocycler (Px2). All reactions were performed in a final volume of 100 μ l containing 200 nM each primer, 200 μ M dNTPs, 2 μ l template, and 1 U polymerase in 0.5 $\times$ buffer (20 mM Tricine-KOH, pH 8.0, 8 mM KCl, 1.75 mM MgCl₂, and 1.87 μ g/ml bovine serum albumin (BSA)). The quality of the DNA was first ascertained by PCR (45 cycles) with the β -globin primers. The program used for β -globin amplification consisted of initial denaturation at 94°C for 5 min followed by denaturation at 94°C for 30 s, annealing at 43°C for 30 s, and elongation at

72°C for 1 min, followed by final elongation at 72°C for 7 min. For early region amplification, two rounds of 30 cycles each were used. The template for second-round amplification consisted of 30 µl product from the first-round amplification. The program consisted of initial denaturation for 5 min at 94°C followed by denaturation at 94°C for 30 s, annealing at 47°C for 30 s, and elongation at 72°C for 1 min, followed by final elongation at 72°C for 7 min. For regulatory region amplification, each reaction was increased to 40 cycles and 1 × buffer was used. The program consisted of initial denaturation for 5 min at 94°C followed by denaturation at 94°C for 30 s, annealing for 30 s, and elongation at 72°C for 1 min, followed by final elongation at 72°C for 7 min. The annealing temperature for primers BKRLT1/BKRLT6 was 52°C and BRP1/BRP2 was 47°C. The negative control tube contained all components except DNA template. Plasmid DNA controls were not amplified at the same time as tissue samples, to avoid cross contamination.

Southern blotting

A 9-µl aliquot of the PCR reaction, or plasmid DNA, was electrophoresed on a 1.5% agarose gel in 1 × TAE buffer (40 mM Tris acetate and 1 mM EDTA, pH 7.8), stained with EtBr, and photographed before transfer. The probes were labeled with terminal transferase using the DIG oligonucleotide tailing kit (Roche). The gel was washed sequentially before transfer in 250 mM HCl, 10 min; ddH₂O, 5 min; denaturation solution (0.5 M NaOH and 1.5 M HCl), 2 × 15 min; ddH₂O, 5 min; neutralization solution (0.5 M Tris-HCl, pH 7.5 and 1.5 M NaCl), 2 × 15 min; and 20 × SSC (0.3 M NaCl and 0.03 M sodium citrate, pH 7.03), 10 min. The gel was transferred overnight to nylon membrane (Roche) in 20 × SSC using the capillary method. The membrane was washed for 2 × 15 min in 2 × SSC before baking at 110°C for 30 min to fix the DNA to the membrane, followed by incubation in prehybridization buffer (DIG Easy Hyb buffer; Roche) containing 0.1 mg/ml PolyA for 30 min at 45°C. This was followed by hybridization with denatured DIG-labeled probe (1 pmol/ml for PCR products; 10 pmol/ml for plasmid DNA) for 2 h at 45°C in DIG Easy buffer containing 0.1 mg/ml PolyA. After hybridization, the membrane was washed with 2 × SSC containing 0.1% sodium dodecyl sulfate (SDS) at room temperature for 2 × 15 min followed by 0.1 × SSC containing 0.1% SDS for 30 min at 55°C. All subsequent incubations were at room temperature. The membrane was incubated with 1 × maleic acid (Roche) for 10 min, blocked with blocking buffer (Roche) for 30 min, and incubated with alkaline phosphatase-conjugated anti-DIG antibody (Roche) for 30 min. The anti-DIG antibody was diluted 1:7000 for ECL detection of PCR products or 1:1000 for colorimetric detection of plasmid DNA. After four 15 min washes, the membranes were incubated with either BCIP/NBT (5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium) substrate for colorimetric detection or CSPD substrate (Roche) for ECL detection, which was subsequently exposed to film.

Sequence analysis

PCR products were separated by agarose-gel electrophoresis, extracted (Qiaquick gel extraction kit, Qiagen), and sent for sequencing by the DNA Sequencing Core at the University of Michigan. Samples that did not yield interpretable results were cloned into pGEM-T easy (Promega) and sequenced. The sequences were analysed using the ClustalW server alignment (NCBI Server).

Cell fixation for in situ studies

Cells in culture were fixed using a slightly modified published protocol (Berger, 1986). BSC-1 (African green monkey kidney cells) and BSC-BKT cells were grown on sterilized charged Silane prep™ slides (Sigma) in Dulbecco's modified Eagle's medium (GIBCO), supplemented with 100 U/ml penicillin, 100 µg/ml streptomycin, and 10% fetal bovine serum, at 37°C in a 5% CO₂ incubator, and fixed by 20 min incubation in 4% paraformaldehyde solution (16% stock obtained from Electron Microscopy Sciences) at room temperature. After fixation, the cells were rinsed in phosphate-buffered saline (PBS), first with 3 × PBS (5 min) and then 1 × PBS (3 × 2 min). The slides were dehydrated using increasing concentrations of ethanol (50, 70, and 95%; 5 min each at room temperature), air dried, and stored at -80°C.

In situ DNA hybridization (ISH)

ISH was performed using a modified protocol (Singer et al., 1986; Nuovo, 1996; Wiedorn et al., 1999), with digoxigenin (DIG)-labeled 45 bp probes. The probes were labeled with terminal transferase using the DIG oligonucleotide tailing kit (Roche). Formalin-fixed paraffin-embedded sections (4 µm), mounted on AES charged slides, were deparaffinized by immersing in xylene solution (2 × 5 min) and then rehydrated by incubation with decreasing concentration of ethanol (95, 75, and 50%; 1 min each). The sections were incubated in Tris-buffered saline (TBS; 0.1 M Tris-HCl, pH 7.4, and 0.15 M NaCl) for 10 min before being treated with proteinase K (4.5 µg/ml in TBS) at 37°C for 25 min for permeabilization. The proteinase K treatment was stopped by washing in TBS rigorously for 10 min, with a change of buffer after 5 min. This was followed by two 10 min washes, first with ddH₂O then with ethanol, and air-drying. For hybridization, the sections were prehybridized in 10% formamide, 10% dextran sulfate, and 2 × SSC for 15 min at 45°C. The probe (100 pmol/ml) was added to the prehybridization mixture and denatured for 10 min at 92°C after application on the sections, and hybridization was performed for 2 h at 45°C in a humidified chamber. The sections were then washed with 2 × SSC containing 0.2% BSA fraction V (Roche) at room temperature for 15 min followed by 0.1 × SSC containing 0.2% BSA fraction V for 15 min at 54°C. All subsequent incubations were at room temperature. The sections were blocked with BSA fraction V (50 mg/ml) in TBS for 30 min then incubated with alkaline phosphatase-conjugated anti-DIG antibody (Roche) for 30 min in a humidified chamber. The anti-DIG antibody was diluted 1:65 in conjugate dilution buffer (0.1 M Tris-HCl pH 7.3, 0.15 M MgCl₂, 10 mg/ml BSA fraction V). After two 10 min washes with TBS, the sections were incubated with BCIP/NBT substrate, mounted with crystal mount (Biomedica Corp), and visualized using an Olympus BX41 microscope.

Immunohistochemistry (IHC)

IHC was performed using a modified protocol (Brown et al., 1996; Shi et al., 2001). Paraffin-embedded 4 µm sections were deparaffinized and rehydrated as above. Nonenzymatic antigen retrieval was performed by treatment with 0.01 M sodium citrate, pH 6.0, at 92°C for 50 min with a change of buffer after 20 min. The sections were cooled for 25 min, rinsed with PBS (3 × 5 min), treated with 3% H₂O₂ for 10 min at room temperature to quench endogenous peroxidase, and rinsed in PBS for 2 × 5 min. The sections were blocked with 1.5% horse serum (Vector Labs) in PBS for 45 min and incubated with primary antibody in PBS containing 1.5% horse serum for 45 min at room temperature in a humidified chamber. We tried five different antigen retrieval

techniques and the results were identical. The anti-TAG antibody was a mouse monoclonal antibody originally raised against SV40 TAG (Harlow *et al.*, 1981), which crossreacts with BKV TAG (PAb416; 1:200 dilution in blocking buffer; isotype IgG_{2a}). The anti-p53 was a mouse monoclonal antibody, p53Ab-6 (Vojtesek *et al.*, 1992) (Clone DO-1; 1:20 dilution in blocking buffer; isotype IgG_{2a}; Lab Vision). The anti-androgen receptor was a mouse monoclonal antibody, ARAb-1 ((Ruizeveld de Winter *et al.*, 1991) (Clone AR 441; 1:10 dilution in blocking buffer; Lab Vision)), and the anti-breast carcinoma 90 kDa glycoprotein was a mouse monoclonal antibody, ((Colcher *et al.*, 1981) (Clone B6.2; 1:50 dilution in blocking buffer; BioGenex). Purified mouse IgG_{2a} was used as a negative control (Sigma; 1:20 dilution in blocking buffer). After incubation with primary antibody, sections were washed with PBS for 3 × 5 min and incubated with biotinylated secondary anti-mouse immunoglobulin antibody (ABC kit; Vector labs) at 1:200 dilution in blocking buffer for 45 min at room temperature. Sections were washed with PBS for 3 × 10 min and incubated with avidin/biotin-conjugated peroxidase (1:50 dilution in PBS) for 45 min at room temperature. The sections were subsequently washed with PBS for 10 min, developed with diaminobenzidine tetra hydrochloride substrate (Sigma), counterstained with hematoxylin, mounted with crystal mount (Biomedica Corp), and visualized.

Double-labeling immunofluorescence

For double-labeling immunofluorescence, deparaffinization and antigen retrieval were performed as described above.

References

- Arthur RR, Shah KV, Baust SJ, Santos GW and Saral R. (1986). *N. Engl. J. Med.*, **315**, 230–234.
- Abate-Shen C and Shen MM. (2000). *Genes Dev.*, **14**, 2410–2434.
- Bauer JJ, Sesterhenn IA, Mostofi KF, McLeod DG, Srivastava S and Moul JW. (1995). *Clin. Cancer Res.*, **1**, 1295–1300.
- Berg DT, Walls JD and Grinnell BW. (1988). *Nucleic Acids Res.*, **16**, 9057.
- Berger CN. (1986). *EMBO J.*, **5**, 85–93.
- Bersagel DJ, Finegold MJ, Butel JS, Kupsky WJ and Garcea RL. (1992). *N. Engl. J. Med.*, **326**, 988–993.
- Brewster SF, Oxley JD, Trivella M, Abbott CD and Gilherin DA. (1999). *J. Urol.*, **161**, 1238–1243.
- Brown D, Lydon J, McLaughlin M, Stuart-Tilley A, Tyszkowski R and Alper S. (1996). *Histochemistry*, **105**, 261–267.
- Bookstein R, MacGrogan D, Hilsenbeck SG, Sharkey F and Allred DC. (1993). *Cancer Res.*, **53**, 3369–3373.
- Bookstein R, Rio P, Madreperla SA, Hong F, Allred C, Grizzle WE and Lee WH. (1990). *Proc. Natl. Acad. Sci. USA*, **87**, 7762–7766.
- Bosari S, Viale G, Bossi P, Maggioni M, Coggi G, Murray JJ and Lee AKC. (1994). *J. Natl. Cancer Inst.*, **86**, 681–687.
- Bosari S, Viale G, Roncalli M, Graziani D, Borsani G, Lee AKC and Coggi G. (1995). *Am. J. Pathol.*, **147**, 790–798.
- Bostwick DG. (1995). *Cancer*, **75**, 1823–1836.
- Bostwick DG and Brawer MK. (1987). *Cancer*, **59**, 788–794.
- Castresana JS, Rubio MP, Vazquez JJ, Idoate M, Sober AJ, Seizinger BR and Barnhill RL. (1993). *Int. J. Cancer*, **55**, 562–565.
- Cattoretti G, Rilke F, Andreola S, D'Amato L and Delia D. (1988). *Int. J. Cancer*, **41**, 178–183.
- Colcher D, Hand PH, Nuti M and Schlom JA. (1981). *Proc. Natl. Acad. Sci. USA*, **78**, 3199–3203.
- Cuzik J. (1995). *Cancer Surv.*, **23**, 91–95.
- Dalrymple SA and Beemon KL. (1990). *J. Virol.*, **64**, 1182–1191.
- De Marzo AM, Marchi VL, Epstein JI and Nelson WG. (1999). *Am. J. Pathol.*, **155**, 1985–1992.
- De Marzo AM, Meeker AK, Zha S, Luo J, Nakayama M, Platz EA, Issacs WB and Nelson WG. (2003). *Urology*, **62**, 55–62.
- De Mattei M, Martini F, Corallini A, Gerosa M, Scotlandi K, Carinci P, Barbanti-Brodano G and Tognon M. (1995). *Int. J. Cancer*, **61**, 756–760.
- Dyson N, Buchkovich K, White P and Harlow E. (1989). *Cell*, **58**, 249–255.
- Dyson NR, Bernards SH, Friend LR, Gooding JA, Hassell EQM, Pipas JM, Vandyke T and Harlow E. (1990). *J. Virol.*, **64**, 1353–1356.
- Dörries K, Vogel E, Gunther S and Czub S. (1994). *Virology*, **198**, 59–70.
- Flaegstad T, Andresen PA, Johnsen JI, Asomani SK, Jorgensen GE, Vignarajan S, Kjuul A, Kogner P and Traavik T. (1999). *Cancer Res.*, **59**, 1160–1163.
- Franks LM. (1954). *J. Pathol. Bacteriol.*, **68**, 617–621.
- Frisque RJ, Bream GL and Cannella MT. (1984). *J. Virol.*, **51**, 458–469.
- Gardner SD. (1973). *BMJ*, **1**, 77–78.
- Gardner SD, Field AM, Coleman DV and Hulme B. (1971). *Lancet*, **1**, 1253–1257.
- Gardner Jr WA and Bennett BD. (1992). *The Prostate-Overview: Recent Insights and Speculations*. Weinstein RS and Gardner Jr WA (eds). Williams and Wilkins: Baltimore, MD, pp. 129–148.
- Gardner Jr WA and Culbertson DE. (1987). *J. Urol.*, **137**, 53–56.
- Geetha D, Tong BC, Racusen L, Markowitz JS and Westra WH. (2002). *Transplantation*, **73**, 1933–1936.

- Gonzalzo ML and Isaacs WB. (2003). *J. Urol.*, **170**, 2444-2452.
- Harris KF, Chang E, Christensen JB and Imperiale MJ. (1998a). *Dev. Biol. Stand.*, **94**, 81-91.
- Harris KF, Christensen JB and Imperiale MJ. (1996). *J. Virol.*, **70**, 2378-2386.
- Harris KF, Christensen JB, Radany EH and Imperiale MJ. (1998b). *Mol. Cell. Biol.*, **18**, 1746-1756.
- Harlow E, Crawford LV, Pim DC and Williamson NM. (1981). *J. Virol.*, **39**, 861-869.
- Hiraoka A, Ishikawa J, Kitayama H, Yamagami T, Teshima H, Nakamura H, Shibata H, Masaoka T, Ishigami S and Taguchi F. (1991). *Bone Marrow Transplant.*, **7**, 107-111.
- Hirsch HH and Steiger J. (2003). *Lancet Infect. Dis.*, **3**, 611-623.
- Ibrahim GK, Gravitt PE, Dittrich KL, Ibrahim SN, Melhus O, Anderson SM and Robertson CN. (1992). *J. Urol.*, **148**, 1822-1826.
- Imperiale MJ. (2000). *Virology*, **267**, 1-7.
- Imperiale MJ. (2001). *Oncogene*, **20**, 7917-7923.
- Iwasaka T, Oh-uchida M, Matsuo N, Yokoyama M, Fukuda K, Hara K, Fukuyama K, Hori K and Sugimori H. (1993). *Gynecol. Oncol.*, **48**, 104-109.
- Jemal A, Murray T, Samuels A, Ghafoor A, Ward and Thun MJ. (2003). *CA Cancer J. Clin.*, **53**, 5-26.
- Jiang Z, Woda BA, Rock KL, Xu Y, Savas L, Khan A, Pihan G, Cai F, Babcook JS, Rathanaswami P, Reed SG, Xu J and Fanger G. (2001). *Am. J. Surg. Pathol.*, **25**, 1397-1404.
- Jorgensen GEK, Johnsen JJ, Ponthan F, Kogner P, Flaegstad T and Traavik T. (2000). *Med. Pediatr. Oncol.*, **35**, 593-596.
- Kang S and Folk WR. (1992). *Virology*, **191**, 754-764.
- Knippschild U, Oren M and Deppert W. (1996). *Oncogene*, **12**, 1755-1765.
- Kuefer R, Varambally S, Zhou M, Lucas PC, Loeffler M, Wolter H, Mattfeldt T, Hautmann RE, Gschwend JE, Barrette TR, Dunn RL, Chinnaiyan AM and Rubin MA. (2002). *Am. J. Pathol.*, **161**, 841-848.
- Lane DP and Crawford LV. (1979). *Nature*, **278**, 261-263.
- Li RM, Mannon RB, Kleiner D, Tsokos M, Bynum M, Kirk AD and Kopp JB. (2002). *Transplantation*, **74**, 1497-1504.
- Lusky M and Botchan MR. (1981). *Nature*, **293**, 79-81.
- McNeal JE. (1988). *Am. J. Surg. Pathol.*, **12**, 619-633.
- McNicol PJ and Dodd JG. (1991). *J. Urol.*, **145**, 850-853.
- Melamed J, Einhorn JM and Ittmann MM. (1997). *Clin. Cancer Res.*, **10**, 1867-1872.
- Moens U and Rekvig OP. (2001). *Human Polyomaviruses: Molecular and Clinical Perspectives* Khalili K and Stoner GL (eds). Wiley-Liss: New York, pp. 215-226.
- Moll UM, LaQuaglia M, Benard J and Riou G. (1995). *Proc. Natl. Acad. Sci. USA*, **92**, 4407-4411.
- Moll UM, Ostermeyer AG, Haladay R, Winkfield B, Frazier M and Zambetti G. (1996). *Mol. Cell. Biol.*, **16**, 1126-1137.
- Moll UM, Riou G and Levine AJ. (1992). *Proc. Natl. Acad. Sci. USA*, **89**, 7262-7266.
- Monini P, Rotola A, DeLellis L, Corallini A, Secchiero P, Albini A, Benelli R, Parravicini C, Barbanti-Brodano G and Cassai E. (1996). *Int. J. Cancer*, **66**, 717-722.
- Monini P, Rotola D, DiLuca D, DeLellis L, Chiari E, Corallini A and Cassai E. (1995). *Virology*, **214**, 273-279.
- Myers RB, Srivastava S, Oelschlagel DK and Grizzle WE. (1994). *J. Natl. Cancer Inst.*, **86**, 1140-1145.
- Nakashatri H, Pater MM and Pater A. (1988). *J. Virol.*, **62**, 4613-4621.
- Negrini M, Sabbioni S, Arthur RR, Castagnoli A and Barbanti-Brodano G. (1991). *J. Virol.*, **65**, 5092-5095.
- Nelson WG, De Marzo AM and Isaacs WB. (2003). *N. Engl. J. Med.*, **349**, 366-381.
- Nikolaev AY, Li M, Puskas N, Qin J and Gu W. (2003). *Cell*, **112**, 29-40.
- Noda T, Sasagawa T, Dong Y, Fuse H, Namiki M and Inoue M. (1998). *Urol. Res.*, **26**, 165-169.
- Nuovo GJ. (1996). *PCR In Situ Hybridization: Protocols and Applications*. Raven Press: New York.
- Nupponen N and Visakorpi T. (1999). *Eur. Urol.*, **35**, 351-354.
- Osman I, Drodjak M, Fazzari M, Ferrara J, Scher HI and Cordon-Cardo C. (1999). *Clin. Cancer Res.*, **5**, 2082-2088.
- Padgett B. (1981). *Human Papovaviruses* Tooze J (ed). Cold Spring Harbor Laboratory: Cold Spring Harbor, NY, pp. 339-370.
- Pappo O, Demetris AJ, Raikow RB and Randhawa PS. (1996). *Mod. Pathol.*, **9**, 105-109.
- Pater A. and Pater MM. (1986). *J. Virol.*, **58**, 680-683.
- Pipas JM. (1992). *J. Virol.*, **66**, 3979-3985.
- Pipas JM and Levine AJ. (2001). *Cancer Biol.*, **11**, 23-30.
- Portolani M, Barbanti-Brodano G and Placa ML. (1975). *J. Virol.*, **15**, 420-422.
- Putzi MJ and De Marzo AM. (2000). *Urology*, **56**, 828-832.
- Randhawa PS, Finkelstein S, Scantlebury V, Shapiro R, Vivas C, Jordan M, Picken MM and Demetris AJ. (1999). *Transplantation*, **67**, 103-109.
- Replogue MD, Storch GA and Clifford DB. (2001). *Clin. Infect. Dis.*, **33**, 191-202.
- Rotola A, Monini P, Luca D, Savioli A, Sione R, Secchiero P, Reggiani A and Cassai E. (1992). *Int. J. Cancer*, **52**, 359-365.
- Rubin MA, Zhou M, Dhanasekaran SM, Varambally S, Barrette TR, Sanda MG, Pienta KJ, Ghosh D and Chinnaiyan AM. (2002). *JAMA*, **287**, 1662-1670.
- Rubinstein R, Pare N and Harley EH. (1987). *J. Virol.*, **61**, 1747-1750.
- Ruska KM, Sauvageot J and Epstein JI. (1998). *Am. J. Surg. Pathol.*, **22**, 1073-1077.
- Ruizeveld de Winter JA, Trapman J, Vermey M, Mulder E, Zegers ND and Van der Kwast TH. (1991). *J. Histochem. Cytochem.*, **39**, 927-936.
- Sarkar FH, Sakr W, Li Y-W, Macoska J, Ball DE and Crissman JD. (1992). *Prostate*, **21**, 145-152.
- Scheffner M, Munger K, Byrne JC and Howley PM. (1991). *Proc. Natl. Acad. Sci. USA*, **88**, 5523-5527.
- Seif I, Khoury G and Dhar R. (1979). *Cell*, **18**, 963-977.
- Shah KV. (1996). *Fields Virology* Fields BN, Knipe DM and Howley PM (eds). Lippincott-Raven: Philadelphia, pp. 2027-2043.
- Shah R, Mucci NR, Amin A, Macoska JA and Rubin MA. (2001). *Am. J. Pathol.*, **158**, 1767-1773.
- Shi SR, Cote RJ and Taylor CR. (2001). *J. Histochem. Cytochem.*, **49**, 931-937.
- Shinohara T, Matsuda M, Cheng SH, Marshall J, Fujita M and Nagashima K. (1993). *J. Med. Virol.*, **41**, 301-305.
- Singer RH, Lawrence JB and Villnave C. (1986). *BioTechniques*, **4**, 230-250.
- Simmons DT. (1995). *Oncogenic mechanisms* Barbanti-Brodano G, Bendinelli M and Friedman F (eds). Plenum Press: New York, pp. 27-50.
- Small JA, Khoury G, Jay G, Howey PM and Scangos GA. (1986). *Proc. Natl. Acad. Sci. USA*, **83**, 8288-8292.
- Staib C, Pesch J, Gerwig R, Gerber JK, Brehm U, Stangl A and Grummt F. (1996). *Virology*, **219**, 237-246.
- Sundsford A, Johansen T, Flaegstad T, Moens U, Villand P, Subramani S and Traavik T. (1990). *J. Virol.*, **64**, 3864-3871.
- Theile M and Grabowski G. (1990). *Arch. Virol.*, **113**, 221-233.

- Tognon M, Casalone R, Martini F, De Mattei M, Granata P, Minelli E, Arcuri C, Collini P and Bocchini V. (1996). *Cancer Genet. Cytogenet.*, **90**, 17–23.
- Tognon M, Corallini A, Martini F, Negrini M and Barbanti-Brodano G. (2003). *Oncogene*, **22**, 5192–5200.
- Trabanelli C, Corallini A, Gruppioni R, Sensi A, Bonfatti A, Campioni D, Merlin M, Calza N, Possati L and Barbanti-Brodano G. (1998). *Virology*, **243**, 492–496.
- Varambally S, Dhanasekaran SM, Zhou M, Barrette TR, Sinha CK, Sanda MG, Ghosh D, Pienta KJ, Sewalt RGAB, Otte AP, Rubin MA and Chinnaiyan AM. (2002). *Nature*, **419**, 572–573.
- Vilchez RA, Kozinetz CA, Jorgensen JL, Kroll MH and Butel JS. (2002). *AIDS Res. Hum. Retroviruses*, **18**, 237–242.
- Vivaldi A, Pacini F, Martini F, Iaccheri L, Pezzetti F, Elisei R, Pinchera A, Faviana P, Basolo F and Tognon M. (2003). *J. Clin. Endocrinol. Metab.*, **88**, 892–899.
- Vojtesek B, Bartek J, Midgley CA and Lane DP. (1992). *J. Immunol. Methods*, **151**, 237–244.
- Völter C, zurHansen H, Alber D and DeVilliers EM. (1997). *Virology*, **237**, 389–396.
- Watanabe S and Yoshiike K. (1989). *Virology*, **169**, 204–212.
- Weiss J, Heine M, Korner B, Pilch H and Jung EG. (1995). *Br. J. Dermatol.*, **133**, 23–31.
- Wiedorn KH, Kuhl H, Galle J, Caselitz J and Vollmer E. (1999). *Histochem. Cell Biol.*, **111**, 89–95.
- Yang RC and Wu R. (1979). *Science*, **206**, 456–462.
- Yang XJ, Wu CL, Woda BA, Dresser K, Tretiakova M, Fanger GR and Jiang Z. (2002). *Am. J. Surg. Pathol.*, **26**, 921–925.
- Zambrano A, Kalantari M, Simoneau A, Jensen JL and Villarreal LP. (2002). *Prostate*, **53**, 263–276.
- Zambrano A and Villarreal LP. (2002). *Oncogene*, **37**, 5725–5732.



Rapid Communication

BKV and SV40 infection of human kidney tubular epithelial cells in vitro

Jonathan Low,^a H. David Humes,^{b,c} Mark Szczypka,^c and Michael Imperiale^{a,d,*}

^aDepartment of Microbiology and Immunology, University of Michigan Medical School, Ann Arbor, MI 48109, USA

^bDepartment of Internal Medicine, University of Michigan Medical School, Ann Arbor, MI 48109, USA

^cNephros Therapeutics, Ann Arbor, MI 48108, USA

^dComprehensive Cancer Center, University of Michigan Medical School, Ann Arbor, MI 48109, USA

Received 4 February 2004; returned to author for revision 9 March 2004; accepted 26 March 2004

Available online 4 May 2004

Abstract

The interaction of BKV with its natural target cells, human kidney epithelial cells, has not been studied. In vitro infections of human primary kidney epithelial cells were performed to investigate a BKV infection in its natural host cell. BKV undergoes a lytic replication cycle in this system: high levels of T antigen expression were first detected at 36 h postinfection, while viral DNA replication, capsid protein expression, and progeny virus were observed at 48 h postinfection. It was observed that the related polyomavirus SV40 is incapable of infecting human kidney epithelium except in the presence of the GM1 ganglioside, recently reported to be an SV40 receptor.

© 2004 Elsevier Inc. All rights reserved.

Keywords: BKV; SV40; Kidney; Proximal tubule

Introduction

The process by which BKV infects its host, completes an acute replication cycle, and then remains subclinical for the lifetime of the host or becomes active upon immunosuppressive conditions is currently poorly understood. Therefore, we wanted to develop a model system in which BKV could be studied in vitro under conditions similar to those it might encounter in vivo. Early studies on BKV were performed using monkey-derived Vero cells, human embryonic kidney cells, or human embryonic fibroblast cells, which are not the normal target cells for the virus in vivo (Maraldi et al., 1975; Seehafer et al., 1975). We have chosen to characterize a BKV infection in human kidney epithelial cells due to the known ability of BKV to infect tubular epithelium in humans (Randhawa et al., 1999). Human proximal tubule epithelial cells (HPTE cells) are an important part of the kidney nephron. These cells

recover essential non-waste blood products and return them to circulation, produce vitamin precursors, maintain blood pressure and blood volume, and communicate with the host immune system through the release of cytokines when they sense infections or toxic molecules (Briggs et al., 2001; Daha and van Kooten, 2000). A system has been developed in which HPTE cells are able to maintain their differentiated state in vitro and function in a renal assist device. This device was developed to perform the nutrient recovery, blood pressure, and blood volume functions for patients with failing kidneys (Humes et al., 2002). The HPTE cells used in the renal assist device are capable of being passaged in culture up to six times. Using these cells, we were able to observe viral protein expression, DNA replication, and growth over the course of infection of a known target cell type for BKV. We find that BKV is capable of replicating efficiently and producing infectious virus particles within these cells. Interestingly, SV40 cannot infect these cells except in the presence of the GM1 ganglioside. The characterization of BKV infection of HPTE cells sets the groundwork for further studies to better understand how BKV is able to persist within the human host and reactivate under immunosuppressed conditions.

* Corresponding author. Department of Microbiology and Immunology, University of Michigan Medical School, 1500 East Medical Center Drive, 6304 Cancer Center, Ann Arbor, MI 48109-0942. Fax: +1-734-615-6560.

E-mail address: imperial@umich.edu (M. Imperiale).

Results

We began these studies by asking whether BKV could infect HPTE cells *in vitro*, and what the outcome of an infection would be. We first examined expression of large T antigen (TAg) as a marker for early gene expression. The expression of TAg and its interaction with p53 and the retinoblastoma family of tumor suppressors are critical for a successful infection by polyomaviruses. We infected HPTE cells with BKV at an MOI of 5 fluorescence forming units (FFU)/cell, isolated total protein over the course of 11 days, and assayed TAg expression by immunoblotting (Fig. 1A). We first detected TAg expression between 12 and 24 h postinfection. Much higher levels of TAg expression were detected between 24 and 36 h postinfection. The expression of TAg peaked at 7 days postinfection, and remained high throughout the rest of the time course.

The next step in the BKV life cycle is replication of its DNA. To assess this step, we isolated low molecular weight DNA from HPTE cells infected with BKV and digested with *EcoRI* to linearize the BKV genome. We analyzed the amount of viral DNA present by Southern blotting using the BKV genome as a probe. We detected the presence of low levels of viral DNA starting at 0 h postinfection (Fig. 1B). Due to the low and constant level of viral DNA seen during the first 36 h of the infection, we believe that this is DNA from input virus. The first increase in viral DNA occurred at 36 h postinfection, which is expected due to the onset of expression of TAg 12 h earlier. The level of viral DNA increased throughout the course of infection, peaking at 9 days postinfection. Metabolic labeling experiments using tritiated thymidine confirmed that viral DNA synthesis did not begin until 48 h postinfection (data not shown).

After determining when early viral proteins were expressed and viral DNA was replicated, we went on to establish when the major late viral protein VP1 was expressed. Total cellular protein was isolated from HPTE

cells infected with BKV over the course of 11 days. Immunoblot analysis revealed the presence of VP1 at all time points from 0 h postinfection to 11 days postinfection (Fig. 1A). As was the case for viral DNA, the VP1 present at 0 h postinfection is probably from the input virus. The first time that VP1 was expressed *de novo* was at 36 h postinfection, when there was a marked increase in its steady-state levels. High levels of VP1 expression were seen 12 h later. The amount of VP1 increased throughout the remainder of the infection and was at its highest level when the time course ended at 11 days postinfection.

When BKV infects a human kidney, it establishes a subclinical infection unless the immune system has been compromised. Under immunocompromised conditions or immunosuppression, usually resulting from organ or bone marrow transplantation, pregnancy, or HIV infection, high levels of BKV replication are observed in the kidneys and urinary tract, which can lead to hemorrhagic cystitis and polyomavirus nephropathy (PVN) (Binet et al., 1999). Three of the morphologic hallmarks of PVN are intranuclear viral inclusions, focal necrosis of tubular cells, and cells rounding up from the epithelial layer and extruding into the tubular lumen (Nickel et al., 1999). We examined the morphology of BKV-infected HPTE cells. As early as 4 days postinfection, we detected morphological characteristics in HPTE cells that are also commonly observed in PVN. The HPTE cells rounded up from the cell monolayer, likely due to cell death, and contained enlarged nuclei and what appeared to be inclusions (Fig. 2). These characteristics were more evident at 9 days postinfection. Thus, BKV apparently undergoes a lytic infection in HPTE cells, producing morphologic changes similar to those seen in PVN.

The key event to which all other viral events lead is progeny virus production. We determined the viral yield over the course of an 11-day infection (Fig. 3). At 24 h postinfection, we detected the eclipse phase of infection, when fewer mature virions were present. We then observed




Fig. 1. Expression of BKV early and late proteins, and DNA replication. (A) Total cellular protein (25 μ g) isolated at the indicated times postinfection was analyzed as described in Materials and methods. (B) *EcoRI*-digested low molecular weight DNA was separated on an agarose gel, transferred to PVDF, and probed for BKV sequences.




Fig. 2. Morphology of BKV-infected HPTE cells. (A and B) HPTE cells were mock infected or infected with BKV at an MOI of 5 FFU/cell. Cells were fixed and stained with hematoxylin at 4 days postinfection. (C) Kidney biopsy from a patient with PVN stained with hematoxylin. Arrows identify enlarged nuclei and inclusion bodies in proximal tubule cells. Magnification is $\times 400$.

a relatively rapid rise in the amount of virus that peaked at 4 days postinfection, concomitant with robust viral DNA and late protein synthesis. After this early peak, the release of progeny virus remained relatively constant for the remainder of the infection.

We were also interested in whether SV40 can infect HPTE cells. SV40 is the best studied of the polyomavirus family and a close relative of BKV. An examination of SV40 infection of human kidney cells is particularly interesting because of accumulating evidence that SV40 infects




Fig. 3. Growth of BKV in HPTE cells. Triplicate wells of HPTE cells were infected with BKV at an MOI of 5 FFU/cell. Viral lysates were prepared at the indicated days pi, and the titer determined. The mean and standard deviation are shown.

humans (Butel and Lednicky, 1999; Garcea and Imperiale, 2003). SV40 is known to infect proximal tubule cells in the monkey kidney, causing a disease similar to that seen in humans (Sheffield et al., 1980). Upon infecting HPTE cells and BSC-1 cells, an established line of monkey kidney cells, with SV40 we expected similar morphologies to those seen in BKV-infected HPTE cells. However, at 11 days postinfection, cytopathic effects were evident in SV40-infected BSC-1 cells, but not in HPTE cells. We looked for SV40 Tag as a marker of early protein expression. We detected high levels of Tag expression in BSC-1 cells, but none at all in HPTE cells, even at 13 days postinfection (Fig. 4A). To determine if SV40 had entered the HPTE cells, we infected HPTE and BSC-1 cells, extracted low molecular weight DNA, and probed for the presence of SV40 DNA using a Southern blot. In BSC-1 cells, the level of SV40 DNA began to increase at 24 h postinfection, but in HPTE cells no SV40 DNA was detected even at 4 days postinfection (Fig. 4B). These results suggest that SV40 is unable to infect HPTE cells.

It has been reported that the cellular receptor for SV40 is the major histocompatibility complex (MHC) class I molecule (Atwood and Norkin, 1989). Immunoblotting of protein lysates from the HPTE cells revealed that these cells express this molecule at levels similar to human fibroblasts susceptible to SV40 infection (data not shown). More recently, however, Tsai et al. (2003) have reported that the ganglioside GM1 is required for entry of SV40 into mammalian cells. We considered the possibility that a lack of GM1 on the proximal tubule cells was the cause for the lack of




Fig. 4. SV40 infection of HPTE and BSC-1 cells. Cells were infected with SV40 at an MOI of 5 PFU/cell. Total cellular protein (25 μ g; A) or equal amounts of DNA (B) isolated at the indicated days pi were analyzed as in Fig. 1, except that an SV40 probe was used for the Southern blot. (C) Cells were incubated with the indicated ganglioside for 18 h before infection with BKV or SV40. Protein lysates were prepared at 0 and 96 h postinfection and analyzed as in Fig. 1A.

infectivity by SV40. HPTE cells were incubated with either GM1 or GD1a, reported as a receptor for mouse polyomavirus (Py), and infected with SV40 or BKV (Tsai et al., 2003). Immunoblots were used to detect the expression of TAg. SV40 was able to infect HPTE cells preincubated with GM1, confirming that the GM1 ganglioside is a receptor for SV40 and indicating that the inability of SV40 to infect HPTE cells is due to a lack of this ganglioside (Fig. 4C). BKV was able to infect HPTE cells that had been preincubated with either GM1 or GD1a, illustrating that GD1a does not have an inhibitory effect on viral entry into these cells.

Discussion

The experiments described in this paper define a system in which BKV can be studied *in vitro* using a cell type that it infects *in vivo*. While others have examined BKV infections in cell culture, none of these previous studies have been performed at a molecular level in a cell type infected *in vivo* (Flaegstad and Traavik, 1987; Maraldi et al., 1975). We have shown that BKV causes morphological changes *in vitro* which are similar to those seen in BKV-induced nephropathy *in vivo* and that a productive infection occurs. While descriptive, the characterization of BKV infection of

these cells is a critical first step to understanding how BKV causes disease in the human host.

We first observed expression of high levels of TAg at 36 h postinfection. This 36-h delay from infection to expression of abundant TAg corresponds to work by others who observed that in human fibroblasts, nuclear uncoating of BKV occurs as late as between 24 and 30 h postinfection (Maraldi et al., 1975). We also observed high levels of VP1 expression and DNA replication beginning at 48 h postinfection, 12 h after TAg was strongly expressed. Two days after the components of the mature virion were first synthesized, we observed a large peak in mature virion production.

We do not have any indication that BKV can establish a latent or persistent infection in tubular epithelial cells *in vitro*. No cells survived the infection despite repeated attempts to passage them. One major difference between HPTE cells infected *in vitro* and *in vivo* is the lack of an immune system. Therefore, our system reproduces what likely occurs during a BKV infection in an immunosuppressed individual. Because BKV is reactivated upon immunosuppression, it is clear that the body keeps viral replication in check without destroying the entire population of infected cells. A component of the immune system controlling BKV infection may be chemokines or cytokines.

For example, it has been demonstrated that cytokines are capable of inhibiting promoter activity of other viruses: IFN- α inhibits murine CMV immediate early gene expression and IFN- γ inhibits gene expression of constructs under control of the human CMV or SV40 promoters (Gribaudo et al., 1995; Harms and Splitter, 1995; Qin et al., 1997). Further study is required to determine under which conditions BKV gene expression and replication might be repressed. The immune system may not be the only factor playing a role in activation of BKV from its latent or persistent state. In studies of Py, it has been reported that mouse kidneys become permissive to lytic infection upon cellular damage and subsequent tissue regeneration (Atencio et al., 1993). It may be that a similar situation exists in the transplant setting, in which BKV replication is stimulated due to either ischemia or kidney damage during handling. It is also possible that a change in hormone levels or stress within an individual could cause the reactivation of BKV, such as during pregnancy. PCR analysis of DNA isolated from uninfected HPTE cells revealed that these cells did not harbor any detectable BKV sequences before infection (data not shown). Because it is known that BKV commonly infects the proximal tubule *in vivo*, it may be slightly surprising that we did not detect any viral sequences in our cell lines. It is possible that during an *in vivo* infection, only a few cells are infected, with this infection being kept in check by the immune system and localized to certain areas of the kidney (Chesters et al., 1983). Upon immunosuppression, the infection may become lytic and infect a much larger population of kidney cells.

We have also demonstrated that SV40 does not infect HPTE cells. Because SV40 infects tubular epithelial cells in monkeys, (Sheffield et al., 1980) we expected that our proximal tubule lines would be infected by SV40. The virus apparently does not enter HPTE cells without the addition of the GM1 ganglioside. It remains possible that in the absence of GM1, SV40 is internalized into HPTE cells through a different mechanism, but is unable to traffic to the nucleus and begin lytic replication. Overexposures of our Southern blots demonstrate the presence of input SV40 DNA in HPTE infections in the absence of added gangliosides, but we are unable to determine if the virus has been internalized or remains on the cell surface. It is also possible that in humans SV40 infects a different type of cell within the kidney, retaining its organ tropism, or infects a different tissue entirely. Two groups have reported the detection of SV40 sequences in diseased human kidneys (Butel et al., 1999; Li et al., 2002), so it is possible that SV40 can lytically infect kidney cells under some circumstances. Further study is required to determine why SV40 cannot infect proximal tubule epithelium and which human cells, if any, might act as reservoirs for the virus *in vivo*.

We have developed a system in which we can study lytic BKV replication in phenotypically normal human kidney epithelial cells. Given the prevalence of BKV in the human population and its known associations with human disease,

we believe that this system is an important first step towards better understanding how BKV interacts with the host. We are left with at least two major areas of study to pursue. First, there are likely many cellular proteins and processes that are influenced during infection. Second, we wish to develop a model of BKV persistent infection and determine what causes BKV to reactivate from its subclinical state.

Materials and methods

Cell culture

Human kidney proximal tubule epithelial cells, isolated from cadaveric tissue, were cultured as previously described (Humes et al., 2002). The experiments described in this report were performed on cells isolated from two separate donors with no difference in the results.

Viruses

BKV stocks were initially propagated in 293 cells from virus obtained from the American Type Culture Collection (VR-837). Viral lysates were made through three cycles of freezing the infected cells at -80°C and thawing at 37°C . Once we determined that HPTE cells could be lytically infected, one subsequent passage was performed on HPTE cells. The regulatory region of these stocks, as determined by sequencing of PCR products, was identical to the TU strain (Sundsford et al., 1990). For the SV40 studies we used strain 776.

Viral titers

Viral titer was determined using a fluorescent focus assay, rather than hemagglutination inhibition, due to its ability to detect infectious virus. Seventy percent confluent HPTE cells in 12-well dishes were infected with 10-fold dilutions of BKV for 4 days at 37°C . The cells were fixed for 30 min using 70% ethanol/30% PBS (Liu and Atwood, 2001) PAb416 (Harlow et al., 1981) followed by a fluorescein-conjugated anti-mouse antibody (Sigma) used to detect TAg. Titer, measured in fluorescence forming units (FFU), was determined by counting five random fields. The titer of SV40 was determined on BSC-1 monolayers by the agar overlay method (Brockman et al., 1973).

Infections

Cells were grown in 6-well dishes to 70% confluence, and infected with BKV or SV40 at an MOI of 5 FFU/cell. To stain with hematoxylin (Gill III formula), cells were fixed in 4% paraformaldehyde, washed with PBS, and rehydrated with ethanol. For infections to which gangliosides were added, cells were preincubated with either media, 3.2 μM GM1, or 2.7 μM GD1a (gift of B. Tsai) for 18 h at 37°C ,

then washed three times with media before infection (Tsai et al., 2003).

Immunoblotting

Total cell protein was extracted using E1A lysis buffer (Harlow et al., 1986) containing 0.05 M NaF and Complete EDTA-free protein inhibitors (Roche). Protein concentration was determined using the Bio-Rad protein assay, and equal amounts of protein were electrophoresed on an 8% SDS–polyacrylamide gel. PAb416 was used to detect TAG expression, P5G6 (gift of D. Galloway) for VP1 expression, and GAPDH Ab8245 (Abcam) for glyceraldehyde-3-phosphate-dehydrogenase expression. After washing, blots were probed with a horseradish peroxidase-conjugated secondary antibody (Sigma). Antibody complexes were detected using ECL+ reagent (Amersham) and exposure to film.

DNA replication assays

Low molecular weight DNA was isolated from infected cells using the protocol described by Hirt (1967). One quarter of the DNA isolated from each sample was digested with *EcoRI*, electrophoresed, and transferred to a PVDF membrane. A random-labeled probe was synthesized from a plasmid containing either the BKV or SV40 genome. The probe was allowed to hybridize for 16 h at 65 °C, after which the membrane was washed twice with 2× SSC and once with 2× SSC and 1%. The membrane was then exposed to film.

Virus growth assay

Viral lysates, prepared by three freeze–thaw cycles as described above (Knowles, 2001), were collected in triplicate from HPTE cells infected with BKV at an MOI of 5 FFU/cell. The viral lysates were prepared by three freeze–thaw cycles as described above. The titer of the virus in these lysates was determined using the fluorescent focus assay described above.

Acknowledgments

We thank the members of the Imperiale laboratory for help with this work, Denise Galloway for anti-BKV VP1 antibodies, Bill Tsai for GM1 and GD1a gangliosides, Diane Cibrik for providing biopsy specimens, and Kathy Spindler and Kathy Collins for useful discussions and suggestions. This work was supported by Department of Defense CDMRP grant PC001595.

References

- Atencio, I.A., Shadan, F.F., Zhou, X.J., Vaziri, N.D., Villarreal, L.P., 1993. Adult mouse kidneys become permissive to acute polyomavirus infection and reactivate persistent infections in response to cellular damage and regeneration. *J. Virol.* 67, 1424–1432.
- Atwood, W.J., Norkin, L.C., 1989. Class I major histocompatibility proteins as cell surface receptors for simian virus 40. *J. Virol.* 63 (10), 4474–4477.
- Binet, I., Nickleleit, V., Hirsch, H.H., Prince, O., Dalquen, P., Gudat, F., Mihatsch, M.J., Thiel, G., 1999. Polyomavirus disease under new immunosuppressive drugs: a cause of renal graft dysfunction and graft loss. *Transplantation* 67 (6), 918–922.
- Briggs, J.P., Kriz, W., Schnermann, J.B., 2001. Overview of renal function structure. In: Greenberg, A. (Ed.), *Primer on Kidney Diseases*, 3rd ed. Academic Press, New York, pp. 3–19.
- Brockman, W.W., Lee, T.N., Nathans, D., 1973. The evolution of new species of viral DNA during serial passage of simian virus 40 at high multiplicity. *Virology* 54 (2), 384–397.
- Butel, J.S., Lednicky, J.A., 1999. Cell and molecular biology of simian virus 40: implications for human infections and disease. *J. Natl. Cancer Inst.* 91, 119–134.
- Butel, J.S., Arrington, A.S., Wong, C., Lednicky, J.A., Finegold, M.J., 1999. Molecular evidence of simian virus 40 infections in children. *J. Infect. Dis.* 180 (3), 884–887.
- Chesters, P.M., Heritage, J., McCance, D.J., 1983. Persistence of DNA sequences of BK virus and JC virus in normal human tissues and in diseased tissues. *J. Infect. Dis.* 147, 676–684.
- Daha, M.R., van Kooten, C., 2000. Is the proximal tubular cell a proinflammatory cell? *Nephrol., Dial., Transplant.* 15 (Suppl. 6), 41–43.
- Flaegstad, T., Traavik, T., 1987. BK virus in cell culture: infectivity quantitation and sequential expression of antigens detected by immunoperoxidase staining. *J. Virol. Methods* 16 (1–2), 139–146.
- Garcea, R.L., Imperiale, M.J., 2003. Simian virus 40 infection of humans. *J. Virol.* 77 (9), 5039–5045.
- Gribaudo, G., Ravaglia, S., Gaboli, M., Gariglio, M., Cavallo, R., Landolfo, S., 1995. Interferon-alpha inhibits the murine cytomegalovirus immediate-early gene expression by down-regulating NF-kappa B activity. *Virology* 211 (1), 251–260.
- Harlow, E., Crawford, L.V., Pim, D.C., Williamson, N.M., 1981. Monoclonal antibodies specific for simian virus 40 tumor antigens. *J. Virol.* 39, 861–869.
- Harlow, E., Whyte, P., Franza Jr., B.R., Schley, C., 1986. Association of adenovirus early-region 1A proteins with cellular polypeptides. *Mol. Cell. Biol.* 6, 1579–1589.
- Hams, J.S., Splitter, G.A., 1995. Interferon-gamma inhibits transgene expression driven by SV40 or CMV promoters but augments expression driven by the mammalian MHC I promoter. *Hum. Gene Ther.* 6 (10), 1291–1297.
- Hirt, B., 1967. Selective extraction of polyoma DNA from infected mouse cell cultures. *J. Mol. Biol.* 26, 365–369.
- Humes, H.D., Fissell, W.H., Weitzel, W.F., Buffington, D.A., Westover, A.J., MacKay, S.M., Gutierrez, J.M., 2002. Metabolic replacement of kidney function in uremic animals with a bioartificial kidney containing human cells. *Am. J. Kidney Dis.* 39 (5), 1078–1087.
- Knowles, W.A., 2001. Propagation and assay of BK virus. *Methods Mol. Biol.* 165, 19–31.
- Li, R.M., Branton, M.H., Tanawattanacharoen, S., Falk, R.A., Jennette, J.C., Kopp, J.B., 2002. Molecular identification of SV40 infection in human subjects and possible association with kidney disease. *J. Am. Soc. Nephrol.* 13 (9), 2320–2330.
- Liu, C.K., Atwood, W.J., 2001. Propagation assay of the JC virus. In: Raptis, L.H. (Ed.), *Methods Mol. Biol.*, vol. 165. Humana Press, Totowa, NJ, pp. 9–17.
- Maraldi, N.M., Barbanti-Brodano, G., Portolani, M., La Placa, M., 1975. Ultrastructural aspects of BK virus uptake and replication in human fibroblasts. *J. Gen. Virol.* 27 (1), 71–80.
- Nickleleit, V., Hirsch, H.H., Binet, I.F., Gudat, F., Prince, O., Dalquen, P., Thiel, G., Mihatsch, M.J., 1999. Polyomavirus infection of renal allograft recipients: from latent infection to manifest disease. *J. Am. Soc. Nephrol.* 10 (5), 1080–1089.

- Qin, L., Ding, Y., Pahud, D.R., Chang, E., Imperiale, M.J., Bromberg, J.S., 1997. Promoter attenuation in gene therapy: interferon-gamma and tumor necrosis factor-alpha inhibit transgene expression. *Hum. Gene Ther.* 8, 2019–2029.
- Randhawa, P.S., Finkelstein, S., Scantlebury, V., Shapiro, R., Vivas, C., Jordan, M., Picken, M.M., Demetris, A.J., 1999. Human polyoma virus-associated interstitial nephritis in the allograft kidney. *Transplantation* 67 (1), 103–109.
- Seehafer, J., Salmi, A., Scraba, D.G., Colter, J.S., 1975. A comparative study of BK and polyoma viruses. *Virology* 66 (1), 192–205.
- Sheffield, W.D., Strandberg, J.D., Braun, L., Shah, K., Kalter, S.S., 1980. Simian virus 40-associated fatal interstitial pneumonia and renal tubular necrosis in a rhesus monkey. *J. Infect. Dis.* 142 (4), 618–622.
- Sundsford, A., Johansen, T., Flaegstad, T., Moens, U., Volland, P., Subramani, S., Traavik, T., 1990. At least two types of control regions can be found among naturally occurring BK virus strains. *J. Virol.* 64 (8), 3864–3871.
- Tsai, B., Gilbert, J.M., Stehle, T., Lencer, W., Benjamin, T.L., Rapoport, T.A., 2003. Gangliosides are receptors for murine polyoma virus and SV40. *EMBO J.* 22 (17), 4346–4355.