

GAS CHROMATOGRAPHIC ANALYSIS OF THE STEREOISOMERS OF THE CHEMICAL WARFARE AGENT GF

Jan E. Kolakowski, Steven P. Harvey and Louis P. Reiff

U.S. Army Edgewood Chemical Biological Center
Aberdeen Proving Ground, MD 21010

ABSTRACT

Synthesis of the nerve agent cyclohexyl methylphosphonofluoridate (GF) yields a racemic mixture of two stereoisomers due to the presence of an asymmetric phosphorus atom. These two enantiomers, designated as P(+) and P(-), were separated in 8 minutes by chiral gas chromatography and quantified using flame photometric detection. Plots of peak area versus concentration of the individual enantiomers resulted in correlation coefficients greater than 0.99. The enantiomers were present in racemic GF in the ratio of 3:2 as P(-):P(+). The P(-) enantiomer was isolated using a method whereby the P(+) isomer was removed from the racemic mixture via preferential enzymatic catalyzed hydrolysis. The specific rotation of the P(-) enantiomer at 589 nm and 25 °C in methylene chloride was calculated to be -19.3° . The P(-) enantiomer also inhibited acetylcholinesterase more strongly than the racemic mixture, consistent with the findings for other optically active nerve agents.

INTRODUCTION

The chemical warfare agents GA, GB, GD, GF, and VX all have asymmetric phosphorus atoms resulting in pairs of P(+) and P(-) stereoisomers. GD also has an asymmetric carbon, which gives it four stereoisomers. Previously reported data on GA, GB, GD, and VX show that most of the toxicity of these compounds, as measured by inhibition rate of acetylcholinesterase (AChE), is due to the P(-) isomers.¹⁻⁵ No data have yet been published on the toxicities of the individual GF enantiomers. The enantiomers for all four of the G-agents have been separated by chiral gas chromatography (GC).⁶ VX enantiomers were resolved using a chiral liquid chromatography method.⁷ Separation of the GF enantiomers proved to be the most challenging, with retention times in excess of 80 minutes.⁸ A chiral GC method was developed to perform rapid quantitative measurements of the enantiomers of GF. The work described here was part of an investigation into the stereospecificity of nerve agent degrading enzymes.^{9,10}

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 00 JAN 2002	2. REPORT TYPE N/A	3. DATES COVERED -	
4. TITLE AND SUBTITLE Gas Chromatographic Analysis Of The Stereoisomers Of The Chemical Warfare Agent Gf		5a. CONTRACT NUMBER	
		5b. GRANT NUMBER	
		5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)		5d. PROJECT NUMBER	
		5e. TASK NUMBER	
		5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Edgewood Chemical Biological Center Aberdeen Proving Ground, MD 21010		8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)	
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited			
13. SUPPLEMENTARY NOTES This article is from ADA409494 Proceedings of the 2001 ECBC Scientific Conference on Chemical and Biological Defense Research, 6-8 March , Marriott's Hunt Valley Inn, Hunt Valley, MD.			
14. ABSTRACT			
15. SUBJECT TERMS			
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified	UU
			18. NUMBER OF PAGES 5
			19a. NAME OF RESPONSIBLE PERSON

EXPERIMENTAL

The gas chromatograph was a Hewlett-Packard model 6890 equipped with a flame photometric detector and an automatic sampler. The conditions were as follows:

Column: 25m X 250 um id X 0.12 um Chrompack Chirasil-Val-L.

Injection volume: 1 microliter, 100:1 split.

Inlet temperature: 200 °C.

Oven temperature: 90 °C (isothermal).

FPD temperature: 200 °C (phosphorus mode).

Carrier (helium) flow rate: 1 ml/min.


A chromatographically pure preparation of a single isomer of GF was made by degrading the isomer on which the enzyme had the greater activity. The enzymatic reaction was run at 15°C and pH 7.0 in order to minimize spontaneous hydrolysis. Just past the midpoint deflection, the reaction was extracted with methylene chloride and the extract was analyzed by GC using the above conditions. A single peak was observed representing a single GF isomer. When NaF was shaken with a solution of this isomer, subsequent GC analysis showed two peaks, consistent with NaF catalyzed racemization of GF. Since the racemization occurs in the absence of enzyme, the direct effect of NaF on GF appears to be at least one means by which GF is racemized.

The specific rotation of the single GF isomer was then calculated based on measurements of the observed rotation made at 589 nm (sodium line) and 25 °C using a Perkin Elmer 141 Electronic Polarimeter and a sample cell with a path length of 10 cm.


RESULTS AND DISCUSSION

The enantiomers of GF were consistently separated by 0.15 minutes at retention times near 8 minutes. A typical chromatogram is shown in figure 1a. Hydrolysis of the racemic mixture yielded the single isomer shown in figure 1b. Optical rotation measurements later revealed this to be the P(-) isomer, the specific rotation at 589 nm and 25 °C being calculated as -19.3°. The isomers were consistently present in the ratio P(-):P(+) of 3:2 based on peak area.

GA, GB, GD, and VX are known to stereospecifically bind acetylcholinesterase (AChE) resulting in a significant difference in toxicity between enantiomers.¹⁻⁵ Nerve agents and enzymes all exert their effects in a biological, chiral environment, so it should be expected that those effects would be stereoselective. For these agents, the P(-) isomers are all more toxic than the P(+) isomers. In the case of GA, the (-) isomer is 7 times more toxic than the (+) isomer.³ Racemic GB is half as toxic as (-) GB, indicating that essentially all the toxicity is derived from the P(-) isomer.^{1,5} Essentially all the toxicity of GD is derived from the two P(-) isomers.² The (-) isomer of VX is 13 times more toxic than the (+) isomer.⁴ Figure 2 shows the results of an AChE inhibition assay in the presence of racemic and (-) GF as compared to spontaneous inhibition. The decrease in product yield is a result of the degree of binding of GF to AChE, or the increased inhibition of the enzyme. The rate of product formed is approximately halved in the



Peak #	RetTime [min]	Type	Width [min]	Area 150 pA*s	Height [150 pA]	Area %
1	7.845	VV	0.0851	746.98712	117.62490	37.40245
2	8.003	VV	0.1160	1250.17371	131.37010	62.59755



Peak #	RetTime [min]	Type	Width [min]	Area 150 pA*s	Height [150 pA]	Area %
1	7.981	VV	0.1160	1100.49902	122.31263	1.000e2

Figure 1. Gas Chromatograms of Racemic GF and (-)GF.




Figure 2. AChE Inhibition by GF.




Figure 3. Standard Curves of GF Enantiomers.

presence of (-)GF when compared to racemic GF. This shows that pure (-)GF is more toxic than the racemic mixture.

Although baseline separation was not achieved, the separation between (-) and (+) GF was sufficient to quantitatively measure these isomers. Attempts to achieve baseline resolution of the enantiomers by modifying the GC conditions proved unsuccessful. Figure 3 shows sample plots of peak area versus concentration of the stereoisomers, as calculated from the ratio of peak area of the individual enantiomers to that of the racemic agent. Linear regression analyses of these data yielded correlation coefficients greater than 0.99.

CONCLUSIONS

A rapid and quantitative chiral GC method was developed to measure the individual stereoisomers of GF with retention times near 8 minutes. The P(-) enantiomer was successfully purified from the racemic agent and its specific rotation at 589 nm and 25 °C in methylene chloride was calculated to be -19.3° . The P(-) isomer of GF was shown to be more toxic than the racemic mixture and, consequently, much more toxic than the P(+) isomer. This is consistent with the findings for other nerve agents with optically active isomers.

REFERENCES

1. Boter, H.L. and Van Dijk, C, *Biochem. Pharmacol.* 18, 2403 (1969).
1. Benschop, H.P., et. al., *Toxicol. Appl. Pharmacol.* 72, 61 (1984).
2. Degenhardt, C.E.A.M., et. al., *J. Am. Chem. Soc.* 108, 8290 (1986).
3. Hall, C.R., et. al., *J. Pharm. Pharmacol.* 29, 574 (1977).
4. Van DeMeent, et. al., TNO, unpublished results (1987).
5. Smith, J.R. and Schlager, J.J., *J. High Resol. Chromatogr.* 19, 151 (1986).
6. Smith, J.R. and Shih, M.L., Analysis of the Stereoisomers of the Organophosphorus Chemical Warfare Agent VX Using Chiral LC and APCI Mass Spectrometry, Proceedings of the 47th ASMS Conference on Mass Spectrometry and Allied Topics, Dallas, Texas, June 1999.
7. Smith, J.R. and Schlager, J.J., Proceedings of the 1994 ERDEC Scientific Conference on Chemical and Biological Defense Research, ERDEC-SP-036, May 1996, p. 697.
8. Harvey, S.P., et. al., Catalytic Decontamination: Recent Developments, presented at the Army Research Office Decon Meeting, Jackson Hole, Wyoming, September 2000.
9. Harvey, S.P., et. al., Phosphotriesterase Mutants with Enhanced VX and G-agent Activity, presented at the 22nd Army Science Conference, Baltimore, Maryland, December 2000.