



# **Interval Estimates for Probabilities of Non-Perforation Derived From a Generalized Pivotal Quantity**

**by David W. Webb**

**ARL-TR-3596**

**September 2005**

## **NOTICES**

### **Disclaimers**

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

Citation of manufacturer's or trade names does not constitute an official endorsement or approval of the use thereof.

**DESTRUCTION NOTICE**—Destroy this report when it is no longer needed. Do not return it to the originator.

# **Army Research Laboratory**

Aberdeen Proving Ground, MD 21005-5066

---

**ARL-TR-3596****September 2005**

---

## **Interval Estimates for Probabilities of Non-Perforation Derived From a Generalized Pivotal Quantity**

**by David W. Webb**  
**Weapons and Materials Research Directorate, ARL**

| REPORT DOCUMENTATION PAGE                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |                             |                              |                                          | Form Approved<br>OMB No. 0704-0188                             |                                                           |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------|------------------------------|------------------------------------------|----------------------------------------------------------------|-----------------------------------------------------------|
| <p>Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.</p> <p><b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b></p> |                             |                              |                                          |                                                                |                                                           |
| 1. REPORT DATE (DD-MM-YYYY)<br>September 2005                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |                             | 2. REPORT TYPE<br>Final      |                                          | 3. DATES COVERED (From - To)<br>March 2004 to August 2005      |                                                           |
| 4. TITLE AND SUBTITLE<br><br>Interval Estimates for Probabilities of Non-Perforation Derived From a Generalized Pivotal Quantity                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |                             |                              |                                          | 5a. CONTRACT NUMBER                                            |                                                           |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                             |                              |                                          | 5b. GRANT NUMBER                                               |                                                           |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                             |                              |                                          | 5c. PROGRAM ELEMENT NUMBER                                     |                                                           |
| 6. AUTHOR(S)<br><br>David W. Webb (ARL)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                             |                              |                                          | 5d. PROJECT NUMBER<br>1L162618AH80                             |                                                           |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                             |                              |                                          | 5e. TASK NUMBER                                                |                                                           |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                             |                              |                                          | 5f. WORK UNIT NUMBER                                           |                                                           |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)<br>U.S. Army Research Laboratory<br>Weapons and Materials Research Directorate<br>Aberdeen Proving Ground, MD 21005-5066                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |                             |                              |                                          | 8. PERFORMING ORGANIZATION<br>REPORT NUMBER<br><br>ARL-TR-3596 |                                                           |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                             |                              |                                          | 10. SPONSOR/MONITOR'S ACRONYM(S)                               |                                                           |
|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                             |                              |                                          | 11. SPONSOR/MONITOR'S REPORT<br>NUMBER(S)                      |                                                           |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT<br><br>Approved for public release; distribution is unlimited.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                              |                             |                              |                                          |                                                                |                                                           |
| 13. SUPPLEMENTARY NOTES                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |                             |                              |                                          |                                                                |                                                           |
| 14. ABSTRACT<br><br>A generalized pivotal quantity is developed that yields exact confidence intervals for the cumulative distribution function (CDF) at a specific value when the underlying distribution is assumed to be normal. This problem is similar to the development of a tolerance interval, and unsurprisingly, its solution involves the non-central "t" distribution. Confidence bands for a normal CDF follow easily. Military applications include vulnerability and lethality assessment (e.g., interval estimation for the probability of non-perforation against homogeneous armored targets).                                                                                                                                                                                                                                                                                                                                                   |                             |                              |                                          |                                                                |                                                           |
| 15. SUBJECT TERMS<br><br>confidence interval; confidence limit; generalized pivotal quantity; non-central t distribution; perforation                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |                             |                              |                                          |                                                                |                                                           |
| 16. SECURITY CLASSIFICATION OF:                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |                             |                              | 17. LIMITATION<br>OF ABSTRACT<br><br>SAR | 18. NUMBER<br>OF PAGES<br><br>40                               | 19a. NAME OF RESPONSIBLE PERSON<br>David W. Webb          |
| a. REPORT<br>Unclassified                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | b. ABSTRACT<br>Unclassified | c. THIS PAGE<br>Unclassified |                                          |                                                                | 19b. TELEPHONE NUMBER (Include area code)<br>410-278-7014 |

---

## Contents

---

|                                                                               |           |
|-------------------------------------------------------------------------------|-----------|
| <b>List of Figures</b>                                                        | <b>iv</b> |
| <b>List of Tables</b>                                                         | <b>iv</b> |
| <b>Acknowledgments</b>                                                        | <b>v</b>  |
| <b>1. Introduction</b>                                                        | <b>1</b>  |
| <b>2. General Solution</b>                                                    | <b>2</b>  |
| <b>3. Lower Confidence Bound (LCB)</b>                                        | <b>6</b>  |
| <b>4. Upper Confidence Bound (UCB)</b>                                        | <b>7</b>  |
| <b>5. Two-Sided Confidence Interval</b>                                       | <b>8</b>  |
| <b>6. Iterative Solutions for the Confidence Bounds and Confidence Limits</b> | <b>8</b>  |
| <b>7. Confidence Bands</b>                                                    | <b>10</b> |
| <b>8. An Application</b>                                                      | <b>10</b> |
| <b>9. Summary</b>                                                             | <b>12</b> |
| <b>10. References</b>                                                         | <b>13</b> |
| <b>Appendix A. Proof of the Monotonicity of the Function H</b>                | <b>15</b> |
| <b>Appendix B. MATLAB Code for Probability of Non-perforation LCB</b>         | <b>17</b> |
| <b>Appendix C. Mathematica Code for Probability of Non-perforation LCB</b>    | <b>19</b> |
| <b>Appendix D. MATLAB Code for Probability of Non-perforation UCB</b>         | <b>21</b> |
| <b>Appendix E. Mathematica Code for Probability of Non-perforation UCB</b>    | <b>23</b> |

|                                                                                                  |           |
|--------------------------------------------------------------------------------------------------|-----------|
| <b>Appendix F. MATLAB Code for Probability of Perforation Two-Sided Confidence Interval</b>      | <b>25</b> |
| <b>Appendix G. Mathematica Code for Probability of Perforation Two-Sided Confidence Interval</b> | <b>27</b> |
| <b>Appendix H. MATLAB Code for Probability of Non-perforation CDF and Lower Confidence Band</b>  | <b>29</b> |
| <b>List of Symbols Used</b>                                                                      | <b>31</b> |
| <b>Distribution List</b>                                                                         | <b>33</b> |

---

## List of Figures

---

|                                                                                                                                              |    |
|----------------------------------------------------------------------------------------------------------------------------------------------|----|
| Figure 1. Two non-central t density functions, $f_1$ and $f_2$ , with respective non-centrality parameters $\lambda_1$ and $\lambda_2$ ..... | 9  |
| Figure 2. Estimated probability of non-perforation and 95% lower confidence boundary as a function of armor thickness .....                  | 11 |

---

## List of Tables

---

|                                                                    |    |
|--------------------------------------------------------------------|----|
| Table 1. Penetration depths into armor (no units specified). ..... | 10 |
|--------------------------------------------------------------------|----|

---

## Acknowledgments

---

The author wishes to thank the following people:

Messrs. Thomas Havel and Michael Zoltoski of the U.S. Army Research Laboratory (ARL), who brought this problem to my attention;

Professor Thomas Mathew, University of Maryland, Baltimore County, for pointing out Weerahandi's generalized pivotal quantity for  $\frac{\mu}{\sigma}$ ;

Major William H. Kaczynski and Commandant Michael A. Faries, United States Military Academy, who are responsible for the Mathematica code that appears in appendix B; and

Dr. Gene Cooper, ARL, for his many hours spent fine tuning the Mathematica programs and his review of the original manuscript.

INTENTIONALLY LEFT BLANK

---

## 1. Introduction

---

The estimation of non-perforation probabilities is of significant interest to the developers of armor systems, whose objective is to provide protection for vehicles and personnel against enemy threats. In particular, the following questions may arise:

1. What is the probability that homogenous armor of a given thickness will not be perforated by a specific enemy threat?
2. How thick should the armor be if one is to be highly confident that the probability of non-perforation with this threat is high?

Conversely, these probabilities are of interest to the developers of armor-piercing projectiles, whose objective is to perforate enemy armor. From this point of view, one might ask

3. What is the probability that a projectile will be able to perforate an enemy armor of a given thickness?

For a particular projectile, non-perforation<sup>1</sup> occurs if its penetration depth, denoted by  $x$ , is less than the thickness of the armor, denoted by  $x_0$ . Now consider the random variable  $X$ , which represents the penetration depth of a randomly selected projectile. The probability of non-perforation is an unknown, fixed constant within the interval  $[0, 1]$  and is written as  $P(X \leq x_0)$ . An estimate for the true non-perforation probability will be based on sample data and as such, is subject to sampling variation. However, a point estimate for the non-perforation probability is a single value and therefore does not give any sense of the uncertainty associated with it. Preferable to a point estimate is an interval estimate that contains the non-perforation probability with high probability, i.e., a confidence interval. The width of the interval provides a quantitative measure of the estimation error.

The problem of estimating  $P(X \leq x_0)$  has its origins in acceptance sampling plans for statistical quality control in the 1950's. Lieberman and Resnikoff (1955) and Barton (1961) yielded expressions for the minimum variance estimator. It was not until the 1970's that confidence interval estimators surfaced. In their 1977 paper, Owen and Hua derived lower confidence limits for  $P(X \geq x_0)$  and described how to obtain one-sided and two-sided limits for  $P(X \leq x_0)$ .

However, their derivation contains several errors in notation and as a result, is quite difficult to follow. Owen and Hua's paper also provided tables that assist in the calculation of lower confidence limits for  $P(X \geq x_0)$ ; these tables were dramatically expanded in Odeh and Owen

---

<sup>1</sup> For the remainder of this report, we only discuss non-perforation events, their probabilities of occurrence, and interval estimates for these probabilities. Analogous results for the probability of perforation,  $P(X > x_0)$ , are easily derived if we use the axiomatic property  $P(X > x_0) = 1 - P(X \leq x_0)$ .

(1980, tables 7.1 through 7.7). Odeh and Owen's tables are cited in the more recent engineering statistics text by Hahn and Meeker (1991, section 4.5).

In this report, the recent concept of generalized inference is used to develop one- and two-sided confidence intervals for the probability of non-perforation when it can be assumed that the penetrations follow a normal distribution. Although this may be seen as an attempt to "reinvent the wheel," it is meant to show the wide applicability of the generalized approach to confidence interval construction. As pointed out by Weerahandi (2004, section 1.7), generalized inference may be used to obtain confidence intervals for *any* function of the normal distribution parameters.

Solutions for the limits of the intervals require iterative calculation of percentiles of non-central  $t$  distributions. As an alternative to the tables by Odeh and Owen and to avoid the interpolation errors inherent in their usage, software code is provided in the appendices so that the practitioner can easily obtain exact solutions for the interval limits.

---

## 2. General Solution

---

Suppose that  $n$  depths of penetration  $x_1, x_2, \dots, x_n$  are collected in an experiment for which the observations are assumed to come from a normal distribution with mean  $\mu$  and standard deviation  $\sigma$ . Historically, this assumption has been found to be tenable when the projectiles are of the same ammunition type with (approximately) equal muzzle velocities, obliquities, etc., and the target is a homogeneous armor. For the purpose of this report, we assume that the armor is thick enough to guarantee that no projectiles will perforate. From this sample, an estimate is desired for the probability of non-perforation, i.e., the probability that a future penetration,  $X$ , from the same population of projectiles will be less than or equal to some specified depth,  $x_0$ . This probability is written as  $P(X \leq x_0)$  and is known as the cumulative distribution function (CDF) evaluated at  $x_0$ , or  $F_X(x_0)$ . Notice that

$$P(X \leq x_0) = P\left(\frac{X - \mu}{\sigma} \leq \frac{x_0 - \mu}{\sigma}\right) = P\left(Z \leq \frac{x_0 - \mu}{\sigma}\right) = \Phi\left(\frac{x_0 - \mu}{\sigma}\right), \quad (1)$$

in which  $Z$  is a standard normal random variable and  $\Phi(\cdot)$  is the standard normal cumulative distribution function (CDF) commonly tabulated in statistics texts and available in many software packages. Therefore, calculating a simple point estimate for  $P(X \leq x_0)$  is

straightforward: one computes the standard normal CDF at an estimate of  $\frac{x_0 - \mu}{\sigma}$ . We obtain

this estimate by substituting the sample mean ( $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$ ) and the sample standard deviation

( $s = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}$ ) for the population mean ( $\mu$ ) and the population standard deviation ( $\sigma$ ), respectively<sup>2</sup>. That is,

$$\hat{F}_X(x_0) = \hat{P}(X \leq x_0) = \hat{P}\left(Z \leq \frac{x_0 - \mu}{\sigma}\right) \approx P\left(Z \leq \frac{x_0 - \bar{x}}{s}\right) = \Phi\left(\frac{x_0 - \bar{x}}{s}\right). \quad (2)$$

However, any simple point estimate for  $F_X(x_0)$  does not give information about its precision. What we would prefer to construct is a confidence interval or perhaps a confidence bound for  $F_X(x_0)$  that by its width gives some sense of the degree of uncertainty associated with it.<sup>3</sup>

We begin by recalling from equation 1 that  $F_X(x_0) = \Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ . Since  $\Phi(\cdot)$  is a strictly monotonic function, we can easily attain a confidence interval for  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$  by deriving a confidence interval for  $\frac{x_0 - \mu}{\sigma}$  and then applying the function  $\Phi(\cdot)$  to the resulting confidence limits (see Mood, Graybill, and Boes, 1974, page 378).

The following derivation of a confidence interval for  $\theta = \frac{x_0 - \mu}{\sigma}$  is based on a method by Weerahandi (1995) in which he uses a generalized pivotal quantity to obtain a confidence interval for the coefficient of variation,  $\frac{\sigma}{\mu}$ , for a normal population with mean  $\mu$  and standard deviation  $\sigma$ . The construction of a generalized pivotal quantity requires one to find a function  $R$  with arguments

1.  $\bar{X}$  and  $S^2$  (independent and sufficient statistics for the random sample  $X_1, \dots, X_n$ ),
2.  $\bar{x}$  and  $s^2$  (the observed values of  $\bar{X}$  and  $S^2$ ),
3.  $\theta$ , the parameter of interest, and
4. perhaps a vector,  $\xi$ , of additional unknown (nuisance) parameters.

---

<sup>2</sup>Although this estimate is easily obtained, it does not have the desirable property of being unbiased. An unbiased estimator with minimum variance is presented in either Lieberman and Resnikoff (1955) or Barton (1961).

<sup>3</sup>Note the distinction between the problems of solving for a confidence interval for  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$  and solving for a tolerance interval. With the former, the percentile value ( $x_0$ ) is specified and an interval for its associated CDF value,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ , is desired. On the other hand, with tolerance interval construction, the CDF value ( $p$ ) is specified while an interval for the associated percentile,  $\mu + z_p \sigma$ , is desired.

This function, denoted in full by  $R(\bar{X}, S^2; \bar{x}, s^2, \theta, \xi)$ , must satisfy the following two conditions:

Condition 1:  $R(\bar{X}, S^2; \bar{x}, s^2, \theta, \xi)$  has a distribution that is free of any unknown parameters.

Condition 2: The observed value of  $R$ , i.e.,  $r = R(\bar{x}, s^2; \bar{x}, s^2, \theta, \xi)$ , is equal to  $\theta$ .

If such a function can be found, then it is a generalized pivotal quantity for  $\theta$ , and its percentiles can be used to obtain confidence intervals for  $\theta$ . Since their introduction in the 1980's, generalized pivotal quantities have been regarded as quite challenging to derive. Even Weerahandi (1993) states "... the construction of pivots requires some intuition." In essence, one obtains the pivotal quantity by working backwards from the expression for  $\theta$ . The mathematical "tricks" of adding 0 and/or multiplying by 1 are adroitly employed to link all unknown parameters in  $\theta$  with functions of the sufficient statistics whose distributions are parameter free. Once the linking is complete, all remaining random variables are converted to their observed values.

Step 1: The sufficient statistics for a normally distributed random sample are  $\bar{X}$  and  $S^2$ .

Random variables based on these statistics whose distributions are free of unknown parameters include

1.  $Y = \frac{\bar{X} - \mu}{\sigma}$ , distributed as a normal random variable with mean zero and variance  $\frac{1}{n}$ ; and
2.  $V = (n-1)\frac{S^2}{\sigma^2}$ , distributed as a chi-square random variable with  $n-1$  degrees of freedom.

Step 2: We attempt to construct a random variable from  $\theta$  that involves  $Y$  and/or  $V$ . First, notice that by adding zero in the form of  $(-\bar{X} + \bar{X})$  to the numerator,  $\theta$  can be rewritten in the following manner:

$$\theta = \frac{x_0 - \mu}{\sigma} = \frac{x_0 - \bar{X} + \bar{X} - \mu}{\sigma} = \frac{x_0 - \bar{X}}{\sigma} + \frac{\bar{X} - \mu}{\sigma}. \quad (3)$$

Now the parameter  $\mu$  is linked to the random variable  $\frac{\bar{X} - \mu}{\sigma}$  which was previously noted as

having a parameter-free distribution. When we make the substitution  $Y = \frac{\bar{X} - \mu}{\sigma}$ , the parameter  $\mu$  is removed from  $\theta$ :

$$\theta = \frac{x_0 - \bar{X}}{\sigma} + Y. \quad (4)$$

It remains to manipulate the left addend of equation 4 so that the unknown parameter  $\sigma$  is linked to a random variable whose distribution is free of unknown parameters. We can achieve this by multiplying the left addend by 1, in the form of  $\sqrt{\frac{(n-1)S^2}{(n-1)S^2}}$ :

$$\theta = \frac{x_0 - \bar{X}}{\sigma} + Y = \frac{x_0 - \bar{X}}{\sigma} \sqrt{\frac{(n-1)S^2}{(n-1)S^2}} + Y = \frac{x_0 - \bar{X}}{S\sqrt{n-1}} \sqrt{\frac{(n-1)S^2}{\sigma^2}} + Y. \quad (5)$$

Now  $\sigma$  is linked to  $\frac{(n-1)S^2}{\sigma^2}$  which was noted in step 1 as having a parameter-free distribution.

After  $V = (n-1)\frac{S^2}{\sigma^2}$  is substituted in equation 5, all unknown parameters are removed from  $\theta$ :

$$\theta = \frac{x_0 - \bar{X}}{S\sqrt{n-1}} \sqrt{\frac{(n-1)S^2}{\sigma^2}} + Y = \frac{x_0 - \bar{X}}{S\sqrt{n-1}} \sqrt{V} + Y. \quad (6)$$

Step 3: At this point, we replace the remaining sufficient statistics in equation 6 with their observed values; that is,  $\frac{x_0 - \bar{x}}{s\sqrt{n-1}} \sqrt{V} + Y$ .

Step 4: Finally, we rewrite the random variables  $V$  and  $Y$  in their original form (as functions of the sufficient statistics) to obtain the generalized pivotal quantity:

$$R(\bar{X}, S^2; \bar{x}, s^2, \mu, \sigma) = \frac{x_0 - \bar{x}}{s\sqrt{n-1}} \sqrt{\frac{(n-1)S^2}{\sigma^2}} + \frac{\bar{X} - \mu}{\sigma}. \quad (7)$$

As was demonstrated in the development of  $R$ , its distribution is free of any unknown parameters. Thus, condition 1 is satisfied. Furthermore, the observed value of  $R$  is

$$r = R(\bar{x}, s^2; \bar{x}, s^2, \mu, \sigma) = \frac{x_0 - \bar{x}}{s} \sqrt{\frac{s^2}{\sigma^2}} + \frac{\bar{x} - \mu}{\sigma} = \theta,$$

thus satisfying condition 2. Therefore,  $R$  is a generalized pivotal quantity whose percentiles can be used to obtain a confidence interval for  $\theta$ . Using the normal cumulative distribution,  $\Phi(\cdot)$ , we then ultimately obtain a desired confidence interval for the probability of non-perforation.

---

### 3. Lower Confidence Bound (LCB)

---

Since effective armor offers a high probability of non-perforation, engineers and management will most often be interested in a lower confidence bound, since this would represent a “pessimistic” bound on the true probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ .

A  $(1 - \alpha)100\%$  LCB for  $\theta = \frac{x_0 - \mu}{\sigma}$  is that value  $B_L$  for which

$$1 - \alpha = P(B_L \leq R) \quad (8)$$

is satisfied.<sup>4</sup> We can estimate a generalized LCB for  $\theta$  by randomly generating a large number of observations of  $R$  and selecting a value for which equation 5 is empirically satisfied. For instance, with  $\alpha = .05$ , one could start by generating 100,000 observations of  $R$ , denoted by  $r_1, r_2, \dots, r_{100000}$ . If the ordered observations are written as  $r_{(1)} < r_{(2)} < \dots < r_{(100000)}$ , then a 95%

LCB for  $\theta$  would be any real value  $B_L$  so that  $r_{(5000)} < B_L < r_{(5001)}$ , such as  $B_L = \frac{r_{(5000)} + r_{(5001)}}{2}$ .

However, equation 8 can be expanded to get

$$1 - \alpha = P\left(B_L \leq \frac{x_0 - \bar{x}}{s} \sqrt{\frac{S^2}{\sigma^2}} + \frac{\bar{X} - \mu}{\sigma}\right) = P\left(B_L \leq \frac{x_0 - \bar{x}}{s} \sqrt{\frac{V}{n-1}} + Y\right). \quad (9)$$

Notice that the random variable  $Y$  in equation 9 is equal in probability to  $\frac{Z}{\sqrt{n}}$ , in which  $Z$  is a standard normal random variable (having mean 0 and variance 1). By using this fact and rearranging the terms in the last probability statement, we have

$$\begin{aligned} 1 - \alpha &= P\left(\frac{x_0 - \bar{x}}{s} \sqrt{\frac{V}{n-1}} + \frac{Z}{\sqrt{n}} \geq B_L\right) \\ &= P\left(\frac{x_0 - \bar{x}}{s} \sqrt{\frac{n}{n-1}} \sqrt{V} + Z \geq \sqrt{n} B_L\right) \\ &= P\left(\frac{Z - \sqrt{n} B_L}{\sqrt{\frac{V}{n-1}}} \geq \frac{\bar{x} - x_0}{s} \sqrt{n}\right) \\ &= P\left(T_{n-1, -\sqrt{n} B_L} \geq \frac{\bar{x} - x_0}{s / \sqrt{n}}\right). \end{aligned} \quad (10)$$

---

<sup>4</sup>It is implicit that the upper bound equals 1 in the probability statement of equation 8, i.e.,  $1 - \alpha = P(B_L \leq R \leq 1)$ .

In equation 10,  $T_{n-1, -\sqrt{n}B_L}$  is a non-central t random variable with  $n-1$  degrees of freedom and non-centrality parameter  $-\sqrt{n}B_L$  (see for example, Casella and Berger, 1990). However, a non-central t random variable with non-centrality parameter  $-\sqrt{n}B_L$  is the mirror image of a non-central t random variable with non-centrality parameter  $\sqrt{n}B_L$  (Johnson and Kotz, 1970).

Therefore,

$$1 - \alpha = P\left(T_{n-1, -\sqrt{n}B_L} \geq \frac{\bar{x} - x_0}{s/\sqrt{n}}\right) = P\left(T_{n-1, \sqrt{n}B_L} \leq \frac{x_0 - \bar{x}}{s/\sqrt{n}}\right). \quad (11)$$

The final probability expression of equation 11 is the CDF of a non-central t random variable with  $n-1$  degrees of freedom and non-centrality parameter  $\sqrt{n}B_L$ , i.e.,

$$1 - \alpha = G_{n-1, \sqrt{n}B_L}\left(\frac{x_0 - \bar{x}}{s/\sqrt{n}}\right). \quad (12)$$

A lower confidence bound for  $\theta = \frac{x_0 - \mu}{\sigma}$ , is  $B_L$ , the solution to equation 12. When we exploit the strict monotonicity of the normal distribution function, a  $(1 - \alpha)100\%$  LCB for the probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ , is  $\Phi(B_L)$ .

## 4. Upper Confidence Bound (UCB)

A UCB would represent an “optimistic” bound on the true probability of non-perforation. As such, it is of little practical value and rarely calculated. However, its derivation is briefly discussed here for completeness and as a precursor to the development of a two-sided confidence interval.

Following a similar progression to that of the previous section, a  $(1 - \alpha)100\%$  UCB for  $\theta = \frac{x_0 - \mu}{\sigma}$  is that value  $B_U$  satisfying

$$1 - \alpha = P(R \leq B_U) = P\left(T_{n-1, \sqrt{n}B_U} \geq \frac{x_0 - \bar{x}}{s/\sqrt{n}}\right).$$

Therefore,

$$\alpha = G_{n-1, \sqrt{n}B_U}\left(\frac{x_0 - \bar{x}}{s/\sqrt{n}}\right), \quad (13)$$

and a  $(1-\alpha)100\%$  UCB for the probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ , is  $\Phi(B_U)$ .

---

## 5. Two-Sided Confidence Interval

---

A two-sided confidence interval might be of interest to the researcher simultaneously wanting lower and upper limits for the probability of non-perforation, neither of which is automatically set to their extreme value of 0 or 1, respectively.

A  $(1-\alpha)100\%$  two-sided confidence interval for  $\theta = \frac{x_0 - \mu}{\sigma}$  is given by values  $C_L$  and  $C_U$  satisfying

$$\frac{\alpha}{2} = P(R \leq C_L) \text{ and } \frac{\alpha}{2} = P(R \geq C_U)$$

Following steps similar to those used in determining one-sided confidence bounds, two-sided confidence limits for  $\theta = \frac{x_0 - \mu}{\sigma}$  are solutions to

$$1 - \frac{\alpha}{2} = G_{n-1, \sqrt{n}C_L} \left( \frac{x_0 - \bar{x}}{s/\sqrt{n}} \right), \quad (14)$$

and

$$\frac{\alpha}{2} = G_{n-1, \sqrt{n}C_U} \left( \frac{x_0 - \bar{x}}{s/\sqrt{n}} \right). \quad (15)$$

Thus, a  $(1-\alpha)100\%$  two-sided confidence interval for the probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ , is  $(\Phi(C_L), \Phi(C_U))$ .

---

## 6. Iterative Solutions for the Confidence Bounds and Confidence Limits

---

As an example, consider solving equation 12 for the LCB of  $\theta = \frac{x_0 - \mu}{\sigma}$ . We start by rewriting the equation so that the right side equals 0:

$$G_{n-1, \sqrt{n}B_L} \left( \frac{x_0 - \bar{x}}{s/\sqrt{n}} \right) + \alpha - 1 = 0. \quad (16)$$

In equation 16,  $n$ ,  $x_0$ ,  $\bar{x}$ ,  $s$ , and  $\alpha$  are fixed constants, and the only unknown is  $B_L$ . Therefore,

if we let  $H(B_L; n, x_0, \bar{x}, s, \alpha) = G_{n-1, \sqrt{n}B_L} \left( \frac{x_0 - \bar{x}}{s/\sqrt{n}} \right) + \alpha - 1$ , the problem becomes one of solving for

the root of the function  $H$ . We can show  $H$  to be a monotonic decreasing function in  $B_L$  by recognizing that the non-central t distribution function is monotonic decreasing in the non-centrality parameter (see figure 1) or by formal proof (see appendix A). Therefore, one can use the bisection method or other root-finding algorithm to solve for  $B_L$ . Finally, the LCB for the probability of non-perforation is  $\Phi(B_L)$ .

The bisection method can be easily programmed into most mathematical software packages. MATLAB<sup>5</sup> and Mathematica<sup>6</sup> programs appear in appendices B through G for calculating an LCB, a UCB, and a two-sided confidence interval for  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ .




Figure 1. Two non-central t density functions,  $f_1$  and  $f_2$ , with respective non-centrality parameters  $\lambda_1$  and  $\lambda_2$ . (As the non-centrality parameter increases, the area to the right of  $x_0$  under the non-central t curve decreases.)

<sup>5</sup>MATLAB is a registered trademark of The MathWorks.

<sup>6</sup>Mathematica is a registered trademark of Wolfram Research, Inc.

---

## 7. Confidence Bands

---

By plotting the function  $\Phi\left(\frac{x-\bar{x}}{s}\right)$  for all real numbers  $x$ , one obtains a smooth, monotonically increasing function that is a simple estimate for the entire CDF. A lower confidence band for the CDF is a plot of the function  $\Phi(B_L(x))$ . This band lies below the estimated CDF and enables one to see the region above the band in which the entire actual CDF lies (with the stated level of confidence). If one fixes the probability level on the vertical axis, then the corresponding x-axis value from the lower confidence band is an upper tolerance bound. In the context of armor design, the upper tolerance bound is very important as it states with a specified level of confidence what armor thickness is needed to stop a (high) percentage of projectiles from perforating the materiel. A MATLAB program for generating a lower confidence band appears in appendix H.

Upper confidence bands and two-sided confidence bands are constructed in a similar fashion; however, they are likely to be of much less interest to the armor designer. Although MATLAB programs for their construction are not included here, the interested reader could easily tailor the program in appendix H to accomplish this.

---

## 8. An Application

---

Table 1 gives the penetrations of 14 projectiles into an extended armor pack. Engineers plan to use 115 units as the armor thickness. To determine if this thickness will provide enough protection, an estimate is desired for the probability that the next projectile fired will not penetrate deeper than 115 units. Letting  $X$  be the penetration of this next projectile, we seek an estimate for  $P(X \leq 115)$ .

Table 1. Penetration depths into armor (no units specified).

|    |    |    |     |     |
|----|----|----|-----|-----|
| 47 | 81 | 89 | 99  | 114 |
| 59 | 86 | 90 | 100 | 118 |
| 80 | 88 | 90 | 113 |     |

The data are first checked for normality by Lilliefors Test (see Conover, 1980). The value of the test statistic is 0.1713, which corresponds to a P-value greater than 0.20. Therefore, the assumption of normality is not rejected.

Summary statistics for the sample data are  $\bar{x} = 89.571$  and  $s^2 = 390.725$ . Equation 2 is used to obtain a point estimate for  $P(X \leq 115)$ . The solution is

$$\hat{P}(X \leq 115) = \Phi\left(\frac{115 - 89.571}{\sqrt{390.725}}\right) = \Phi(1.286) = 0.901.$$

To achieve a 95% LCB for  $P(X \leq 115)$ , we start by using equation 12:

$$.95 = G_{13, \sqrt{14}B_L}(4.813).$$

Using the bisection method, we obtain a solution of  $B_L = 0.6672$ . Therefore, a 95% LCB for  $P(X \leq 115)$  is  $\Phi(0.6672) = .7477$ . That is, for an armor of thickness 115 units, one can be 95% confident that the probability of non-perforation is at least 74.77%.

At this point, a reasonable question might be “With 95% confidence, what armor thickness would offer at least 90% protection against perforation?” The lower confidence band for  $\Phi(B_L)$ , shown in figure 2, will help determine this. We begin by extending a horizontal (dashed) line from the “y”-axis at  $P(\text{non-perforation}) = .90$  to the lower confidence band and then dropping to the x-axis to find the desired armor thickness. This thickness of 131.25 is the 90% upper tolerance bound.




Figure 2. Estimated probability of non-perforation and 95% LCB as a function of armor thickness. (The line formed by the bounds is referred to as the lower confidence band. From the band, one is able to determine what armor thickness yields a certain minimum probability of non-perforation, e.g., with 95% confidence, one can state that a thickness of 131.25 will stop at least 90% of projectiles from perforating the material.)

A 95% two-sided confidence interval for  $P(X \leq 115)$  is calculated from equations 12 and 13. It has a lower limit given by  $\Phi(C_L)$  in which  $C_L$  is the solution to  $1 - \frac{\alpha}{2} = G_{n-1, \sqrt{n}C_L} \left( \frac{x_0 - \bar{x}}{s/\sqrt{n}} \right)$ , or  $.975 = G_{13, \sqrt{14}C_L} (4.813)$ . The upper limit is given by  $\Phi(C_U)$  in which  $C_U$  is the solution to  $\frac{\alpha}{2} = G_{n-1, \sqrt{n}C_U} \left( \frac{x_0 - \bar{x}}{s/\sqrt{n}} \right)$ , or  $.025 = G_{13, \sqrt{14}C_U} (4.813)$ . Using the bisection method again, we obtain  $C_L = .5566$  and  $C_U = 1.9901$ . Therefore, the lower confidence limit is  $\Phi(.5566) = .7111$ , and the upper confidence limit is  $\Phi(1.9901) = .9767$ . We can be 95% confident that the true probability of non-perforation is between 71.11% and 97.67%.

---

## 9. Summary

---

Reporting an interval estimate along with a point estimate gives designers a sense of the error of estimation which is a function of the sample size used in the study and the inherent variability of each projectile-armor interaction. Using a generalized pivotal quantity, we have achieved an interval estimate for the probability that a homogeneous armor plate of specified thickness will successfully stop a projectile from perforating. In the derivation of this interval estimate, we have assumed that the penetration depths are random and that they follow a normal distribution of unknown mean and variance.

The interval limit (or limits if a two-sided interval is desired) is a function of the sample mean and sample standard deviation of penetration depths, the number of shots fired, the thickness of the armor and percentiles of a non-central t distribution. However, since the non-centrality parameter associated with this distribution is a function of the interval limit, numerical methods are required to obtain the final solution. MATLAB and Mathematica codes to perform these calculations are provided in appendices B through G.

Confidence bands that graphically display the entire the relationship between armor width and bounds on the probability of non-perforation follow naturally. These bands may be of help to armor designers in selecting an armor width that will provide a high degree of protection against specific enemy threats.

---

## 10. References

---

- Barton, D.E. Unbiased estimation of a set of probabilities. *Biometrika* **1961**, 48, 227–229.
- Casella, G.; Berger, R.L. *Statistical Inference*, 2nd Ed., Duxbury Press: Belmont, CA, 1990.
- Conover, W. J. *Practical Nonparametric Statistics*, 2nd ed.; John Wiley & Sons: New York, 1980.
- Hahn G. J.; Meeker, W. Q. *Statistical Intervals: A Guide for Practitioners*, John Wiley & Sons: New York, 1991.
- Johnson, N. L.; Kotz, S. *Distributions in Statistics --- Continuous Univariate Distributions - 2*, John Wiley & Sons: New York, 1970.
- Lieberman, G. J.; Resnikoff, G. J. Sampling plans for inspection by variables. *Journal of the American Statistical Association* **1955**, 50, 457–516.
- Mood, A.; Graybill, F.; Boes, D. *Introduction to the Theory of Statistics*, 3rd ed.; McGraw-Hill: New York, 1974.
- Odeh, R.E.; Owen, D.B. *Tables for Normal Tolerance Limits, Sampling Plans, and Screening*, Marcel Dekker, Inc.: New York, 1980.
- Owen, D. B.; Hua, T. A. *Tables of Confidence Limits on the Tail Area of the Normal Distribution, Communications in Statistics – Simulation and Computation B* **1977**, 6, 285–311.
- Weerahandi, S. Generalized Confidence Intervals, *Journal of the American Statistical Association* **1993**, 88, 899–905.
- Weerahandi, S. *Exact Statistical Methods for Data Analysis*, Springer: New York, 1995.
- Weerahandi, S. *General Inference in Repeated Measures*, John Wiley & Sons: New York, 2004.

INTENTIONALLY LEFT BLANK

---

## Appendix A. Proof of the Monotonicity of the Function H

---

Show that  $H(\cdot)$  is a monotonic decreasing function. That is, show that if  $\lambda_1 < \lambda_2$ , then  $H(\lambda_1) > H(\lambda_2)$ .

Proof:

Let  $n, x_0, \bar{x}, s, \alpha, \lambda_1$ , and  $\lambda_2$  be fixed constants with  $\lambda_1 < \lambda_2$ . Let  $X_1$  and  $Y$  be independent random variables,  $X_1$  being normally distributed with mean  $\lambda_1$  and variance 1, and  $Y$  following a chi-squared distribution with  $n-1$  degrees of freedom. The support of  $X_1$  is the set of all real numbers, and the support of  $Y$  is the set of all positive reals.

Now define  $W = \sqrt{\frac{nY}{n-1}} \frac{x_0 - \bar{x}}{s}$ . The support of  $W$  is either the set of all positive real numbers (if  $x_0 \geq \bar{x}$ ) or the set of all non-positive real numbers (if  $x_0 < \bar{x}$ ). However, in general, we denote the support of  $W$  by  $\Theta$ .

$P(X_1 < W)$  can be expressed as a double integral of the joint density of  $X_1$  and  $W$ , namely,

$$P(X_1 < W) = \int_{\Theta} \int_{-\infty}^w f_{X_1}(x) f_W(w) dx dw. \quad (\text{A-1})$$

The double integral of equation A-1, can be rewritten as the sum of two double integrals:

$$P(X_1 < W) = \int_{\Theta} \int_{-\infty}^{w-(\lambda_2-\lambda_1)} f_{X_1}(x) f_W(w) dx dw + \int_{\Theta} \int_{w-(\lambda_2-\lambda_1)}^w f_{X_1}(x) f_W(w) dx dw. \quad (\text{A-2})$$

Each of these double integrals is positive, so by dropping the latter of the two, we obtain the inequality,

$$P(X_1 < W) > \int_{\Theta} \int_{-\infty}^{w-(\lambda_2-\lambda_1)} f_{X_1}(x) f_W(w) dx dw. \quad (\text{A-3})$$

However, the double integral in equation A-3, represents a probability, namely,  $P(X_1 < W - (\lambda_2 - \lambda_1))$ , which in turn equals  $P(X_1 + \lambda_2 - \lambda_1 < W)$ . So,

$$P(X_1 < W) > P(X_1 + \lambda_2 - \lambda_1 < W). \quad (\text{A-4})$$

Now define  $X_2 = X_1 + (\lambda_2 - \lambda_1)$ ; then  $X_2$  is normally distributed with mean  $\lambda_2$  and variance 1. Therefore,

$$P(X_1 < W) > P(X_2 < W), \quad (\text{A-5})$$

Consider the standard normal random variable,  $Z$ , having mean 0 and variance 1. Employing the equalities  $X_1 = Z + \lambda_1$  and  $X_2 = Z + \lambda_2$  and the definition of  $W$ , we expand the above result to obtain

$$P\left(Z + \lambda_1 < \sqrt{\frac{nY}{n-1}} \frac{x_0 - \bar{x}}{s}\right) > P\left(Z + \lambda_2 < \sqrt{\frac{nY}{n-1}} \frac{x_0 - \bar{x}}{s}\right). \quad (\text{A-6})$$

Rearranging the expressions inside each of the probability statements,

$$P\left(\frac{Z + \lambda_1}{\sqrt{Y/n-1}} < \frac{x_0 - \bar{x}}{s/\sqrt{n}}\right) > P\left(\frac{Z + \lambda_2}{\sqrt{Y/n-1}} < \frac{x_0 - \bar{x}}{s/\sqrt{n}}\right). \quad (\text{A-7})$$

By the definition of a non-central t random variable,

$$P\left(T_{n-1, \lambda_1} < \frac{x_0 - \bar{x}}{s/\sqrt{n}}\right) > P\left(T_{n-1, \lambda_2} < \frac{x_0 - \bar{x}}{s/\sqrt{n}}\right). \quad (\text{A-8})$$

Each of these probabilities is the cumulative non-central t distribution function evaluated at  $\frac{x_0 - \bar{x}}{s/\sqrt{n}}$ :

$$G_{n-1, \lambda_1}\left(\frac{x_0 - \bar{x}}{s/\sqrt{n}}\right) > G_{n-1, \lambda_2}\left(\frac{x_0 - \bar{x}}{s/\sqrt{n}}\right). \quad (\text{A-9})$$

Next, we add the constant  $\alpha - 1$  to both sides of the inequality:

$$G_{n-1, \lambda_1}\left(\frac{x_0 - \bar{x}}{s/\sqrt{n}}\right) + \alpha - 1 > G_{n-1, \lambda_2}\left(\frac{x_0 - \bar{x}}{s/\sqrt{n}}\right) + \alpha - 1. \quad (\text{A-10})$$

Using the definition of the function  $H(\cdot)$ , we obtain the desired result.

$$H(\lambda_1) > H(\lambda_2). \quad (\text{A-11})$$

---

## Appendix B. MATLAB Code for Probability of Non-perforation LCB

---

MATLAB Release 14, Version 7, code for calculating an LCB for the probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ . The user may supply his or her own data and values for alpha, x0, and epsilon.

```
% Define variables %
alpha=.05;    x0=115;    epsilon=.000001;

% Declare data %
data=[47 59 80 81 86 88 89 90 90 99 100 113 114 118]';

% Calculate summary statistics and define K %
n=length(data);    xbar=mean(data);    sigma=std(data);
K=sqrt(n)*(x0- xbar)/sigma;

% Initialize flag and bounds on non-centrality parameter %
flag=0;    lononcent=xbar-10*sigma;    hinoncent=xbar+10*sigma;

% Execute bisection method %
noncent=(lononcent+hinoncent)/2;
while flag==0
    if hinoncent-lononcent < epsilon
        flag=1;
    elseif nctcdf(K,n-1,noncent)+alpha-1 < 0
        hinoncent=noncent;    noncent=(noncent+lononcent)/2;
    else
        lononcent=noncent;    noncent=(noncent+hinoncent)/2;
    end
end

%Print lower confidence bound for probability of non-perforation %
fprintf('%3.0f%% LCB for P(non-perforation) = %6.4f\n',...
    100*(1-alpha),normcdf(noncent/sqrt(n)))
```

INTENTIONALLY LEFT BLANK

---

## Appendix C. Mathematica Code for Probability of Non-perforation LCB

---

Mathematica Version 5.1 code for calculating an LCB for the probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ . The user may supply his or her own data and values for alpha, x0, and epsilon.

```
(* Remove warning messages that do not affect results but clutter output *)
Off[General::spell]; Off[General::spell1];
Off[NIntegrate::slwcon]; Off[NIntegrate::ncvb];

(* Enable use of statistical tools *)
<<Statistics`ContinuousDistributions`
<<Statistics`NormalDistribution`

(* Clear variables for use *)
Clear[data,n,noncent,x0,a,epsilon,lononcent,hinoncent,flag,K,ndist]

(* Define variables *)
alpha=.05; x0=115; epsilon=.000001;

(* Declare data *)
data={47,59,80,81,86,88,89,90,90,99,100,113,114,118};

(* Calculate summary statistics and define K *)
n=Length[data]; xbar=Mean[data]/N; sigma=StandardDeviation[data]/N;
K=(x0-xbar)/(sigma/Sqrt[n])/N;

(* Initialize flag and bounds on non-centrality parameter *)
flag=0; lononcent=xbar-10sigma/N; hinoncent=xbar+10sigma/N;

(* Execute bisection method *)
H:=CDF[NoncentralStudentTDistribution[n-1,noncent],K]+alpha-1
While[flag==0,noncent=Mean[{lononcent,hinoncent}];
  If[hinoncent-lononcent<epsilon,flag=1,
  If[H<0,hinoncent=noncent,lononcent=noncent]]]

(* Print lower confidence bound for probability of non-perforation *)
Print[TraditionalForm[StringForm["``% LCB for P(non-perforation) = ``",
  100 (1-alpha),CDF[NormalDistribution[0,1],noncent/Sqrt[n]]]]]
```

INTENTIONALLY LEFT BLANK

---

## Appendix D. MATLAB Code for Probability of Non-perforation UCB

---

MATLAB Release 14, Version 7, code for calculating a UCB for the probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ . The user may supply his or her own data and values for alpha, x0, and epsilon.

```
% Define variables %
alpha=.05;    x0=115;    epsilon=.000001;

% Declare data %
data=[47 59 80 81 86 88 89 90 90 99 100 113 114 118]';

% Calculate summary statistics and define K %
n=length(data);    xbar=mean(data);    sigma=std(data);
K=sqrt(n)*(x0- xbar)/sigma;

% Initialize flag and bounds on non-centrality parameter %
flag=0;    lononcent=xbar-10*sigma;    hinoncent=xbar+10*sigma;

% Execute bisection method %
noncent=(lononcent+hinoncent)/2;
while flag==0
    if hinoncent-lononcent < epsilon
        flag=1;
    elseif nctcdf(K,n-1,noncent)-alpha < 0
        hinoncent=noncent;    noncent=(noncent+lononcent)/2;
    else
        lononcent=noncent;    noncent=(noncent+hinoncent)/2;
    end
end

% Print upper confidence bound for probability of non-perforation %
fprintf('%3.0f%% UCB for P(non-perforation) = %6.4f\n',...
    100*(1-alpha),normcdf(noncent/sqrt(n)))
```

INTENTIONALLY LEFT BLANK

---

## Appendix E. Mathematica Code for Probability of Non-perforation UCB

---

Mathematica Version 5.1 code for calculating a UCB for the probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ . The user may supply his or her own data and values for alpha, x0, and epsilon.

```
(* Remove warning messages that do not affect results but clutter output *)
Off[General::spell]; Off[General::spell1];
Off[NIntegrate::slwcon]; Off[NIntegrate::ncvb];

(* Enable use of statistical tools *)
<<Statistics`ContinuousDistributions`
<<Statistics`NormalDistribution`

(* Clear variables for use *)
Clear[data,n,noncent,x0,a,epsilon,lononcent,hinoncent,flag,K,ndist]

(* Define variables *)
alpha=.05; x0=115; epsilon=.000001;

(* Declare data *)
data={47,59,80,81,86,88,89,90,90,99,100,113,114,118};

(* Calculate summary statistics and define K *)
n=Length[data]; xbar=Mean[data]/N; sigma=StandardDeviation[data]/N;
K=(x0-xbar)/(sigma/Sqrt[n])/N;

(* Initialize flag and bounds on non-centrality parameter *)
flag=0; lononcent=xbar-10sigma/N; hinoncent=xbar+10sigma/N;

(* Execute bisection method *)
H:=CDF[NoncentralStudentTDistribution[n-1,noncent],K]-alpha
While[flag==0,noncent=Mean[{lononcent,hinoncent}];
  If[hinoncent-lononcent<epsilon,flag=1,
    If[H<0,hinoncent=noncent,lononcent=noncent]]]

(* Print upper confidence bound for probability of non-perforation *)
Print[TraditionalForm[StringForm["``% UCB for P(non-perforation) = ``",
  100 (1-alpha),CDF[NormalDistribution[0,1],noncent/Sqrt[n]]]]]
```

INTENTIONALLY LEFT BLANK

---

## Appendix F. MATLAB Code for Probability of Perforation Two-Sided Confidence Interval

---

MATLAB Release 14, Version 7, code for calculating a two-sided confidence interval for the probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ . The user may supply his or her own data and values for alpha, x0, and epsilon.

```
% Define variables %
alpha=.05;    x0=115;    epsilon=.000001;

% Declare data %
data=[47 59 80 81 86 88 89 90 90 99 100 113 114 118]';

% Calculate summary statistics and define K %
n=length(data);    xbar=mean(data);    sigma=std(data);
K=sqrt(n)*(x0- xbar)/sigma;

% Initialize flag and bounds on non-centrality parameter %
flag=0;    lononcent=xbar-10*sigma;    hinoncent= xbar+10*sigma;

% Execute bisection method for lower limit %
noncent=(lononcent+hinoncent)/2;
while flag==0
    if hinoncent-lononcent < epsilon
        flag=1;
    elseif nctcdf(K,n-1,noncent)+alpha/2-1 < 0
        hinoncent=noncent;    noncent=(noncent+lononcent)/2;
    else
        lononcent=noncent;    noncent=(noncent+hinoncent)/2;
    end
end

% Print lower confidence limit for probability of non-perforation %
fprintf('%4.0f%% CI for P(non-perforation) = (%6.4f, ',...
        100*(1-alpha),normcdf(noncent/sqrt(n)))

% Re-initialize flag and bounds on non-centrality parameter %
flag=0;    lononcent=xbar-10*sigma;    hinoncent= xbar+10*sigma;
```

```

% Execute bisection method for upper limit %
noncent=(lononcent+hinoncent)/2;
while flag==0
    if hinoncent-lononcent < epsilon
        flag=1;
    elseif nctcdf(K,n-1,noncent)-alpha/2 < 0
        hinoncent=noncent; noncent=(noncent+lononcent)/2;
    else
        lononcent=noncent; noncent=(noncent+hinoncent)/2;
    end
end

% Print upper confidence limit for probability of non-perforation %
fprintf('%6.4f)\n',normcdf(noncent/sqrt(n)))

```

---

## Appendix G. Mathematica Code for Probability of Perforation Two-Sided Confidence Interval

---

Mathematica Version 5.1 code for calculating a two-sided confidence interval for the probability of non-perforation,  $\Phi\left(\frac{x_0 - \mu}{\sigma}\right)$ . The user may supply his or her own data and values for alpha, x0, and epsilon.

```
(* Remove warning messages that do not affect results but clutter output *)
Off[General::spell]; Off[General::spell1];
Off[NIntegrate::slwcon]; Off[NIntegrate::ncvb];

(* Enable use of statistical tools *)
<<Statistics`ContinuousDistributions`
<<Statistics`NormalDistribution`

(* Clear variables for use *)
Clear[data,n,noncent,x0,a,epsilon,lononcent,hinoncent,flag,K,
      ndist,lowerlimit,upperlimit]

(* Define variables *)
alpha=.05; x0=115; epsilon=.000001;

(* Declare data *)
data={47,59,80,81,86,88,89,90,90,99,100,113,114,118};

(* Calculate summary statistics and define K *)
n=Length[data]; xbar=Mean[data]/N; sigma=StandardDeviation[data]/N;
K=(x0-xbar)/(sigma/Sqrt[n])/N;

(* Initialize flag and bounds on non-centrality parameter *)
flag=0; lononcent=xbar-10sigma/N; hinoncent=xbar+10sigma/N;

(* Execute bisection method for lower limit *)
H:=CDF[NoncentralStudentTDistribution[n-1,noncent],K]+alpha/2-1
While[flag==0,noncent=Mean[{lononcent,hinoncent}];
      If[hinoncent-lononcent<epsilon,flag=1,
```

```

    If[H<0,hinoncent=noncent,lononcent=noncent]]]
lowerlimit = N[CDF[NormalDistribution[0,1],noncent/Sqrt[n]]];

(* Re-initialize flag and bounds on non-centrality parameter *)
flag=0; lononcent=xbar-10sigma//N; hinoncent=xbar+10sigma//N;

(* Execute bisection method for upper limit *)
H:=CDF[NoncentralStudentTDistribution[n-1,noncent],K]-alpha/2
While[flag==0,noncent=Mean[{lononcent,hinoncent}]];
    If[hinoncent-lononcent<epsilon,flag=1,
        If[H<0,hinoncent=noncent,lononcent=noncent]]]
upperlimit = N[CDF[NormalDistribution[0,1],noncent/Sqrt[n]]];

(* Print confidence interval for probability of non-perforation *)
Print[TraditionalForm[StringForm["``0% CI for P(non-perforation) = (``,``)",
    100 (1-alpha),lowerlimit,upperlimit]]]

```

---

## Appendix H. MATLAB Code for Probability of Non-perforation CDF and Lower Confidence Band

---

MATLAB Release 14, Version 7, code for plotting an estimated CDF and lower confidence band. The user may supply his or her own data and values for alpha, x0, and epsilon.

```
% Define variables %
alpha=.05;    epsilon=.0001;

% Declare data %
data=[47 59 80 81 86 88 89 90 90 99 100 113 114 118]';

% Calculate summary statistics %
n=length(data);  Xbar=mean(data);  SDev=std(data);

% Initialize counter for points along the X-axis %
i=0;

% Declare armor thickness %
for x0=Xbar-3*SDev:SDev/10:Xbar+4*SDev
    i=i+1;  x(i)=x0;

    % Calculate plug-in estimate for probability of non-perforation %
    pointest(i)=normcdf((x0-Xbar)/SDev);

    % Define K %
    K=sqrt(n)*(x0-Xbar)/SDev;

    % Initialize flag and bounds on non-centrality parameter %
    flag=0;  lononcent=-1000;  hinoncent=1000;

    % Execute bisection method for lower confidence bound %
    noncent=(lononcent+hinoncent)/2;
    while flag==0
        if hinoncent-lononcent < epsilon
            flag=1;
        elseif nctcdf(K,n-1,noncent)-1+alpha < 0
```

```

        hinoncent=noncent;  noncent=(noncent+lononcent)/2;
    else
        lononcent=noncent;  noncent=(noncent+hinoncent)/2;
    end
end
lcb(i)=normcdf(noncent/sqrt(n));
end

% as a function of armor thickness, plot non-perforation probability curve %
% in blue, and lower confidence band for probability of non-perforation in %
% red %
plot(x,pointest,'b-',x,lcb,'r-')
xlabel('ARMOR THICKNESS'); ylabel('P(NON-PERFORATION)')
text(Xbar-.25*SDev,.5,'Estimated','Color','b','Rotation',69,...
    'HorizontalAlignment','Center')
text(Xbar+.75*SDev,.5,'LC Band','Color','r','Rotation',67,...
    'HorizontalAlignment','Center')

```

---

## List of Symbols Used

---

### Scalar Values

|           |                                                                                                                            |
|-----------|----------------------------------------------------------------------------------------------------------------------------|
| $n$       | number of observations in sample data                                                                                      |
| $r_i$     | the $i^{th}$ observed value of the random variable $R$                                                                     |
| $r_{(i)}$ | the $i^{th}$ ordered, observed value of the random variable $R$                                                            |
| $s$       | observed sample standard deviation                                                                                         |
| $x_i$     | the $i^{th}$ observed value from a data set; in this paper, the observed depth of penetration from the $i^{th}$ projectile |
| $x_0$     | a specified thickness of armor                                                                                             |
| $\bar{x}$ | observed sample mean                                                                                                       |
| $z_p$     | the value from the standard normal distribution having an area of $p$ to its right under the density curve                 |
| $\alpha$  | level of significance                                                                                                      |
| $\sigma$  | population standard deviation for a normally distributed random variable                                                   |
| $\theta$  | parameter for which a confidence interval is constructed; may actually be a function of one or more parameters             |
| $\mu$     | population mean for a normally distributed random variable                                                                 |
| $\xi$     | a set of one or more nuisance parameters                                                                                   |

### Random Variables

|           |                                                                                                                                  |
|-----------|----------------------------------------------------------------------------------------------------------------------------------|
| $R( )$    | generalized test variable; a function of random data, observed data, parameter of interest and perhaps other nuisance parameters |
| $S$       | sample standard deviation of depths of penetration                                                                               |
| $T$       | student's $t$ random variable                                                                                                    |
| $V$       | chi-square random variable                                                                                                       |
| $X$       | depth of penetration for a randomly selected projectile                                                                          |
| $\bar{X}$ | sample mean depth of penetration                                                                                                 |

$Y$  a normal random variable with mean zero and variance  $\frac{1}{n}$

$Z$  a standard normal random variable, with mean zero and variance one

### Functions

$F( )$  cumulative distribution function

$\hat{F}( )$  estimated cumulative distribution function

$G( )$  student's t cumulative distribution function

$\Phi( )$  standard normal cumulative distribution function

### Miscellaneous

$B_L$  lower confidence bound

$B_U$  upper confidence bound

$C_L$  lower confidence limit

$C_U$  upper confidence limit

$P( )$  probability of the parenthesized expression

$\hat{P}( )$  estimated probability of the parenthesized expression

$\Sigma$  summation

$\wedge$  estimate of

NO. OF  
COPIES ORGANIZATION

- \* ADMINISTRATOR  
DEFENSE TECHNICAL INFO CTR  
ATTN DTIC OCA  
8725 JOHN J KINGMAN RD STE 0944  
FT BELVOIR VA 22060-6218  
\*pdf file only
- 1 DIRECTOR  
US ARMY RSCH LABORATORY  
ATTN IMNE ALC IMS MAIL & REC MGMT  
2800 POWDER MILL RD  
ADELPHI MD 20783-1197
- 1 DIRECTOR  
US ARMY RSCH LABORATORY  
ATTN AMSRD ARL CI OK TL TECH LIB  
2800 POWDER MILL RD  
ADELPHI MD 20783-1197
- 1 US ARMY RESEARCH OFFICE  
MATHEMATICS DIVISION  
ATTN H CHANG  
PX BOX 12211  
RSCH TRIANGLE PARK NC 27709-1221

ABERDEEN PROVING GROUND

- 1 DIRECTOR  
US ARMY RSCH LABORATORY  
ATTN AMSRD ARL CI OK (TECH LIB)  
BLDG 4600
- 1 US ARMY EVALUATION CTR  
ATTN N DUNN  
BLDG 4120
- 1 DIRECTOR  
US ARMY RSCH LABORATORY  
ATTN AMSRD ARL CI CT B BODT  
BLDG 321
- 1 DIRECTOR  
US ARMY RSCH LABORATORY  
ATTN AMSRD ARL WM B R COATES  
BLDG 309
- 1 DIRECTOR  
US ARMY RSCH LABORATORY  
ATTN AMSRD ARL WM M ZOLTOSKI  
BLDG 4600

NO. OF  
COPIES ORGANIZATION

- 7 DIRECTOR  
US ARMY RSCH LABORATORY  
ATTN AMSRD ARL WM TA T HAVEL (4 CYS)  
D KLEPONIS J RUNYON  
S SCHOENFELD  
BLDG 393
- 1 DIRECTOR  
US ARMY RSCH LABORATORY  
ATTN AMSRD ARL WM TC L MAGNESS  
BLDG 309