

REPORT DOCUMENTATION PAGEForm Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY) 2005		2. REPORT TYPE Open Literature		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE A Convenient Fluorometric Method to Study Sulfur Mustard-induced Apoptosis in Human Epidermal Keratinocytes Monolayer Microplate Culture				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Ray, R, Hauck, S, Kramer, R, Benton, B				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) US Army Medical Research Institute of Chemical Defense ATTN: MCMR-UV-PB 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5400				8. PERFORMING ORGANIZATION REPORT NUMBER USAMRICD-P04-018	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) US Army Medical Research Institute of Chemical Defense ATTN: MCMR-UV-RC 3100 Ricketts Point Road Aberdeen Proving Ground, MD 21010-5400				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Published in Drug and Chemical Toxicology, 1, 105-116, 2005.					
14. ABSTRACT See reprint.					
15. SUBJECT TERMS Apoptosis, fluorometric apoptosis assay, human epidermal keratinocytes, monolayer culture, sulfur mustard					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UNLIMITED	18. NUMBER OF PAGES 12	19a. NAME OF RESPONSIBLE PERSON Radharaman Ray
a. REPORT UNLIMITED	b. ABSTRACT UNLIMITED	c. THIS PAGE UNLIMITED			19b. TELEPHONE NUMBER (include area code) 410-436-3074

A Convenient Fluorometric Method to Study Sulfur Mustard-Induced Apoptosis in Human Epidermal Keratinocytes Monolayer Microplate Culture

Radharaman Ray, Stephanie Hauck, Rachel Kramer,
and Betty Benton

Biochemical Pharmacology Branch, Pharmacology Division, U.S. Army Medical Research Institute of Chemical Defense, Aberdeen Proving Ground, Maryland, USA

Sulfur mustard [SM; bis-(2-chloroethyl) sulfide], which causes skin blistering or vesication [(1991). Histo- and cytopathology of acute epithelial lesions. In: Papirmeister, B., Feister, A. J., Robinson, S. I., Ford, R. D., eds. *Medical Defense Against Mustard Gas: Toxic Mechanisms and Pharmacological Implications*. Boca Raton: CRC Press, pp. 43–78.], is a chemical warfare agent as well as a potential terrorism agent. SM-induced skin blistering is believed to be due to epidermal–dermal detachment as a result of epidermal basal cell death via apoptosis and/or necrosis. Regarding the role of apoptosis in SM pathology in animal skin, the results obtained in several laboratories, including ours, suggest the following: 1) cell death due to SM begins via apoptosis that proceeds to necrosis via an apoptotic–necrotic continuum and 2) inhibiting apoptosis decreases SM-induced microvesication in vivo. To study the mechanisms of SM-induced apoptosis and its prevention in vitro, we have established a convenient fluorometric apoptosis assay using monolayer human epidermal keratinocytes (HEK) adaptable for multiwell plates (24-, 96-, or 384-well) and high-throughput applications. This assay allows replication and multiple types of experimental manipulation in

Disclaimer: The opinions or assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the U.S. Army or the Department of Defense.

Address correspondence to Radharaman Ray, Biochemical Pharmacology Branch, Pharmacology Division, U.S. Army Medical Research Institute of Chemical Defense, Aberdeen Proving Ground, MD 21010-5400, USA; Fax: (410) 436-1960; E-mail: radharaman.ray@apg.amedd.army.mil

20060126 061

sister cultures so that the apoptotic mechanisms and the effects of test compounds can be compared statistically. SM affects diverse cellular mechanisms, including endoplasmic reticulum (ER) Ca^{2+} homeostasis, mitochondrial functions, energy metabolism, and death receptors, each of which can independently trigger apoptosis. However, the biochemical pathway in any of these apoptotic mechanisms is characterized by a pathway-specific sequence of caspases, among which caspase-3 is a key member. Therefore, we exposed 80–90% confluent HEK cultures to SM and monitored apoptosis by measuring the fluorescence generated due to hydrolysis of a fluorogenic caspase-3 substrate (acetyl- or benzyl oxycarbonyl-Asp-Glu-Val-Asp-fluorochrome, also designated as AC-or Z-DEVD- fluorochrome) added to the assay medium. Fluorescence was measured using a plate reader. We used two types of substrates, one (Sigma-Aldrich, CASP-3-F) required cell disruption and the other (Beckman-Coulter CellProbe HT Caspase-3/7 Whole Cell Assay Kit) was cell permeable. The latter substrate was useful in experiments such as determining the time-course of apoptosis immediately following SM exposure without disruption (e.g., due to cell processing). In SM-exposed HEK, fluorescence generated from the fluorogenic caspase-3 substrate hydrolysis increased in a time (0–24 h) and concentration (0.05, 0.1, 0.15, 0.2, 0.3, 0.5 mM) dependent manner. SM caused maximum fluorescence at about 0.5 mM. However, at 2 mM SM, fluorescence decreased compared with 0.5 mM, which remains to be explained. Following 0.3 mM SM exposure, which is considered to be the *in vitro* equivalent of a vesicating dose *in vivo* (Smith, W. J., Sanders, K. M., Ruddle, S. E., Gross, C. L. (1993). Cytometric analysis of DNA changes induced by sulfur mustard. *J. Toxicol.-Cut. Ocular Toxicol.* 12(4):337–347.), a small fluorescence increase was observed at 6 to 8 h, which was markedly higher at 12 h. At 24 h, all SM concentrations increased fluorescence. Fluorescence increase due to SM was prevented 100% by a caspase-3–specific peptide inhibitor AC-DEVD-CHO (acetyl-Asp-Glu-Val-Asp-aldehyde, 0.1 mM), but less effectively by a general caspase inhibitor Z-VAD-FMK (benzyl oxycarbonyl-Val-Ala-Asp-fluoromethylketone, 0.01 mM), indicating that the fluorescence increase was due to caspase-3–mediated apoptosis. These results suggest potential applications of this method to study apoptosis mechanisms involving caspase-3 substrates and possibly those involving other caspase substrates.

Keywords Apoptosis, Fluorometric apoptosis assay, Human epidermal keratinocytes, Monolayer culture, Sulfur mustard.

INTRODUCTION

The term apoptosis is used to describe cell death via modulation of a defined set of biochemical parameters in a cascade of events leading to apoptotic morphological expressions. The major morphological features are cellular shrinkage, nuclear and cytoplasmic condensation, and cellular disintegration followed by prolific budding to produce apoptotic bodies (Kerr et al., 1972; Trump and Berezsky, 1998). As cited in the literature (Kerr et al., 1972; Trump and Berezsky, 1998) and as revealed in our studies (Ray et al., 1999), extensive budding of the cytoplasm is a characteristic morphological feature of

cells undergoing apoptosis. Budding of the cytoplasm results in the formation of apoptotic bodies, which are further characterized, morphologically, as containing membrane-bound nuclear chromatin fragments. Cellular shrinkage, cytoplasmic budding, nuclear fragmentation, and apoptotic bodies are pathognomonic of apoptosis. Confirmation by transmission electron microscopy of cytoplasmic budding bearing nuclear fragments is essential in the differential diagnosis of cellular apoptosis (Kan et al., 2003). Apoptosis contributes to the mechanisms of removal of unwanted cells in normal physiological processes (e.g., during organ development, sloughing of the endometrium in early menstruation period, immune response, etc.). Apoptosis also is the mechanism of cell death responsible for pathological conditions such as cancer, Alzheimer disease, ischemic cardiac damage, oxidative stress, autoimmune syndromes (Hollinger, 2002), or chemical toxicity, for example, due to the highly reactive alkylating compound sulfur mustard (SM) (Rosenthal et al., 1998). SM reacts with multiple targets in the cell and, therefore, causes toxicity via different mechanisms such as DNA damage, disturbance in intracellular calcium homeostasis, abnormal cellular energy metabolism to include oxidative stress, and upregulation of death receptors (Fas) and Fas ligand (Ray et al., 2002). Any of these mechanisms may independently contribute to apoptosis by initially activating a pathway that is characteristic of the mechanism involved, but later converging to a common pathway culminating into the final apoptotic manifestations. There are reports showing that in cultured human epidermal keratinocytes (HEK), SM induces apoptosis via at least two separate mechanisms: 1) an intrinsic mechanism involving mitochondria, Ca^{2+} -calmodulin (Ca^{2+} -CaM), and caspase-3/poly (ADP-ribose) polymerase (PARP), and 2) an extrinsic mechanism involving Fas (death) receptors and Fas ligand (FasL) (Rosenthal et al., 1998, 2003). These reports also show that inhibiting the Ca^{2+} -CaM or the Fas response attenuates the biochemical markers of apoptosis and pathology due to SM. However, it is possible that other mechanisms, for example, involving 1) endoplasmic reticulum (ER) Ca^{2+} and 2) abnormal energy metabolism, may also contribute to SM-induced apoptosis. Our goal is to understand thoroughly the mechanisms of apoptosis due to SM via each of the pathways discussed above and to use the knowledge in developing medical countermeasures against SM toxicity and pathology.

In apoptosis, a consistent feature is the induction of a series of cytosolic proteases, which specifically cleave their substrates and lead to apoptotic morphology (Keppler-Hafkemeyer et al., 1998; Kumar, 1999; Porter and Janicke, 1999; Stennicke and Salvesen, 1997). These proteases are designated as caspases because they are cysteine proteases and cleave their substrates after aspartic acid residues (cysteine aspartase). Caspases remain dormant in the cell, but when activated by an apoptotic stimulus, they act by

proteolytic processing, thus modulating the function of their substrates. The substrate can be the activated caspase itself (autocatalysis) or another caspase. This way, caspases function not only as proteases, but also as specific signaling molecules in apoptosis. Different types of caspases initiate different apoptotic pathways. However, caspase-3, also called apopain, or CPP32, has been shown to be a key mediator of the final biochemical and morphological steps in apoptosis (Kumar, 1999). In apoptosis, the roles of different caspases have been shown to be pathway-specific as well as sequential in the steps involved in the pathway (Ray et al., 2002). Therefore, it seems possible to understand the mechanisms of apoptosis (e.g., due to SM) by studying the caspases involved. At present, this approach is facilitated by the availability of 1) some fluorogenic peptide substrates, which are specific for different caspases, and 2) some peptide caspase inhibitors, specific for different caspases. It should be noted that besides being useful to study apoptotic mechanisms, these inhibitors might also serve as prospective antiapoptotic medical countermeasures. There are reports in the literature showing that systemic injections of a broad-spectrum caspase inhibitor Z-VAD-FMK into mice protect the animals from caspase-3 activation and apoptosis in the liver and the lung due to anti-Fas antibody or lipopolysaccharide, respectively, and subsequent lethality (Kawasaki et al., 2000; Rodriguez et al., 1996). In this report, we demonstrate the usefulness of a convenient fluorometric caspase-3 assay to study SM-induced apoptosis in monolayer human epidermal keratinocytes (HEK) grown in multiwell plates.

MATERIALS AND METHODS

Chemicals and Cells

SM (>98% pure) was obtained from the US Army Research, Development and Engineering Command, Aberdeen Proving Ground (Maryland, USA). Frozen stock normal human epidermal keratinocyte (HEK) culture and keratinocyte growth medium (KGM) were from CAMBREX (Walkersville, MD, USA). The caspase-3 assay kits containing fluorogenic caspase-3 substrates were obtained from two separate sources. One substrate required cell lysis and was purchased from Sigma-Aldrich (Saint Louis, MO, USA; Product code: CASP-3-F). The other cell-permeable substrate did not require cell lysis and was procured from Beckman-Coulter (Fullerton, CA, USA; Product code: Beckman-Coulter CellProbe HT Caspase-3/7 Whole Cell Assay Kit). The caspase-3-specific inhibitor AC-DEVD-CHO was included in the caspase-3 assay kits from both sources. The general caspase inhibitor

Z-VAD-FMK was purchased from BD Biosciences (San Diego, CA, USA). All other chemicals were of the purest grade available.

Cell Culture

Frozen stock HEK (passage 2) were cultured in 150 cm² tissue culture flasks in KGM according to company instructions to initiate the culture. When these monolayer cells became 70–80% confluent, they were subcultured to 80–90% confluency in multiwell plates (96- or 24-well) and used in the experiments. To maintain consistency, HEK from a single donor and subcultured only to passage 3 were used.

Exposure of HEK to SM

Frozen stock of SM in KGM (Broomfield and Gross, 1989) was thawed and vortexed to obtain a 4 mM solution. Exposure of HEK to SM was done as described previously (Ray et al., 1995). The stock 4 mM SM solution was diluted in multiwell plates containing monolayer HEK in KGM for specified concentrations and kept for times as indicated.

Caspase-3 Assay

The caspase-3 fluorometric assays were done according to the instructions provided with the assay kits, but with necessary modifications required for developing the method described here. This enzyme assay was based on the hydrolysis of the caspase-3 peptide substrate (acetyl-Asp-Glu-Val-Asp or AC-DEVD) conjugated to a fluorochrome at the C-terminal Asp, resulting in the release of the fluorescent moiety. The fluorescence (absolute units) was measured using the CytoFluor Multi-Well Plate Reader Series 4000 spectrofluorometer from PerSeptive Biosystems (Framingham, MA, USA).

Caspases are present in the cytosol. In the assay kit from Sigma-Aldrich, the substrate AC-DEVD-AMC (7-amido-4-methylcoumarin) is not cell permeable and, therefore, the use of this substrate requires cell lysis. Cell lysis was accomplished by sonication of monolayer cultures in 1 × lysis buffer added to the wells of the culture plate as described by Gross et al. (2004). We used the Misonix Sonicator 3000 purchased from Misononix (Farmingdale, NY, USA). The sonicator was equipped with a microplate horn housed inside an acoustic enclosure. During sonication, the microplate horn was chilled by circulating water at 0–2°C using a temperature-controlled water circulator from Thermo Neslab (Portsmouth, NH, USA). To avoid any contamination of cultures by cooling water, the microplate containing monolayer cultures in lysis buffer was tightly sealed using an adhesive plastic sheet. Sonication of cells was done for 30-s intervals at the highest power setting on the sonicator dial followed by

a 60-s cooling period; the sonication/cooling cycles were repeated four times. At the end, cell disruption was confirmed by light microscopic observation.

The substrate in the Beckman-Coulter proprietary caspase-3/7 assay kit is cell permeable and, therefore, no cell lysis was required. This assay is based on the cleavage of the bisamide substrate Z-DEVD-R110 [bis-(N-CBZ-L-aspartyl-L-glutamyl-L-Valyl-L-aspartic acid amide derivative of rhodamine 110). Caspase-3 or caspase-7 cleaves the bond between aspartic acid and rhodamine molecules, releasing the fluorescent rhodamine molecule. The extent of fluorescence is, therefore, a measure of caspase-3/7 activity.

It should be noted that the use of the Sigma-Aldrich substrate would require a cell-processing (lysis) time at the end of each time interval following cell exposure to an apoptosis-inducing agent such as SM. On the contrary, the Beckman-Coulter substrate could be added along with the apoptosis inducer, and the time-course of caspase-3/7 activation could be monitored beginning immediately after SM addition and continuing without interruption.

All assays were done using monolayer cells grown in 96-well plates. Exposure of cells to different dilute SM concentrations was done according to approved procedures, which included exposing cells to SM at room temperature inside a total exhaust chemical/biological hood and then keeping the exposed cells inside this hood for at least 1 h for venting of volatile gases.

In assays using the Beckman-Coulter substrate, cells were exposed to SM in a total volume of 100 μ L, which also contained 20 μ L substrate solution added just prior to SM. Cells were incubated for specified times, and fluorescence was read at excitation and emission wavelengths of 485 nm and 530 nm, respectively.

In assays using the Sigma-Aldrich substrate, cells were exposed to desired SM concentrations, with appropriate controls, in fresh KGM in a total volume of 80 μ L. SM-exposed cells were then incubated inside a cell culture incubator overnight for 18–24 h. Following this incubation, 20 μ L of $5 \times$ lysis buffer was added to each well, and the cells were disrupted by sonication as described above. The substrate was then added to a total volume of 150 μ L, and fluorescence was read at increasing times at excitation and emission wavelengths of 360 nm and 460 nm, respectively.

Inhibitor Study

The effects of caspase inhibitors on SM-induced caspase-3 activation were studied by using the Beckman-Coulter substrate to avoid any possible effect of cell sonication on caspase-3 activity. Cells were first incubated with the inhibitors in KGM at 37°C inside a cell culture incubator for 30 min prior to SM addition. These SM-exposed cells were further incubated inside a cell culture incubator overnight, the caspase-3 substrate was added, and fluorescence was monitored at specified times after substrate addition.

Data Analysis

The results of assays performed in replicates of eight were averaged and expressed as the mean $\pm$ standard error of the mean (SEM). For each assay type, the differences between treatment groups at each time point were determined using a one-way analysis of variance (ANOVA) followed by post hoc comparison of treatment group means using the Tukey's test. In addition, for each assay type, the differences between observation time points within each treatment group were compared using a mixed-model ANOVA with time as a fixed factor and wells as a random factor followed by post hoc comparison of observation time means using the Tukey's test. Statistical significance was defined as $p < 0.05$.

RESULTS AND DISCUSSION

Experiments were designed to study the caspase-3 activation (apoptosis) due to SM in monolayer HEK cultures grown in multiwell plates, using the fluorogenic caspase-3 substrates AC-DEVD-AMC or Z-DEVD-R110 described above. The




Figure 1: Time-course of caspase-3 activation in HEK following exposure to increasing SM concentrations. Monolayer HEK cultures in a 96-well plate were exposed to different SM concentrations with the cell-permeable caspase-3 substrate Z-DEVD-R110 added just prior to SM. SM exposures were done inside a total exhaust chemical/biological hood at room temperature according to approved procedures. After 1 h inside the hood, the cells were transferred to a 37°C cell culture incubator. Fluorescence generated due to substrate hydrolysis was measured at specified times shown in the figure using a plate reader/spectrofluorometer as described under "Materials and Methods." Percent fluorescence values in experimental samples compared with SM-unexposed controls as 100% are mean $\pm$ SEM of eight replicate determinations at each SM concentration. The absence of SEM bars means that the values are too small to be visible. Statistical significance was defined as $p < 0.05$ versus SM-unexposed control group as described under "Materials and Methods."




Figure 2: Effects of peptide caspase-3 inhibitors on 0.3 mM SM-induced caspase-3 activation. HEK cultures in KGM were first incubated inside a cell culture incubator without or with the caspase-3-specific inhibitor AC-DEVD-CHO (0.1 mM) or the general caspase inhibitor Z-VAD-FMK (0.01 mM) for 30 min prior to 0.3 mM SM exposure inside a total exhaust chemical/biological hood at room temperature according to approved procedures. SM-exposed cells were then incubated overnight, and caspase-3 assays were conducted by adding the cell-permeable caspase-3 substrate Z-DEVD-R110 to cultures as shown. Fluorescence was measured at specified times after substrate addition. The results shown as absolute fluorescence units are mean $\pm$ SEM of eight replicate determinations. The absence of SEM bars means that the values are too small to be visible. Statistical significance was defined as $p < 0.05$ versus corresponding control values obtained in experiments without substrate.

time-course of enzyme activation, the effects of caspase inhibitors, and the SM concentration dependence are shown in Figures 1, 2, and 3, respectively.

The data shown in Figure 1 were generated in an experiment in which replicate HEK cultures were exposed to increasing concentrations of SM (0.05 to 0.3 mM) in KGM and the caspase-3 substrate Z-DEVD-R110 (Beckman-Coulter) had been added to the cells just prior to SM. Fluorescence increase following SM exposure was monitored at specified times throughout the day until 12 h and at 24 h next morning. The results are shown as percent fluorescence relative to SM-unexposed controls. In SM-exposed HEK, fluorescence generated from the fluorogenic caspase-3 substrate hydrolysis increased in a time (0–24 h) and concentration (0.05, 0.1, 0.15, 0.2, 0.3, mM) dependent manner. Following 0.3 mM SM exposure, which is considered to be the *in vitro* equivalent of a vesicating dose *in vivo*, a very small but statistically significant increase in fluorescence was observed at 6 to 8 h, which was markedly higher at 12 h. At 24 h, all SM concentrations increased fluorescence. These results are in agreement with those reported by Rosenthal et al. (1998) on the concentration and time dependence of SM-induced caspase-3 processing (conversion of the inactive 32 kDa procaspase-3




Figure 3: SM concentration dependence of caspase-3 activation in HEK. Monolayer HEK cultures in a 96-well plate were exposed to different SM concentrations inside a total exhaust chemical/biological hood at room temperature according to approved procedures as described in "Materials and Methods." SM-exposed cells were incubated inside a 37°C cell culture incubator overnight for 18 h. Following this incubation overnight, the cells were lysed in lysis buffer by sonication, and the caspase-3 substrate AC-DEVD-AMC was added to monitor the kinetics of substrate hydrolysis as measured by increase in fluorescence with time. The cells were placed inside the 37°C cell culture incubator in between readings. Percent fluorescence increase with time compared with SM-unexposed controls as 100% are shown as mean $\pm$ SEM of eight replicate determinations at each SM concentration. The absence of SEM bars means that the values are too small to be visible. Statistical significance was defined as $p < 0.05$ versus SM-unexposed control group as described under "Materials and Methods."

to the active 17 kDa caspase-3) and activation as well as other apoptosis indicators in HEK studied by the Western blotting method. The time-course and concentration dependence of SM-induced caspase-3 activation in HEK observed by our fluorometric method, therefore, appear to be valid indicators of apoptosis. These results also show that in 0.3 mM SM-exposed HEK, an overnight incubation is needed using this method to see a high level of caspase-3 activity.

The effects of some peptide caspase inhibitors on 0.3 mM SM-induced hydrolysis of the caspase-3 substrate AC-DEVD-R110 (Beckman-Coulter) in HEK are shown in Figure 2. In these experiments, as described in "Materials and Methods," the cells in KGM were first incubated inside a cell culture incubator without or with the caspase-3 specific inhibitor AC-DEVD-CHO or the general caspase inhibitor Z-VAD-FMK for 30 min prior to 0.3 mM SM exposure. The cells were then incubated overnight, and caspase-3 assays were conducted. The results (Figure 2) are shown as absolute fluorescence units $\pm$ SEM. SM-unexposed control cells incubated in KGM overnight had some caspase-3 activity, indicating spontaneous apoptosis in HEK cultures

without stimulation. Fluorescence increase due to 0.3 mM SM was prevented 100% by the caspase-3-specific peptide inhibitor (AC-DEVD-CHO), but less effectively by the general caspase inhibitor (Z-VAD-FMK), indicating that the fluorescence increase was due to caspase-3-mediated apoptosis.

We also examined the SM concentration dependence of caspase-3 activation at concentrations higher than 0.3 mM using 0.5, 1, and 2 mM. In these experiments, cells were exposed to different SM concentrations in KGM according to approved procedures and then incubated overnight inside a cell culture incubator for 18 h. The cells were lysed in lysis buffer by sonication as described under "Materials and Methods," and then the caspase-3 substrate AC-DEVD-AMC (Sigma-Aldrich) was added to cultures to monitor the kinetics of substrate hydrolysis between 0 to 8 h. The results (Figure 3) presented as percent fluorescence relative to SM-unexposed controls showed that when caspase-3 was almost fully activated during overnight incubation of HEK cultures following exposure to specified SM concentrations, there was a time-dependent increase in substrate hydrolysis that reached a plateau at around 4 h. Fluorogenic substrate hydrolysis due to caspase-3 activity increased progressively, being maximum at 0.5 mM. The extent and the kinetics of substrate hydrolysis at 1 mM SM were almost identical to those at 0.5 mM SM, and in some experiments, 1 mM SM produced values slightly lower than those at 0.5 mM SM as shown in Figure 3. However, at 2 mM SM, fluorescence decreased compared with 0.5 mM, which remains to be explained. One possible reason could be that at a SM concentration as high as 2 mM, the cells were metabolically compromised.

CONCLUSION

The results presented in this report support the following conclusions. The fluorometric method described here to study chemically induced apoptosis using fluorogenic caspase-3 substrates and monolayer cell cultures in multiwell plates is convenient, adaptable for high-throughput applications, if needed, and suitable for evaluations of test compounds as inhibitors of apoptosis. This assay clearly demonstrates apoptosis, as indicated by caspase-3 activation, in cultured HEK due to SM in a SM concentration- and time-dependent manner. Because apoptosis is believed to be an important process in many cases of chemical toxicity (e.g., due to SM) and a number of diseases including cancer, this method provides a valuable means to conduct research on the role of apoptosis in respective situations and drug development. These results also suggest potential applications of this method to study the roles in apoptosis of caspases other than caspase-3, provided 1) suitable fluorogenic substrates are available and 2) the subject caspases are stimulated enough to generate measurable fluorescent signals.

REFERENCES

- Broomfield, C. A., Gross, C. L. (1989). Stability of sulfur mustard in a two-phase storage configuration. In: Proceedings 1989 Medical Defense Bioscience Review, AD B139550. Aberdeen Proving Ground, MD: United States Army Medical Research Institute of Chemical Defense, pp. 433–436.
- Gross, C. L., Clark, O. E., Nealley, E. W., Nipwoda, M. T., Smith, W. J. (2004). Lysis of adherent human epidermal keratinocytes in situ by a misonix tissue culture plate sonicator. *Toxicologist* 78(1-S):265.
- Hollinger, M. A. (2002). Series foreword. In: Davis, M. A., ed. *Apoptosis Methods in Pharmacology and Toxicology—Approaches to Measurement and Quantification*. Totowa, NJ: Humana Press, p. 5.
- Kan, R. K., Pleva, C. A., Hamilton, T. A., Anderson, D. A., Petralli, J. P. (2003). Sulfur mustard-induced apoptosis in hairless guinea pig skin. *Toxicol. Pathol.* 31(2):185–190.
- Kawasaki, M., Kuwana, K., Hagimoto, N., Matsuba, T., Kunitake, R., Tanaka, T., Maeyama, T., Hara, N. (2000). Protection from lethal apoptosis in lipopolysaccharide-induced acute lung injury in mice by a caspase inhibitor. *Am. J. Pathol.* 157(2):597–603.
- Keppeler-Hafkemeyer, A., Brinkman, U., Pastan, I. (1998). Role of caspases in immunotoxin-induced apoptosis of cancer cells. *Biochemistry* 37(48):16934–16942.
- Kerr, J. F. R., Wyllie, A. H., Currie, A. R. (1972). Apoptosis: a basic biological phenomenon with wide-ranging implications in tissue kinetics. *Br. J. Cancer* 26:239–257.
- Kumar, S. (1999). Regulation of caspase activation in apoptosis: implications in pathogenesis and treatment of disease. *Clin. Exp. Pharmacol. Physiol.* 26(4):295–303.
- Porter, A. G., Janicke, R. U. (1999). Emerging role of caspase-3 in apoptosis. *Cell Death Differ.* 6(2):99–104.
- Ray, R., Benton, B. J., Anderson, D. R., Brobst, D. E., Rosenthal, D. S., Ray, P. (1999). Poly(ADP-ribose) polymerase (PARP) mediated mechanisms of sulfur mustard (SM) toxicity and its protection. *FASEB J.* 13(7):A1341.
- Ray, R., Bhat, K. R., Benton, B. J., Burke, M. E., Dierking, E., Roman, J. M., Anderson, D. R., Petralli, J. P., Smith, W. J., Rosenthal, D. S., Ray, P. (December 2–6, 2002). In: *Mechanism-Based Intervention of Vesicant Toxicity*, Proceedings of the 3rd Singapore International Symposium on Protection Against Toxic Substances (SISPAT), DSO National Laboratories, Singapore.
- Ray, R., Legere, R. H., Majerus, B. J., Petralli, J. P. (1995). Sulfur mustard-induced increase in intracellular free calcium level and arachidonic acid release from cell membrane. *Toxicol. Appl. Pharmacol.* 131:44–52.
- Rodriguez, I., Matsuura, K., Ody, C., Nagata, S., Vassali, P. (1996). Systemic injection of a tripeptide inhibits the intracellular activation of CPP32-like proteases in vivo and fully protects mice against Fas-mediated fulminant liver destruction and death. *J. Exp. Med.* 184:2067–2072.
- Rosenthal, D. S., Rosenthal, C. M. G., Iyer, S., Smith, W., Ray, R., Smulson, M. E. (1998). Sulfur mustard induces terminal differentiation and apoptosis in keratinocytes via a Ca²⁺-calmodulin and apopain-dependent pathway. *J. Invest. Dermatol.* 111(1):64–71.
- Rosenthal, D. S., Valena, A., Choi, F.-P., Schlege, I. R., Ray, R., Benton, B., Anderson,

- D., Smith, W. J., Simbulan-Rosenthal, C. M. (2003). Expression of dominant-negative Fas-associated death domain blocks human keratinocyte apoptosis and vesication induced by sulfur mustard. *J. Biol. Chem.* 278(10):8531–8540.
- Stennicke, H. R., Salvesen, G. S. (1997). Biochemical characteristics of caspases-3, -6, -7, and -8. *J. Biol. Chem.* 272(41):25719–25723.
- Trump, B. F., Berezesky, I. K. (1998). The reactions of cells to lethal injury: oncosis and necrosis—the role of calcium. In: Lockshin, R. A., Zakeri, Z., Tilly, J. L., eds. *When Cells Die*. New York: Wiley-Liss, pp. 57–96.