

Hyperbaric Oxygen Therapy – An Adjunct to Optimal Combat Trauma Management

Brett B. Hart, CDR, MC, USN
Department of Hyperbaric Medicine
Naval Medical Center
Portsmouth, VA 23708

The views expressed in this article are those of the author and do not reflect the official policy or position of the Department of the Navy, Department of Defense, or the United States Government.

INTRODUCTION

Although wound statistics from Operation Iraqi Freedom are still being analyzed, a preliminary analysis of casualty data demonstrates that approximately 70% of the injured combatants sustained trauma to their extremities.(Malcolm 2004) During the acute management of these injuries at Echelon II and III medical facilities, approximately 75 - 100 major vascular reconstructions and 180 limb amputations were performed. Anecdotal reports reveal that a number of the wounded combatants required tissue grafting, flap reconstruction, revision amputation or treatment for resistant infection subsequent to their evacuation to tertiary care facilities. Indeed, of the 560 surgical procedures performed for combat related injuries aboard the USNS Comfort, approximately 31% were noted to be sufficiently complicated by persistent tissue necrosis, wound infection, graft failure or delayed wound healing that further surgical management was required.(Helmert 2004) Residual ischemia has been shown to play a role in each of these processes. Hyperbaric Oxygen (HBO₂) has been shown to be an effective in reversing ischemia and limiting wound healing complications in hospital-based, clinical care settings. It is proposed that HBO₂ therapy could be applied in Echelon II and III operational environments to similarly limit the extent of surgical debridements, improve tissue flap and graft survival, decrease wound infection rates, speed complex wound healing and, ultimately, reduce the morbidity experienced by wounded combatants. This paper discusses the physiologic mechanisms underlying HBO₂ therapy's clinically beneficial effects, examines potential roles for HBO₂ in the optimal management of combat-related trauma, and provides practical suggestions for HBO₂ treatment chamber deployment into operational environments.

MECHANISMS OF HBO₂ THERAPY

Reversal of Acute Ischemia

All wounds are characterized by a state of relative hypoxia.(Hunt, Twomey et al. 1967) Local oxygen tensions in the vicinity of a wound are approximately half the values observed in normal, non-wounded tissue.(Sheffield 1998) Musculoskeletal injuries secondary to crush, blast and penetrating trauma each produce local tissue ischemia, hypoxic gradients from zones of necrotic to healthy tissue and, when inadequately treated, the potential for propagation of ischemic injury into adjacent healthy tissues.(Warriner and Hopf 2003) Clinically, HBO₂ therapy can be used to correct tissue hypoxia. This is accomplished when a patient breathes 100% oxygen at elevated atmospheric pressure. Physiologically, a directly proportional

Paper presented at the RTO HFM Symposium on "Combat Casualty Care in Ground Based Tactical Situations: Trauma Technology and Emergency Medical Procedures", held in St. Pete Beach, USA, 16-18 August 2004, and published in RTO-MP-HFM-109.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 01 SEP 2004		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Hyperbaric Oxygen Therapy An Adjunct to Optimal Combat Trauma Management				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Department of Hyperbaric Medicine Naval Medical Center Portsmouth, VA 23708				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001795, Combat Casualty Care in Ground-Based Tactical Situations: Trauma Technology and Emergency Medical Procedures (Soins aux blessés au combat dans des situations tactiques : echnologies des traumatismes et procédures médicales d'urgence), The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 12	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

increase in plasma oxygen tensions is produced and, at 2 - 2.5 atmospheres absolute pressure, arterial PO₂ elevations in excess of 1500 mmHg are achieved.(Warriner and Hopf 2003) As a result, tissue oxygen tensions are also elevated and the diffusion of oxygen into areas of relative hypoxia markedly enhanced.(Strauss 2003; Niinikoski 2004) This significant level of hyperoxygenation allows for the reversal of localized tissue hypoxia and corrects the pathophysiology related to oxygen deficiency. This improved tissue oxygenation translates into the ability to limit the duration and progression of ischemic tissue necrosis(Zamboni, Roth et al. 1989; Bouachour, Cronier et al. 1996; Ramon, Abramovich et al. 1998), improve demarcation between necrotic and viable tissue(Isakov Iu, Atroshchenko et al. 1979; Rosenthal, Benderly et al. 1985; von Schroeder and Botte 1998; Murphy, Banwell et al. 2000), temporize against exceptional blood loss anemia(Hart 1974; Sherman, Sennik et al. 1989; Bitterman, Reissman et al. 1991; Greensmith 2000; MacFarlane, Cronje et al. 2000; Stark, Coatesworth et al. 2003) and enhance the survival of reconstructive tissue grafts and flaps.(Tai, Birely et al. 1992; Gampper, Zhang et al. 2002; Kalani, Jorreskog et al. 2002; Ulkur, Yuksel et al. 2002; Richards, Lineaweaver et al. 2003)

Reduction of Reactive Edema Formation

Post-traumatic vasogenic edema develops as a direct consequence of acute soft tissue injury and, as the injured, hypoxic tissues lose their ability to regulate intracellular water, it becomes accentuated by the formation of cytogenic edema.(Strauss 2003) The resultant extracellular fluid accumulation increases diffusion distances between oxygen-carrying capillaries and surrounding cells, reducing total oxygen delivery and perpetuating cytogenic edema formation. In closed tissue compartments, increases in interstitial pressure cause collapse of the microcirculation, further exacerbating existing ischemia and threatening previously uninvolved, healthy tissues. Clinically, this pathologic process is recognized as compartment syndrome. HBO₂ therapy has been shown to be an effective adjunct in the management of compartment syndrome. Three distinct processes contribute to this clinical effectiveness. First, by reversing acute tissue hypoxia, HBO₂ breaks the cycle between cellular ischemia and progressive edema formation, limits the total volume of soft tissue necrosis.(Nylander 1986; Skyhar, Hargens et al. 1986; Nylander, Nordstrom et al. 1988; Zamboni, Roth et al. 1993) Second, HBO₂-induced vasoconstriction produces a 20% decrease in arterial blood flow.(Bird and Telfer 1965; Bird and Telfer 1966) Edema reduction occurs because filtration of capillary fluid is decreased, while vascular outflow and improved oxygen delivery are maintained. Finally, the affected soft tissues are protected from reperfusion injury, a topic covered more fully in the next section.(Haapaniemi, Sirsjo et al. 1995) From the clinical standpoint, HBO₂ therapy can be used to prophylax against suspected or impending compartment syndrome.(Myers 2000; Strauss 2003) Where a compartment syndrome is already established, HBO₂ can be used to decrease the extent of required surgical fasciotomy, accelerate resolution of residual edema, and speed primary wound closure.(Bouachour, Cronier et al. 1996; Fitzpatrick, Murphy et al. 1998; Lindstrom, Gullichsen et al. 1998; Assenza, Borromeo et al. 2001; Van Poucke, Leenders et al. 2001; Gold, Barish et al. 2003)

Optimizing Host Antibacterial Defenses

Beyond the occurrence of frank necrosis, pathologic levels of hypoxia are correlated with increased rates of wound infection and, consequently, delayed wound healing.(Niinikoski 1969; Silver 1977) HBO₂ therapy has been shown to be effective in limiting the incidence of wound infection, speeding resolution of refractory infections, and decreasing the morbidity and mortality associated with malignant infections. Several discrete mechanisms account for the beneficial effects of HBO₂ in controlling infection. First, neutrophils required tissue oxygen tensions of 30-40 mmHg to destroy bacteria by oxidative killing mechanisms.(Mandell 1974; Hohn 1977) Leukocyte-mediated killing of aerobic gram-positive and anaerobic gram-negative organisms is restored when the pathologically low tissue oxygen tensions characteristic of wounded and infected tissues are

increased to physiologic or supraphysiologic levels by HBO₂ treatment.(Mader 1987; Knighton, Fiegel et al. 1990) Second, aminoglycoside and cephalosporin antibiotic transport across the bacterial cell wall does not occur if tissue oxygen tensions are below 20-30 mmHg. Therefore, HBO₂ therapy may enhance transport and augment antibiotic efficacy.(Park, Muhvich et al. 1991; Hirn 1993; Mendel, Reichert et al. 1999) Third, hyperoxygenation of the tissues surrounding areas of malignant infection may be of significance in preventing the extension of invading microorganisms.(Korhonen, Hirn et al. 1998; Korhonen 2000) Clinically, the most dramatic benefits are seen in reducing the morbidity and mortality of necrotizing fasciitis and gas gangrene, where treatment with HBO₂ results in clinical improvement even when standard measures have failed.(Schreiner, Tonjum et al. 1974; Bakker 1985; Riseman, Zamboni et al. 1990) Such benefits have been demonstrated for combat casualties.(Shupak, Halpern et al. 1984; Paillet and Labeau 1986) Clinical improvements have also been noted in the management of peritonitis.(Bogomolova and Bol'shakov 1996) HBO₂ therapy is useful as a rescue treatment of refractory bone infections.(Aitasalo, Niinikoski et al. 1998; Maynor, Moon et al. 1998; Chen, Shih et al. 2003) In addition, HBO₂ is effective in reducing the need for re-operations in neurosurgical procedures complicated by infected grafts.(Eltorai, Hart et al. 1984; Larsson, Engstrom et al. 2002)

Prophylaxis Against Reperfusion Injury

The body of scientific literature continues to elucidate the multiple biochemical mechanisms behind HBO₂ therapy's ability to limit and prophylax against ischemia-reperfusion (I/R) injury. These include upregulation of TGF-beta1 which ameliorates reperfusion injury by up-regulating bcl-2 and inhibiting TNF-alpha production (Yang, Bosco et al. 2001; Grunenfelder, Miniati et al. 2002), catalase induction (Kim, Choi et al. 2001), inhibition of intracellular adhesion molecule (ICAM-1) formation (Buras, Stahl et al. 2000) and stimulation of the fibrinolytic enzymes tissue plasminogen activator (t-PA), urokinase plasminogen activator (uPA) and plasminogen activator inhibitor type one (PAI-1).(Tjarnstrom, Holmdahl et al. 2001) While it is not possible to say at this point which of these mechanisms is most predominant in the beneficial effects of HBO₂ in limiting I/R injury, the preponderance of models consistently demonstrate this effect.(Buras 2000) Such inducible I/R protection has been demonstrated even when HBO₂ treatment was delayed two to eight hours.(Murakami, Horinouchi et al. 2001; Tjarnstrom, Holmdahl et al. 2001; Agir, Mersa et al. 2003) Pretreatment with HBO₂ has also been shown to be protective against subsequent periods of ischemia in neural and musculoskeletal tissues.(Murakami, Horinouchi et al. 2001; Dong, Xiong et al. 2002) Clinically, prophylaxis against I/R injury could be used to help improve post-surgical outcomes in patients undergoing peripheral vascular repairs, flap and grafting, and primary closure of large tissue defects.(Chen, Chen et al. 1998; Mazariegos, O'Toole et al. 1999; Myers 2000)

Improving Tissue Repair Rates

As noted previously, the hypoxic nature of all wounds has been demonstrated and, when pathologically increased, is correlated with impaired wound healing.(Hunt, Twomey et al. 1967; Niinikoski 1969; Gottrup 2004) Fibroblast replication, collagen deposition and angiogenesis are all oxygen sensitive responses that are necessary to proper wound healing.(Hunt and Pai 1972; Knighton, Silver et al. 1981; LaVan and Hunt 1990; Ishii, Miyanaga et al. 1999) Induction of several oxygen dependent growth factors has been elucidated in this process. Specifically, the production of nitric oxide (NO), vascular endothelial growth factor (VEGF), platelet derived growth factor (PDGF), basic fibroblast growth factor (bFGF), angiopoietin-2 (ANG-2) and their respective tissue receptors are all upregulated in the presence of HBO₂.(Zhao, Davidson et al. 1994; Bonomo, Davidson et al. 1998; Boykin 2000; Sheikh, Gibson et al. 2000; Lin, Shyu et al. 2002; Kang, Gorti et al. 2004) These effects have been shown to persist after patient removal from the hyperbaric environment.(Siddiqui, Davidson et al. 1997) Additionally, the degree of wound healing induced by HBO₂ therapy appears to be

either superior to or synergistic with topically applied growth factors.(Bonomo, Davidson et al. 2000; Boykin 2000; Chen, Lai et al. 2002) In animal models, acceleration of wound healing has been noted for both wounds in both regionally ischemic and normal tissue beds.(Uhl, Sirsjo et al. 1994) Improvements in healing have also been demonstrated for orthopedic and neurosurgical procedures.(Atesalp, Komurcu et al. 2002; Chen, Lai et al. 2002) Clinically, HBO₂ therapy has been shown to improve rates of wound healing and reduce amputation rates in patients sustaining injuries in both civilian and combat environments.(Bialik, Fishman et al. 1987; LaVan and Hunt 1990; Radonic, Baric et al. 1995; Porcellini, Bernardo et al. 1997; Atesalp, Komurcu et al. 2002; Wang, Li et al. 2002; Warriner and Hopf 2003; Zamboni, Browder et al. 2003)

Hyperbaric Effects Summary

For hypoxic, complex wounds such as those induced by combat trauma, the net effect of serial hyperbaric exposures is improved local host immune response, increased clearance of infection, enhanced tissue growth, angiogenesis and wound epithelialization. When applied as an adjunct to surgical interventions, antibiotics and other clinically indicated therapies, HBO therapy is effective adjunct in the management of severe and refractory musculoskeletal problems. Indeed, the timely application of hyperbaric oxygen can be a limb- and life-saving therapy. (Wang, Li et al. 2002) More practically, adding HBO₂ treatment to other standard of care therapies can reduce the number and extent of required surgical procedures, decrease the duration of hospitalization and recovery time, improve post recovery function and, ultimately, enhance the wounded patient's prospects for full recovery.

Combat-Related Treatment Indications:

	Reduce Acute Tissue Ischemia	Minimize Edema Formation	Reperfusion Injury Prophylaxis	Improve Host Antibacterial Defenses	Improve Tissue Healing Rates
Musculoskeletal Trauma	Indicated Practical	Indicated Practical	Indicated Practical	Indicated Practical	Indicated Practical
Compartment Syndrome	Indicated Practical	Indicated Practical	Indicated Practical	Indicated Practical	Indicated Practical
Compromised Grafts / Flaps	Indicated Practical	Indicated Practical	Indicated Practical	Indicated Practical	Indicated Practical
Major Vascular Injury	Indicated Practical	Indicated Practical	Indicated Practical	N/A	N/A
Necrotizing Infections	Indicated Practical	Indicated Practical	Unknown	Indicated Practical	Indicated Practical
Antibiotic Resistant Infection	N/A	N/A	N/A	Indicated Practical	Indicated Practical
Acute Blood Loss Anemia	Indicated Not Practical	N/A	Unknown	N/A	N/A

Operational Deployment of HBO₂ Chambers:

The use of HBO₂ therapy for the management of combat related trauma has been suggested by a number of authors.(Cramer 1985; Spichev and Gostev Iu 1987; Rudge 1993; Broome 1997; Radonic, Baric et al. 1997; Fitzpatrick, Murphy et al. 1998; MacFarlane, Cronje et al. 2000) The primary difficulty with employment of this therapeutic modality is providing ready access of combat casualties to hyperbaric chamber facilities. As noted from the mechanistic discussions above, the acute benefits of HBO₂ therapy are best achieved when the therapy is provided within a few hours of injury. However, for the diseases and injuries presented, treatment with HBO₂ is generally considered to be an adjunct to appropriate antibiotic therapy and standard surgical management. Thus, whenever possible, initial treatment with HBO₂ should be timed to occur immediately after definitive surgical interventions have taken place, between planned surgical procedures, or when lack of OR availability allows HBO₂ therapy to be used as a temporizing measure. Unfortunately, logistical issues complicate deployment of hyperbaric chambers into the combat environment. Indeed, size, safety and mobility constraints render co-location of hyperbaric chambers at Echelon I treatment levels impractical. In contrast, at least two hyperbaric systems are available for use at Echelon II level treatment facilities. The most portable of these systems is the single person (i.e. a monoplace chamber) folding hyperbaric stretcher. This foldable, lightweight chamber, called the Emergency Evacuation Hyperbaric Stretcher (EEHS), has received certification for use within DoD. While the majority of these systems were initially intended for use in the remote treatment of decompression sickness and submarine escape and rescue operations, their potential application to combat trauma cases has been noted.(Locklear 2002) Deployment of U.S. Navy's Transportable Recompression Chamber System (TRCS), as was used by the Special Medical Response Team in August 2002 to treat miners trapped in a Pennsylvania coal mine. This later chamber has the advantage of being large enough to allow a medical attendant to accompany the patient during treatment (i.e. a multiplace chamber). At Echelon III level treatment facilities, multiple hyperbaric treatment options could be employed. These could include those chamber systems already mentioned as well as larger chamber systems such as the Fly Away Recompression Chamber (FARC). For maritime operations, a multiplace hyperbaric chamber could be stationed aboard amphibious ships or designated hospital ships, such as USNS Comfort and USNS Mercy. HBO₂ therapy used in support of reconstructive procedures and complex wound management can be arranged by medevac of select patients to regional Echelon IV treatment facilities that have fixed multiplace and monoplace chamber facilities. Such hospital-associated HBO₂ treatment capabilities can be found in most industrialized nations. Many of the chamber complexes are associated with existing tertiary care, military treatment facilities.

Agir, H., B. Mersa, et al. (2003). "[Histologic effects of hyperbaric oxygen therapy administered immediately after or two hours after ischemia-reperfusion injury: a rat abdominal skin flap model]." Kulak Burun Bogaz Ihtis Derg **10**(1): 18-24.

Aitasalo, K., J. Niinikoski, et al. (1998). "A modified protocol for early treatment of osteomyelitis and osteoradionecrosis of the mandible." Head Neck **20**(5): 411-7.

Assenza, M., C. Borromeo, et al. (2001). "[Compartment syndrome or gas gangrene? A case report]." G Chir **22**(10): 345-7.

Atesalp, A. S., M. Komurcu, et al. (2002). "The treatment of infected tibial nonunion with aggressive debridement and internal bone transport." Mil Med **167**(12): 978-81.

Bakker, D. J. (1985). "Pure and mixed aerobic and anaerobic soft tissue infections." HBO Rev **6**: 65-96.

- Bialik, V., J. Fishman, et al. (1987). "The management of severe compound war injuries of the extremities in paediatric patients." Z Kinderchir **42**(1): 57-9.
- Bird, A. D. and A. B. Telfer (1965). "Effect of Hyperbaric Oxygen on Limb Circulation." Lancet **13**: 355-6.
- Bird, A. D. and A. B. Telfer (1966). "The effect of oxygen at 1 and 2 atmospheres on resting forearm blood flow." Surg Gynecol Obstet **123**(2): 260-8.
- Bitterman, H., P. Reissman, et al. (1991). "Oxygen therapy in hemorrhagic shock." Circ Shock **33**(3): 183-91.
- Bogomolova, N. S. and L. V. Bol'shakov (1996). "[Anaerobic infection in abdominal surgery]." Vestn Ross Akad Med Nauk(2): 30-3.
- Bonomo, S. R., J. D. Davidson, et al. (2000). "Enhancement of wound healing by hyperbaric oxygen and transforming growth factor beta3 in a new chronic wound model in aged rabbits." Arch Surg **135**(10): 1148-53.
- Bonomo, S. R., J. D. Davidson, et al. (1998). "Hyperbaric oxygen as a signal transducer: upregulation of platelet derived growth factor-beta receptor in the presence of HBO2 and PDGF." Undersea Hyperb Med **25**(4): 211-6.
- Bouachour, G., P. Cronier, et al. (1996). "Hyperbaric oxygen therapy in the management of crush injuries: a randomized double-blind placebo-controlled clinical trial." J Trauma **41**(2): 333-9.
- Boykin, J. V., Jr. (2000). "The nitric oxide connection: hyperbaric oxygen therapy, becaplermin, and diabetic ulcer management." Adv Skin Wound Care **13**(4 Pt 1): 169-74.
- Broome, J. R. (1997). "The Sir James Watt Hyperbaric Medicine Unit--an asset to the Royal Hospital Haslar and the defence medical services." J R Nav Med Serv **83**(1): 14-5.
- Buras, J. (2000). "Basic mechanisms of hyperbaric oxygen in the treatment of ischemia-reperfusion injury." Int Anesthesiol Clin **38**(1): 91-109.
- Buras, J. A., G. L. Stahl, et al. (2000). "Hyperbaric oxygen downregulates ICAM-1 expression induced by hypoxia and hypoglycemia: the role of NOS." Am J Physiol Cell Physiol **278**(2): C292-302.
- Chen, C. E., S. T. Shih, et al. (2003). "Hyperbaric oxygen therapy in the treatment of chronic refractory osteomyelitis: a preliminary report." Chang Gung Med J **26**(2): 114-21.
- Chen, M. F., H. M. Chen, et al. (1998). "Hyperbaric oxygen pretreatment attenuates hepatic reperfusion injury." Liver **18**(2): 110-6.
- Chen, W. J., P. L. Lai, et al. (2002). "The effect of hyperbaric oxygen therapy on spinal fusion: using the model of posterolateral intertransverse fusion in rabbits." J Trauma **52**(2): 333-8.
- Cramer, F. S. (1985). "Care of the injured soldier: a medical readiness role for clinical hyperbaric oxygen therapy." Mil Med **150**(7): 372-5.
- Dong, H., L. Xiong, et al. (2002). "Preconditioning with hyperbaric oxygen and hyperoxia induces tolerance against spinal cord ischemia in rabbits." Anesthesiology **96**(4): 907-12.

- Eltorai, I., G. B. Hart, et al. (1984). "Osteomyelitis in the spinal cord injured: a review and a preliminary report on the use of hyperbaric oxygen therapy." Paraplegia **22**(1): 17-24.
- Fitzpatrick, D. T., P. T. Murphy, et al. (1998). "Adjunctive treatment of compartment syndrome with hyperbaric oxygen." Mil Med **163**(8): 577-9.
- Gampper, T. J., F. Zhang, et al. (2002). "Beneficial effect of hyperbaric oxygen on island flaps subjected to secondary venous ischemia." Microsurgery **22**(2): 49-52.
- Gold, B. S., R. A. Barish, et al. (2003). "Resolution of compartment syndrome after rattlesnake envenomation utilizing non-invasive measures." J Emerg Med **24**(3): 285-8.
- Gottrup, F. (2004). "Oxygen in wound healing and infection." World J Surg **28**(3): 312-5.
- Greensmith, J. E. (2000). "Hyperbaric oxygen reverses organ dysfunction in severe anemia." Anesthesiology **93**(4): 1149-52.
- Grunenfelder, J., D. N. Miniati, et al. (2002). "Up-regulation of Bcl-2 through hyperbaric pressure transfection of TGF-beta1 ameliorates ischemia-reperfusion injury in rat cardiac allografts." J Heart Lung Transplant **21**(2): 244-50.
- Haapaniemi, T., A. Sirsjo, et al. (1995). "Hyperbaric oxygen treatment attenuates glutathione depletion and improves metabolic restitution in postischemic skeletal muscle." Free Radic Res **23**(2): 91-101.
- Hart, G. B. (1974). "Exceptional blood loss anemia. Treatment with hyperbaric oxygen." Jama **228**(8): 1028-9.
- Helmerts, S. (2004). Personal Communication. June 04.
- Hirn, M. (1993). "Hyperbaric oxygen in the treatment of gas gangrene and perineal necrotizing fasciitis. A clinical and experimental study." Eur J Surg Suppl(570): 1-36.
- Hohn, D. C. (1977). Oxygen and leukocyte microbial killing. Hyperbaric Oxygen Therapy. J. C. Davis and T. K. Hunt. Bethesda, Undersea Medical Society: 101-110.
- Hunt, T. K. and M. P. Pai (1972). "The effect of varying ambient oxygen tensions on wound metabolism and collagen synthesis." Surg Gynecol Obstet **135**(4): 561-7.
- Hunt, T. K., P. Twomey, et al. (1967). "Respiratory gas tensions and pH in healing wounds." Am J Surg **114**(2): 302-7.
- Isakov Iu, V., Z. B. Atroshchenko, et al. (1979). "[Hyperbaric oxygenation in the prevention of wound infection in open injury to the locomotor apparatus]." Vestn Khir Im I I Grek **123**(10): 117-21.
- Ishii, Y., Y. Miyanaga, et al. (1999). "Effects of hyperbaric oxygen on procollagen messenger RNA levels and collagen synthesis in the healing of rat tendon laceration." Tissue Eng **5**(3): 279-86.
- Kalani, M., G. Jorneskog, et al. (2002). "Hyperbaric oxygen (HBO) therapy in treatment of diabetic foot ulcers. Long-term follow-up." J Diabetes Complications **16**(2): 153-8.

- Kang, T. S., G. K. Gorti, et al. (2004). "Effect of hyperbaric oxygen on the growth factor profile of fibroblasts." Arch Facial Plast Surg **6**(1): 31-5.
- Kim, C. H., H. Choi, et al. (2001). "Hyperbaric oxygenation pretreatment induces catalase and reduces infarct size in ischemic rat myocardium." Pflugers Arch **442**(4): 519-25.
- Knighton, D. R., V. D. Fiegel, et al. (1990). "Oxygen as an antibiotic. The effect of inspired oxygen on bacterial clearance." Arch Surg **125**(1): 97-100.
- Knighton, D. R., I. A. Silver, et al. (1981). "Regulation of wound-healing angiogenesis-effect of oxygen gradients and inspired oxygen concentration." Surgery **90**(2): 262-70.
- Korhonen, K. (2000). "Hyperbaric oxygen therapy in acute necrotizing infections. With a special reference to the effects on tissue gas tensions." Ann Chir Gynaecol **89 Suppl 214**: 7-36.
- Korhonen, K., M. Hirn, et al. (1998). "Hyperbaric oxygen in the treatment of Fournier's gangrene." Eur J Surg **164**(4): 251-5.
- Larsson, A., M. Engstrom, et al. (2002). "Hyperbaric oxygen treatment of postoperative neurosurgical infections." Neurosurgery **50**(2): 287-95; discussion 295-6.
- LaVan, F. B. and T. K. Hunt (1990). "Oxygen and wound healing." Clin Plast Surg **17**(3): 463-72.
- Lin, S., K. G. Shyu, et al. (2002). "Hyperbaric oxygen selectively induces angiopoietin-2 in human umbilical vein endothelial cells." Biochem Biophys Res Commun **296**(3): 710-5.
- Lindstrom, T., E. Gullichsen, et al. (1998). "Effects of hyperbaric oxygen therapy on perfusion parameters and transcutaneous oxygen measurements in patients with intramedullary nailed tibial shaft fractures." Undersea Hyperb Med **25**(2): 87-91.
- Locklear, K. R. (2002). U.S. Department of Defense Certifies the Hyperlite for Operational Use. Hyperbaric Medicine Today. **1**: 35.
- MacFarlane, C., F. J. Cronje, et al. (2000). "Hyperbaric oxygen in trauma and surgical emergencies." J R Army Med Corps **146**(3): 185-90.
- Mader, J. T. (1987). Potential of tobramycin by hyperbaric oxygen in experimental Pseudomonas aeruginosa osteomyelitis. 27th Interscience Conference on Antimicrobial Agents and Chemotherapy.
- Malcolm, R. D. (2004). Personal Communication.
- Mandell, G. L. (1974). "Bactericidal activity of aerobic and anaerobic polymorphonuclear neutrophils." Infect Immun **9**(2): 337-41.
- Maynor, M. L., R. E. Moon, et al. (1998). "Chronic osteomyelitis of the tibia: treatment with hyperbaric oxygen and autogenous microsurgical muscle transplantation." J South Orthop Assoc **7**(1): 43-57.
- Mazariegos, G. V., K. O'Toole, et al. (1999). "Hyperbaric oxygen therapy for hepatic artery thrombosis after liver transplantation in children." Liver Transpl Surg **5**(5): 429-36.

- Mendel, V., B. Reichert, et al. (1999). "Therapy with hyperbaric oxygen and cefazolin for experimental osteomyelitis due to *Staphylococcus aureus* in rats." Undersea Hyperb Med **26**(3): 169-74.
- Murakami, N., T. Horinouchi, et al. (2001). "Hyperbaric oxygen therapy given 30 minutes after spinal cord ischemia attenuates selective motor neuron death in rabbits." Crit Care Med **29**(4): 814-8.
- Murphy, J. V., P. E. Banwell, et al. (2000). "Frostbite: pathogenesis and treatment." J Trauma **48**(1): 171-8.
- Myers, R. A. (2000). "Hyperbaric oxygen therapy for trauma: crush injury, compartment syndrome, and other acute traumatic peripheral ischemias." Int Anesthesiol Clin **38**(1): 139-51.
- Niinikoski, J. (1969). "Effect of oxygen supply on wound healing and formation of experimental granulation tissue." Acta Physiol Scand Suppl **334**: 1-72.
- Niinikoski, J. H. (2004). "Clinical hyperbaric oxygen therapy, wound perfusion, and transcutaneous oximetry." World J Surg **28**(3): 307-11.
- Nylander, G. (1986). "Tissue ischemia and hyperbaric oxygen treatment: an experimental study." Acta Chir Scand Suppl **533**: 1-109.
- Nylander, G., H. Nordstrom, et al. (1988). "Effects of hyperbaric oxygen treatment in post-ischemic muscle. A quantitative morphological study." Scand J Plast Reconstr Surg Hand Surg **22**(1): 31-9.
- Pailler, J. L. and F. Labeau (1986). "[Gas gangrene: a military disease?]." Acta Chir Belg **86**(2): 63-71.
- Park, M. K., K. H. Muhvich, et al. (1991). "Hyperoxia prolongs the aminoglycoside-induced postantibiotic effect in *Pseudomonas aeruginosa*." Antimicrob Agents Chemother **35**(4): 691-5.
- Porcellini, M., B. Bernardo, et al. (1997). "Combined vascular injuries and limb fractures." Minerva Cardioangiol **45**(4): 131-8.
- Radonic, V., D. Baric, et al. (1997). "War injuries of the femoral artery and vein: a report on 67 cases." Cardiovasc Surg **5**(6): 641-7.
- Radonic, V., D. Baric, et al. (1995). "War injuries of the crural arteries." Br J Surg **82**(6): 777-83.
- Ramon, Y., A. Abramovich, et al. (1998). "Effect of hyperbaric oxygen on a rat transverse rectus abdominis myocutaneous flap model." Plast Reconstr Surg **102**(2): 416-22.
- Richards, L., W. C. Lineaweaver, et al. (2003). "Effect of hyperbaric oxygen therapy on the tubed pedicle flap survival in a rat model." Ann Plast Surg **50**(1): 51-6.
- Riseman, J. A., W. A. Zamboni, et al. (1990). "Hyperbaric oxygen therapy for necrotizing fasciitis reduces mortality and the need for debridements." Surgery **108**(5): 847-50.
- Rosenthal, E., A. Benderly, et al. (1985). "Hyperbaric oxygenation in peripheral ischaemic lesions in infants." Arch Dis Child **60**(4): 372-4.
- Rudge, F. W. (1993). "The role of hyperbaric oxygenation in the treatment of clostridial myonecrosis." Mil Med **158**(2): 80-3.

- Schreiner, A., S. Tonjum, et al. (1974). "Hyperbaric oxygen therapy in bacteroides infections." Acta Chir Scand **140**(1): 73-6.
- Sheffield, P. J. (1998). "Measuring tissue oxygen tension: a review." Undersea Hyperb Med **25**(3): 179-88.
- Sheikh, A. Y., J. J. Gibson, et al. (2000). "Effect of hyperoxia on vascular endothelial growth factor levels in a wound model." Arch Surg **135**(11): 1293-7.
- Sherman, D. M., V. T. Sennik, et al. (1989). "[The efficacy of hyperbaric oxygenation in experimental and hemorrhagic shock]." Anesteziol Reanimatol(2): 37-9.
- Shupak, A., P. Halpern, et al. (1984). "Hyperbaric oxygen therapy for gas gangrene casualties in the Lebanon War, 1982." Isr J Med Sci **20**(4): 323-6.
- Siddiqui, A., J. D. Davidson, et al. (1997). "Ischemic tissue oxygen capacitance after hyperbaric oxygen therapy: a new physiologic concept." Plast Reconstr Surg **99**(1): 148-55.
- Silver, I. A. (1977). "Tissue PO₂ changes in acute inflammation." Adv Exp Med Biol **94**: 769-74.
- Skyhar, M. J., A. R. Hargens, et al. (1986). "Hyperbaric oxygen reduces edema and necrosis of skeletal muscle in compartment syndromes associated with hemorrhagic hypotension." J Bone Joint Surg Am **68**(8): 1218-24.
- Spichev, V. P. and P. Gostev Iu (1987). "[Organization and experience in using hyperbaric oxygenation in the medical installations of a military district]." Voen Med Zh(9): 29-31.
- Stark, R. D., A. P. Coatesworth, et al. (2003). "Management of life-threatening epistaxis in a Jehovah's Witness." Int J Clin Pract **57**(1): 60-1.
- Strauss, M. B. (2003). Hyperbaric Oxygen Therapy Committee Report 2003 - Crush Injuries / Skeletal Muscle-Compartment Syndrome. Hyperbaric Oxygen 2003: Indications and Results: The Hyperbaric Oxygen Therapy Committee Report. J. J. Feldmeier. Kensington, Undersea Hyperbaric Medical Society: 27-34.
- Tai, Y. J., B. C. Birely, et al. (1992). "The use of hyperbaric oxygen for preservation of free flaps." Ann Plast Surg **28**(3): 284-7.
- Tjarnstrom, J., L. Holmdahl, et al. (2001). "Effects of hyperbaric oxygen on expression of fibrinolytic factors of human endothelium in a simulated ischaemia/reperfusion situation." Scand J Clin Lab Invest **61**(7): 539-45.
- Uhl, E., A. Sirsjo, et al. (1994). "Hyperbaric oxygen improves wound healing in normal and ischemic skin tissue." Plast Reconstr Surg **93**(4): 835-41.
- Ulkur, E., F. Yuksel, et al. (2002). "Effect of hyperbaric oxygen on pedicle flaps with compromised circulation." Microsurgery **22**(1): 16-20.
- Van Poucke, S., T. Leenders, et al. (2001). "Hyperbaric oxygen (HBO) as useful, adjunctive therapeutic modality in compartment syndrome." Acta Chir Belg **101**(2): 73-4.
- von Schroeder, H. P. and M. J. Botte (1998). "Crush syndrome of the upper extremity." Hand Clin **14**(3): 451-6.

Wang, J., F. Li, et al. (2002). "The role and effectiveness of adjunctive hyperbaric oxygen therapy in the management of musculoskeletal disorders." J Postgrad Med **48**(3): 226-31.

Warriner, R. A. and H. W. Hopf (2003). Hyperbaric Oxygen Therapy Committee Report 2003 - Healing in Problem Wounds. Hyperbaric Oxygen 2003: Indications and Results: The Hyperbaric Oxygen Therapy Committee Report. J. J. Feldmeier. Kensington, Undersea Hyperbaric Medical Society: 41-55.

Yang, Z. J., G. Bosco, et al. (2001). "Hyperbaric O₂ reduces intestinal ischemia-reperfusion-induced TNF- α production and lung neutrophil sequestration." Eur J Appl Physiol **85**(1-2): 96-103.

Zamboni, W. A., L. K. Browder, et al. (2003). "Hyperbaric oxygen and wound healing." Clin Plast Surg **30**(1): 67-75.

Zamboni, W. A., A. C. Roth, et al. (1993). "Morphologic analysis of the microcirculation during reperfusion of ischemic skeletal muscle and the effect of hyperbaric oxygen." Plast Reconstr Surg **91**(6): 1110-23.

Zamboni, W. A., A. C. Roth, et al. (1989). "The effect of acute hyperbaric oxygen therapy on axial pattern skin flap survival when administered during and after total ischemia." J Reconstr Microsurg **5**(4): 343-7; discussion 349-50.

Zhao, L. L., J. D. Davidson, et al. (1994). "Effect of hyperbaric oxygen and growth factors on rabbit ear ischemic ulcers." Arch Surg **129**(10): 1043-9.

