

AD _____

Award Number: W81XWH-04-1-0806

TITLE: Elucidating the Mechanism of p27 Inactivation by the Bcr-Abl Tyrosine Kinase

PRINCIPAL INVESTIGATOR: Michael W. Deininger, Ph.D.

CONTRACTING ORGANIZATION: Oregon Health and Science University
Portland, OR 97239-0398

REPORT DATE: October 2005

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 01-11-2005		2. REPORT TYPE Final		3. DATES COVERED (From - To) 30 Sep 2004 – 29 Sep 2005	
4. TITLE AND SUBTITLE Elucidating the Mechanism of p27 Inactivation by the Bcr-Abl Tyrosine Kinase				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-04-1-0806	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Michael W. Deininger, Ph.D. E-mail: deininge@ohsu.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon Health and Science University Portland, OR 97239-0398				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Inhibition of Bcr-Abl kinase activity in Mo7e – P210BCR-ABL cells induces accumulation of cells in G0/1. This is associated with (a) increased total levels of p27, (b) accumulation of p27 in the nucleus but not the cytoplasm, and (c) decrease of nuclear but not cytoplasmic Cdk2/cylin E kinase activity. Despite the decrease of Cdk2/cyclin E activity, most of the nuclear p27 is phosphorylated on threonine 187, consistent with diminished nuclear degradation. In accordance with this, the expression of KPC (subunit 1 and 2), the ubiquitin ligase responsible for targeting cytoplasmic p27 for degradation, was not affected by inhibition of Bcr-Abl. The reduced degradation of nuclear p27 is likely due to downregulation of Skp2, the F-box protein of the SCFSKP2 complex that specifically recognizes T187 phosphorylated p27 and targets it for proteasomal degradation. Although no major differences in lineage marker expression were seen in Skp2 knockout mice compared to wildtype mice, there appears to have reduced clonogenicity in response to growth factors and upon expression of Bcr-Abl. This may lead to reduced leukomogenicity of Bcr-Abl in a Skp2 -/- background.					
15. SUBJECT TERMS Bcr-Abl, p27, Skp2, chronic myeloid leukemia (CML)					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	22	19b. TELEPHONE NUMBER (include area code)

Table of Contents

Cover.....	1
SF 298	2
Introduction.....	4
Body.....	4
Key Research Accomplishments	12
Reportable Outcomes.....	12
Conclusions.....	12
References.....	12
Appendix.....	14

INTRODUCTION

Bcr-Abl is a constitutively active tyrosine kinase that is responsible for the pathogenesis of chronic myeloid leukemia (CML) and some cases of acute lymphoblastic leukemia (ALL)(1). Imatinib, a specific inhibitor of Bcr-Abl is effective in patients with CML and Bcr-Abl-positive ALL, however relapse after an initial response is common in patients with advanced disease(2,3). Therefore new therapeutic approaches are required. p27 is an inhibitor of cyclin dependent kinases but has additional functions. It directly inhibits the activity of cdk2/cyclin E and cdk2/cyclin A complexes but it also serves as an assembly factor to generate these complexes(4-6). p27 regulation is complex, involving transcriptional, posttranslational as well as topological levels of control. In Bcr-Abl expressing cell lines p27 is downregulated, and re-expression from ectopic vectors induces cell cycle arrest(7,8). In primary CML cells there is mislocalization to the cytoplasm, where the protein is unable to exert its inhibitory function towards cdk2(9). The precise mechanisms responsible for the deranged regulation of p27 in Bcr-Abl expressing cells are unknown. The nuclear/cytoplasmic distribution of p27 is regulated by shuttle proteins as well as phosphorylation on specific sites, including serine 10(10) and threonine 157 (in humans)(11,12). Proteasomal degradation occurs via distinct nuclear and cytoplasmic pathways. The key ubiquitin ligase responsible for nuclear degradation is the SCF^{Skp2} complex, in which Skp2 plays the key role of the F-box protein recognizing p27 after the latter has been phosphorylated on T187 by cyclin E/cdk2(13,14). Cytoplasmic proteasomal degradation occurs via a novel ubiquitin ligases termed KPC(15).

In many solid tumors cytoplasmic mislocalization is observed at the adenoma stage while more advanced tumors are characterized by down-regulation of overall protein levels. Low levels of p27 tend to correlate with a poor prognosis, and there is an inverse relationship between Skp2 and p27 levels, suggesting that Skp2 may be responsible for the low levels of p27(16,17). In CML, the mechanisms underlying downregulation and mislocalization of p27 are not well understood; however the fact that low p27 levels have adverse prognostic significance in solid tumors suggests that reversing these abnormalities may be therapeutically beneficial. Based on these considerations 2 specific aims were formulated:

- Specific aim 1.** To determine, whether Bcr-Abl regulates the activity of the SCFSKP complex, and whether Skp2 is required for induction of leukemia in a murine model of CML.
- Specific aim 2.** To determine the cellular localization of p27, whether it is associated with phosphorylation of specific residues and whether it influences the degradation of the protein.


BODY

- Task 1.* To determine the ubiquitination activity toward p27 in Mo7e leukemia cells expressing the Bcr-Abl oncoprotein
- Generate a GST-p27 construct and test ubiquitination capacity toward recombinant p27 in cytoplasmic and nuclear extracts from Mo7-p210^{Bcr-Abl} cells (treated and untreated with STI571 [imatinib]). (Months 1-2)
 - Optional follow-up studies: Determine the association of p27 with cyclin D3 and Grb-3-3 by co-immunoprecipitation. (Month 3)

We have generated FLAG-tagged p27 constructs for ubiquitination assays, including wild type p27 as well as S10A, T157A, and T187A mutants. Due to the fact that Dr. Lanker has left OHSU, the technical expertise to perform ubiquitination assays of p27 in nuclear and cytoplasmic lysates is not longer available on site. We have therefore started to set up these assays in our laboratory but have not yet been able to generate consistent results. In addition, with the data shown below it is very likely that cytoplasmic p27 stability is not regulated by Bcr-Abl kinase activity, rendering the ubiquitination experiments a lesser priority.

Task 2. To determine the expression of Skp2 and possibly other SCF^{SKP2} (such as Cks1) in a panel of BCR-ABL-positive leukemia cell lines. (Month 3, in parallel with task 1b)


We tested several Bcr-Abl expressing cell lines, including Mo7e – p210^{BCR-ABL}, K562 and BaF3-p210^{BCR-ABL} cells for expression of Skp2. We find that Skp2 is consistently expressed in BCR-ABL-positive cell lines and that the level of expression decreases with inhibition of Bcr-Abl kinase activity (Figure 1). This is consistent with the notion that Skp2 expression is regulated by Bcr-Abl kinase activity and identifies Skp2 as a downstream target of Bcr-Abl. In order to examine whether these effects were specific to Bcr-Abl expressing cells or were a general phenomenon associated with the expression of an oncogenic tyrosine kinase we have also started to investigate cell lines transformed by different tyrosine kinases, including BaF/3 cells expressing D816V mutant Kit and human lines expressing FLT3-ITD and V617F mutant Jak2. Thus far, results have not been conclusive.



Task 3. To determine, if ectopic Skp2 expression rescues p27 expression and cell cycle progression in Mo7-p210^{BCR-ABL} cell treated with imatinib. (Month 4)

Several considerations led us to change strategy. **(a)** One of the major translational questions to be answered in this project is whether Skp2 may be a therapeutic target – for example, it may be possible to develop specific small molecule inhibitors that interfere with the recognition of p27 by Skp2. This in mind, we realized that it would be more critical to devise a system that permits assessing Skp2's relevance in a direct manner rather than by the experiments originally planned, which would have been based on over-expressing the protein while endogenous protein is downregulated by inhibition of Bcr-Abl. **(b)** The induction of apoptosis along with the G1/0 arrest in Mo7-p210^{BCR-ABL} cells treated with imatinib limits the time frame of the experiments to maximal 20 hours, at which time the cells apoptose. We therefore decided to down-regulate Skp2 expression using shRNA using a lentivirus-based inducible system. The additional advantage of this system is that it can be used in primary cells as the integration of the shRNA construct is stable. As the system is inducible, the expected effect of inhibiting Skp2 expression, namely up-regulation of p27 with subsequent cell cycle arrest can be avoided. Until now we have generated Mo7e and Mo7-p210^{BCR-ABL} cells expressing the tetracycline responsive

transcriptional regulator (Figure 2A), and we were able to show down-regulation of Skp2 in K562 and COS-7 cells using Skp2 shRNA (Figure 2B).



Task 4. To test the requirement of Skp2 for leukemia induction by Bcr-Abl.

- Determine the capacity of *BCR-ABL* to transform Skp2 ^{-/-} murine embryonal fibroblasts compared to wildtype fibroblasts and B cells (Months 5-6)
- Determine the capacity of *BCR-ABL* to induce a myeloproliferative syndrome in Skp2 ^{-/-} mice compared to wildtype mice (Month 5-9, partially in parallel with task 4a)

We have focused on the question whether Bcr-Abl is capable of transforming B-cells and induce a myeloproliferative disease in Skp ^{-/-} compared to Skp2 ^{+/+} mice. Unexpectedly and in contrast to published data we find that Skp2 ^{-/-} mice are born at much lower than Mendelian frequency (Table 1). In addition most Skp ^{-/-} mice do not breed, further limiting the availability of KO mice for experimentation. In initial experiments we determined whether Skp2 ^{-/-} bone marrow cells exhibit a major defect in the distribution of immunophenotypically defined subsets of cells (Table 2).


Litter Group	Parents		Number of pups born	Pups Genotype		
	Father	Mother		WT	KO	HET
1.	HET	HET	7	2	-	5
2.	HET	HET	4	2	-	2
3.	HET	HET	7	3	1	3
4.	HET	HET	6	-	-	6
5.	HET	HET	4	2	-	2

Table 1. Distribution of genotypes in 5 litters using Skp2 ^{+/+} mice. A total 28 mice were born from the 5 heterozygote-heterozygote breedings, but only 1 rather were ^{-/-} compared to 7 mice as expected by mendelian frequency.

Marker	Skp2 ^{+/+}	Skp2 ^{-/-}
Erythroid	4.9%	20.0%
Gr-1	90.2%	86.4%
CD11b	92.1%	85.9%
B220	52.4%	49.2%
Thy12	43.2%	45.5%
CD 133	18.0%	21.1%
CD 45	91.5%	97.6%
CD34	9.1%	13.4%
Thy11	8.3%	5.3%
Sca-1	10.9%	13.1 %
CD117	5.1%	4.5%

Table 2. Immunophenotyping of bone marrow cells derived from Skp2^{+/+} and Skp2^{-/-} mice. No consistent differences in the expression of a panel of lineage specific and early progenitor cell markers were observed

No consistent differences were observed, suggesting that any difference vs. wildtype would be primarily attributable to the role of Skp2 in Bcr-Abl-induced leukemogenesis rather than to differences in the number of target cells available for transformation. In a first set of experiments we compared B-cell transformation and cytokine-independent myeloid colony formation in Skp^{-/-} vs. Skp2^{+/+} bone marrow cells transduced with Bcr-Abl. We find no consistent difference in B-cell transformation (Figure 3), while the formation of myeloid colonies in the absence of cytokines was slightly reduced in Skp2^{-/-} bone marrow (Figure 4). Similarly, myeloid colony formation in response to cytokines was also reduced in Skp2^{-/-} marrow. Given the low numbers of Skp^{-/-} mice, only one bone marrow transplantation experiment could be carried out thus far (Table 3). Since due to unknown reasons the viability of cells prior to injection into the recipients was very low in this experiment we do not consider these data to be reliable. After intense efforts we have now 4 knockout animals available (Table 4). This will allow for transplantation of BCR-ABL transduced bone marrow cells into 8 recipients.






Figure 4. Comparison of myeloid colony formation between *Skp2* $+/+$ and *Skp2* $-/-$ bone marrow cells. Bone marrow cells were transduced with BCR-ABL or empty vector and plated in methylcellulose in the presence and absence of cytokines. The histograms show the average numbers of colonies from triplicate assays. No colonies were recovered from cells transduced with empty vector and grown in the absence of cytokines (not shown).

Sex	HET	WT	KO
Male	22	11	2
Female	12	7	2


Table 4. Size of current *Skp2* mouse colony

Blood counts 13 days after bone marrow transplantation								
Experimental Group	WBC (10e3/mm3)	RBC (10e6/mm3)	HGB (g/dl)	HCT (%)	PLT (10e3/mm3)	LYM (%)	MON (%)	GRA (%)
WT+Bcr-Abl* (n=3)	18.0 $\pm$ 2.6	12.6 $\pm$ 0.3	20.0 $\pm$ 0.7	64.3 $\pm$ 1.9	387.3 $\pm$ 85.4	28.3 $\pm$ 4.5	8.6 $\pm$ 0.7	63.1 $\pm$ 3.8
WT control (n=4)	4.7 $\pm$ 2.1	9.7 $\pm$ 0.9	15.7 $\pm$ 1.3	49.7 $\pm$ 4.3	457.0 $\pm$ 19	26.4 $\pm$ 6.1	11.7 $\pm$ 2.1	61.7 $\pm$ 7.5
KO+Bcr-Abl (n=4)	1.5 $\pm$ 0.12	8.3 $\pm$ 0.5	13.8 $\pm$ 0.7	43.0 $\pm$ 1.68	107.0 $\pm$ 49.7	-	-	-
KO control** (n=3)	0.6 $\pm$ 0.2	4.0 $\pm$ 0.7	6.2 $\pm$ 1.1	17.2 $\pm$ 2.7	30.0 $\pm$ 11.1	-	-	-
Blood counts 20 days after bone marrow transplantation								
WT+Bcr-Abl* (n=1)	32.4	14.6	22.4	74.0	704.0	26.0	5.7	68.3
WT control (n=4)	4.3 $\pm$ 2.3	10.4 $\pm$ 2.6	16.9 $\pm$ 4.1	46.4 $\pm$ 1.3	492.5 $\pm$ 173.5	51.3 $\pm$ 4.2	10.9 $\pm$ 0.8	37.7 $\pm$ 3.9
KO+Bcr-Abl (n=4)	3.1 $\pm$ 1.2	7.4 $\pm$ 1.5	12.8 $\pm$ 2.2	39.12 $\pm$ 5.9	355.0 $\pm$ 335.7	40.2 $\pm$ 3.2	11.6 $\pm$ 0.8	48.2 $\pm$ 2.6
KO control** (n=3)	Sacrificed after 13 days due to poor health condition							

Table 3. Comparison of Bcr-Abl-induced leukemia in *Skp2* $+/+$ vs. *Skp2* $-/-$ mice. Bone marrow cells were transduced with a BCR-ABL retrovirus or the empty construct and injected into lethally irradiated recipients. Blood counts are shown on day 13 and 20 after transplant. * Of the *Skp2* $+/+$ mice transplanted with BCR-ABL-transduced marrow, one died on day 13, other three died on day 17, 20 and 22. ** Of the *Skp2* $-/-$ mice transplanted with empty construct-transduced marrow, one died on day 13. Due to low viability of the cells after retroviral transduction, we consider the data derived from this experiment unreliable, although they may suggest the leukemia induction is delayed in *Skp2* $-/-$ mice.



- Task 5. To determine the intracellular localization, stability and phosphorylation of p27 in Mo7-p210^{BCR-ABL} treated and untreated with imatinib.
- Determine the p27 expression and protein turnover (pulse chase) in cytoplasmic and nuclear lysates (Month 10)
 - Optional follow-up studies (association of p27 with cyclin D, etc.) (Month 11)
 - Studies using leptomycin B to inhibit nuclear export of p27 (Month 11)
 - Determine the phosphorylation of Thr-157 and Ser-10 of p27 and its dependence on Akt and/or hKIS (Months 11-12)


We have determined the time course of p27 upregulation and cell cycle arrest in Mo7-p210^{BCR-ABL} cells upon inhibition of Bcr-Abl kinase activity (Figure 5). We clearly demonstrate that the upregulation of p27 precedes the accumulation of cells in G0/1, suggesting that it is the primary event. Analysis of nuclear and cytoplasmic lysates indicated that the accumulation is almost exclusively the result of increased nuclear levels, while cytoplasmic levels remain practically constant (Figure 5).



While this work was in progress, Kamura et al(15) reported on the identification of KPC, a novel ubiquitin ligase that targets cytoplasmic p27 for degradation. We obtained antibodies against the two components of the KPC complex (KPC1 and KPC2) and demonstrated that levels are not regulated by Bcr-Abl, in agreement with the stable cytoplasmic p27 levels. This corroborates the notion that in the Mo7-p210^{BCR-ABL} cells cytoplasmic p27 is not regulated by Bcr-Abl (Figure 6). In contrast we found nuclear p27 levels strongly increase upon

inhibition of Bcr-Abl and that most of this p27 is phosphorylated on T187, despite the fact that cyclinE/cdk2 activity is decreased (Figure 7). These data concur with the data on Skp2 down-regulation and suggest that Skp2 is the critical mediator of p27 degradation in Bcr-Abl-positive cells.






We were unable to show any differences in S10 p27 phosphorylation, although these experiments were hampered by the poor quality of available antibodies. It remains possible that phosphorylation of T157 is also relevant, as the phosphorylation of this residue by a PI3 kinase dependent pathway has been shown to lead to cytoplasmic relocation of p27, and the inhibition of this phosphorylation could also lead to nuclear retention of the protein. As cytoplasmic levels remained constant this is less likely. Nonetheless, we have generated an anti phospho - T157 antibody that is currently undergoing quality testing (Figure 8). As mentioned above we have generated a Flag-tagged T157A mutant of p27 and plan to use this construct to test the specificity of this antibody. If specific for phosphorylated T157, this reagent will allow us to clarify this question. We have also established and optimized immuno-cytochemistry for p27 to help in analyzing intracellular distribution (Figure 9).

Personnel

The following personnel worked on and received commensurate salary support from this grant:

- Michael Deininger, MD PhD – Principal Investigator
- Paul Yoshihara, PhD – Senior Research Associate
- Taiping Jia – Research Assistant
- Stephanie Willis –Research Associate



KEY RESEARCH ACCOMPLISHMENTS

- Identification of Skp2 as the most likely critical downstream target of Bcr-Abl kinase activity that mediates degradation of p27
- Identification of the nuclear p27 pool as the Bcr-Abl target
- Preliminary results suggesting that myeloid transformation may be delayed in Skp -/- cells

REPORTABLE OUTCOMES

- A manuscript summarizing the main findings is in preparation and will be completed as soon as the bone marrow transplant experiments have been carried out.

CONCLUSIONS

Using this hypothesis development award, we have:

- Identified Skp2 as the most likely candidate mediating the downregulation of p27 by the Bcr-Abl tyrosine kinase. If confirmed in subsequent experiments, our results would imply Skp2 as a therapeutic target in BCR-ABL-positive leukemia;
- Generated a number of reagents that will be invaluable in further and more precisely addressing the question of p27 regulation in BCR-ABL-positive cells, including primary cells from patients with CML and BCR-ABL-positive ALL.

REFERENCES

1. Deininger, M. W.; Goldman, J. M.; Melo, J. V. *Blood* 2000, 96(10), 3343-3356.
2. Druker, B. J.; Sawyers, C. L.; Kantarjian, H.; Resta, D. J.; Reese, S. F.; Ford, J. M.; Capdeville, R.; Talpaz, M. *N.Engl.J.Med.* 2001, 344(14), 1038-1042.
3. Druker, B. J.; Talpaz, M.; Resta, D. J.; Peng, B.; Buchdunger, E.; Ford, J. M.; Lydon, N. B.; Kantarjian, H.; Capdeville, R.; Ohno-Jones, S.; Sawyers, C. L. *N.Engl.J.Med.* 2001, 344(14), 1031-1037.
4. Blain, S. W.; Montalvo, E.; Massague, J. *J.Biol.Chem.* 1997, 272(41), 25863-25872.
5. Sheaff, R. J.; Groudine, M.; Gordon, M.; Roberts, J. M.; Clurman, B. E. *Genes Dev.* 1997, 11(11), 1464-1478.
6. Cheng, M.; Sexl, V.; Sherr, C. J.; Roussel, M. F. *Proc.Natl.Acad.Sci.U.S.A.* 1998, 95(3), 1091-1096.
7. Jonuleit, T.; van der, K. H.; Miething, C.; Michels, H.; Hallek, M.; Duyster, J.; Aulitzky, W. E. *Blood* 2000, 96(5), 1933-1939.
8. Gesbert, F.; Sellers, W. R.; Signoretti, S.; Loda, M.; Griffin, J. D. *J.Biol.Chem.* 2000, 275(50), 39223-39230.
9. Jiang, Y.; Zhao, R. C.; Verfaillie, C. M. *Proc.Natl.Acad.Sci.U.S.A* 2000, 97(19), 10538-10543.
10. Rodier, G.; Montagnoli, A.; Di Marcotullio, L.; Coulombe, P.; Draetta, G. F.; Pagano, M.; Meloche, S. *EMBO J.* 2001, 20(23), 6672-6682.
11. Viglietto, G.; Motti, M. L.; Bruni, P.; Melillo, R. M.; D'Alessio, A.; Califano, D.; Vinci, F.; Chiappetta, G.; Tschlis, P.; Bellacosa, A.; Fusco, A.; Santoro, M. *Nat.Med.* 2002, 8(10), 1136-1144.
12. Viglietto, G.; Motti, M. L.; Fusco, A. *Cell Cycle* 2002, 1(6), 394-400.
13. Carrano, A. C.; Eytan, E.; Hershko, A.; Pagano, M. *Nat.Cell Biol.* 1999, 1(4), 193-199.
14. Pagano, M. *Mol.Cell* 2004, 14(4), 414-416.

15. Kamura, T.; Hara, T.; Matsumoto, M.; Ishida, N.; Okumura, F.; Hatakeyama, S.; Yoshida, M.; Nakayama, K.; Nakayama, K. I. *Nat.Cell Biol.* 2004, 6(12), 1229-1235.
16. Gstaiger, M.; Jordan, R.; Lim, M.; Catzavelos, C.; Mestan, J.; Slingerland, J.; Krek, W. *Proc.Natl.Acad.Sci.U.S.A* 2001, 98(9), 5043-5048.
17. Lim, M. S.; Adamson, A.; Lin, Z.; Perez-Ordenez, B.; Jordan, R. C.; Tripp, S.; Perkins, S. L.; Elenitoba-Johnson, K. S. *Blood* 2002, 100(8), 2950-2956.

APPENDIX

- CV

MICHAEL WERNER NIKOLAUS DEININGER

3181 SW Sam Jackson Park Road L592 • Portland, Oregon 97239

503-494-1603 • fax 503-494-3688 • deininge@ohsu.edu

DOB: April 3, 1963

EDUCATION

Primary School: Schwarzenbach (Germany)	1969-73
Secondary School: Jean-Paul-Gymnasium Hof (Germany)	1973-82
Best final exam in the state of Bavaria (867/900 possible points)	

MEDICAL SCHOOL

University of Regensburg	1982-84
University of Würzburg	1984-90
- clinical clerkships abroad: Glasgow (2 months), London (6 months), Vienna (1 months);	
- final year abroad: Madison/US (3 months), Krakow (3 months), Moscow (5 months)	

QUALIFICATIONS

1990	Graduation
1993	US Medical Licensing Examination, Part II
1994	US Medical Licensing Examination, Part I
1995	M.D. (University of Würzburg)
2000	Ph.D. (University of London)
2001	German Board Certification for Internal Medicine
2002	Habilitation

ADDITIONAL QUALIFICATION

1986	Russicum (university degree, Russian as a foreign language)
------	---

PRESENT APPOINTMENT

Assistant Professor, Division of Hematology/Oncology Oregon Health & Science University, Portland OR	04/02 - present
---	-----------------

PREVIOUS APPOINTMENTS

Consultant, Department of Hematology University of Leipzig, Germany	2001 - 2002
Clinical Fellow, Department of Hematology University of Leipzig	1999 - 2001
Research Fellow, Department of Haematology Hammersmith Hospital, London, UK	1995 - 1999
Resident, Institute of Medical Oncology and Haematology Nürnberg General Hospital, Germany	1992 - 1995
Intern, Marienhospital Aachen, Germany	1990 - 1992

PROFESSIONAL MEMBERSHIPS

1998	German Society of Internal Medicine
------	-------------------------------------

1999 East German Study Group Hematology/Oncology
 2000 Chronic Leukemia Working Party, EBMT
 2000 American Society of Hematology

AWARDS/HONORS

1987 Study grant (Studienstiftung des Deutschen Volkes/German Scholarship Foundation)
 1988 Study grant (Studienstiftung des Deutschen Volkes) to study medicine at the University of London
 1989 12 months grant for final year at the 2. Moscow Medical Institute (DAAD/German Academic Exchange Service)
 1995 1 year fellowship (Deutsche Krebshilfe) to study CML pathogenesis (Imperial College, London)
 1996 1 year fellowship (Deutsche Krebshilfe) to study CML pathogenesis (Imperial College, London)
 1997 Travel grant to attend the ASH meeting (British Society of Hematology)
 1998 Travel grant to attend the Wilsede meeting (Leukemia Research Fund, UK)
 1998 1 year fellowship (Dr Ernst und Anita Bauer Stitung, Germany) to continue Ph.D. at Imperial College
 1998 1 year fellowship (Leuka2000, London) to continue Ph.D. at Imperial College
 1999 Research Grant (Leukemia Research Fund): “*Characterisation of cell lines resistant to STI571*” (co-applicant with Dr Junia Melo) (\$100,000).
 1999 Research Grant (University of Leipzig): “*Role of cyclin D2 in the pathogenesis of CML*” (\$10,000).
 2000 Research Grant (Novartis): “*Prediction of response to STI571 in patients with myeloid blast crisis of CML*” (\$30,000).
 2001 Research grant (Deutsche Krebshilfe): “*Charcaterisation of 2 Bcr-Abl-dependent genes*” (\$ 60,000).
 2001 Research grant (IZKF Leipzig): “*Role of cyclin D2 in the pathogenesis of CML*” (\$75,000).
 2001 Research grant (IZKF Leipzig): “*Gene array analysis to predict response to imatinib in CML patients*” (\$15,000).
 2002 Research grant (Medical Research Foundation of Oregon): “*Clonality analysis of CML patients with complete cytogenetic response to imatinib*” (\$30,000).
 2004 Fellow of the American Society of Hematology

OTHER SCIENTIFIC ACTIVITIES

Reviewer of Scientific Journals

1998- Bone Marrow Transplantation
 2000- Leukemia
 2002- Oncogene
 2002- Experimental Hematology
 2002- Hematologica
 2002- Blood
 2002- Cancer Research
 2003- The Lancet
 2003- Cancer Cell
 2003- Nature Genetics

Reviewer of Grants

2002- Leukemia Research Fund

Membership in Scientific Committees

2000- Molecular diagnostics (German competence network acute and chronic leukemias)
 2001- Website “Chronische Myeloische Leukämie” (German)

(Co-) Principal Investigator in Clinical Studies

- “A phase II study to determine the efficacy and safety of STI571 in patients with chronic myeloid leukemia in myeloid blast crisis” (Novartis)
- “A phase II study to determine the efficacy and safety of STI571 in patients with chronic myeloid leukemia in accelerated phase” (Novartis)
- “A phase II study to determine the efficacy and safety of STI571 in patients with CML resistant or refractory to interferon-alpha” (Novartis)
- “Treatment of CML patients with imatinib + hydroxyurea- a phase I/II study” (East German Study Group Hematology/Oncology [OSHO])
- “Treatment of patients with newly diagnosed CML with pegylated interferon-alpha (pegasys[®]) – a phase I/II study” (OSHO)

Co-Investigator in Clinical Studies

- “A Phase I/II Dose-Finding Study to Determine the Safety, Tolerability, and Anti-Leukemic Effects of STI571 in Combination with Interferon-alpha in Patients with Chronic Myelogenous Leukemia in Chronic Phase” (National Institute of Health [NIH])
- “A Phase II Study to Determine the Anti-Leukemic Effects of STI571 in Combination with Ara-C in Patients with Chronic Myelogenous Leukemia in Chronic Phase” (NIH)
- “A Phase I/II Study to Determine the Safety, Tolerability, and Anti-Leukemic Effects of Trisenox (Arsenic Trioxide) in Combination with Gleevec (STI571) in Patients with Resistant Chronic Myelogenous Leukemia in Chronic Phase” (NIH)
- “Phase II Study Of AG-858 Plus Gleevec[™] In Patients With Chronic Myelogenous Leukemia (CML) In Chronic Phase Who Are Cytogenetically Positive After Treatment With Gleevec[™]”

CLINICAL EXPERIENCE***General Medicine***

Covered all areas of internal medicine including critical care. Experience in the setting of a small community hospital as well as tertiary referral center. Extensive experience in diagnostic procedures (echocardiography, sonography, gastroscopy).

Hematology/Oncology

Extensive experience in the diagnosis and management of patients with hematological malignancies and solid tumors. Several years of experience in high-dose therapy and autologous stem cell transplantation (mainly for lymphoma and germ cell tumors). Almost 3 years experience in allogeneic bone marrow transplantation, including reduced intensity conditioning.

Hematological Diagnostics

Standard morphology. Supervised molecular diagnostics in the Department of Hematology/Oncology at the University of Leipzig. Established or contributed to the establishment of qualitative and quantitative RT-PCR assays and Fluorescence in situ hybridization assays for molecular leukaemia diagnosis. Established data base and specimen repository.

RESEARCH EXPERIENCE***Clinical Research***

- Co - investigator in several Novartis-sponsored international multicenter trials that tested imatinib at various stages of CML and Ph-positive ALL.
- Co - PI in 2 multicenter trials of the East German Study Group Hematology/Oncology (OSHO) that test pegylated interferon and imatinib combined with hydrea
- Co-investigator in several NIH-sponsored multicenter trials that test imatinib in combination with conventional alpha-interferon, cytarabine and arsenic trioxide.

- Habilitation: “Hemmung der *Bcr-Abl*-Tyrosinkinase durch Imatinib, einen selektiven pharmakologischen Inhibitor: Präklinische und klinische Evaluation und Studien zur Pathogenese der chronischen myeloischen Leukämie (CML)”

Laboratory

- MD thesis: “The value of bone marrow biopsies in clinical routine diagnostics: statistical and histological studies”.
- PhD thesis: “STI571, a novel tyrosine kinase inhibitor: pre-clinical evaluation and application to identify downstream targets of Bcr-Abl”
- Extensive experience in molecular biology techniques, including RT-PCR, Northern, Southern, Western blotting, expression cloning and protein expression, Fluorescence in-situ hybridization, cell culture, progenitor cell assays, FACS.

TEACHING EXPERIENCE

Basic/Laboratory

- Extensive experience in supervising research projects.
- Co-supervised PhD thesis of Dr Sara Vieira at Imperial College, London
- Supervised MD theses of Thomas Bumm and Uwe Reinhold at the University of Leipzig
- Co-supervised PhD thesis of Allie Grossmann (Oregon Health & Science University) (submitted June, 2003)
- Lecturer in Biochemistry (graduate course for Ph.D. students at Oregon Health & Science University)

Clinical

- Supervised several final year medical students (Nürnberg General Hospital; University of Leipzig).
- Held lectures during in the main course “Internal Medicine” at Leipzig University
- Lecturer in Clinical Hematology (Graduate course for medical students at Oregon Health & Science University).

RESEARCH SUPPORT

Ongoing Research Support

No # (Deininger)	7/1/04-6/30/06
ASH Scholar Award	51%
	\$75,000

Epidemiology of clonal chromosomal abnormalities, analysis of clonality and assessment of DNA repair capacity in Ph-negative hematopoietic progenitor cells from CML patients with a complete cytogenetic response to imatinib
The major goals of this project are to determine 1) the incidence, prognosis and associated risk factors for the development of karyotypic abnormalities in Ph-negative cells of CML patients treated with imatinib; 2) the incidence of clonal hematopoiesis in patients with a complete cytogenetic response to imatinib and if the Philadelphia-positive cell clone and the abnormal Ph-negative clone(s) originate from a common ancestral cell; and 3) the DNA repair capacity of Ph-negative progenitor cells from CML patient, with and without chromosomal abnormalities in Ph-negative cells, in the presence and absence of imatinib.

Role: PI

7393-06: Specialized Center of Research (Druker)	10/1/05 – 9/30/10
The Leukemia and Lymphoma Society	11%
	\$137,500

Molecularly Targeted Therapy for Leukemia (consortium with Dr. Charles Sawyers, UCLA, and Dr. John Kuriyan, UC Berkeley)

The major goals of Project 1 are to 1) fully characterize the biochemical and biological properties of the most commonly observed BCR-ABL mutants; and 2) define the mechanism of disease persistence in CML patients who are refractory to imatinib at the cytogenetic or molecular level, despite a complete hematologic remission.

Role: Co-PI, Project 1 – Evaluation of Mechanisms of Resistance to Imatinib in CML

1R01 HL082978-01 (Deininger)	9/20/05-8/31/09
------------------------------	-----------------

NIH:NHLBI
Biology of Imatinib-Resistant Mutants of BCR-ABL

23%
\$225,000

The major goals of this project are to 1) determine differences in the transformation potencies of clinically detected BCR-ABL kinase domain (KD) mutants in comparison with wild type BCR-ABL; and 2) compare intrinsic kinase activity between wild type and mutant BCR-ABL and identify critical switches in signaling pathways that cause differences in transformation potency

Role: PI

Completed Research Support

Novartis Pharmaceuticals – Deininger (PI) 2000

Definition of mechanisms of resistance to imatinib in the myeloid
blast crisis of CML

Role: PI

Interdisciplinary Center for Clinical Research – Deininger (PI) 2001

Prediction of cytogenetic response to imatinib using microarray technology

Role: PI

Interdisciplinary Center for Clinical Research – Deininger (PI) 2001-2002

Role of Cyclin D2 for the transformation of lymphoblasts by BCR-ABL

Role: PI

German Cancer Aid (Mildred Scheel Foundation) – Deininger (PI) 2001-2002

Molecular cloning and characterization of 2 BCR-ABL-regulated genes

Role: PI

(No #) – Deininger (PI) 6/1/03-5/31/04

5%

Medical Research Foundation of Oregon – Deininger (PI) \$30,000

Analysis of clonality in complete cytogenetic responders to imatinib

The major goals of this project were to 1) determine if the Bcr-Abl translocation is the only genetic lesion in the chronic phase of CML (i.e. is CML a two step genetic process); 2) determine if treatment with Gleevec will restore polyclonality; and 3) determine if Bcr-Abl positive cells and chromosomally abnormal Ph-negative cells arise from a common ancestor.

Role: PI

CM030024: Hypothesis Development – Deininger (PI) 9/30/04-9/30/05

15%

Department of Defense Chronic Myelogenous Leukemia Research Program \$66,225

Elucidating the Mechanism of p27 Inactivation by the Bcr-Abl Tyrosine Kinase

The major goals of this project are 1) to determine whether Bcr-Abl regulates the activity of the SCFSKP complex and whether Skp2 is required for induction of leukemia in a murine model of CML and 2) to determine the cellular localization of p27, whether it is associated with phosphorylation of specific residues and whether it influences the degradation of the protein.

Role: PI

ADMINISTRATIVE EXPERIENCE

Responsible for recruitment of scientific personnel (laboratory of Professor Brian Druker).

BIBLIOGRAPHY

1. Deininger, M.W., et al., The tyrosine kinase inhibitor CGP57148B selectively inhibits the growth of BCR-ABL-positive cells. *Blood*, 1997. 90(9): p. 3691-8.
2. Bose, S., et al., The presence of typical and atypical BCR-ABL fusion genes in leukocytes of normal individuals: biologic significance and implications for the assessment of minimal residual disease. *Blood*, 1998. 92(9): p. 3362-7.
3. Deininger, M.W. and J.M. Goldman, Chronic myeloid leukemia. *Curr Opin Hematol*, 1998. 5(4): p. 302-8.
4. Deininger, M.W., et al., Selective induction of leukemia-associated fusion genes by high-dose ionizing radiation. *Cancer Res*, 1998. 58(3): p. 421-5.
5. Gora-Tybor, J., et al., The susceptibility of Philadelphia chromosome positive cells to FAS-mediated apoptosis is not linked to the tyrosine kinase activity of BCR-ABL. *Br J Haematol*, 1998. 103(3): p. 716-20.
6. Deininger, M.W., J.M. Goldman, and J.V. Melo, The molecular biology of chronic myeloid leukemia. *Blood*, 2000. 96(10): p. 3343-56.
7. Deininger, M., et al., No evidence for persistence of BCR-ABL-positive cells in patients in molecular remission after conventional allogeneic transplantation for chronic myeloid leukemia. *Blood*, 2000. 96(2): p. 779-80.
8. Deininger, M.W., et al., BCR-ABL tyrosine kinase activity regulates the expression of multiple genes implicated in the pathogenesis of chronic myeloid leukemia. *Cancer Res*, 2000. 60(7): p. 2049-55.
9. Mahon, F.X., et al., Selection and characterization of BCR-ABL positive cell lines with differential sensitivity to the tyrosine kinase inhibitor STI571: diverse mechanisms of resistance. *Blood*, 2000. 96(3): p. 1070-9.
10. Marley, S.B., et al., The tyrosine kinase inhibitor STI571, like interferon-alpha, preferentially reduces the capacity for amplification of granulocyte-macrophage progenitors from patients with chronic myeloid leukemia. *Exp Hematol*, 2000. 28(5): p. 551-7.
11. Deininger, M.W., et al., Direct relation between BCR-ABL tyrosine kinase activity and cyclin D2 expression in lymphoblasts. *Cancer Res*, 2001. 61(21): p. 8005-13.
12. Deininger, M., et al., Chemotherapy for mobilisation of Ph-negative progenitor cells from patients with CML: impact of different mobilisation regimens. *Bone Marrow Transplant*, 2001. 27(11): p. 1125-32.
13. Vieira, S.A., et al., Transcription factor BACH2 is transcriptionally regulated by the BCR/ABL oncogene. *Genes Chromosomes Cancer*, 2001. 32(4): p. 353-63.
14. Al-Ali, H.K., et al., CML with an e1a3 BCR-ABL fusion: rare, benign, and a potential diagnostic pitfall. *Blood*, 2002. 100(3): p. 1092-3.
15. Deininger, M. and D. Niederwieser, [Molecular therapy of chronic myeloid leukemia--a new era in the treatment of malignant diseases]. *Med Klin (Munich)*, 2002. 97 Suppl 1: p. 1.
16. Huntly, B.J., et al., Derivative chromosome 9 deletions in chronic myeloid leukemia: poor prognosis is not associated with loss of ABL-BCR expression, elevated BCR-ABL levels, or karyotypic instability. *Blood*, 2002. 99(12): p. 4547-53.
17. Kohler, T., et al., High Bax and Bcl-2 mRNA expression correlate with negative outcome in acute myeloid leukemia (AML). *Leukemia*, 2002. 16(1): p. 22-9.
18. La Rosee, P., et al., Activity of the Bcr-Abl kinase inhibitor PD180970 against clinically relevant Bcr-Abl isoforms that cause resistance to imatinib mesylate (Gleevec, STI571). *Cancer Res*, 2002. 62(24): p. 7149-53.
19. Muller, C., et al., The BCR/ABL-extra signal fluorescence in situ hybridization system reliably detects deletions upstream of the ABL locus: implications for reporting of results and followup of chronic myelogenous leukemia patients. *Cancer Genet Cytogenet*, 2002. 136(2): p. 149-50.
20. Ottmann, O.G., et al., A phase 2 study of imatinib in patients with relapsed or refractory Philadelphia chromosome-positive acute lymphoid leukemias. *Blood*, 2002. 100(6): p. 1965-71.
21. Reid, A.G., et al., Deletions of the derivative chromosome 9 do not account for the poor prognosis associated with Philadelphia-positive acute lymphoblastic leukemia. *Blood*, 2002. 99(6): p. 2274-5.
22. Sawyers, C.L., et al., Imatinib induces hematologic and cytogenetic responses in patients with chronic myelogenous leukemia in myeloid blast crisis: results of a phase II study. *Blood*, 2002. 99(10): p. 3530-9.
23. Talpaz, M., et al., Imatinib induces durable hematologic and cytogenetic responses in patients with accelerated phase chronic myeloid leukemia: results of a phase 2 study. *Blood*, 2002. 99(6): p. 1928-37.
24. Bumm, T., et al., Emergence of clonal cytogenetic abnormalities in Ph- cells in some CML patients in cytogenetic remission to imatinib but restoration of polyclonal hematopoiesis in the majority. *Blood*, 2003. 101(5): p. 1941-9.
25. Corbin, A.S., et al., Several Bcr-Abl kinase domain mutants associated with imatinib mesylate resistance remain sensitive to imatinib. *Blood*, 2003. 101(11): p. 4611-4.

26. Deininger, M.W. and B.J. Druker, Specific targeted therapy of chronic myelogenous leukemia with imatinib. *Pharmacol Rev*, 2003. 55(3): p. 401-23.
27. Deininger, M.W., Cytogenetic studies in patients on imatinib. *Semin Hematol*, 2003. 40(2 Suppl 2): p. 50-5.
28. Deininger, M.W., et al., Practical management of patients with chronic myeloid leukemia receiving imatinib. *J Clin Oncol*, 2003. 21(8): p. 1637-47.
29. Dinulescu, D.M., et al., c-CBL is not required for leukemia induction by Bcr-Abl in mice. *Oncogene*, 2003. 22(55): p. 8852-60.
30. Huntly, B.J., et al., Imatinib improves but may not fully reverse the poor prognosis of patients with CML with derivative chromosome 9 deletions. *Blood*, 2003. 102(6): p. 2205-12.
31. La Rosee, P., et al., No correlation between the proliferative status of Bcr-Abl positive cell lines and the proapoptotic activity of imatinib mesylate (Gleevec/Glivec). *Hematol J*, 2003. 4(6): p. 413-9.
32. Lange, T., D.W. Niederwieser, and M.W. Deininger, Residual disease in chronic myeloid leukemia after induction of molecular remission. *N Engl J Med*, 2003. 349(15): p. 1483-4.
33. Lange, T., et al., High levels of BAX, low levels of MRP-1, and high platelets are independent predictors of response to imatinib in myeloid blast crisis of CML. *Blood*, 2003. 101(6): p. 2152-5.
34. Muller, M.C., et al., Dynamics of BCR-ABL mRNA expression in first-line therapy of chronic myelogenous leukemia patients with imatinib or interferon alpha/ara-C. *Leukemia*, 2003. 17(12): p. 2392-400.
35. Niederwieser, D., et al., Low-dose total body irradiation (TBI) and fludarabine followed by hematopoietic cell transplantation (HCT) from HLA-matched or mismatched unrelated donors and postgrafting immunosuppression with cyclosporine and mycophenolate mofetil (MMF) can induce durable complete chimerism and sustained remissions in patients with hematological diseases. *Blood*, 2003. 101(4): p. 1620-9.
36. O'Brien, S.G. and M.W. Deininger, Imatinib in patients with newly diagnosed chronic-phase chronic myeloid leukemia. *Semin Hematol*, 2003. 40(2 Suppl 2): p. 26-30.
37. Olavarria, E., et al., Response to imatinib in patients who relapse after allogeneic stem cell transplantation for chronic myeloid leukemia. *Leukemia*, 2003. 17(9): p. 1707-12.
38. Reinhold, U., et al., FISH for BCR-ABL on interphases of peripheral blood neutrophils but not of unselected white cells correlates with bone marrow cytogenetics in CML patients treated with imatinib. *Leukemia*, 2003. 17(10): p. 1925-9.
39. Rumpel, M., T. Friedrich, and M.W. Deininger, Imatinib normalizes bone marrow vascularity in patients with chronic myeloid leukemia in first chronic phase. *Blood*, 2003. 101(11): p. 4641-3.
40. Tipping, A.J., et al., Comparative gene expression profile of chronic myeloid leukemia cells innately resistant to imatinib mesylate. *Exp Hematol*, 2003. 31(11): p. 1073-80.
41. Wiedmann, M., et al., Imatinib mesylate (STI571; Glivec)--a new approach in the treatment of biliary tract cancer? *Anticancer Drugs*, 2003. 14(9): p. 751-60.
42. Al-Ali, H.K., et al., High incidence of BCR-ABL kinase domain mutations and absence of mutations of the PDGFR and KIT activation loops in CML patients with secondary resistance to imatinib. *Hematol J*, 2004. 5(1): p. 55-60.
43. Corbin, A.S., et al., Sensitivity of oncogenic KIT mutants to the kinase inhibitors MLN518 and PD180970. *Blood*, 2004. 104(12): p. 3754-7.
44. Deininger, M.W. and B.J. Druker, SRCircumventing imatinib resistance. *Cancer Cell*, 2004. 6(2): p. 108-10.
45. Deininger, M., Src kinases in Ph+ lymphoblastic leukemia. *Nat Genet*, 2004. 36(5): p. 440-1.
46. Deininger, M.W., et al., Detection of ABL kinase domain mutations with denaturing high-performance liquid chromatography. *Leukemia*, 2004. 18(4): p. 864-71.
47. Deininger, M.W., Basic science going clinical: molecularly targeted therapy of chronic myelogenous leukemia. *J Cancer Res Clin Oncol*, 2004. 130(2): p. 59-72.
48. Gaston, I., et al., Coexistence of phosphotyrosine-dependent and -independent interactions between Cbl and Bcr-Abl. *Exp Hematol*, 2004. 32(1): p. 113-21.
49. Griswold, I.J., et al., Effects of MLN518, a dual FLT3 and KIT inhibitor, on normal and malignant hematopoiesis. *Blood*, 2004. 104(9): p. 2912-8.
50. Grossmann, A.H., et al., Catalytic domains of tyrosine kinases determine the phosphorylation sites within c-Cbl. *FEBS Lett*, 2004. 577(3): p. 555-62.
51. La Rosee, P., et al., In vitro efficacy of combined treatment depends on the underlying mechanism of resistance in imatinib-resistant Bcr-Abl-positive cell lines. *Blood*, 2004. 103(1): p. 208-15.

52. Lange, T., et al., BCR-ABL transcripts are early predictors for hematological relapse in chronic myeloid leukemia after hematopoietic cell transplantation with reduced intensity conditioning. *Leukemia*, 2004. 18(9): p. 1468-75.
53. Lange, T., et al., Quantitative reverse transcription polymerase chain reaction should not replace conventional cytogenetics for monitoring patients with chronic myeloid leukemia during early phase of imatinib therapy. *Haematologica*, 2004. 89(1): p. 49-57.
54. Loriaux, M. and M. Deininger, Clonal cytogenetic abnormalities in Philadelphia chromosome negative cells in chronic myeloid leukemia patients treated with imatinib. *Leuk Lymphoma*, 2004. 45(11): p. 2197-203.
55. Mauro, M.J., M. Loriaux, and M.W. Deininger, Ph-positive and -negative myeloproliferative syndromes may co-exist. *Leukemia*, 2004. 18(7): p. 1305-7.
56. Melo, J.V. and M.W. Deininger, Biology of chronic myelogenous leukemia--signaling pathways of initiation and transformation. *Hematol Oncol Clin North Am*, 2004. 18(3): p. 545-68, vii-viii.
57. O'Hare, T., et al., Inhibition of wild-type and mutant Bcr-Abl by AP23464, a potent ATP-based oncogenic protein kinase inhibitor: implications for CML. *Blood*, 2004. 104(8): p. 2532-9.
58. Silver, R.T., et al., The second international congress on myeloproliferative and myelodysplastic syndromes. *Leuk Res*, 2004. 28(9): p. 979-85.
59. Corbin, A.S., et al., In vitro and in vivo activity of ATP-based kinase inhibitors AP23464 and AP23848 against activation loop mutants of Kit. *Blood*, 2005.
60. Deininger, M.W. and T.L. Holyoake, Can we afford to let sleeping dogs lie? *Blood*, 2005. 105(5): p. 1840-1.
61. Deininger, M., E. Buchdunger, and B.J. Druker, The development of imatinib as a therapeutic agent for chronic myeloid leukemia. *Blood*, 2005. 105(7): p. 2640-53.
62. Demehri, S., et al., e8a2 BCR-ABL: more frequent than other atypical BCR-ABL variants? *Leukemia*, 2005.
63. Mayerhofer, M., et al., Identification of mTOR as a novel bifunctional target in chronic myeloid leukemia: dissection of growth-inhibitory and VEGF-suppressive effects of rapamycin in leukemic cells. *Faseb J*, 2005.
64. O'Hare, T., et al., AMN101: tightening the grip of imatinib. *Cancer Cell*, 2005. 7(2): p. 117-9.
65. Walters, D.K., et al., RNAi-induced down-regulation of FLT3 expression in AML cell lines increases sensitivity to MLN518. *Blood*, 2005. 105(7): p. 2952-4.
66. Crossman, L.C., et al., A single nucleotide polymorphism in the coding region of ABL and its effects on sensitivity to imatinib. *Leukemia*, 2005.
67. Chuah, C., et al., Zoledronate inhibits proliferation and induces apoptosis of imatinib-resistant chronic myeloid leukaemia cells. *Leukemia*, 2005.
68. O'Hare, T., et al., Combined Abl inhibitor therapy for minimizing drug resistance in chronic myeloid leukemia: Src/Abl inhibitors are compatible with imatinib. *Clin Cancer Res*, 2005. 11(19 Pt 1): p. 6987-93.
69. Levine, R.L., et al., The JAK2V617F activating mutation occurs in chronic myelomonocytic leukemia and acute myeloid leukemia, but not in acute lymphoblastic leukemia or chronic lymphocytic leukemia. *Blood*, 2005.
70. Crossman, L.C., et al., hOCT 1 and resistance to imatinib. *Blood*, 2005. 106(3): p. 1133-4; author reply 1134.
71. O'Hare, T., et al., In vitro activity of Bcr-Abl inhibitors AMN107 and BMS-354825 against clinically relevant imatinib-resistant Abl kinase domain mutants. *Cancer Res*, 2005. 65(11): p. 4500-5.
72. Willis, S.G., et al., High-sensitivity detection of BCR-ABL kinase domain mutations in imatinib-naive patients: correlation with clonal cytogenetic evolution but not response to therapy. *Blood*, 2005. 106(6): p. 2128-37.
73. Lange, T., et al., Durability of molecular remission in chronic myeloid leukemia patients treated with imatinib vs allogeneic stem cell transplantation. *Leukemia*, 2005. 19(7): p. 1262-5.
74. Crossman, L.C., et al., In chronic myeloid leukemia white cells from cytogenetic responders and non-responders to imatinib have very similar gene expression signatures. *Haematologica*, 2005. 90(4): p. 459-64.
75. Corbin, A.S., et al., In vitro and in vivo activity of ATP-based kinase inhibitors AP23464 and AP23848 against activation loop mutants of Kit. *Blood*, 2005.

Book Chapters

1. Deininger M. Grundlagen der Therapie mit Imatinib Mesilat in: Fischer T (ed.) Der Signalhemmer Imatinib Mesilat (STI571). Uni-Med Verlag, Bremen 2002 (German)
2. Deininger MW. Clinical applications in: Stein G and Pardee A (eds.). Cell cycle and growth control: biomolecular regulation and cancer. Wiley&Sons, N.Y., 2003 (*in press*)

Invited Lectures at International Conferences

- Identification of Bcr-Abl target genes using differential display (Biarritz, 1999)
- Mechanisms of defective cell cycle regulation in CML cells (Rapallo, 2000)
- Molecular pathogenesis of CML (Porto Allegre, 2001)
- Role of imatinib prior to allografting (Paris, 2001)
- Targeted therapy of CML with imatinib (Krakow, 2001)
- Imatinib prior to allogeneic transplantation (Madrid, 2002)
- Cytogenetics in CML patients on imatinib (Bordeaux, 2002)
- Role of imatinib for remission induction prior to allografting (Houston, 2002)
- Targeted therapy of CML with imatinib: a paradigm shift in cancer therapy? (Berlin, 2003)