

DNA Microarray Analysis of Human Monocytes Early Response Genes upon Infection with *Rickettsia rickettsii*

Chien-Chung Chao

Rickettsiae Diseases Department

Infectious Diseases Directorate

Naval Medical Research Center

Silver Spring, MD

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 15 NOV 2004		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE DNA Microarray Analysis of Human Monocytes Early Response Genes upon Infection with Rickettsia rickettsii				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Rickettsiae Diseases Department Infectious Diseases Directorate Naval Medical Research Center Silver Spring, MD				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001849, 2004 Scientific Conference on Chemical and Biological Defense Research. Held in Hunt Valley, Maryland on 15-17 November 2004 . , The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 18	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Rickettsiae

- Gram negative coccobacillary bacteria
- Obligate intracellular organisms
- Arthropod-borne
- Cause febrile diseases (mild to life threatening)
- Military importance
 - Epidemic Typhus: Peloponnesian war, Napoleon, WWI, WWII
 - Trench fever: WWI and WWII
 - Scrub typhus: WWII, Vietnam, Camp Fuji
 - Ehrlichiosis: (HME Ft Chaffee; Quantico, Ft Campbell)
 - Spotted Fever: (Ft Chaffee, Ft Bragg, Botswana)

Typhus Group (TG) Rickettsiae

- Epidemic (louse-borne) Typhus: *Rickettsia prowazekii*
- Murine (flea-borne) Typhus: *R. typhi*

Spotted Fever Group (SFG)

- Rocky Mountain spotted fever: *Rickettsia rickettsii*
- Mediterranean spotted fever: *R. conorii*
- African tick bite fever: *R. africae*
- North Asian/Siberian tick typhus: *R. sibirica*

Rocky Mountain Spotted Fever

- Etiologic agent: *R. rickettsii*
- Reported in USA, Canada, Mexico, Costa Rica, Panama, Columbia, and Brazil.
- Most severe SFG rickettsial disease and is the most commonly fatal tick-borne disease in the United States. Case-fatality rates up to 30% were reported in the pre-antibiotic era but rates have remained between 2-10 % since the 1950's.
- Recent Brazilian outbreak, 66% of cases were fatal.
- Seroconversions were observed for military units that trained in Arkansas (38%) and more than 40% of these individuals received medical treatment.


Diagnosis of rickettsial disease

- Generally based on clinical presentation and exposure history of a patient.
- Clinical characterizations are sudden onset of fever, primary eschars, severe headache, myalgia, arthralgia, malaise, and skin rashes.
- Differentiating rickettsial diseases from other acute tropical febrile illnesses can be difficult because of the similarities in signs and symptoms.

Currently available laboratory diagnosis of rickettsial diseases

- serodiagnostic assays such as the indirect immunoperoxidase (IIP) assay and indirect immunofluorescent (IFA) tests (weeks)
- nucleic acid detection by PCR amplification of rickettsial genes (days)

General Reaction Scheme



Hypothesis


Infection of human monocytes with *R. rickettsii* causes specific mRNA expression pattern. This pattern could be used to distinguish different types of infection and to identify novel genes involved in host response.

Experimental Design


- Human monocytes (THP-1) were infected with *R. rickettsii* for 45 min in 5% CO₂ incubator at 35°C with gentle rotation.
- Non-associated *R. rickettsii* were removed by centrifugation.
- Infected and control THP-1 cells were left in the incubator for additional 1, 4, 8 and 18 h.
- At indicated time, cells were centrifuged, washed with PBS and RNA was extracted with trizol.
- Trizol extracted RNA was further cleaned with a RNeasy kit.

Experimental Design (continued)


- Reverse transcription reaction was carried out using poly dT and Superscript kit in the presence of Cy3-dUTP (for labeling sample RNA) and Cy5-dUTP (for labeling human reference RNA).
- Labeled RNA was purified and hybridized at 42°C for overnight onto a DNA microarray slide with 7680 human genes.
- The fluorescence images were scanned and visualized using GenePix Pro 4.0 (Axon Lab).
- Data was analyzed using web-based analysis software from NCI (<http://nciarray.nci.nih.gov>).




ScatterPlot log2 Ratios; $r = 0.797$



Up-regulated genes



Down-regulated genes



Comparison of gene lists from various infectious agents

Infectious agents	<i>Orientia tsutsugamushi</i>	<i>R. prowazekii</i>	<i>R. rickettsii</i>	8 others (Nau et. al.) ^a
<i>Orientia tsutsugamushi</i>	20/2 ^b	1 ^c	0	3
<i>R. prowazekii</i>	---	24/7	0	6
<i>R. rickettsii</i>	---	---	60/25	2
8 others	---	---	---	139/62

^a. Data taken from Nau et. al., PNAS, 2002, 99(3), 1503-1508.

^b. Presented as the number of up/down regulated genes.

^c. Numbers represent genes identified in both infectious agents.

Conclusions

- Infection of human monocytes with *R. rickettsii* resulted in 60 up-regulated and 25 down-regulated genes. These affected genes may be important for the design of diagnostic markers.
- Comparison with microarray results from other infectious agents indicated list of unique genes responsive to *R. rickettsii* infection (Poster by Ge et. al, in Diagnostics session).

Future work

- Confirmation of up- and down-regulated genes.
- Replicates and multiple time points are necessary to identify genes consistently regulated by *R. rickettsii* infection.
- Statistical analysis is needed to better identify genes significantly regulated by *R. rickettsii* infection.

Acknowledgement

RDD/IDD/NMRC

Dr. Wei-Mei Ching

Dr. Allen Richards

T. C. Chan

NCI/NIH

Dr. Eric Y. Chuang

Shuping Zhao

Funding

MIDRP Task Area L