



United States Army Medical Research Institute of Infectious Diseases

Increased Stability of a Designed, Single Domain Ricin A-chain Vaccine

John H. Carra

November 17, 2003

Toxinology and Aerobiology Division
USAMRIID
Fort Detrick, Maryland

Report Documentation Page

Form Approved
OMB No. 0704-0188

Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

1. REPORT DATE 01 OCT 2005		2. REPORT TYPE N/A		3. DATES COVERED -	
4. TITLE AND SUBTITLE Increased Stability of a Designed, Single Domain Ricin A-chain Vaccine				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Toxinology and Aerobiology Division USAMRIID Fort Detrick, Maryland				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM001851, Proceedings of the 2003 Joint Service Scientific Conference on Chemical & Biological Defense Research, 17-20 November 2003. , The original document contains color images.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Goal: Create a soluble and stable platform for safe presentation of a neutralizing epitope against ricin A-chain.

Limitations of earlier vaccine candidates:


- Residual catalytic activity
- Poor solubility -- aggregation
- Relatively unstable

Solution: Design a stable, monomeric, single-domain derivative of RTA by protein engineering.

- Mark Olson modeling
- Virginia Roxas cloning
- Leonard Smith, Robert Wannemacher toxicology, immunology
- many others ... team effort

The A-chain of Ricin (267 a.a.)

(1rtc.pdb, Mlsna et al., 1993, *Protein Sci.* 2:429-435)



Vaccine candidates created: RTA 1-33/44-198
RTA 1-198

Design considerations:

1. Conservation of structure from WT RTA
 - Maintain epitope in natural conformation.
2. Soluble and monomeric
 - No aggregation.
3. High stability
 - Avoid degradation during production and storage.
 - Avoid need for stabilizing excipients in formulation, the "cold chain."
4. No toxicity, adequate neutralizing antigenicity
 - *Verified in mouse model*

Secondary structure contents close to that of RTA
are predicted for the mutants,
if the N-terminal domain is not altered by the residue deletions


Table 1. Predicted Secondary Structure Content^a

Protein	% β sheet	% α helix	% Other
RTA	16	32	52
RTA 1-198	16	34	50
RTA 1-33/44-198	17	36	47

^a Percentage of amino acids predicted in each type of secondary structure, based on the known structure of RTA (Mlsna et al., 1993).


The RTA derivatives are folded and have secondary structure contents similar to recombinant and natural wild-type RTA

Far-UV Circular Dichroism Spectra



rRTA ———
1-33/44-198 - - -
1-198
nRTA ●●●●●

Fourier-transform infrared spectra in the amide I' region also indicate conservation of structure



rRTA

—

1-33/44-198

—

RTA 1-33/44-198 is a monodisperse monomer in solution

Table 2. Dynamic Light Scattering Measurements

Protein	R_h^a	hydrodynamic MW_r^b	actual MW_r^c
RTA	2.53	29.5×10^3	30.0×10^3
RTA 1-198	2.16	20.4×10^3	22.5×10^3
RTA 1-33/44-198	2.11	19.3×10^3	21.3×10^3


^a The hydrodynamic radius of the protein in solution as measured by DLS, in nanometers.

^b Molecular weight calculated using the Dynamics software (Protein Solutions).

^c Molecular weight predicted from the amino acid sequence.

0.8 mg/ml protein in PBS buffer
PBS buffer was used in all experiments because this
is the expected formulation condition.


RTA 1-33/44-198 and RTA 1-198 have increased thermostability





- 1 rRTA
- 2 nRTA
- 3 1-198
- 4 1-33/44-198

RTA 1-33/44-198 is more stable to long-term incubation at 37 °C


rRTA and RTA 1-33/44-198 were mixed together and incubated at 37 °C for a period of time, then precipitated protein was removed by centrifugation and the soluble fraction run on a gel.



% Solubility of proteins after extended incubation at 37 °C
RTA 1-33/44-198 remains in solution longest



Denaturation by guanidine-HCl followed by CD at 225 nm
RTA 1-33/44-198 is the most stable to chemical denaturation



- rRTA
- ▲ 1-198
- 1-33/44-198
- ◆ nRTA

Stabilities of wild-type and mutant RTA molecules

Table 3. Stability parameters of RTA molecules.

Protein	T_d^a	C_m^b	% soluble ^c
rRTA	46.6 ±0.1	1.56 ±0.022	6
rRTA 1-198	50.8 ±0.5	2.29 ±0.004	25
rRTA 1-33/44-198	53.1 ±0.1	2.53 ±0.041	58
nRTA	45.6 ±0.1	1.39 ±0.020	31

^a Temperature of onset of denaturation in thermal melting in °C as monitored by CD at 205 nm. T_d was taken as the point at which the signal had diverged from that of the native state by 5%. Results are the average and standard error of three measurements.

^b The midpoint of denaturation in Molar [GuHCl] as measured by CD at 225 nm.

^c The percent of protein remaining soluble after incubation at 37 °C for 106 hours.

Errors are shown in Figure 5.

Demonstrated:


1. Conservation of secondary structure from wild-type RTA
 - Circular dichroism and FTIR data
2. Soluble and monomeric
 - Dynamic light scattering
3. High stability
 - Melting curves
 - Extended incubation at physiological temperature

**What is the origin of the increased stability
of RTA 1-33/44-198 and 1-198?**


Three-state unfolding of RTA through an intermediate

12 Tyr


1 Trp
2 Tyr



Denaturation of wild-type rRTA by guanidine-HCl proceeds through a partially unfolded intermediate




- ▲ Fluor ex 295 em 320
- ▲ CD 225 nm
- ▲ Fluor ex 280 em 303
- ANS Fluor 480 nm




Anilinonaphthalene sulfonic acid (ANS)

Denaturation of wild-type rRTA by guanidine-HCl




Followed by fluorescence emission with excitation at 280 nm

Denaturation of the truncated RTA derivatives by guanidine-HCl shows no intermediate -- a two-state process



- 1-33/44-198 CD 225
- 1-198 CD 225
- 1-33/44-198 Fluor
(ex 280, em 303)
- 1-198 Fluor
- ▲ 1-33/44-198 ANS
- ▲ 1-198 ANS

Denaturation of RTA1-33/44-198 by guanidine-HCl




Followed by tyrosine fluorescence emission with excitation at 280 nm


Three-state unfolding of RTA through an intermediate

12 Tyr

1 Trp
2 Tyr



Comparison of tryptophan vs. tyrosine fluorescence emission data and the absence of the intermediate for the vaccine candidates indicate that the C-terminal domain of RTA is relatively unstable, unfolding before the N-terminal domain.



Intracellular translocation of RTA uses pathways for export from the ER of misfolded proteins.

-- A physiological role for an intermediate state?

[Argent et al., 2000, *JBC* 275:9263-9269, Day et al., 2002, *Biochemistry* 41:2836-2843, Sandvig and van Deurs, 2002, *FEBS Letts.* 529:49-53]

Main conclusions

1. The RTA vaccine candidates are soluble, folded, monomeric, and more stable than WT RTA.
 - Superior biophysical properties for a vaccine.
2. Unfolding of RTA involves a partially unfolded intermediate state with a disrupted C-terminal domain.
3. Mutant derivatives lacking the C-domain do not adopt the intermediate state, which helps explain their greater stability.

The native state of RTA is limited in its stability by a propensity to transform to a partially unfolded state.

Acknowledgements

Colleen McHugh
Rowena Schokman
Christine Johnston

Ralph Tammariello
LTC Charles Millard