

AD _____

Award Number: DAMD17-01-1-0733

TITLE: Eliciting Autoimmunity to Ovarian Tumors in Mice by Genetic Disruption of
T Cell Tolerance Mechanisms

PRINCIPAL INVESTIGATOR: Brad H. Nelson, Ph.D.

CONTRACTING ORGANIZATION: British Columbia Cancer Agency
Victoria, British Columbia, Canada V8R 6V5

REPORT DATE: August 2006

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 01-08-2006		2. REPORT TYPE Final		3. DATES COVERED 1 Aug 2001 – 31 Jul 2006	
4. TITLE AND SUBTITLE Eliciting Autoimmunity to Ovarian Tumors in Mice by Genetic Disruption of T Cell Tolerance Mechanisms				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER DAMD17-01-1-0733	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Brad H. Nelson, Ph.D.				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) British Columbia Cancer Agency Victoria, British Columbia, Canada V8R 6V5				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES Original contains colored plates: ALL DTIC reproductions will be in black and white.					
14. ABSTRACT We have developed a mouse model for ovarian cancer that allows monitoring of tumor-specific T cell clones as they encounter ovarian tumors in vivo. We "tagged" the neu oncogene with two defined T cell epitopes so as to confer recognition by available T cell receptor (TCR) transgenic T cells. When expressed in the murine ovarian tumor cell line ID8, epitope-tagged neu (designated neuOT1/OT2) induces the formation of aggressive ovarian adenocarcinomas that express the epitope tags and hence are recognizable by adoptively transferred TCR transgenic T cells. We successfully made the neuOT1/OT2 expression construct and stably expressed it in an aggressive subclone of the ID8 cell line, designated ID8-G7, which was derived by serial in vivo passage of the original ID8 line. When injected intraperitoneally into syngenic mice, ID8-G7 cells expressing neuOT1/OT-II give rise within one month to disseminated ovarian cancer with extensive ascites (Aim 1). CD8+ (OT-I) T cells specific for neuOT1/OT-II proliferate extensively after adoptive transfer into tumour-bearing hosts and, remarkably, induce complete tumour regression within 10 days in a dose-dependent manner (Aim 2). In the next year, we will test whether the dose-dependency of this response can be mitigated by use of autoimmune-prone Cbl-b-deficient CD8+ T cells (Aim 3).					
15. SUBJECT TERMS Tumor immunology, immunotherapy, animal models, CD4+ and CD8+ T 15. Number of Pages (count all pages including appendices) cells, HER2/neu, tumor antigens					
16. SECURITY CLASSIFICATION OF:			UU	18. NUMBER OF PAGES 15	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

Introduction.....	4
Body.....	4
Key Research Accomplishments.....	6
Reportable Outcomes.....	7
Conclusions.....	8
References.....	8
Appendices.....	8

DAMD17-01-1-0733 Annual Progress Report 2006

PI: Brad H. Nelson, Ph.D.

Title of Project: Eliciting Autoimmunity to Ovarian Tumors in Mice by Genetic Disruption of T Cell Tolerance Mechanisms

Introduction:

Research in the fields of basic immunology and autoimmunity has identified several distinct mechanisms through which immune tolerance is established and maintained in the normal host, and additional mechanisms will likely be identified in future. We hypothesize that ovarian tumors are recognized in an antigen-specific manner by T cells but induce immunologic tolerance through one or more of these homeostatic mechanisms, which have evolved to protect the host from autoimmune attack. We further hypothesize that tolerance to ovarian tumors can be overcome by disrupting critical components of tolerogenic pathways through genetic manipulation of T cells. To test this hypothesis, we proposed to develop a murine model for ovarian cancer that will allow, for the first time, precise monitoring of the functional responses of naïve, tumor-specific CD4+ and CD8+ T cell clones to ovarian tumors. Multiple properties of tumor-reactive T cells will be assessed *in vivo*, including their localization, activation, anergic status, proliferation and apoptosis. Differential responses and anti-tumor activities of the CD4+ and CD8+ T cell subsets will be investigated. Finally, the model will be used to evaluate the functional responses of tumor-specific CD4+ and CD8+ T cells that are genetically pre-disposed to autoimmune activity. The first tolerogenic pathway tested will be that involving the Cbl-b gene, as T cells lacking Cbl-b have a greatly reduced requirement for CD28 co-stimulation and demonstrate hyperactivity *in vivo* with profound autoimmune sequelae. The specific aims of this proposal are:

- Aim 1. To generate an ovarian tumor cell line that is recognized by antigen-specific CD4+ and CD8+ T cell clones from TCR transgenic mice.
- Aim 2. To define the mechanisms by which ID8 ovarian tumors evade rejection by tumor-specific CD4+ and CD8+ T cells.
- Aim 3. To determine whether tumor-specific CD4+ and CD8+ T cells lacking the Cbl-b gene show enhanced functional responses to ovarian tumors.

Body:

Aim 1: To generate an ovarian tumor cell line that is recognized by antigen-specific CD4+ and CD8+ T cell clones from TCR transgenic mice.

As described in the previous progress report, we had to generate a more aggressive ovarian tumor cell line for our experiments, as we encountered problems with spontaneous rejection of the original ID8 cell line when it was made to express epitope-tagged *neu* (*neu*^{OT-I/OT-II}). We generated a more aggressive subclone by serial transplantation of ID8 cells in syngeneic host mice. The new subclone (ID8-G7) induces tumors in just 30-40 days, as opposed to the 120-day latency of the original ID8 cell line. We have successfully transfected the new subclone with the *neu*^{OT-I/OT-II} construct under the control of the β -actin promoter (ID8-NOO) and achieved stable expression (Figure 1). However, the transfected cells are still rejected by wild-type C57/BL6 hosts due to expression of the highly immunogenic OT-I and OT-II epitopes. Fortunately, for another DOD-funded project, we created transgenic mice that express *neu*^{OT-I/OT-II} in mammary epithelium under the control of the MMTV promoter (MMTV/NTOO mice). These mice are tolerant to *neu*^{OT-I/OT-II} as it is essentially a self protein. MMTV/NTOO mice do not

develop mammary tumours until >12 months of age, therefore they can serve as hosts for ID8-G7-induced ovarian tumours at younger ages. Indeed, when MMTV/NTOO mice are injected i.p. with the ID8-G7/*neu*^{OT-I/OT-II} cell line, disseminated ovarian cancer forms within one month in 100% of cases and is associated with extensive ascites. Thus, we now have a fully operational ovarian cancer model involving epitope-tagged *neu*.

Aim 2: To define the mechanisms by which ID8 ovarian tumors evade rejection by tumor-specific CD4+ and CD8+ T cells.

We obtained from other labs CD8+ and CD4+ T cells expressing TCR transgenes specific for the OT-I and OT-II epitopes on *neu*^{OT-I/OT-II}, respectively. To evaluate the CD8+ OT-I T cell response to ovarian tumors in our model, we adoptively transferred 1×10^7 million lymphocytes from TCR transgenic OT-I donor mice into MMTV/NTOO transgenic mice bearing established ovarian tumors. Control mice received a similar dose of CD8+ T cells expressing an irrelevant TCR (obtained from P14 donor mice). The OT-I T cells (but not the negative control P14 T cells) began to proliferate within 3 days of adoptive transfer and subsequently underwent a massive proliferative response such that by Day 9 they constituted as much as 65% of the circulating CD8+ T cell population in blood and 96% of CD8+ T cells in ascites (Figure 2 and data not shown). Whereas all control mice (6/6) receiving P14 T cells had to be euthanized by day 10 after adoptive transfer due to progressive tumour growth, 75% of mice (15/20) receiving OT-I T cells achieved complete tumor regression and fully recovered by day 10 post-adoptive transfer. Recurrences have been observed in all animals, typically commencing ~30 days after adoptive transfer, a period in which OT-I T cells are still abundant in peripheral blood. Some recurrent tumours no longer express *neu*^{OT-I/OT-II}, whereas others remain positive for *neu*^{OT-I/OT-II} and MHC Class I, suggesting ID8 tumours can escape CD8+ T cells by more than one means.

To investigate how large an OT-I T cell population was required to induce tumour regression, we reduced the dose of OT-I cells to 6×10^6 or 1×10^6 . Donor T cells still underwent activation, proliferation and infiltration of the peritoneal cavity and tumor site (Figure 3). However, tumor regressions were never observed at these lower T cell doses (Table 1). In principle, if 6×10^6 T cells were to undergo only 1 additional round of proliferation, they could theoretically reach equivalent numbers as achieved with the 1×10^7 dose. This lead us to question what factors may limit the expansion or persistence of CD8+ T cells in the ovarian tumour microenvironment.

In this regard, we observed that activated OT-I T cells express CD25 (the alpha subunit of the IL-2 receptor) when they infiltrate ovarian tumors, yet are CD25-negative in blood, lymph node and ascites (Figure 4 and data not shown). This suggests that OT-I cells may undergo local, IL-2-induced proliferation within the tumour bed, which may be an important determinant of the final population size. Indeed, OT-I T cells rendered genetically deficient for the IL-2 receptor beta subunit show reduced numbers in ascites after adoptive transfer into tumour-bearing hosts (Figure 5). Similar results were seen with OT-I cells rendered genetically deficient for the IL-2 receptor alpha chain (not shown). This indicates IL-2 signaling plays an important role in sustaining antigen-specific T cell proliferation and tumor cell killing.

Finally, we investigated whether the massive proliferative response demonstrated by OT-I cells in response to ovarian tumours required antigen-expression by the tumour, or whether it was driven non-specifically by cytokines in the tumour microenvironment. OT-I cells were co-transferred with an equal number of CD8+ P14 cells, which express a T cell receptor specific for an antigen not present in the tumour (gp33 from LCMV). P14 cells were activated by subcutaneous immunization with cognate antigen (gp33 peptide). Although they mounted a normal proliferative response in blood (Fig. 6A), P14 cells failed to achieve the exceptionally high numbers seen with OT-I cells in blood, ascites or solid tumour nodules (Fig. 6A, B). Thus, the massive proliferation and accumulation of OT-I cells is not only triggered by antigen in

draining lymph nodes, but requires re-encounter with antigen at the tumour site as well. This suggests that effective eradication of ovarian tumours depends on antigen- and IL-2-driven CD8+ T cell proliferation at the tumour site.

Aim 3: To determine whether tumor-specific CD4+ and CD8+ T cells lacking the Cbl-b gene show enhanced functional responses to ovarian tumors.

As described in previous progress reports, the Cbl-b ^{-/-} mice we received from Dr. Josef Penninger's lab were not on a pure B6 background. Therefore, we have had to backcross the mice onto the B6 background. We completed 10 generations of backcrossing to obtain Cbl-b deficient OT-I cells on a pure C57Bl6 background. To date, we have performed one experiment in which these cells (1×10^7) were adoptively transferred into wild type mice bearing ID8-G7 ovarian tumours. Compared to wild-type OT-I cells, Cbl-b-deficient OT-I cells mounted a 25-50% greater proliferative response in peripheral blood, which was followed by tumour regression in all cases. This supports our hypothesis that CD8+ T cells lacking the Cbl-b gene show enhanced functional responses to ovarian tumors. Current experiments are determining Cbl-b-deficient OT-I cells outperform wild type OT-I cells when infused at limiting doses (i.e., 1×10^6 cells). In the near future, we will test the functional activity of Cbl-b-deficient CD4+ OT-II cells in our model.

Key Research Accomplishments:

The following items have been completed or are underway:

Task 1. To generate an ovarian tumor cell line that is recognized by antigen-specific CD4+ and CD8+ T cell clones from TCR transgenic mice (July 2003-Dec 2003).

- a. Evaluate signaling and transforming properties of epitope-tagged and untagged version of *neu* in cell lines; if problems noted, modify epitopes as needed. ***completed**
- b. Generate ID8 cell subclones that stably express *neu*^{OT-I/OT-II}; perform in vitro assays to evaluate recognition of OT-I and OT-II epitopes by CD4+ and CD8+ T cells from TCR-transgenic mice. ***completed**
- c. Inject ID8/*neu*^{OT-I/OT-II} cells intraperitoneally into MMTV/NTOO transgenic mice to establish ovarian cancer (July 2003-Dec2003). ***completed**

Task 2. To define the mechanisms by which ID8 ovarian tumors evade rejection by tumor-specific CD4+ and CD8+ T cells (Jan 2004-July 2005).

- a. Generate sufficient numbers of mice bearing tumors expressing *neu*^{OT-I/OT-II}. ***completed**
- b. Perform immunological studies of adoptively transferred OT-I- and OT-II-specific T cells and control T cells in mice bearing ovarian tumors expressing *neu*^{OT-I/OT-II}, as per Aim 2. ***completed**

Task 3. To determine whether tumor-specific CD4+ and CD8+ T cells lacking the Cbl-b gene show enhanced functional responses to ovarian tumors (July 2005-June 2006).

- a. Breed OT-I, OT-II, TEa and P14 TCR transgenes onto the Cbl-b background (Months 1-12). ***completed**

- b. Generate sufficient numbers of mice bearing tumors expressing *neu*^{OT-I/OT-II} (Months 1-12). ***completed**
- c. Perform immunological studies of adoptively transferred Cbl-b-deficient OT-I- and OT-II-specific T cells and control T cells in mice bearing ovarian tumors expressing *neu*^{OT-I/OT-II} (Months 4-12). ***in progress**

Reportable Outcomes:

Peer-reviewed manuscripts:

Regression of Advanced Ovarian Cancers after Adoptive Transfer of CD8+ T Cells in a Novel Murine Model. Taimei Yang, Erika M. Wall, Katy Milne, Pathy Theiss and **Brad H. Nelson**. *In preparation*.

Conference proceedings:

Proliferation and Differentiation of CD8+ T Cells in the Absence of IL-2/IL-15 Receptor beta Chain Expression or STAT5 Activation. Ryan M. Teague, Richard M. Tempero, Sunil Thomas, Murali-Krishna Kaja and **Brad H. Nelson**. Annual Meeting of the American Association for Cancer Research, Orlando, March 2004.

Proliferation and Differentiation of CD8+ T Cells in the Absence of IL-2/IL-15 Receptor beta Chain Expression or STAT5 Activation. Ryan M. Teague, Richard M. Tempero, Sunil Thomas, Murali-Krishna Kaja and **Brad H. Nelson**. 12th International Congress of Immunology and 4th Annual Conference of the Federation of Clinical Immunology Societies (FOCIS), Montreal, July 2004, Publication Number: (76PM) W8.29

Mapping and Manipulating the Immune Response to Ovarian Cancer. **Brad H. Nelson**, Brad C. Stone and Cassian Yee. Second Canadian Conference on Ovarian Cancer Research, Ottawa, May 2004.

Monitoring the T cell response to spontaneous mammary tumors using a novel transgenic mouse model. Erika M. Wall, Katy Milne, **Brad H. Nelson**. Annual Meeting of the American Association for Cancer Research, Anaheim, April 2005.

Monitoring the T cell response to spontaneous mammary tumors using a novel transgenic mouse model. Erika M. Wall, Katy Milne, **Brad H. Nelson**. Canadian Society for Immunology Annual Meeting, Whistler BC, April 2005.

Regression of Advanced Ovarian Cancers After Adoptive Transfer of CD8+ T Cells in a Novel Murine Model. Taimei Yang, Erika M. Wall, Katy Milne and **Brad H. Nelson**. NCI/SPORE Annual Meeting, Washington DC July 2005.

Evaluation of the T Cell Response to Mammary Tumours Using a Novel Transgenic Mouse Model. Elaine K. Wong, Erika M. Wall, Katy Milne and **Brad H. Nelson**. Translational Research in Radiation Oncology Symposium, San Francisco, CA, August 2005.

Invited presentations:

Mapping and Manipulating the Immune Response to Ovarian Cancer. **Brad H. Nelson**, Brad C. Stone and Cassian Yee. Second Canadian Conference on Ovarian Cancer Research, Ottawa, May 2004.

Mapping and Manipulating the Immune Response to Cancer. **Brad H. Nelson**, Canada's Michael Smith Genome Sciences Centre, Vancouver, Sept. 2004.

The Immune Response to Ovarian Cancer. **Brad H. Nelson**. National Ovarian Cancer Association, Educational Session, Victoria BC, Nov. 20, 2004.

The Immune Response to Ovarian Cancer. **Brad H. Nelson**. BC Cancer Agency's Annual Meeting, Pharmacy Session, Vancouver, Nov. 2004.

Mapping and Manipulating the Immune Response to Cancer. **Brad H. Nelson**. University of Victoria, Feb. 4 2005.

The Immune Response to Ovarian Cancer. **Brad H. Nelson**. OBGYN Grand Rounds, Vancouver General Hospital, Feb.16 2005.

Mapping and Manipulating the Immune Response to Cancer. **Brad H. Nelson**. Canadian Association of Medical Oncologists Annual Meeting, Montreal, March 2005.

Tracking and Manipulating the Immune Response to Cancer. **Brad H. Nelson**. McMaster University, Hamilton Ontario, Oct. 2005.

Career advancement:

The PI, Brad Nelson, has been appointed Director of the Research Laboratories for the British Columbia Cancer Agency's Vancouver Island Centre (Victoria, BC). His work in this animal tumor model was integral to his success in this competition. He moved there on July 1, 2003. (The DOD has already been informed of this move.)

Conclusions:

We have created a novel mouse model for ovarian cancer that allows precise analysis of CD4+ and CD8+ T cell responses in the tumour microenvironment. After adoptive transfer, CD8+ OT-I T cells undergo massive proliferation in response to ovarian tumours, and are able to induce tumor regression in a dose-dependent manner. This response is dependent on both IL-2 signaling and re-encounter with antigen at the tumour site. Genetic deletion of the negative regulator Cbl-b from OT-I cells leads to an enhanced proliferative response, which is followed by tumour regression. Current studies are investigating whether Cbl-b-deficient OT-I cells outperform wild type OT-I cells by inducing tumour regression at limiting cell doses. If so, then genetic manipulation of human CD8+ T cells to disrupt Cbl-b function may represent an attractive, novel approach to enhancing the efficacy of T cell therapy against ovarian cancer.

References:

None.

Appendices:

See accompanying Figures 1-6 and Table 1.




Figure 1. Cell surface expression of *neu*^{OT-I/OT-II}, MHC Class I and MHC Class II on the ovarian tumour cell line ID8-G7.




Figure 2. Expansion of adoptively transferred CD8+ OT-I T cells in peripheral blood in response to *neu*OT-I/OT-II expressing ovarian tumours, and correlation to tumor regression as assessed by ascites scores. Note that tumours recur after about 30 days despite the continued presence of OT-I cells.







Figure 3. Percentage of tumour-infiltrating OT-I T cells relative to total CD8+ T cells after adoptive transfer of 1×10^6 versus 1×10^7 lymphocytes from OT-I transgenic mice into ovarian tumor-bearing hosts.






Isotype control —



CD25 staining —



Ascites



Solid tumor

Figure 4. CD25 expression on OT-I donor T cells in ascites and solid tumor after adoptive transfer into ovarian tumour-bearing mice.




Figure 5. CD8+ OT-I T cells deficient in the IL-2 receptor beta subunit (an essential component of the IL-2 and IL-15 receptors) show reduced accumulation in the ascites of ovarian tumour-bearing mice (Day 7 after adoptive transfer).



A**B**

Figure 6. The massive proliferative response demonstrated by OT-I cells in response to ovarian tumours is antigen dependent. OT-I cells were co-transferred with an equal number of CD8+ P14 cells, which express a T cell receptor specific for an antigen not present in the tumour (gp33 from LCMV). P14 cells were activated by immunization with cognate antigen (gp33 peptide). Although they proliferated in blood (panel A), they failed to achieve the high numbers seen with OT-I cells in blood, ascites or tumour nodules (panels A and B).

Dose of CD8+ T cells	Regression	Non-regression
10 x 10 ⁶ OT-I	15	5
6 x 10 ⁶ OT-I	0	6
1 x 10 ⁶ OT-I	0	19
10 x 10 ⁶ P14 (irrelevant antigen receptor)	0	6

Table1. Dose-dependent regression of advanced ovarian cancers by adoptively transferred CD8+ T cells.