

AD _____

Award Number: W81XWH-04-1-0540

TITLE: Role of Notch/VEGF-Receptor 3 in Breast Tumor Angiogenesis and Lymphangiogenesis

PRINCIPAL INVESTIGATOR: Jan K. Kitajewski, Ph.D.

CONTRACTING ORGANIZATION: Columbia University
New York, NY 10032

REPORT DATE: May 2006

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 01-05-2006		2. REPORT TYPE Annual		3. DATES COVERED 1 May 2005 – 30 Apr 2006	
4. TITLE AND SUBTITLE Role of Notch/VEGF-Receptor 3 in Breast Tumor Angiogenesis and Lymphangiogenesis				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-04-1-0540	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Jan K. Kitajewski, Ph.D.				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Columbia University New York, NY 10032				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES Original contains colored plates: ALL DTIC reproductions will be in black and white.					
14. ABSTRACT The overall objective is to define the interaction between Notch and VEGFR-3 signaling in breast cancer. We are examining a role for Notch in breast tumor vessels and attempting to block Notch and VEGFR-3 activity in breast tumors grown in mice. We proposed two aims: 1) studies of Notch/Dll4 function in murine mammary tumorigenesis and 2) studies of the inhibitory effects of a Notch antagonist (Notch decoy) in a murine mammary tumor model. In aim 1, to study the role for notch in murine mammary tumorigenesis, progress has been made in developing two new transgenic lines that will allow for conditional activation or inactivation of Notch specifically within the endothelium. The first mouse line, EF1-Notch1IC can be manipulated in a conditional fashion, as demonstrated by lethality if activated in embryonic endothelium. The second mouse line, EF1-Notch1ECD/Fc, has been generated and is being further tested. We have carried out experiments to demonstrate that breast tumor xenograft growth is inhibited by the Notch decoy, an antagonist made up of the Notch1 extracellular domain fused to the Fc protein, Notch1ECD/Fc. This block appears to be a result of reduced tumor angiogenesis. This strategy, now shown to inhibit mammary tumor growth in our mouse models, is proving to be a promising area for therapeutic intervention in breast cancer.					
15. SUBJECT TERMS Notch, VEGF, Breast Cancer, Angiogenesis					
16. SECURITY CLASSIFICATION OF:			UU	18. NUMBER OF PAGES 11	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

TABLE OF CONTENTS

Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	10
Reportable Outcomes.....	10
Conclusions.....	10
References.....	11
Appendices.....	-

INTRODUCTION

Expression studies have shown that the angiogenic/lymphangiogenic factor VEGFR-3, its ligand VEGF-C and the Notch ligand, Dll4 are up-regulated in the invading blood and lymphatic vessels in human breast cancer (1, 2). Using mouse models, Notch signaling has been found to be essential for angiogenesis to progress to completion in the developing embryo (3). However, the exact mechanism(s) by which Notch signaling regulates angiogenesis and/or lymphangiogenesis is not well understood. It is known that Notch modulates cell-fate decisions by regulating the expression of tissue specific genes. As many angiogenic regulators have been identified, we have focused on defining the interactions between Notch and that of known angiogenic regulators. Using quantitative RT-PCR analysis, we found that VEGFR-3 was induced in response to an activated form of Notch4 (Notch4/int-3) in three human primary endothelial cells (HUVEC-umbilical vein, HUAEC-umbilical artery, and HMVEC-dermal microvascular). Taken together these data suggested a relationship between Notch signaling and VEGFR-3. Thus, we **hypothesized** that Notch may regulate blood and/or lymphatic vessel development via its induction of VEGFR-3 during physiological angiogenesis, as well as pathological angiogenesis and/or lymphangiogenesis in breast cancer. The **overall objective** of this proposal is to define the interaction between Notch and VEGFR-3 signaling in breast cancer.

Our original study design was as follows: We will investigate the relationship between Notch and VEGFR-3 signaling in gain-of-function and loss-of-function mouse models. We have developed two mouse models in which Notch signaling can be specifically induced in the vascular endothelium. Murine mammary tumor cells will be implanted into these mice and the affect(s) of Notch signaling on tumor angiogenesis/lymphangiogenesis determined. Expression of Dll4 and VEGFR-3 and its ligands, VEGF-C and VEGF-D will also be monitored. We will also determine if blocking endogenous Notch signaling will perturb breast tumor angiogenesis/lymphangiogenesis. We have generated Notch antagonists, the Notch decoys, which encode the extracellular domains of the four Notch proteins but not the signaling domain. Human mammary tumor cell lines will be generated that express the different Notch antagonists and xenografted into immuno-compromised mice. The affects of the Notch1 antagonist on tumor angiogenesis and/or lymphangiogenesis will be examined, as well as the affects on Dll4 and VEGFR-3 expression.

In year two we made progress on three of our designated tasks. In **Task 1** (Months 1-3) we set out to optimize conditions for tamoxifen treatment of mice to conditionally activate Notch. We have made progress in establishing that Notch1C can be conditionally activated using our altered strategy to develop a new transgenic Notch1C line and a transgenic Notch decoy (inhibitor) line and our progress will be described. In **Task 2** (Months 4-24) we set out to induce Notch activity in adult mice and determine the consequences for breast tumor cell growth. The mice needed for this task, EF1-Notch1C and EF1-Notch1ECD/Fc are now generated and validated. In **Task 4** (Months 6-36) analysis of VEGFR-3 expression in tumors expressing Notch antagonists. An initial goal in this task was to develop breast cancer cell lines over-expressing Notch1C or Notch decoy in order to manipulate Notch activity in the cancer cell. Ultimately, we wanted to create line that secrete decoy to assess the affects on tumor cell/endothelial interaction. What we found was that Notch activity affected the cancer cell. Activation of Notch suppressed breast cancer cell growth and expression of Notch decoy promoted growth. This result was unexpected and will be discussed in the body, below. However, we now have worked with a mouse mammary tumor line, Mm5MT, which is not affected by the Notch decoy when analyzed in vitro. When analyzing the consequences of Notch decoy expression in xenografted breast cancer cells, we found that tumor growth was indeed inhibited by the Notch decoy.

BODY

Specific Aims 1: Notch and Dll4 function in murine mammary tumorigenesis.

Task 1 (Months 1-3) Optimizing conditions for TM treatment of N1ICD and Dll4 double transgenic mice

Experiments using both gain-of-function and loss-of-function mouse breast tumor models will aid our understanding of the role of Notch in tumor angiogenesis. The receptors for Dll4, Notch1 and Notch4 are endogenously expressed in endothelial cells (4, 5). Notch4 activation in embryonic endothelium causes vascular remodeling defects in murine embryos (6). Notch4 nullizygous mice are viable and fertile (7). However, we noticed that the adults have vascular patterning defects. Notch4 is primarily expressed within the vascular endothelium during development and this expression is maintained in the adult (8). In our original proposal, we pursued a system to conditionally express N1ICD, an activated form of Notch1, in the adult vasculature. We have generated mice doubly transgenic for a Flk1-Cre-ERT transgene and an N1ICD CAG-CAT transgene in the C57BL/6J background. The Flk1-Cre-ERT transgene encodes the Flk1 (VEGFR-2) regulatory region upstream of a Cre-estrogen receptor fusion gene. In the adult mouse, Flk1 expression is restricted the vascular endothelium. Thus, this transgene limits the expression of the Cre-ERT fusion protein to the vascular endothelial


cells. The Cre-ERT fusion protein has a tamoxifen(TM)-responsive estrogen receptor binding component and is latent until TM is administered. The CAG-CAT transgene contains a non-functional gene (CAT) with its associated stop codon, flanked by lox P sites followed by either N1ICD or Dll4 gene. In the absence of Cre expression, the CAT gene is expressed. When the CAT gene is excised by Cre, the N1ICD or Dll4 gene is placed behind the CAG promoter and expressed. The CAG promoter consists of a CMV enhancer-chicken β -actin hybrid protomoter that is strongly expressed in endothelial cells. Thus, when treated with tamoxifen these double transgenic mice activate Notch signaling specifically within the vascular endothelium (data not shown).

New conditional mouse models to regulate Notch (EF-1-flox)

As discussed above, we have successfully developed one transgenic line to conditionally regulate Notch1C expression (CAG-CAT-Notch1C). To enhance our chances of success in conditionally manipulating Notch in

tumor endothelium, we have pursued another model for conditional Notch regulation, which we will refer to as EF1-flox-Notch (Figure 1A, 1B). In this new strategy, we also include a mouse line designed to inactivate Notch signaling in tumor endothelium that incorporates the Notch decoy. Although these approaches were not originally proposed, we feel that inclusion enhances our ability to understand Notch function in tumor endothelium by either activating or inactivating Notch in tumor vessels.

The EF1-flox-Notch constructs use the strong promoter from the eukaryotic Elongation Factor 1 (EF1) gene to drive expression (Figure 1A). Using homologous recombination, we have inserted a flox-stop-Notch1C construct and a flox-stop-Notch1ECDFc (Notch decoy) construct into the EF1 locus (Figure 1B). The proper targeting of ES cells has been confirmed and a mouse has already been generated for expression of Notch1C (EF1-flox-stop-Notch1C). This will be used in the same fashion as CAG-CAT-Notch1C. To date, we have crossed EF1-flox-stop-Notch1C mice with Rosa-cre mice and this leads to embryonic lethality, as expected. The mice have also been crossed to the endothelial specific Flk1-cre mouse and this also leads to embryonic lethality. Thus, abnormal activation of Notch, either globally or in an endothelial specific fashion, is not compatible with proper embryonic development.



In order to further test the EF1-Notch1IC mice, we crossed to a mouse line that expresses cre recombinase in vascular smooth muscle cells, SM22cre. This cross did not produce viable doubly transgenic offspring, SM22cre; EF1-N1IC (Figure 2). Thus, expression of N1IC via either an endothelial- or vascular smooth muscle-specific cre leads to lethality, validating the use of this conditional line for the study of Notch in tumor vasculature.

Figure 2. Result of cross of SM22 cre-recombinase mice and EF1-Notch1IC

Cross: SM22 ^{Cre/+} x EF1 ^{N1IC/+}				
	WT ; EF1 ^{N1IC/+}	SM22 ^{Cre/+} ; WT	WT ; WT	SM22 ^{Cre/+} ; EF1 ^{N1IC/+}
Live births				
Expected	25%	25%	25%	25%
Actual	31%	19%	50%	0%
	n = 11/36	n = 7/36	n = 18/36	n = 0/36

Generation of EF-Notch Decoy (Notch1ECD^ΔEc) mice.

The ES cells with the EF1-flox-stop Notch1ECD^ΔEc have been used to generate mouse lines. We now have two chimeric mice lines that are being bred to develop a line with EF1-flox-stop Notch decoy.


Specific Aims 2: Analysis of Notch antagonists in a murine mammary tumor model


Task 4 (Months 6-36) Analysis of VEGFR-3 expression in tumors expressing Notch antagonist

In previous studies of murine xenografts of human mammary tumor cells, Dll4 expression was found to be enhanced within the tumor vessels (2). Moreover, ectopic expression of the VEGFR-3 ligand, VEGF-C, in mammary xenografts induced lymphangiogenesis and nodal spread of the breast cancer cell (9, 10). Taken together with the observation that Notch signaling induces VEGFR-3 expression in cultured endothelial cells, we want to test whether blocking Notch activity suppressed tumor growth and pathological angiogenesis/lymphangiogenesis. We have generated Notch antagonists that are composed of the signal peptide and EGF-like repeats of Notch1 and Notch2 fused in frame with Fc fragment of human IgG. Using an *in vitro* co-culture assay, we have found that the Notch antagonists N1ECDFc and N2ECDFc perturb ligand-activated signaling of Notch1, Notch2 and Notch4 (data not shown). Thus, we proposed to generate mammary tumor cell lines expressing these Notch antagonists and transplant them into immunocompromised mice.

Analysis of VEGFR-3 expression in tumors expressing Notch antagonists. In previous studies of murine xenografts of human mammary tumor cells, Dll4 expression was found to be enhanced within the tumor vessels. Delta4, an endothelial specific notch ligand expressed at sites of physiological and tumor angiogenesis (2). Moreover, ectopic expression of the VEGFR-3 ligand, VEGF-C, in mammary xenografts induced lymphangiogenesis and nodal spread of the breast cancer cells (9). VEGF-C-mediated lymphangiogenesis promoted tumor metastasis and VEGF-C induced lymphangiogenesis is associated with lymph node metastasis in orthotopic MCF-7 tumors (10). Taken together with the observation that Notch signaling induces VEGFR-3 expression in cultured endothelial cells, we want to test whether blocking Notch activity suppressed tumor growth and pathological angiogenesis/lymphangiogenesis. We have generated Notch antagonists that are composed of the signal peptide and EGF-like repeats of Notch1 and Notch2 fused in frame with Fc fragment of human IgG. Using an *in vitro* co-culture assay, we have found that the Notch antagonists N1ECDFc and N2ECDFc perturb ligand-activated signaling of Notch1, Notch2 and Notch4 (data not shown). Thus, we proposed to generate mammary tumor cells expressing these Notch antagonists and transplant them into mice.


Our expression studies of human mammary tissue demonstrated that Notch1, Notch4 and Dll4 were expressed in the normal and malignant ductal epithelial cells. This suggested to us that Notch signaling functions in normal as well as malignant ductal epithelium. Therefore, we decided to determine the affects of Notch signal activation on growth and tumorigenicity of human MDA-MB-231 breast cancer cell line and FGF expressing mouse Mm5MT mammary tumor cell line. MDA-MB-231 and FGF expressing Mm5MT cells were retrovirally transduced with empty virus or viruses expressing activated forms of Notch1 (N1IC) or Notch4 (N4/int-3). Cell populations were selected in hygromycin. Consistent with Notch signaling being activated, cell populations expressing N1IC or N4/int-3 transactivated luciferase reporters encoding 6 Notch/CSL binding sites relative to the mock infected controls (Figure 3A). Next, we determined the ability of the mock and Notch expressing mammary tumor populations to form colonies in soft agar (Figure 3B). Notch signal activation inhibited adhesion independent growth of the transformed mammary tumor lines. In MDA-MB-231, Notch1 signaling suppressed cell growth greater than 20 fold relative to the mock-infected cells (Figure 3C). The inhibition of cell growth was also observed in monolayer cultures and thus, they could not be subcutaneously transplanted in immunocompromised mice.





Next, we determined the affect of expressing the Notch antagonist, N1ECDFc on the tumorigenicity of mammary tumor cells. FGF-expressing Mm5MT cells were retrovirally infected with empty virus, Fc (control) or N1ECDFc expressing viruses and cell populations generated by hygromycin selection. In soft agar assays, expression of the Notch antagonists had a nominal affect on soft agar growth as determined by WST-8 assay (Figure 4). 1×10^6 mock, Fc, or N1ECDFc expressing cells were injected subcutaneously into 10 C3H syngeneic mice each. Three independent trials were monitored for tumor growth for 3 weeks (Figure 5). In the three

trials, expression of N1ECDFc significantly inhibited tumor growth. This affect of the Notch1 antagonist may be on either the tumor epithelial cells, tumor angiogenesis or both. To discriminate between these possibilities, immunohistochemistry will be performed to analyze tumor angiogenesis, lymphangiogenesis, epithelial apoptosis and epithelial proliferation.






Figure 6 Expression of blood and lymphatic endothelial cell markers in micropapillary breast carcinomas. A) CD34 expression in the blood endothelium. Expression of the lymphatic markers B) LYVE-1 and C) podoplanin.

Notch1 and Notch4 are co-expressed in the breast cancer lymphatic endothelium.




Figure 7 Notch1 and Notch4 are expressed in the lymphatics of micropapillary breast carcinomas. A, B) LYVE-1, C) Notch1, D) Notch4 double immunohistochemistry of breast tumor tissue. E) Notch1 and LYVE-1 co-expression in the lymphatic vessel. F) Notch4 and LYVE-1 co-expression in the lymphatic vessel. G, H) Magnified view of lymphatic vessels from panels E and F, respectively.

We observed that Notch1, Notch4, Dll4 and VEGFR-3 were all expressed in the vasculature of human breast carcinoma (data not shown). We also observed that Jagged1 is expressed with the vasculature of normal breast tissue (data not shown). Whether these proteins are expressed in both the blood and lymphatic endothelium remains to be determined. To discriminate between the blood and lymphatic vessels, we performed immunohistochemistry on human micropapillary breast carcinomas using antibodies against CD34, as a marker for blood endothelium and LYVE-1 and podoplanin as markers for lymphatic endothelium (Fig. 6).

We then performed double immunohistochemistry with antibodies against LYVE-1 and Notch1 or Notch4 (Fig. 7). We found that both Notch1 and Notch4 are expressed in the lymphatic endothelium of human micropapillary breast carcinomas. Notch1 and




Figure 8 Notch1 is activated in a lymphatic endothelium of micropapillary breast carcinomas. A) Notch1 Val, and B) Podoplanin double immunohistochemistry of breast tumor tissue. C) DAPI and Notch1 Val image demonstrating perinuclear and nuclear expression of activated Notch1. D) Triple stain, Notch1 Val, podoplanin and DAPI.

Notch4 were co-expressed with LYVE-1 in the lymphatic endothelial cell bodies. Interestingly, Notch1 and Notch4 proteins were also detected in the nuclei of the lymphatic endothelial cells suggesting that Notch signaling was activated in these cells (Fig. 7 G & H). To further validate this observation, we co-immunostained the micropapillary breast carcinoma tissue with Notch1Val, an antibody that only recognizes the activated Notch1 peptide and the lymphatic marker, podoplanin (Fig. 8). Activated Notch1 peptide was detected in a subset of the tumor lymphatic endothelium.

KEY RESEARCH ACCOMPLISHMENTS

- Developed a EF-1-flox-Notch1C mouse line that expresses an activated form of Notch1 within the murine vasculature when crossed with either an endothelial specific driven Cre or a vascular smooth muscle-specific cre.
- Generated EF-1-flox-Notch decoy mice for conditional loss-of-function Notch mice.
- Found that the expression of activated Notch constructs or Notch antagonists altered the growth of murine and human mammary tumor cell lines in soft agar.
- Found that the expression of activated Notch constructs inhibited the growth of murine and human mammary tumor cell lines in soft agar.
- Demonstrated that the Notch1 antagonist inhibited tumor growth of subcutaneous xenografts of mouse mammary tumor cells.
- Demonstrated that Notch1 and Notch4 are expressed and most likely active in the lymphatic endothelium of micropapillary breast carcinomas.

REPORTABLE OUTCOMES (PUBLICATIONS/ABSTRACTS)

Publications:

Vorontchikhina, M.A., Zimmerman, R.C., Shawber, C.J., Tang, H., and **Kitajewski, J.** (2005) Unique patterns of Notch1, Notch4 and Jagged1 expression in ovarian vessels during folliculogenesis and corpus luteum formation. *Gene Expression Patterns*. (5 : 701-9)

Zeng Q, Li S, Chepeha DB, Giordano TJ, Li J, Zhang H, Polverini PJ, Nor J, **Kitajewski J**, Wang CY. (2005) Crosstalk between tumor and endothelial cells promotes tumor angiogenesis by MAPK activation of Notch signaling. *Cancer Cell* **8**:13-23.

Abstract:

Shawber, C.J., Funahashi, Y., Francisco, E., Podgrabinska, S., Kitamura, Y., Vorontchikhina, M., Shiraishi, K., Chawengsaksohak, K., Rossant, J., Accili, D., Skobe, M., and Kitajewski, J. (2006) *VEGFR-3, a direct endothelial cell specific target of Notch signaling in vitro and in vivo*. Molecular Mechanisms in Lymphatic Function and Disease. Gordon Research Conference. Les Diablerets, Switzerland

CONCLUSIONS

The proposal **objective** is to define the interaction between Notch and VEGFR-3 in breast cancer. To study the role for notch in murine mammary tumorigenesis, progress has been made in developing two new transgenic lines that will allow for conditional activation or inactivation of Notch specifically within the endothelium. These will be used to manipulate Notch activity in tumor endothelium of mice. angiogenesis. The first mouse line, EF1-Notch1C can be manipulated in a conditional fashion, as demonstrated by lethality if activated in embryonic endothelium. The second mouse line, EF1-Notch1ECD/Fc, has been generated and is being further tested. We have carried out experiments to demonstrate that breast tumor xenograft growth is inhibited by the Notch decoy, an antagonist made up of the Notch1 extracellular domain fused to the Fc protein, Notch1ECD/Fc. This block appears to be a result of reduced tumor angiogenesis. This strategy, now shown to inhibit mammary tumor growth in our mouse models, is proving to be a promising area for therapeutic intervention in breast cancer.

REFERENCES

1. Valtola, R., Salven, P., Heikkila, P., Taipale, J., Joensuu, H., Rehn, M., Pihlajaniemi, T., Weich, H., deWaal, R., and Alitalo, K. 1999. VEGFR-3 and its ligand VEGF-C are associated with angiogenesis in breast cancer. *Am J Path* 154:1381-1390.
2. Mailhos, C., Modlich, U., Lewis, J., Harris, A., Bicknell, R., and Ish-Horowicz, D. 2001. Delta4, an endothelial specific Notch ligand expressed at sites of physiological and tumor angiogenesis. *Differentiation* 69:135-144.
3. Shawber, C., Kandel, J.J., and Kitajewski, J. 2004. Notch: cell fate determination from vascular development to human vasculopathy. *Drug Discovery Today: Disease Models* 1:351-358.
4. Shawber, C.J., Das, I., Francisco, E., and Kitajewski, J. 2003. Notch signaling in primary endothelial cells. *Ann N Y Acad Sci* 995:1-9.
5. Shawber, C., and Kitajewski, J. 2004. Notch Signaling in the Vasculature; Insights from Zebrafish, Mouse, and Man. *BioEssays* in press.
6. Uyttendaele, H., Ho, J., Rossant, J., and Kitajewski, J. 2001. Vascular patterning defects associated with expression of activated Notch4 in embryonic endothelium. *PNAS* 98:5643-5648.
7. Krebs, L.T., Xue, Y., Norton, C.R., Shutter, J.R., Maguire, M., Sundberg, J.P., Gallahan, D., Closson, V., Kitajewski, J., Callahan, R., et al. 2000. Notch signaling is essential for vascular morphogenesis in mice. *Genes and Development* 14:1343-1352.
8. Uyttendaele, H., Marazzi, G., Wu, G., Yan, Q., Sassoon, D., and Kitajewski, J. 1996. Notch4/int-3, a mammary proto-oncogene, is an endothelial cell-specific mammalian Notch gene. *Development* 122:2251-2259.
9. Mandriota, S.J., Jussila, L., Jeltsch, M., Compagni, A., Baetens, D., Prevo, R., Banerji, S., Huarte, J., Montesano, R., Jackson, D.G., et al. 2001. Vascular endothelial growth factor-C-mediated lymphangiogenesis promotes tumour metastasis. *Embo J* 20:672-682.
10. Mattila, M.M., Ruohola, J.K., Karpah, T., Jackson, D.G., Alitalo, K., and Harkonen, P.L. 2002. VEGF-C induced lymphangiogenesis is associated with lymph node metastasis in orthotopic MCF-7 tumors. *Int J Cancer* 98:945-951.