

HARD X RAYS FROM SUPERNOVA 1993J

M.D. Leising¹, J.D. Kurfess², D.D. Clayton¹, D.A. Grabelsky³, J.E. Grove², W.N. Johnson², G.V. Jung⁴, R.L. Kinzer², R.A. Kroeger², W.R. Purcell³, M.S. Strickman², L.-S. The¹, and M.P. Ulmer³

ABSTRACT

The Oriented Scintillation Spectrometer Experiment (OSSE) on the *Compton Observatory* observed SN 1993J during three intervals centered approximately 12, 30, and 108 days after its outburst. Hard X-ray emission was detected in the first two of these intervals. No emission was seen in the third observation or in two earlier observations in 1991 and 1992. The coincidence of the observed excess with the outburst of SN 1993J and the consistency of the spectra and time evolution with those seen at lower energies by ROSAT and ASCA argue that the observed emission is indeed from SN 1993J. It is probably due the the interaction of the fast supernova ejecta with circumstellar material. The luminosity, 5×10^{40} erg s⁻¹ (50–150 keV) in the first interval, is significantly larger than predicted. Extrapolating the spectrum to a few keV accounts for most or all of the observed emission at low energy. The observed high temperature, 10⁹ K, is easily obtained in the shocked circumstellar matter, but a surprisingly high density is required there to give the observed luminosity and little or no additional X-ray emission from denser shocked supernova ejecta is allowed. The hard emission might also be explained in terms of the shocked supernova ejecta itself with unexpectedly high temperature.

Subject headings: X rays: general – gamma rays: observations – supernovae: individual (SN 1993J) – stars: circumstellar matter

¹Department of Physics and Astronomy, Clemson University, Clemson, SC 29634-1911

²E.O. Hulburt Center for Space Research, Naval Research Lab, Washington, DC 20375-5320

³Department of Physics and Astronomy, Northwestern University, Evanston, IL 60201

⁴Universities Space Research Association, Washington, DC 20024

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1994		2. REPORT TYPE		3. DATES COVERED 00-00-1994 to 00-00-1994	
4. TITLE AND SUBTITLE Hard X Rays from Supernova 1993J				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory,E.O. Hulbert Center for Space Research,4555 Overlook Avenue, SW,Washington,DC,20375				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 11	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

1. INTRODUCTION

The great potential of hard photon diagnostics of supernovae has been well documented. The quantity and location of radioactivity, the nature of the interaction of the fast ejecta with external material, and the presence and character of a compact remnant can all be addressed with X and γ -ray observations. Indeed some of this potential was realized from observations of SN 1987A (Sunyaev et al. 1987; Dotani et al. 1987; Matz et al. 1988). In that case the hard X-ray emission, as well as the γ -ray line emission, can be understood as arising from decay of radioactive ^{56}Co within the ejecta. Supernova 1980K was also detected in soft X rays (Canizares, Kriss, & Feigelson 1982) that were probably produced by the interaction of the ejecta with circumstellar material. Whether this emission was of thermal or nonthermal origin could not be determined.

Supernova 1993J in NGC 3031 (M81) has been extensively observed at many wavelengths and has yielded a wealth of new information about core-collapse supernovae (Wheeler & Filipenko 1994, and references therein). It is a transition object, appearing initially with Type II supernova spectral characteristics and then evolving into a Type Ib object (Filipenko, Matheson, & Ho 1993; Swartz et al. 1993). The light curve was also quite unusual (Schmidt et al. 1993). The observed characteristics are well modelled as the result of a core collapse and subsequent explosion in a red supergiant that had lost almost all of its hydrogen-rich envelope (Nomoto et al. 1993; Podsiadlowski et al. 1993; Wheeler et al. 1993; Schmidt et al. 1993; Hoffich, Langer, & Duschinger 1993; Woosley, Eastman, & Weaver 1994), probably due to interactions with a binary companion.

Soft X rays were detected from SN 1993J beginning only six days after the explosion, and subsequent X-ray observations showed a slowly declining flux, with decay timescales of a few months (Zimmermann et al. 1994; Tanaka 1993). All X-ray spectra were quite flat, with photon power-law indices near -1.0 . For thermal spectral models only lower limits on the electron temperature, near $kT_e=10$ keV, could be established. These data have been interpreted in terms of thermal bremsstrahlung emission from shock-heated electrons due to the interaction of the expanding supernova ejecta with circumstellar material, presumably the presupernova stellar wind (Suzuki et al. 1993; Chevalier & Fransson 1994). In principle, significant X-ray emission can come from both the wind material shocked to high temperatures ($\sim 10^9$ K) just ahead of the ejecta (Fransson 1984), and from the ejecta (Chevalier 1982) if the overpressure resulting from the primary shock drives a “reverse” shock into the ejecta (McKee 1974). Despite the lower temperature in the shocked ejecta, it could dominate the soft X-ray luminosity because of its higher free-free emissivity due to its higher density. Hard X-ray observations are required to assess the contribution of the shocked wind itself, and to better constrain the electron temperature, if the emission is

Table 1: SN 1993J/M81 Observation Summary

CGRO Period	Dates	Days from Explosion ¹	Accumulation Time ² (10^5 s)
216	1993 April 6–12	9.9–15.4	0.84
218	1993 April 20–May 3	23.7–36.9	1.49
227/228	1993 June 29–July 27	93.6–121.5	4.53
7	1991 Aug 9–15	...	0.80
18	1992 Jan. 10–23	...	1.80

¹Taken to be 1993 March 28.0.

²Total time of source observations with four detectors.

indeed thermal. Combined hard and soft X-ray measurements can in principle distinguish between thermal and nonthermal emission mechanisms. Early detections of radio emission from SN 1993J also imply circumstellar interactions (Pooley & Green 1993; Van Dyk et al. 1993; Phillips & Kulkarni 1993).

2. Observations and Data Analysis

The Oriented Scintillation Spectrometer Experiment (OSSE) is one of four experiments on NASA’s *Compton Gamma-Ray Observatory* satellite (Johnson et al. 1993). OSSE was designed to undertake observations of astrophysical sources in the 0.05 – 10 MeV range. It consists of four identical detector systems with independent, single-axis orientation controls that provide a range of 192 degrees. The primary detecting element for each detector is a large-area NaI(Tl) scintillation crystal (13 in. diameter by 4 in. thick) shielded in the rear by an optically coupled 3-in. thick CsI(Na) scintillation crystal in a “phoswich” configuration. Each phoswich is actively shielded by a 3.3-in. thick annulus of NaI(Tl) scintillation crystals and passively collimated by a tungsten slit collimator that defines a $3.8^\circ \times 11.4^\circ$ full width at half maximum (FWHM) γ -ray aperture. The total aperture area of the four detectors is 2620 cm², with an effective photopeak area of 2000 cm² at 511 keV.

Shortly after the discovery of SN 1993J the nature of the explosion was unclear. Mainly in view of the possibility of early leakage of ⁵⁶Ni γ rays, the *Compton Observatory* was reoriented to observe the supernova as soon as possible. A 6-day observation was followed by a full two-week observation after 8 intervening days. These two observations are the main focus of this work. Later, when the probability of detecting ⁵⁶Co γ rays was

near maximum, OSSE observed SN 1993J for four weeks. The three observation intervals are summarized in Table 2.. The accumulation times listed are total on-source times for 4 detectors. Comparable time was spent on background observations. Two previous observations, nominally of M82 but also including M81 in the field of view (they are separated by $\sim 1^\circ$) are also listed.

OSSE observations consist of a sequence of two-minute observations of a source field alternated with two-minute offset-pointed background measurements. The two-minute duration is selected to be short relative to the typical orbital background variations. When possible, source-free fields 4.5° offset on both sides of the source position along the detector scan plane are used for background observations.

Data screening selected only the highest quality data of both source and background observations. Background estimation was achieved by performing a quadratic fit to three or (where possible) four two-minute background accumulations within 6 minutes on either side of every two-minute source accumulation. The interpolated background count rate spectrum is subtracted from each source spectrum. The differences are screened for anomalies and then averaged over times appropriate for spectral analysis. Spectra are characterized by defining a photon spectrum model, convolving the model with the instrument response function, comparing the resultant energy loss spectrum with the observed count rate spectrum, and varying the model parameters until the best agreement is obtained. A detailed description of OSSE performance and spectral data analysis procedures can be found in Johnson et al. (1993).

3. Results

The difference spectrum averaged over April 6–12 (days 9.9–15.4) shows a clear excess at energies ≤ 200 keV. The corresponding photon spectrum is shown in Figure 1. Also shown as the solid line is the best-fit thermal bremsstrahlung model used in the model-dependent deconvolution. The fit was performed with the full OSSE energy resolution and the reduced chi-squared $\chi^2_\nu = 0.85$ for 159 degrees of freedom. The intensity at 100 keV is $2.11^{+0.44}_{-0.45} \times 10^{-3} \text{ cm}^{-2} \text{ s}^{-1} \text{ MeV}^{-1}$ and the temperature is $kT_e = 82^{+61}_{-29} \text{ keV}$. These uncertainties are “1- σ ” values derived from unit increase in χ^2 with other parameters allowed to vary freely. More meaningful uncertainties are represented in Figure 2, where we show confidence level contours for the two-parameter fit. The electron temperature is very well constrained on the low side; the spectrum can not cut off below the OSSE energy range. However, very high temperatures are allowed. Thus we do not determine well the degree of equipartition between electrons and ions. Although it is not shown, even the contour corresponding to




Fig. 1.— The SN 1993J photon spectrum observed by OSSE during the interval 9.9 to 15.4 days after the explosion (assumed 28.0 March). Error bars are $1\text{-}\sigma$ and the upper limit is the $2\text{-}\sigma$ limit. The solid line shows the best-fit thermal bremsstrahlung model used in the model-dependent deconvolution.

significance of 10^{-4} does not extend to zero flux; the contour of 3×10^{-5} just does. This statement best reflects the total statistical significance of the measurement.

A power-law model also fits equally well with $\chi^2_\nu = 0.86$. The best-fit power-law amplitude at 100 keV is $1.89^{+0.41}_{-0.42} \times 10^{-3} \text{ cm}^{-2} \text{ s}^{-1} \text{ MeV}^{-1}$, and the photon spectral index is $-2.24^{+0.46}_{-0.48}$. For this model the contour corresponding to significance 1×10^{-5} is the first to extend to zero flux. If the emission is attributed to SN 1993J, either model implies a very large hard X-ray luminosity, near $5.3 \times 10^{40} \text{ erg s}^{-1}$ (50–150 keV) at distance 3.6 Mpc (Freedman et al. 1994).

The next observation, April 20 – May 3, also exhibits some evidence, although statistically weak, for hard X-ray emission. The photon spectrum from this interval is shown in Figure 3, as is the best-fit thermal bremsstrahlung model. The fitted intensity at 100 keV is $9.4^{+3.6}_{-3.8} \times 10^{-4} \text{ cm}^{-2} \text{ s}^{-1} \text{ MeV}^{-1}$, with temperature $kT_e = 74^{+120}_{-40} \text{ keV}$, and $\chi^2_\nu = 1.02$ for 159 degrees of freedom. This is formally less intense than the previous observation,




Fig. 2.— Confidence levels (68%, 90%, and 99%) for various values of the two parameters of the thermal bremsstrahlung model fit to the OSSE spectrum from the interval 9.9 to 15.4 days. The cross shows the position of the best fit. The temperature is well constrained on only the low side. The dashed line shows the combinations of parameter values that extrapolate to the absorption corrected total flux observed from 0.1–2.4 keV by ROSAT on day 12 (Zimmermann et al. 1994). Similarly, the dotted line shows parameter combinations that agree with the reported ASCA total flux (1–10 keV) on day 8 (Tanaka 1993).

although we can not rule out a constant spectrum over the two intervals.

In the last observation, June 29 – July 27, we find no evidence for emission at any energy. Because of the long observation time, the upper limit on the hard X-ray flux is well below the previously observed values. We have also analyzed 1991 and 1992 observations, nominally of NGC 3034 (M82), because they also included M81 in their source fields (see also Bhattacharya et al. 1994). We find no evidence for emission from any object in the source fields at any energy during those two pre-SN 1993J observations. The measured 50–200 keV fluxes, derived using thermal bremsstrahlung fits, for all five observations are plotted in Figure 4.




Fig. 3.— The SN 1993J photon spectrum observed by OSSE during the interval 23.7 to 37.0 days after the explosion (assumed 28.0 March). The solid line shows the best-fit thermal bremsstrahlung model.

4. Discussion

Because of OSSE's large field of view and lack of imaging capability, we can not with certainty attribute the excesses observed in the first two 1993 observations to SN 1993J. Both M82 and the nucleus of M81 are known X-ray sources and are in the OSSE source fields. However, the coincidence of the positive flux with the occurrence of SN 1993J is suggestive. Also, both ROSAT and ASCA definitely detected X rays from SN 1993J just before and during the first OSSE detection. Both observed declining fluxes after that time, with approximate e-folding times of 86 days (Zimmermann et al. 1994) and 60 days (Wheeler & Filipenko 1994), respectively. The OSSE flux variations are, within the large uncertainties, consistent with these declines, as if from the same source. As described below the observed OSSE spectra extrapolate for thermal models, with large uncertainties, very close to the ROSAT (0.1–2.4 keV) and ASCA (1–10 keV) spectra at similar times. Nature could be so perverse as to cause another source to mimic what we might reasonably expect




Fig. 4.— The hard X-ray flux measured by OSSE in five separate observations of the M81/M82 region. The outburst of SN 1993J is shown by the arrow.

to be the spectrum and time variation of emission from SN 1993J, but we view that as less likely than that the emission is from SN 1993J.

Hard X-ray emission of this intensity was not anticipated. It is almost certainly not from scattered γ rays from radioactivity. The optical light curve has quite well revealed the mass of ^{56}Co present, and even if it was mixed through the entire ejecta the hard X-ray flux from it is roughly a factor of 3 below the OSSE sensitivity in these observations, as are the γ -ray line fluxes (Woosley et al. 1994). The observed spectrum is also significantly steeper than that of down-scattered line photons. The low-energy X-ray observations were quite naturally explained in terms of the reverse-shocked supernova ejecta (Suzuki et al. 1993; Chevalier & Fransson 1994), although a contribution from the shocked wind could also be present. The electron temperature we find, 10^9 K, is easily accounted for in the shocked wind for either relatively modest expansion velocities (10^4 km s $^{-1}$) with equal electron and ion temperatures (Fransson 1984), or for higher ejecta velocities in non-equipartition solutions (Suzuki et al. 1993). However, all calculations indicate the ejecta free-free luminosity exceeds that of the lower density wind by at least an order of magnitude. We

can check the possibility that both emission regions are seen by comparing the soft and hard X-ray spectra.

The extrapolation of OSSE spectra to keV energies is very uncertain. We show in Figure 2 as a dashed line the intensity/temperature combinations that extrapolate to the absorption corrected luminosity observed from 0.1–2.4 keV by ROSAT (Zimmermann et al. 1994) during the OSSE observation. The center of the contours extrapolates right to the ROSAT data. All points to the upper left of the dashed line are allowed in the sense that they do not exceed the observed soft flux. The dotted line shows models that extrapolate to the preliminary ASCA 1–10 keV flux (Tanaka 1993).

Although a power-law model also fits the OSSE data, the extrapolated best-fit power-law greatly exceeds the observed soft X-ray fluxes. Only very flat spectra, allowed at a few percent confidence, do not violate the soft X-ray constraint. Still a nonthermal source, or a partial contribution thereof, can not be ruled out by the OSSE data.

The simplest interpretation is that there is a single free-free emission region with kT_e in the range 50 to 100 keV. If this is the shocked wind the required high emissivity demands an unusually high density, of order 10^9 cm^{-3} for pure hydrogen. This in turn implies a high progenitor mass-loss rate, $\geq 10^{-4} M_\odot \text{ yr}^{-1}$ for a constant spherically symmetric wind with a velocity of 10 km s^{-1} . If the emission seen by OSSE is from the forward shock there is little room left for a softer component from the reverse-shocked ejecta between the low-energy X-ray observations and the extrapolation from OSSE data. Also, with the high temperature and relatively high optical depth of the shocked wind, the power-law spectrum from Comptonization of photospheric photons would extend into the soft X-ray range (Fransson 1984). This is apparently not observed. We speculate that high density clumps in the circumstellar matter could possibly avoid some of these problems.

Alternatively, the single emission region observed could be the shocked ejecta, but with an unusually high temperature. The favored model of Suzuki et al. (1993) has initial ejecta velocity of $5 \times 10^4 \text{ km s}^{-1}$ and gives a shocked ejecta temperature of $kT_e = 25 \text{ keV}$. To explain the OSSE data, the temperature and free-free luminosity have to be increased by factors 2–4 over this already rather extreme model. This could be achieved by some combination of increased velocity, shallower ejecta density gradient, and lower ejecta/wind density contrast. Any such model must also be tested against many other related constraints (Fransson, Lundqvist, & Chevalier 1994) such as from the radio and optical light curves and models of the stellar atmosphere.

5. Summary

For the first time a supernova has been detected in >50 keV hard X rays soon after outburst. This emission was observed to decline on the timescale of several weeks. If thermal in nature the temperature was initially $T \simeq 10^9$ K. A contribution of a nonthermal source can not be excluded. The observed emission probably arose from the interaction of the supernova ejecta with circumstellar material. The hard X-ray luminosity 10–15 days after outburst, $\simeq 5 \times 10^{40}$ erg s $^{-1}$ from 50–150 keV, dominates the observed luminosity of the interaction region. It is much larger than was anticipated prior to the event and larger than predicted by models designed to explain the observed <10 keV X-ray emission. The > 50 keV emission might be bremsstrahlung from either the shocked presupernova wind or the reverse-shocked supernova ejecta. The former can easily have the requisite high electron temperature (10^9 K) but was not thought to be dense enough to give high enough luminosity, and there can be additionally very little softer emission from the latter without violating soft X-ray observations. For the ejecta itself to give the observed hard X rays rather extreme models with high expansion velocity are required. Understanding these and other observations of SN 1993J will ultimately lead to a detailed description of the structure of the presupernova star and its wind, and the energy of the explosion.

This work was supported by NASA Grant DPR S-10987C.

REFERENCES

- Bhattacharya, D. et al. 1994 preprint
- Canizares, C. R., Kriss, G. A., & Feigelson, E. D. 1982, *Astrophys. J.*, **253**, L17
- Chevalier, R. A. 1982, *Astrophys. J.*, **259**, 302
- Chevalier, R. A. & Fransson, C. 1994, *Astrophys. J.*, in press
- Dotani, T., Hayashida, K., Inoue, H., Itoh, M., Koyama, K., Makino, F., & Mitsuda, K. 1987, *Nature*, **330**, 230
- Van Dyk, S. D. et al. 1993, *Bull. American Astron. Soc.*, **25**, 1244
- Filippenko, A. V., Matheson, T., & Ho, L. C. 1993, *Astrophys. J.*, **415**, L103
- Fransson, C. 1984, *Astron. Astrophys.*, **133**, 264
- Fransson, C., Lundqvist, P., and Chevalier, R. A. 1994, preprint.
- Freedman, W. L. et al. 1994, *Astrophys. J.*, 427, 628
- Hoflich, P., Langer, N., & Duschinger, M. 1993, *Astron. Astrophys.*, **275**, L29
- Johnson, W. N. et al. 1993, *Astrophys. J. Suppl.*, 86, 693, 23-D3

- Matz, S. M., Share, G. H., Leising, M. D., Chupp, E. L., Vestrand, W. T., Purcell, W. R., Strickman, M. S., & Reppin, C. 1988, *Nature*, **331**, 416
- McKee, C. F. 1974, *Astrophys. J.*, **188**, 335
- Nomoto, K., Suzuki, T., Shigeyama, T., Kumagai, S., Yamaoka, H., & Saio, H. 1993, *Nature*, **364**, 507
- Phillips, J. A. & Kulkarni, S. R. 1993, *IAU Circ.*, **5775**, 1
- Podsiadlowski, P., Hsu, J. L., Joss, P. C., & Ross, R. R. 1993, *Nature*, **364**, 509
- Pooley, G. G. & Green, D. A. 1993, *IAU Circ.*, **5773**, 1
- Schmidt, B. P. et al. 1993, *Nature*, **364**, 600
- Sunyaev, R. et al. 1987, *Nature*, **330**, 227
- Suzuki, T., Kumagai, S., Shigeyama, T., Nomoto, K., Yamaoka, H., & Saio, H. 1993, *Astrophys. J.*, **419**, L73
- Swartz, D. A., Clocchiatti, A., Benjamin, R., Lester, D. F., & Wheeler, J. C. 1993, *Nature*, **365**, 232
- Tanaka, Y. 1993, *IAU Circ.*, **5753**, 1
- Wheeler, J. C. et al. 1993, *Astrophys. J.*, **417**, L71
- Wheeler, J. C. & Filipenko, A. V. 1994, in *Supernovae and Supernova Remnants*, ed. R. McCray, (Cambridge: Cambridge Press), in press
- Woosley, S. E., Eastman, R. G., & Weaver, T. A. 1994, *Astrophys. J.*, in press
- Zimmermann, H.-U. et al. 1994, *Nature*, **367**, 621