

AD_____

Award Number: W81XWH-05-1-0153

TITLE: PSA-Based Screening Outcomes, Dietary Heterocyclic Amine Exposure, and Prostate Cancer Risk in African Americans

PRINCIPAL INVESTIGATOR: Kenneth T. Bogen

CONTRACTING ORGANIZATION: University of California
Livermore, California 94550

REPORT DATE: January 2007

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 01-01-2007		2. REPORT TYPE Annual		3. DATES COVERED (From - To) 1 Jan 2006 – 31 Dec 2006	
4. TITLE AND SUBTITLE PSA-Based Screening Outcomes, Dietary Heterocyclic Amine Exposure, and Prostate Cancer Risk in African Americans				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-05-1-0153	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Kenneth T. Bogen E-Mail: bogen@LLNL.gov				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California Livermore, California 94550				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Prostate cancer (PC) is the second leading cause of male U.S. cancer deaths, with African-Americans having the highest rate of PC mortality worldwide. A 5-year prospective NIH-funded clinic-based study investigated whether prostate-specific antigen (PSA) and digital rectal exam (DRE) screening indicators of PC risk in 500 African-American men 50 to 70 years of age who underwent PC screening in Oakland, CA (East Bay San Francisco area), were associated with estimated dietary exposures to 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP), which forms when meat is overcooked. The DOD-funded study expands that NIH-funded work by adding a new %-free-PSA test for 312 (108 from the NIH-funded study, plus 200 additional) men, results of which will be compared with PSA/ DRE results and PhIP exposures estimated by dietary interviews. For 392 men studied under the NIH protocol, an odds ratio (95% CL) of 32 (3.2, 720) for highly elevated PSA (≥ 20 ng/mL) was observed in the highest 15% vs. the lower 50% of estimated daily PhIP intakes. As of 31-12-06, a total of 220 additional men completed participation using the expanded protocol, for a combined total of 612 men. For 562 of these men studied to date, the corresponding OR was found to be 24 (2.20, 533). This study will help define the potential value of improved screening and dietary/behavioral intervention to reduce PC risk.					
15. SUBJECT TERMS African-American men, cooked-meat carcinogens, epidemiology, PhIP, prostate cancer, screening					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	28	19b. TELEPHONE NUMBER (include area code)

Table of Contents

Introduction.....	4
Body.....	7
Key Research Accomplishments.....	10
Reportable Outcomes.....	10
Conclusions.....	11
References.....	12
Appendix.....	18

Introduction

PhIP is a Dietary Carcinogen that May Pose Heightened Risk to African-American Men

African American (AA) men, who compared to Caucasians die nearly twice as much from prostate cancer (PC), also take in about twice as much of the predominant U.S. dietary heterocyclic amine, 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) (Bogen and Keating, 2001), which occurs primarily in well-cooked chicken and beef. Heterocyclic amines (HAs) are potent mutagens formed in meats, chicken and fish as it is cooked to higher-doneness levels by heat-intensive cooking methods (Thompson *et al.*, 1987; Keating *et al.*, 1999, 2000). HAs also cause cancer at a variety of sites in multiple bioassay animal species/strains/sexes, as well as at multiple sites within many of species/strains/sexes tested (Bogen, 1994). A predominant HA found in cooked and particularly in well-done chicken and beef is 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) (Felton *et al.*, 1984, 1986; Felton and Knize, 1990a-b; Sinha *et al.*, 1995). Dietary exposure to PhIP has been shown to induce colon, intestinal and mammary adenocarcinomas in rats (Ohgaki *et al.*, 1986; Ochiai *et al.*, 1991; Ito *et al.*, 1991; Ito *et al.*, 1997; Ghoshal *et al.*, 1994), as well as prostate cancer in rats (Shirai *et al.*, 1997,1999). HAs are metabolically activated by P450 and N-acetyltransferase enzymes to activated forms that bind covalently to (among other targets) DNA in tissues (including in prostate) where HAs induce cancer in rats (Thorgeirsson *et al.*, 1983; Thorgeirsson, 1984; Rosenkranz and Mermeistein, 1985; Sato *et al.*, 1986; Kato and Yamazoe, 1987; Snyderwine and Battula, 1989; McManus *et al.*, 1990; Turesky *et al.*, 1991; Davis *et al.*, 1993; Kaderlik *et al.*, 1994a-b; Takahashi *et al.*, 1998; Schut and Snyderwine, 1999; Gooderham *et al.*, 2002). In male *lacI* transgenic rats, a diet containing 200 ppm PhIP was shown to be highly mutagenic in prostate tissue (Stuart *et al.*, 2000). In cultured human prostate tissue and primary prostate cells, PhIP is metabolically activated to mutagenic forms that covalently bind to and damage DNA (Williams *et al.*, 2000; Lawson and Kolar, 2002; Martin *et al.*, 2002; Di Paolo *et al.*, 2005).

PC and African-American men

In the U.S., prostate cancer (PC) is a leading cause of cancer death among men, with African Americans having the highest age-specific prostate cancer rate in the world, and a >2-fold higher rate of mortality for PC than white men in the U.S. (Miller *et al.*, 1996; Robbins *et al.*, 1998; Hsing *et al.*, 2000). Although family (particularly father/brother) history of PC is clearly linked to substantially elevated PC risk (Schuurman *et al.*, 1999; Hemminki and Czene, 2002; Zeegers *et al.*, 2003; Hemminki and Chen, 2005) and a human-PC-specific chromosome translocation has been identified (Tomlins *et al.*, 2005), there is (as for most other cancers) no evidence for a predominant heritable factor for PC. Racial-ethnic differences in levels of testosterone-related hormones and related genetic controls on hormone-induced prostate growth have been hypothesized to explain or contribute to corresponding racial-ethnic differences in PC risk, but their role remains unclear (Ross *et al.*, 1998; Pettaway, 1999; Hsing, 2001). Even if hormonally mediated background processes do affect PC risk, dietary exposures to genotoxic HA carcinogens may act independently or interact synergistically with such hormonal processes to further modify PC risk.

HA, Meat, Other Dietary Factors, and PC Risk

Substantial geographical variations in PC incidence indicate the importance of one or more dietary or other environmental factors (Minami *et al.*, 1993; Mettlin, 1997; Angwafo, 1999). Dietary intake of animal or saturated fat is the environmental factor most consistently linked to significantly increased PC risk in previous studies, particularly among African-American men, though these associations appear too weak to explain more than a small fraction of observed racial-ethnic differences in PC risk (Whittemore *et al.*, 1995; Hayes *et al.*, 1999; Kolonel *et al.*, 1999; Daniels *et al.*, 2004), as also appears to be the case for other environmental/dietary factors examined such as calcium, cruciferous vegetables, vitamin D, UV from sunlight, lycopene, and body size (Giovannucci *et al.*, 1997, 1998; Cohen *et al.*, 2000; Chan and Giovannucci, 2001a-b; Kristal and Lampe, 2002; Bodiwala *et al.*, 2003). Because cooked-meat intake is positively associated with total saturated-fat intake (USDA, 1998), previous studies that focused on animal or saturated fat *per se* could have estimated effects due largely or entirely to meat-related HA intakes.

Consumption of cooked meats and associated HAs have been linked to increased risks of colorectal adenoma/adenocarcinoma and of cancer of the stomach, breast, lung and prostate (Mills *et al.*, 1989; Shiffman and Felton, 1990; Gerhardsson de Verdier *et al.*, 1991; Talamini *et al.*, 1992; Lang *et al.*, 1994; De Stefani *et al.*, 1995; Ewings and Bowie, 1996; Probst-Hensch *et al.*, 1997; Ward *et al.*, 1997; Kampman *et al.*, 1999; Norrish *et al.*, 1999; Sinha *et al.*, 1998a, 1999a-b; Zheng *et al.*, 1998, 1999; Murtaugh *et al.*, 2004), while fewer studies found no such associations (Lyon and Mahoney, 1988; Muscat and Wynder, 1994; Augustsson *et al.*, 1999). Positive studies include those in which HA exposure was quantified adjusting for factors expected to determine HA intake—namely, meat type, consumption frequency, cooking method, and cooking doneness. A potential link between PhIP intake, in particular, and the elevated risk of PC experienced by African-American compared to Caucasian men is suggested by relatively greater levels of PhIP and its metabolites detected in urine sampled from the latter vs. the former group (Kidd *et al.*, 1999), although urine analysis only provides a measure of PhIP exposure within 12-24 hours prior to sampling (Malfatti *et al.*, 1999). Such a link is also supported by studies using meat- and cooking-method-specific HA-concentration estimates from multi-laboratory sets of experimental cooking data (Keating and Bogen, 2001) to assess intakes of PhIP and other HAs by >20,000 U.S. (including >3,000 African-American) participants in the nationwide, stratified, random-sample U.S. Continuing Survey of Food Intakes by Individuals (CSFII) (USDA, 1993, 1998). Analyzed by age-, sex, and race/ethnicity, these HA-exposure assessments found at all ages that African-American males consume on average at least twice as much PhIP (and total HA) per kg body weight per day as do U.S. Caucasian males, and that for both groups there is a significant positive (albeit small) correlation between estimated mean intakes of PhIP and those of saturated or total fat (Keating and Bogen, 2001; Bogen and Keating, 2001).

Although the study comparing HA intake (primarily from cooked lamb) and PC risk in New Zealand men by Norrish *et al.* (1999) found no significant PhIP-related

associations involving PC, that study did report a significant association between PC risk and consumption of beefsteak by higher cooking-doneness category (2-sided $p_{\text{trend}} = 0.008$). Several aspects of that study suggest it may have had limited power to assess potential associations between PC and HA or PhIP intake in the U.S. The incidence of PC in New Zealand is only half that of Caucasian men in the U.S. (Hsing et al. 2000). Average meat (primarily lamb) consumption by subjects in the Norrish et al. study (~150 g/d) was well below that of U.S. Caucasian men (260 g/d) and very well below that of African-American men (304 g/d) (USDA, 1998; Table 10A). Moreover, the *minimum* PhIP intake (224 ng/d) in the *top* exposure quartile of Norrish et al. study subjects was well below the estimated *average* PhIP intakes by U.S. Caucasian (390 ng/d) and by African-American (600 ng/d) men (Bogen and Keating 2001).

A randomized, controlled multisite prospective study that ascertained 1,338 PC cases between 1993 and 2001 among 29,361 (including 3.3% African American) men of age 55-74 years in 10 U.S. cities was conducted specifically to determine whether meat intake or meat-related mutagens, including PhIP, was associated with increased PC risk (Cross et al., 2005). In this study, intakes of meat, PhIP, two other heterocyclic amines, and meat-associated Ames-assay mutagenic yield were all estimated using FFQs that were self-administered, but otherwise very similar to those currently administered (by in-person interview) in the clinical study which Project 2 of this application would extend. More than 10 g/day vs. no intake of very well done meat was found to be associated with a 1.7-fold increased risk (95% CI: 1.19-2.40), and the highest vs. lowest PhIP quintile with a 1.3-fold increased risk (95% CI: 1.01-1.61), of PC incidence (Cross et al., 2005). Other intakes were not found to be associated with increased PC risk. No separate analysis was done of study results by race-ethnicity was not reported, nor were genotype or other (e.g., dietary) factors considered that are known to affect PhIP metabolism.

PC Screening and PC

Periodic screening for plasma levels of Prostate-Specific Antigen (PSA) is useful for early detection of PC as well as benign prostatic hyperplasia (BPH) because PSA levels in serum are positively associated with age, prostate volume, and prostatic neoplastic disease (Carter *et al.*, 1992; Oesterling *et al.*, 1993; Etzioni *et al.*, 1999). Significantly higher PSA levels are found in African-Americans than in whites, even after adjustment for age and prostate volume (in men without PC) and for PC grade and stage (in men with PC) (Abdalla *et al.*, 1998a-b; Vijayakumar *et al.*, 1998), due evidently to greater PSA secretion per unit prostate volume by African-American men (Fowler et al., 1999). Although a substantial fraction of “slightly” elevated PSA levels (between 4 and 10 ng/mL) is attributable to BPH or infection, “highly elevated” PSA levels (≥ 20 ng/mL) typically indicate a strong likelihood of localized or metastatic PC, with 80 to 90% positive predictivity and $>99\%$ specificity (Mänttinen et al., 2001; Gerstenbluth et al., 2002; Smith et al., 2004). Likewise, a PSA measure < 4.0 ng/mL often considered within the “normal” range is actually associated with about at 15% of later PC diagnosis (Thompson et al., 2004).

Report Body

Objective/Hypotheses: The study goal is to broaden the scope and power of a ground-breaking study of potential association between dietary HA exposure and screening indicators of PC risk in African-American men by adding a newer %fPSA test to the PSA/DRE protocol now being used. We hypothesize 1) that the added %fPSA test will increase PC detection in our study population, and 2) that the combined screening, follow-up diagnostic and dietary survey data obtained will reveal a positive association between estimated HA intake and screening and diagnostic indicators of increased PC risk in our African American study population.

Specific aims: Our aims are to 1) Add a newer %fPSA test to PSA and DRE screens being done in a current study of potential associations between HA and PC in Oakland, California, area African-American men. 2) Assess potential increased rate of PC identified by including the %fPSA test with PSA and DRE results in light of clinical follow-up diagnoses obtained. 3) Use PSA-related and DRE test results, together with corresponding follow-up diagnostic and dietary survey data, to assess the potential association of HA-related exposure factors and increased PC risk in African-Americans.

Study Design: This is a prospective, clinic-based screening study. For aim 1, 392 participants were solicited from an already-established network of churches, clinics and additional African-American community groups in the Oakland, CA, area. The DOD-funded work continues this effort, adding the new %fPSA biomarker. Detailed data on diet and meat consumption continue to be obtained by in-person interviews using established questionnaires, each followed by PSA-related and DRE screening tests, and follow-up diagnosis—a study design that avoids potential bias due to prior participant/investigator knowledge of PC status. Aims 2 and 3 are being accomplished by statistical data analysis, with aim 3 to be supported by incorporation of similar (other than %fPSA) data obtained for the 392 participants already obtained through the NIH-funded study that ended in 2006, for a total combined study size of 700 men. Our focus on African-Americans provides needed study power, in view of the greater PC risk faced by this specific group.

Progress in Relation to Work Plan

Work Plan Summary

A total of 392 participants were accrued in an ongoing NIH-funded study through December 2004 using dietary interviews, a digital rectal exam (DRE), and a standard prostate serum antigen (PSA) blood test. The DOD Prostate Cancer Research Program support has added a second PSA-related test—the “percent-free PSA” (%fPSA) test—to the protocol already applied to 220 additional participants as of December 2006, and is planned to provide for 88 additional study participants through September of 2008, for whom blood samples will be drawn for both a PSA and a %fPSA test and the dietary interview and DRE will be done, all at the study clinic in Oakland on the same day for each participant.

Delayed Receipt of Funds and HSRRB Approval of Human Subjects Protocol

Initiation of the study was delayed due to delayed receipt of DOD funds by Lawrence Livermore National Laboratory (LLNL). Due to contract language negotiations, funds arrived at LLNL approximately four months after the official start date of the project (Jan. 5, 2005). A further start-up delay occurred due to delayed receipt of U.S. Army Human Subjects Research Review Board (HSRRB) approval of the study human subjects protocol, which had already received approval by the other three Institutional Review Boards (IRBs) involved in this study (those of LLNL, the University of California San Francisco Medical School, and the Summit Alta Bates medical Center in Oakland, CA). A draft human subjects protocol was submitted to the HSRRB in December of 2004, but HSRRB approval was not obtained until mid-May of 2005.

The delayed start-up and extended IRB-related documentation required to start this project caused small, unanticipated shifts in LLNL labor relative to the original budget plan. However, the project is currently planned for completion per the original project schedule.

Study Progress (indicated below in bold)

Task 1. Add %fPSA test to the standard PSA blood test and DRE being done in a current study protocol to assess potential HA/PC associations in Oakland-area African American men (Months 1-27). **Initiated as of June 15, 2005; ongoing.**

1.A. Obtain IRB approvals for modified study protocol (Months 1-3). Accomplishment of this task will be facilitated by the established protocol currently being used in the ongoing parallel NIH-funded study; approvals will be obtained for a modification of this protocol to add the %fPSA test to the PSA and DRE screens currently being used, to extend the study termination date by one year (through January 2008), and to increase the number of participants by 300 to a study total of 700. **Completed in May of 2005.**

1.B. Implement combined PSA-test protocol (PSA + %fPSA) for a total of 400 African American men screened at the Summit Alta-Bates MCEPC clinic in Oakland, California (Months 4-27). The %fPSA, PSA and DRE procedures to be used are all clinical procedures now performed routinely at the study clinic (Markstein Cancer Education and Prevention Center, Alta Bates Summit Medical Center, Oakland, CA). **Initiated as of June 15, 2005; ongoing.**

1.C. We will interview study subjects and edit dietary questionnaires for all study subjects, to include:

1.C.1. 200 participants, comprising 100 DOD-funded participants plus 100 participants funded by NIH, all to be screened during Months 3-15. **All planned NIH (100) and DOD (~100) participants were completed as of December 31, 2005.**

1.C.2. 200 additional DOD-funded participants will be screened during Months 16-27. **A total of about 130 of these planned 200 DOD participants were completed as of January 25, 2007 (total combined study size: ~630 of 700 planned total).**

1.C.3. For all participants to be screened in this study, previously developed dietary survey questionnaires will be used in the same manner they are being used in our ongoing corresponding NIH-funded research study (see Questionnaires, Surveys & Clinical Protocols). **Initiated as of June 15, 2005.**

Task 2. Assess improvement in PC sensitivity and selectivity by addition of %fPSA test to the standard PSA test based on PSA-related test results and follow-up diagnoses in our African American study participants (Months 18-36). **Yet to be done.**

2.A. Data analysis methods to be used will be identical to those in use for the current related NIH-funded study (see Questionnaires, Surveys & Clinical Protocols)

2.B. Preliminary statistical analyses will be conducted during Months 18-27. **Completed as of December 31, 2006, except for %fPSA. See study publication attached.**

2.C. Final statistical analyses will be conducted during Months 27-36

Task 3. Use Combined PSA-related test results, together with dietary survey data, to assess the potential association of HA-related exposure factors and increased PC risk in African Americans (Months 13-36). **See study publication attached.**

3.A. Data analysis methods to be used will be identical to those in use for the current related NIH-funded study. **Analysis of initial data set (n = 392) obtained using the original protocol for NIH-funded work is described in the attached study publication. An updated analysis involving a total of 562 men was completed and will be presented at the upcoming 2007 annual meeting of the American Association for Cancer Research. The updated analysis supports key study hypotheses. (See attached 2007 AACR abstract.) These results set the stage for follow-up analyses that will add new data involving %fPSA test results.**

3.B. Preliminary statistical analyses of the statistical validity of Study Hypotheses 1 and 2 will occur during Months 18-27. **See study publication attached.**

3.C. Final statistical analyses of the statistical validity of Study Hypotheses 1 and 2, as well as manuscript preparation, will occur during Months 27-36. **Yet to be done.**

3.D. We will also, as feasible, test the validity of Study Hypothesis 2 using combine DOD-funded data set (n = 300) with NIH-only-funded data set (n = 400), during Months 27-36. **Yet to be done, except for analysis of expanded interim NIH/DOD data set (n = 562) (see attached 2007 AACR abstract). These results set the stage for follow-up analyses that will add new data involving %fPSA test results.**

Key Research Accomplishments

The NIH-funded P01 work that set the stage for the present expanded study has accomplished the two specific aims it sought to address. We successfully applied methods for estimating HA concentrations in cooked meats based on individually expressed data on meat-specific intakes, cooking methods and doneness preferences to estimate daily PhIP intakes, and we have found these intake estimates to be positively associated with screening indicators of highly elevated PC risk in a prospective clinic-based PC screening study involving nearly 400 African-American men in the San Francisco East Bay area. The observed positive association, which was most significant among men 55 to 70 years of age ($p_{\text{trend}} = 0.00020$), remained statistically significant after adjustments for saturated fat intake, total energy intake and self-reported father/brother history of PC. These study findings were upheld in updated analyses that involved a total of 562 men from the combined NIH- and DOD-supported work (see attached 2007 AACR abstract).

Future work will complete the accrual of all planned participants using the same basic study design, including the expanded screening indicator used to predict PC status (%fPSA test), and assess whether we continue to observe significant PhIP-related associations pertain to combined screening indicator data using a larger study population. In so doing, this study will continue to help define the potential value of improved screening and dietary/behavioral intervention to reduce PC risk, namely, prevention of PhIP intake by avoiding overcooked meats.

Reportable Outcomes

Abstracts/Posters

Bogen, K, W Baker, J Chan, D Nelson, E Holly, G Keating, L Paine, and J Felton. 2005. Prostate cancer screening and dietary HA exposure in African-Americans. UCRL-POST-211476. Poster presented at the University of California Davis Health System Future Fair, May 5, 2005, UC Davis Medical Center, Sacramento, CA.

KT Bogen, GA Keating, EA Holly, J Chan, L Paine, EL Simms, DO Nelson, and J Felton. 2005. Prostate Serum Antigen levels and dietary heterocyclic amines in

African Americans: A prospective clinic-based study. Abstract of poster accepted for presentation at the 97th Annual Meeting of the American Assoc. for Cancer Research, April 1-5, 2006, Washington, DC. UCRL-ABS-217085. Lawrence Livermore National Laboratory, Livermore, CA.

Bogen, KT, J Chan, GA Keating, LJ Paine, EL Simms, EA Holly, and JS Felton. 2006. Prostate-specific antigen levels and dietary PhIP in African Americans: A prospective clinic-based study. [Abstract] 2007 Annual Meeting of the American Association for Cancer Research, 14-18 April 2007, Los Angeles, CA. [UCRL-ABS-226551] [included in the **Appendix** of this report]

Reports and other Publications

Bogen KT. 2006. PSA-Based Screening Outcomes, Dietary Heterocyclic Amine Exposure, and Prostate Cancer Risk in African Americans: Annual Report (Year 1 of 3). UCRL-TR-218258. Lawrence Livermore National Laboratory, Livermore, CA.

Keating GA, K Bogen, and J Chan. 2007. Development of a meat frequency questionnaire for use in diet and cancer studies. *J. Am. Dietetic Assoc.* (in press). [UCRL-JRNL-208025]

Bogen KT, GA Keating II, JM Chan, LJ Paine, EL Simms, DO Nelson, and EA Holly. 2007. Highly elevated PSA and dietary PhIP intake in a prospective clinic-based study among African Americans. *Prostate Cancer and Prostatic Diseases* (in press). [included in the **Appendix** of this report]

Louis ED, Zheng W, Jiang W, Bogen KT, and Keating GA. 2007. Quantification of the neurotoxic β -carboline harmaline in barbecued/grilled meat samples and correlation with level of doneness. *J. Toxicol. Environ. Health* (in press). [UCRL-JRNL-225956] [This publication resulted from a collaboration in which a team from the Columbia University Medical School used the same LLNL meat questionnaire that also is being used in the present DOD-funded work. This publication thus represents an unexpected indirect benefit of the present DOD-funded work.]

Conclusions

Prostate cancer (PC) is the second leading cause of male U.S. cancer deaths, with African-Americans having the highest rate of PC mortality worldwide, as well as more abnormal results from screening tests that correlate with current or eventual PC. Prostate cancer (PC) is the second leading cause of male U.S. cancer deaths, with African-Americans having the highest rate of PC mortality worldwide. A 5-year prospective NIH-funded clinic-based study investigated whether prostate-specific antigen (PSA) and digital rectal exam (DRE) screening indicators of PC risk in 500 African-American men 50 to 70 years of age who underwent PC screening in Oakland,

CA (East Bay San Francisco area), were associated with estimated dietary exposures to 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP), which forms when meat is overcooked. The DOD-funded study expands that NIH-funded work by adding a new %free-PSA test for 312 (108 from the NIH-funded study, plus 200 additional) men, results of which will be compared with PSA/ DRE results and PhIP exposures estimated by dietary interviews. For 562 men studied under the NIH protocol, an odds ratio (95% CL) of 32 (3.2, 720) for highly elevated PSA (≥ 20 ng/mL) was observed in the highest 15% vs. the lower 50% of estimated daily PhIP intakes. As of 31-12-06, a total of about 230 additional men completed participation using the expanded protocol, for a combined total of about 630 men. For 562 of these men studied to date, the corresponding OR was found to be 24 (2.20, 533). This study will help define the potential value of improved screening and dietary/behavioral intervention to reduce PC risk.

Future work will complete the planned accrual of participants in this study using expanded screening indicators to predict PC risk, including the %fPSA test, and to assess whether PhIP-related associations pertain to combined screening indicator data using a larger study population. In so doing, this study will continue to help define the potential value of improved screening and dietary/behavioral intervention to reduce PC risk, namely, prevention of PhIP intake by avoiding overcooked meats.

References

- Abdalla I., Ray P., and Vijayakumar S. Race and serum prostate-specific antigen levels: current status and future directions. *Semin. Urol. Oncol.* 1998a: 16: 207-213.
- Abdalla I., Ray P., Ray V., Vaida F., and Vijayakumar S. Comparison of serum prostate-specific antigen levels and PSA density in African American, white, and Hispanic men without prostate cancer. *Urology* 1998b: 51: 300-305.
- Alavanja, M.C., Sandler D.P., McMaster S.B., Zahm S.H., McDonnell C.J., Lynch C.F., Pennybacker M., Rothman N., Dosemeci M., Bond A.E., and Blair A. The agricultural health study. *Environ. Health Perspect.* 1996: 104: 362-369.
- Block G., Hartman A.M., Dresser C.M., Carroll M.D., Gannon J., and Gardner L. A data-based approach to diet questionnaire design and testing. *Am. J. Epidemiol.* 1986: 124: 453-469.
- Angwafo, F.F. Migration and prostate cancer: an international perspective. *J. Natl. Med. Assoc.* 1998: 90(11 Suppl): S720-S723.
- Augustsson K., Skog K., Jagerstad M., Dickman P.W., and Steineck G. Dietary heterocyclic amines and cancer of the colon, rectum, bladder, and kidney: A population based study. *Lancet* 1999: 353: 703-707.
- Carter H.B., Pearson J.D., Metter E.J., Brant L.J., Chan D.W., Andres R., Fozard J.L., and Walsh P.C. Longitudinal evaluation of prostate-specific antigen levels in men with and without prostate disease. *J. Am. Med. Assoc.* 1992: 267: 2215-2220.
- Bodiwala D., Luscombe C.J., Liu S., Saxby M., French M., Jones P.W., Fryer A.A., Strange R.C. Cancer potencies of heterocyclic amines found in cooked foods. *Cancer Lett.* 2003: 192: 145-149.
- Bogen, K.T. Improved prediction of carcinogenic from mutagenic potencies for chemicals positive in rodents and the Ames test. *Molec. Environ. Mutagen.* 1995: 25: 37-49.
- Bogen K.T. Cancer potencies of heterocyclic amines found in cooked foods. *Food Chem. Toxicol.* 1994: 32: 505-515.
- Bogen K.T., and Keating G.A. U.S. dietary exposures to heterocyclic amines. *J. Exposure Anal. Environ. Epidemiol.* 2001: 11: 155-68.

- Bogen K.T., Keating II G.A., Chan J.M., Paine L.J., Simms E.L., Nelson D.O., and Holly E.A. Highly elevated PSA and dietary PhIP intake in a prospective clinic-based study among African Americans. *Prostate Cancer and Prostatic Diseases* 2007:(in press).
- Breslow N.E., and Day N.E. *Statistical Methods in Cancer Research, Volume II—The Design and Analysis of Cohort Studies*. IARC Scientific Publ. No. 82. International Agency for Cancer Research (IARC), Lyon, France. 1987: 106–118.
- Chan J., and Giovannucci E. Vegetables, fruits, associated micronutrients, and risk of prostate cancer. *Epidemiol. Rev.* 2001a: 23: 82-86.
- Chan J., and Giovannucci E. Dairy products, calcium, and vitamin D, and risk of prostate cancer. *Epidemiol. Rev.* 2001b: 23: 87-92.
- Cohen J.H., Kristal A.R., and Stanford J.L. Fruit and vegetable intakes and prostate cancer. *J. Natl. Cancer Inst.* 2000: 92: 61-68.
- Cross A.J., Peters U., Kirsh V.A., Andriole G.L., Reding D., Hayes R.B., and Sinha R. A prospective study of meat and meat mutagens and prostate cancer risk. *Cancer Res.* 2005: 65:11775-11784.
- Daniels L.K., Snetselaar L.G., Smith B.J., Stewart R.E., and Robbins M.E.C. Problems with the assessment of dietary fat in prostate cancer studies. *Am. J. Epidemiol.* 2004: 160: 436-444.
- De Stefani E., Fierro L., Barrios, E. and Ronco A. Tobacco, alcohol, diet and risk of prostate cancer. *Tumori* 1995: 81: 315-320.
- Di Paolo O.A., Teitel C.H., Nowell S., Coles B.F., and Kadlubar, F.F. Expression of cytochromes P450 and glutathione S-transferases in human prostate, and the potential for activation of heterocyclic amine carcinogens via acetylc-CoA-, PAPS- and ATP-dependent pathways. *Int. J. Cancer* 2005: 117: 8-13.
- Draper N.R., and Smith H. *Applied Regression Analysis*, 2nd ed. John Wiley & Sons: New York, NY, 1981.
- Etzioni R., Cha R., and Cowen M. Serial prostate specific antigen screening for prostate cancer: A computer model evaluates competing strategies. *J. Urology* 1999: 162: 741-748.
- Ewings P., and Bowie C. A case-control study of cancer of the prostate in Somerset and east Devon. *Br. J. Cancer* 1996: 74: 661-666.
- Felton J.S., and Knize M.G. Heterocyclic-amine mutagens/carcinogens in foods. In: Cooper C.S., and Grover P.L. eds. *Handbook of Experimental Pharmacology*, 94/1. Springer-Verlag: Berlin 1990a: 471-502.
- Felton J.S., and Knize M.G. New mutagens from cooked food. In: Pariza M., Aeschbacher H.-U., Felton J.S., and Sato S. eds. *Mutagens and Carcinogens in the Diet*. Wiley-Liss, Inc.: New York 1990b: 19-38.
- Felton J.S., Knize M.G., Shen N.H., Andresen B.D., Bjeldanes L.F., and Hatch F.T. Identification of the mutagens in cooked beef. *Environ. Health Perspect.* 1986: 67: 17-24.
- Felton J.S., Knize M.G., Wood C., Wuebbles B.J., Healy S.K., Stuermer D.H., Bjeldanes L.F., Kimble B.J., and Hatch F.T. Isolation and characterization of new mutagens from fried ground beef. *Carcinogenesis* 1984: 5: 95-102.
- Fowler J.E. Jr., Bigler S.A., Kilambi N.K., and Land S.A. Relationships between prostate-specific antigen and prostate volume in black and white men with benign prostate biopsies. *Urology* 1999: 53:1175-1178.
- Gerhardsson de Verdier M., Hagman U. Peters R.K., Steineck G., and Övervik E. Meat, cooking methods and colorectal cancer: a case-referent study in Stockholm. *Int. J. Cancer* 1991: 49: 520-525.
- Ghoshal A., Preisegger K.-H., Takayama T., Thorgeirsson S., and Snyderwine E.G. Induction of mammary tumors in female Sprague-Dawley rats by the food-derived carcinogen 2-amino-1-methyl-6-phenylimidazo[4, 5,-b]pyridine and effect of dietary fat. *Carcinogenesis* 1994: 15: 2429-2433.
- Giovannucci E., Rimm E.B., Stampfer M.J., Colditz G.A., and Willett W.C. Height, body weight, and risk of prostate cancer. *Cancer Epidemiol., Biomarkers, Prevent.* 1997: 6: 557-563.
- Giovannucci E., Rimm E.B., Wolk A., Ascherio A., Stampfer M.J., Colditz G.A., and Willett W.C. Calcium and fructose intake in relation to risk of prostate cancer. *Cancer Res.* 1998: 58: 442-447.

- Gooderham N.J., Zhu H., Lauber S., Boyce A., and Creton S. 2002. Molecular and genetic toxicology of 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP). *Mutat. Res.* 506-507:91-99.
- Gerstenbluth R.E., Seftel A.D., Hampel N., Oefelein M.G., Resnick M.I. The accuracy of the increased prostate specific antigen level (greater than or equal to 20 ng./ml.) in predicting prostate cancer: is biopsy always required? *J. Urol.* 2002: 168: 1990-1993.
- Hayes R.B., Ziegler R.G., Gridley G., Swanson C., Greenberg R.S., Swanson G.M., Schoenberg J.B., Silverman D.T., Brown L.M., Pottern L.M., Liff J., Schwartz A.G., Fraumeni Jr. J.F., and Hoover R.N. Dietary factors and risks for prostate cancer among blacks and whites in the United States. *Cancer Epidemiol. Biomark. Prev.* 1999: 8: 25-34.
- Hemminki K., and Chen B. Familial association of prostate cancer with other cancers in the Swedish Family-Center Database. *Prostate* 2005: 65: 188-194.
- Hemminki K., and Czene K. Age specific and attributable risks of familial prostate carcinoma from the family-cancer database. *Cancer* 2002: 95: 1346-1353.
- Hsing A.W., Tsao L., and Devesa S.S. International trends and patterns of prostate cancer incidence and mortality. *Int. J. Cancer* 2000: 85: 60-67.
- Hsing A.W. Hormones and prostate cancer: What's next? *Epidemiol. Rev.* 2001: 23: 42-58.
- Ito N., Hasegawa R., Imaida K., Tamano S., Hagiwara A., Hirose M., and Shirai T. Carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) in the rat. *Mutat. Res.* 1997: 376: 107-114.
- Ito N., Hasegawa R., Sano M., Tamano S., Esumi H., Takayama-S., and Sugimura T. A new colon and mammary carcinogen in cooked food, 2-amino-1-methyl-6-phenylimidazo-[4,5-*b*]pyridine (PhIP). *Carcinogenesis* 1991: 12: 1503-1506.
- Kaderlik K.R., Minchin R.F., Mulder G.J., Ilett K.F., Daugaard-Jenson M., Teitel C.H., and Kadlubar F.F. Metabolic activation pathway for the formation of DNA adducts of the carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) in rat extrahepatic tissues. *Carcinogenesis* 1994a: 15: 1703-1709.
- Kaderlik K.R., Mulder G.J., Shaddock J.G., Casciano D.A., Teitel C.H., and Kadlubar F.F. Effect of glutathione depletion and inhibition of glucuronidation and sulfation on 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) metabolism, PhIP-DNA adduct formation and unscheduled DNA synthesis in primary rat hepatocytes. *Carcinogenesis* 1994b: 15: 1711-1716.
- Kampman E., Slattery M.L., Bigler J., Leppert M., Samowitz W., Caan B.J., and Potter J.D. Meat consumption, genetic susceptibility, and colon cancer risk: a United States multicenter case-control study. *Cancer Epidemiol. Biomark. Prev.* 1999: 8: 15-24.
- Keating G.A., and Bogen K.T. Methods to estimate heterocyclic amine concentrations in cooked meats in the U.S. diet. *Food Chem. Toxicol.* 2001: 39: 29-43.
- Keating G.A., Bogen K., and Chan J. . Development of a meat frequency questionnaire for use in diet and cancer studies. *J. Am. Dietetic Assoc.* 2007: (in press).
- Keating G.A., Layton D.W., and Felton J.S. Factors determining dietary intakes of heterocyclic amines in cooked foods. *Mutat. Res.* 1999: 443: 149-156.
- Keating G.A., Sinha, R., Layton, D., Salmon, C.P., Knize, M.G., Bogen, K.T., Lynch, C.F., and Alavanja, M. Comparison of heterocyclic amine levels in home-cooked meats with exposure indicators. *Cancer, Causes and Control* 2000: 11: 731-739.
- Kendall M., and Stuart A. *The Advanced Theory of Statistics, Vol. 2.* MacMillan Publishing Co.: New York, NY, pp. 159-60, 1979.
- Kidd L.R., Stillwell W.G., Yu M.C., Wishnok J.S., Skipper P.L., Ross R.K., Henderson B.E., and Tannenbaum S.R. Urinary excretion of 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) in White, African American, and Asian-American men in Los Angeles County. *Cancer Epidemiol. Biomark. Prev.* 1999: 8: 439-445.
- Kristal A.R., and Lampe J.W. Brassica vegetables and prostate cancer risk: A review of the epidemiological evidence. *Nutr. Cancer* 2002: 42: 1-9.
- Malfatti. M.A., Kulp K.S., Knize M.G., Davis C., Massengill J.P., Williams S., Nowell S., MacLeod S., Dingley K.H., Turteltaub K.W., Lang N.P., and Felton, J.S. The identification of [2-¹⁴C]2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine metabolites in humans. *Carcinogenesis* 1999: 20: 705-713.

- Kolonel L.N., Nomura A.M.Y., and Cooney R.V. Dietary fat and prostate cancer: Current status. *J. Natl. Cancer Inst.* 1999: 91: 414-28.
- Lang N.P., Butler M.A., Massengill J., Lawson M., Stotts R.C., Hauer-Hensen M., and Kadlubar F.F. Rapid metabolic phenotypes for acetyltransferase and cytochrome P4501A2 and putative exposure to food-borne heterocyclic amines increase the risk for colorectal cancer or polyps. *Cancer Epidemiol. Biomark. Prev.* 1994: 8: 675-682.
- Lawson T., and C. Kolar. Human prostate epithelial cells metabolize chemicals of dietary origin to mutagens. *Cancer Lett.* 2002: 175: 141-146.
- Lyon J.L., and Mahoney A.W. Fried foods and the risk of colon cancer. *Am. J. Epidemiol.* 1988: 128: 1000-1006.
- Mänttinen L., Auvinen A., Stenman U.-H., Tammela T., Rannikko S., Aro J., Juusela H., Hakama M., Three-year results of the Finnish Prostate Cancer Screening Trial *J. Natl. Cancer Inst.* 2001: 93: 552-553.
- Martin F.L., Cole K.J., Muir G.H., Kooiman G.G., Williams J.A., Sherwood R.A., Grover P.L., and Phillips D.H. Primary cultures of prostate cells and their ability to activate carcinogens *Prostate Cancer Prostatic Diseases* 2002: 5: 96-104.
- McManus M.E., Burgess W.M., Veronese M.E., Hugget A., Quattrochi L.C., and Tukey R.H. Metabolism of 2-acetylaminofluorene and benzo(a)pyrene and activation of food-derived heterocyclic amine mutagens by cytochromes P450. *Cancer Res.* 1990: 50: 3367-3376.
- Mettlin C. Recent developments in the epidemiology of prostate cancer. *Eur. J. Cancer* 1997: 33: 340-347.
- Miller B.A., Kolonel L.N., Bernstein L., Young Jr. J.L., Swanson G.M., West D., Key C.R., Liff J.M., Glover C.S., Alexander G.A. (eds). *Racial/Ethnic Patterns of Cancer in the United States 1988-1992*. National Cancer Institute. NIH Pub. No. 96-4104: Bethesda, MD, 1996.
- Mills P.K., Beeson W.L., Phillips R.L., and Fraser G.E. Cohort of diet, lifestyle and prostate cancer in Adventist men. *Cancer* 1989: 64: 598-604.
- Minami Y., M.P. Staples, and G.G. Giles. The incidence of colon, breast and prostate cancer in Italian migrants to Victoria, Australia. *Eur. J. Cancer* 1993: 29A: 1735-1740.
- Murtaugh M.A., Ma K., Sweeney C., Caan B.J. and Slattery M.L. Meat consumption patterns and preparation, genetic variants of metabolic enzymes, and their association with rectal cancer in men and women. *J. Nutr.* 2004: 134: 776-784.
- Muscat J.E., and Wynder E.L. The consumption of well-done red meat and the risk of colorectal cancer. *Am. J. Public Health* 1994: 84: 856-858.
- Norrish A.E., Ferguson L.R., Knize M.G., Felton J.S., Sharpe S.J., and Jackson R.T. Heterocyclic amine content of cooked meat and risk of prostate cancer. *J. Natl. Cancer Inst.* 1999: 91: 2038-2044.
- Ochiai M., Ogawa K., Wakabayashi K., Sugimura T., Nagase S., Esumi H., and Nagao M. Induction of intestinal adenocarcinomas by 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine in Nagase albuminemic rats. *Jpn. J. Cancer Res. (Gann)* 1991: 82: 363-366.
- Oesterling J.E., Jacobsen S.J., Chute C.G., Guess H.A., Girman C.J., Panser L.A., and Lieber M.M. Serum prostate-specific antigen in a community-based population of healthy men. *J. Am. Med. Assoc.* 1993: 270: 860-864.
- Ohgaki H., Hasegawa H., Kato T., Suenaga M., Ubakata M., Sato S., Takayama S., and Sugimura T. Carcinogenicity in mice and rats of heterocyclic amines in cooked foods. *Environ. Health Perspect.* 1986: 67: 129-134.
- Pettaway C.A. Racial differences in the adrogen/androgen receptor pathway in prostate cancer. *J. Natl. Med. Assoc.* 1999: 91: 652-660.
- Probst-Hensch N.M., Sinha R., Longnecker M.P., Witte J.S., Ingles S.A., Frankl H.D., Lee E.R., and Haile R.W. Meat preparation and colorectal adenomas in a large sigmoidoscopy-based case-control study in California (United States). *Cancer Causes Control* 1997: 8: 175-183.
- Robbins A.S., Whittemore A.S., and Van Den Eden S.K. Race, prostate cancer survival, and membership in a large health maintenance organization. *J. Natl. Cancer Inst.* 1998: 90: 986-90.

- Rosenkranz H.S., and Mermeistein R. The genotoxicity, metabolism and carcinogenicity of nitrated polycyclic aromatic hydrocarbons. *J. Environ. Sci. Health* 1985: C3: 221-72.
- Ross R.K., Pike M.C., Coetzee G.A., Reichhardt J.K.V., Yu M.C., Feigelson H., Stanczyk F.Z., Kolonel L.N., and Henderson B.E. Androgen metabolism and prostate cancer: Establishing a model of genetic susceptibility. *Cancer Res.* 1998: 58: 4497-504.
- Sato S., Negishi C., Umemoto A., and Sugimura T. Metabolic aspects of pyrolysis mutagens in food. *Environ. Health Perspect.* 1986: 67: 105-9.
- Schurrman A.G., Zeegers P.A., Goldbohm R.A., and van den Brandt P.A. A case-control study on prostate cancer risk in relation to family history of prostate cancer. *Epidemiol.* 1999: 10: 192-195.
- Schut H.A.J., and Snyderwine E.G. DNA adducts of heterocyclic amine food mutagens: implications for mutagenesis and carcinogenesis. *Carcinogenesis* 1999: 20: 353-68.
- Selvin S. *Practical Biostatistical Methods*. Duxbury Press: New York, NY, 1995.
- Shiffman M.H., and Felton J.S. Re "Fried foods and the risk of colon cancer" [letter]. *Am. J. Epidemiol.* 1990: 131: 376-8.
- Shirai T., Cui L., Takahashi S., Futakuchi M., Asamoto M., Kato K., and Ito N. Carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in the rat prostate and induction of invasive carcinomas by subsequent treatment with testosterone propionate. *Cancer Lett.* 1999: 143: 217-21.
- Shirai T., Sano M., Tamano S., Takahashi S., Hirose M., Futakuchi M., Hasegawa R., Imaida K., Matsumoto K., Wakabayashi K., Sugimura T., and Ito N. The prostate: a target for carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) derived from cooked foods. *Cancer Res.* 1997: 57: 195-8.
- Sinha R., Rothman N., Brown E.D., Salmon C.P., Knize M.G., Swanson C.A., Rossi S.C., Mark S.D., Levander O.A., and Felton J.S. High concentrations of the carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine occur in chicken but are dependent on the cooking method. *Cancer Res.* 1995: 55: 4516-4519.
- Sinha R., Kulldorff M., Curtin J., Brown C.C., Alavanja M.C., and Swanson C. Fried, well-done red meat and risk of lung cancer in women (United States). *Cancer Causes Control* 1998a: 9: 621-630.
- Sinha R., Rothman N., Salmon C.P., Knize M.G., Brown E.D., Swanson C.A., Rhodes D., Felton J.S., and Levander O.A. Heterocyclic amine content in beef cooked by different methods and to varying degrees of doneness and gravy made from meat drippings. *Food Chem. Toxicol.* 1998b: 36: 279-287.
- Sinha R., Knize M.G., Salmon C.P., Brown E.D., Rhodes D., Felton J.S., Levander O.A., and Rothman N. Heterocyclic amine content of pork products cooked by different methods and to varying degrees of doneness. *Food Chem. Toxicol.* 1998c: 36: 289-297.
- Sinha R., Chow W.H., Kulldorff M., Denobile J., Butler J., Garcia-Closas M., Weil R., Hoover R.N., and Rothman N. Well-done, grilled red meat increases the risk of colorectal adenomas. *Cancer Res.* 1999a: 59: 4320-4324.
- Sinha R., and Rothman N. Role of well-done, grilled red meat, heterocyclic amines (HCAs) in the etiology of human cancer. *Cancer Lett.* 1999b: 143: 189-194.
- Smith R.P., Malkowicz S.B., Whittington R. VanArsdalen K., Tochner Z., Wein A.J. Identification of clinically significant prostate cancer by prostate-specific antigen screening. *Arch. Intern Med.* 2004: 164: 1227-1230.
- Snyderwine E.G., and Battula N. Selective mutagenic activation by cytochrome P3-450 of carcinogenic arylamines found in foods. *J. Natl. Cancer Inst.* 1989: 81: 223-7.
- Stuart, G. R., Holcroft, J., de Boer, J. G. and Glickman, B. W. (2000). Prostate mutations in rats induced by suspected human carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP). *Cancer Res.* 60:266-268.
- Takahashi S., Tamano S., Hirose M., Kimoto N., Ikeda Y., Sakakibara M., Tada M., Kadlubar F.F., Ito N., and Shirai T. Immunohistochemical demonstration of carcinogen-DNA adducts in tissues of rats given 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP): detection in paraffin-embedded sections and tissue distribution. *Cancer Res.* 1998: 58: 4307-4313.
- Talamini R., Franceschi S., La Vecchia C., Serraino D., and Negri E. Diet and prostatic cancer: a case-control study in northern Italy. *Nutr. Cancer* 1992: 18: 277-286.

- Thompson I.M., Pauler D.K., Goodman P.J., Tangen C.M., Lucia M.S., Parnes H.L., Minasian L.M., Ford L.G., Lippman S.M., Crawford E.D., Crowley J.J., Coltman C.A. Jr. Prevalence of prostate cancer among men with a prostate-specific antigen level ≤ 4.0 ng per milliliter. *N. Engl. J. Med.* 2004; 350: 2239-2246.
- Thompson L.H., Tucker J.D., Stewart S.A., Christensen M.L., Salazar E.P., Carrano A.V., and Felton J.S. Genotoxicity of compounds from cooked beef in repair-deficient CHO cells versus salmonella mutagenicity. *Mutagen.* 1987; 2: 483-487.
- Thorgeirsson S.S. Metabolic determinants in the carcinogenicity of aromatic amines. In: Greim H., Jung R., Kramer M., Marquardt H., and Oesch F. eds. *Biochemical Basis of Chemical Carcinogenesis*. Raven Press: New York 1984: 47-56.
- Thorgeirsson S.S., Glowinski I.B., and McManus M.E. Metabolism, mutagenicity and carcinogenicity of aromatic amines. In: Hodgson E., Bend J.R., and Philpot R.M. eds. *Reviews in Biochemical Toxicology* 5. Elsevier Biomedical: New York-Amsterdam-Oxford 1983: 349-386.
- Tomlins S.A., Rhodes D.R., Perner S., Dhanasekaran M., Mehra R., Sun X.-W., Varambally S., Cao X., Tchinda J., Kuefer R. Lee C., Montie J.E., Shah R.B., Pients K.J., Rubin M.A., Chinnaiyan A.M. Recurrent fusion of *TMPRSS2* and ETS transcription factor genes in prostate cancer. *Science* 2005; 310: 644-648.
- Turesky R.J., Lang N.P., Butler M.A., Teitel C.H., and Kadlubar F.F. Metabolic activation of carcinogenic heterocyclic aromatic amines by human liver and colon. *Carcinogenesis* 1991; 12: 1839-1845.
- U.S. Department of Agriculture (USDA). 1989-91 Continuing Survey of Food Intakes by Individuals and 1989-91 Diet and Health Knowledge Survey (CSFII/DHKS 1989-91 Data Set). CD-ROM. USDA Agricultural Research Service, Beltsville Human Nutrition Research Center, Food Surveys Research Group. 1993.
- U.S. Department of Agriculture (USDA). 1994-96 Continuing Survey of Food Intakes by Individuals and 1994-96 Diet and Health Knowledge Survey (CSFII/DHKS 1994-96 Data Set). CD-ROM. USDA Agricultural Research Service, Beltsville Human Nutrition Research Center, Food Surveys Research Group. 1998.
- Ward M.H., Sinha R., Heinman E.F., Rothman N., Markin R., Weisenburger D.D., Correa P., and Zahm S.H. Risk of adenocarcinoma of the stomach and esophagus with meat cooking method and doneness preference. *Int. J. Cancer* 1997; 71: 14-19.
- Whittemore A.S., Kolonel L.N., Wu A.H., John E.M., Gallagher R.P., Howe G.R., Burch J.D., Hankin J., Dreon D.M., West D.W., Teh C.-Z., and Paffenbarger J., R.S. Prostate cancer in relation to diet, physical activity, and body size in Blacks, Whites, and Asians in the United States and Canada. *J. Natl. Cancer Inst.* 1995; 87: 652-661.
- Williams J.A., Martin F.L., Muir G.H., Hewer A., Grover P.L., Phillips D.H., and Muir G.H. Metabolic activation of carcinogens and expression of various cytochromes P450 in human prostate tissue. *Carcinogenesis* 2000; 21: 1683-1689.
- Wilcox, R.R. Some practical reasons for reconsidering the Kolmogorov-Smirnov test. *Br. J. Math. Stat. Psychol.* 1997; 50: 9-20.
- Wolfram S. *Mathematica Book*, 4th ed. Cambridge University Press: Cambridge, UK, 1999.
- Zeegers M.P., Jellema A., and Ostrer H. Empiric risk of prostate carcinoma for relatives of patients with prostate carcinoma: a meta-analysis. *Cancer* 2003; 97: 1894-1903.
- Zelterman, D. *Models for Discrete Data*. Clarendon Press, Oxford, UK, 1999: 6-24.

Appendix

Bogen, KT, J Chan, GA Keating, LJ Paine, EL Simms, EA Holly, and JS Felton. 2006. Prostate-specific antigen levels and dietary PhIP in African Americans: A prospective clinic-based study. [Abstract] 2007 Annual Meeting of the American Association for Cancer Research, 14-18 April 2007, Los Angeles, CA. [UCRL-ABS-226551]

Bogen KT, GA Keating II, JM Chan, LJ Paine, EL Simms, DO Nelson, and EA Holly. 2007. Highly elevated PSA and dietary PhIP intake in a prospective clinic-based study among African Americans. *Prostate Cancer and Prostatic Diseases* (in press).

[these publications follow]

Prostate-specific antigen levels and dietary PhIP in African Americans: A prospective clinic-based study

Bogen, KT¹, J Chan², GA Keating¹, LJ Paine³, EL Simms³, EA Holly², and JS Felton¹

¹Lawrence Livermore National Laboratory; ²University of California San Francisco Medical Center; ³Alta Bates Summit Medical Center, Oakland, CA

Heterocyclic amines (HAs)—potent mutagens formed as red meat, chicken or fish cooks—cause cancer at multiple sites (including rat prostate) in rodent bioassays, and have been linked to elevated human risk of colon and other cancers. Compared to white men in the U.S., African American (AA) men have about twice the prostate cancer (PC) incidence and about twice the daily intake of 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP), the predominant HA in the U.S. diet. To investigate the hypothesis that dietary PhIP exposure increases PC risk, an ongoing prospective clinic-based study has compared PC-screening outcomes with survey-based estimates of dietary PhIP intake by 40- to 70-year-old AA men in the Oakland, CA, area. Participants with no prior PC diagnosis, recruited through a cancer education center/screening clinic, complete food-frequency and meat-cooking/consumption questionnaires, and have a prostate-specific antigen (PSA) test and digital-rectal exam. Preliminary results for 562 participants indicate that mean (± 1 SD) daily intake of PhIP, the major HA found in cooked meats, in this group is 19 (± 24) ng kg⁻¹ d⁻¹, which is ~2-fold (and ~3-fold) greater than a national estimate of mean PhIP intake for AA (and white U.S.) men of similar age. In the present study, estimated PhIP intakes were found to be attributable mostly (65%) to chicken and positively associated ($R^2 = 0.25$, $p \sim 0$) with estimated saturated fat (SF) intake (a previously hypothesized environmental PC-risk factor). An odds ratio, OR, (and maximum-likelihood 95% confidence limits) of 23.6 (2.20, 533.) for PSA ≥ 20 ng/mL was observed for those in the highest 15% compared to the lower 50% of estimated daily PhIP intakes (≥ 32 vs. ≤ 4.8 ng kg⁻¹ d⁻¹), with a p-value (p_{trend}) of 0.0023 for a chi-square test for trend done across three (including these two) PhIP-intake groups (extended Fisher exact test p-value = 0.0063). This positive trend persisted after separate adjustments for self-reported family (brother or father) history of PC (FH), SF intake, and energy intake ($p_{\text{trend}} = 0.0024$, 0.012, and 0.0032, respectively). PSA measures were found (by Kolmogorov 2-sample tests) to be significantly higher in AA men reporting a positive FH in this study ($p = 0.007$), particularly for those among the highest PSA-measure quartile in each FH group ($p < 0.0002$). These preliminary results are consistent with a positive association between PhIP intake and highly elevated PSA levels, supporting the hypothesis that diet and food preparation interventions may help reduce PC risk in AA and perhaps other groups.

[Work performed under auspices of the U.S. Department of Energy by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48, and the University of California San Francisco, with funding by the National Cancer Institute (NIH grant P01 CA55861-01) and Department of Defense (PCRP grant PC040371).]

To be presented at the 2007 Annual Meeting of the American Association for Cancer Research, 14-18 April 2007, Los Angeles, CA.

ORIGINAL ARTICLE

Highly elevated PSA and dietary PhIP intake in a prospective clinic-based study among African Americans

KT Bogen¹, GA Keating II¹, JM Chan², LJ Paine³, EL Simms³, DO Nelson¹ and EA Holly²

¹Lawrence Livermore National Laboratory, Energy and Environment Directorate, University of California, Livermore, CA, USA;

²Department of Epidemiology and Biostatistics, University of California San Francisco, San Francisco, CA, USA and ³Alta Bates Summit Medical Center, Markstein Cancer Education and Prevention Center, Oakland, CA, USA

African-American men die from prostate cancer (PC) nearly twice as often as white US men and consume about twice as much of the predominant US dietary heterocyclic amine, 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP), a genotoxic rat-prostate carcinogen found primarily in well-cooked chicken and beef. To investigate the hypothesis that PhIP exposure increases PC risk, an ongoing prospective clinic-based study compared PC screening outcomes with survey-based estimates of dietary PhIP intake among 40–70-year-old African-American men with no prior PC in Oakland, CA. They completed food-frequency and meat-cooking/consumption questionnaires and had a prostate-specific antigen (PSA) test and digital-rectal exam. Results for 392 men indicated a 17 (± 17) ng/kg day mean (± 1 s.d.) daily intake of PhIP, about twice that of white US men of similar age. PhIP intake was attributable mostly to chicken (61%) and positively associated ($R^2 = 0.32$, $P < 0.0001$) with saturated fat intake. An odds ratio (95% confidence interval) of 31 (3.1–690) for highly elevated PSA ≥ 20 ng/ml was observed in the highest 15% vs lowest 50% of estimated daily PhIP intake (≥ 30 vs ≤ 10 ng/kg day) among men 50+ years old ($P = 0.0002$ for trend) and remained significant after adjustment for self-reported family history of (brother or father) PC, saturated fat intake and total energy intake. PSA measures were higher in African-American men with positive family history ($P = 0.007$ all men, $P < 0.0001$ highest PSA quartile). These preliminary results are consistent with a positive association between PhIP intake and highly elevated PSA, supporting the hypothesis that dietary intervention may help reduce PC risk.

Prostate Cancer and Prostatic Diseases advance online publication, 16 January 2007; doi:10.1038/sj.pcan.4500941

Keywords: cooked meat; diet; energy; heterocyclic amines; mutagens; race

Introduction

Heterocyclic amines (HAs) are potent mutagens formed in meats, chicken and fish as it is cooked to higher-doneness levels by heat-intensive cooking methods.^{1–3} HAs also cause cancer in a variety of tissue types in multiple bioassays in different animal species, strains and sexes, and so may present a human dietary cancer risk.^{4,5} A predominant HA found particularly in well-done chicken and beef is 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP).^{6–11} Dietary exposure to PhIP has been shown to induce colon, intestinal and mammary adenocarcinomas in rats^{12–18} and prostate cancer in rats.^{19,20} HAs including PhIP are positive in numerous genotoxicity assays.^{21,22} Like other HAs, PhIP is metabolically activated by P450 and N-acetyltransferase enzymes to genotoxic metabolites, which in the case of PhIP mutate both rat and human prostate DNA.^{21,23–40}

PhIP also has potent estrogenic activity that may act synergistically with genotoxic effects to enhance its carcinogenicity.⁴¹

In the US, prostate cancer (PC) is a leading cause of cancer death among men, with African Americans having the highest age-specific prostate cancer rate in the world, and a >2-fold higher rate of mortality for PC than white men in the US.^{42–44} Although family history of PC (particularly father/brother) is clearly linked to substantially elevated PC risk^{45–48} and a human-PC-specific chromosome translocation has been identified,⁴⁹ there is no evidence for a predominant heritable factor for PC. Racial-ethnic differences in testosterone-related hormone and/or receptor levels may also affect racial-ethnic differences in PC risk, but their role remains unclear.^{50–52}

Substantial geographical variations in PC incidence indicate the importance of one or more dietary or other environmental factors.^{53–55} Dietary intake of animal or saturated fat often has been linked to increased PC risk in previous studies, including among African-American men. However, these associations appear too weak to explain more than a small fraction of observed racial-ethnic differences in PC risk.^{56–59} This also

Correspondence: Dr KT Bogen, Lawrence Livermore National Laboratory, Energy and Environment Directorate L-396, University of California, 7000 East Avenue, Livermore, CA 94550-9900, USA.
E-mail: bogen@LLNL.gov
Received 18 October 2006; revised and accepted 6 December 2006

appears to be the case for other environmental/dietary factors examined such as calcium, cruciferous vegetables, vitamin D, UV from sunlight, lycopene and body size.^{60–66} Because cooked-meat intake is positively associated with total saturated fat intake,⁶⁷ previous studies that focused on animal or saturated fat *per se* may have estimated effects due largely or entirely to meat-related HA intake.

Consumption of cooked meats and associated HAs has been linked to increased risks of colorectal adenoma/adenocarcinoma and of cancer of the stomach, breast, lung and prostate,^{68–90} whereas fewer studies have reported no such associations.^{89,91–94} One study⁹⁴ quantified HA-specific intake estimates (including for PhIP), but detected no positive association of consumption of any HA (from white or red meat) with elevated risk of breast cancer, regardless of *N*-acetyltransferase genotype. Other positive studies have tended to include those with HA exposure quantified after adjusting for factors expected to determine HA intake, namely, meat type, consumption frequency, cooking method and cooking doneness. For example, recently published results from a study that used the Block Brief 2000 survey to assess meat-related exposures detected elevated PC risk among 692 African-American men (but not among nearly 65 000 white men) who consumed relatively more cooked red or processed meats.⁸⁹

Periodic screening for serum levels of prostate-specific antigen (PSA) is useful to detect early PC and benign prostatic hyperplasia (BPH) because PSA levels in serum are positively associated with age, prostate volume and prostatic neoplastic disease.^{95–97} Significantly higher PSA levels are found in African Americans than in white subjects, even after adjustment for age and prostate volume in men without PC, and for PC grade and stage in men with PC.^{98–100} This is due to greater PSA secretion per unit prostate volume by African-American men.¹⁰¹ Although a substantial fraction of 'slightly' elevated PSA levels (between 4 and 10 ng/ml) is attributable to BPH or infection, 'highly elevated' PSA levels (≥ 20 ng/ml) typically indicate a strong likelihood of localized or metastatic PC, with $>99\%$ specificity^{102–104} and with $>95\%$ positive predictivity for men of age 50–69 years.¹⁰⁵ Likewise, even among men with a PSA measure <4.0 ng/ml (often considered within the 'normal' range), it is estimated that about 15% later may be diagnosed with PC.¹⁰⁶

To investigate further whether PhIP intake is related to PC risk, we conducted a study to assess the association between highly elevated PSA as a screening indicator for elevated PC risk and estimated dietary exposure to PhIP among African-American men.

Methods

Study design and participants

In an ongoing clinic-based prospective study using Institutional Review Board-approved human participant protocols, we enrolled 392 African-American men from the Oakland, CA area using the following inclusion criteria: (1) African-American men between 40 and 70 years old; (2) no previous PC diagnosis or contra-indicated medical condition; and (3) written informed

consent. Participation was enhanced by a \$30 incentive payment and by >10 years of previous PC-related community outreach undertaken by the study clinic (the Markstein Cancer Education and Prevention Center at Alta Bates Summit Medical Center in Oakland, CA). After providing written informed consent, each participant completed a PC screening medical questionnaire, answered general and detailed meat-related dietary questions and was provided free PC screening including a PSA blood test and a digital-rectal exam (DRE) by a board-certified urologist.

Oakland is a metropolitan city near San Francisco within California's Alameda County, in which prostate cancer is the most common cancer in African-American men with about 920 new cases predicted in 2006.¹⁰⁷ In 2004, Alameda County had a 14.1% African-American population, whereas in Oakland, the site of the medical center where our study was conducted, 35.7% of the population was African American, and African-American populations statewide and nationwide in 2004 were 6.8 and 12.8%, respectively.¹⁰⁸ African-American men in the Alameda County recruitment area have a median age of 34 years, median per capita income of about \$34,700 and a currently married rate of 39%.¹⁰⁸ Outreach was tailored to African-American men aged 40–70 in Alameda County reflecting its current socio-economically diverse African-American population, including a large fraction of unmarried men who do not benefit from positive spousal influences on health-care choices. Outreach was conducted at more sites frequented by men including barbershops catering to African Americans, social service organizations, African-American churches, senior centers and housing units, Veterans Administration outpatient clinics, community health clinics and medical practices of members of the Sinkler-Miller medical association, a professional association of African-American physicians.

Dietary assessment

General dietary intake over the previous year was estimated using the Block-2000 food-frequency questionnaire (FFQ) with portion-size standardized food-model photographs to help each participant select portion sizes.¹⁰⁹ Dietary data on specific meats consumed and preferred cooking methods over the previous year were obtained using an additional questionnaire that includes a validated set of standard meat-doneness descriptors and corresponding set of meat-doneness photographs.^{110–112} All dietary questionnaire data were obtained by in-person interviews administered by trained dietary interviewers.

Data analysis

Combined survey data were used as previously described^{113–115} to estimate annual average dietary PhIP intake from all sources by each participant. Total and basal energy intake in kcal per kg body weight was estimated for each study participant using previously described methods.¹¹³ Standard methods were used to assess the significance of linear associations and outliers where noted, and to assess Pearson product-moment correlations.^{116,117} Approximate significance of differences in mean HA intake was compared using

Welch's *t*-test whereas unequal variance was assessed by corresponding F-tests. Differences in PSA by family history were assessed by the Wilcoxon test.¹¹⁸ Odds ratio (OR) and corresponding 95% confidence interval (CI) estimates obtained by numerical maximum-likelihood procedures are reported together with corresponding χ^2 tests for trend (with or without adjustment for specified factors).¹¹⁹ Reported Fisher's exact *P*-values are for two-tailed tests. Significance *P*-values $\leq 10^{-10}$ were reported as being approximately 0, and values < 0.10 were reported to one significant figure. All calculations were carried out using *Mathematica* 5.2 software.¹²⁰

Results

PhIP intake

Data on 392 African-American men who participated in this study are summarized in Table 1. Corresponding estimated average daily intake of specific meats and total PhIP is summarized in Table 2. The empirical distribution of estimated daily intake of PhIP from all meats (total PhIP) had geometric and arithmetic mean values of 9.6 and 17 ng/kg day, respectively, and a geometric standard deviation (s.d.) of 3.3. A total of 89% of inter-individual variance in PhIP intake could be explained by intake of specific meats and the cooking method of those meats, and doneness-preference data appeared to explain the remaining 11% of inter-individual variation in PhIP intake. Ratios of total to basal daily intake rates of energy per unit body weight estimated for this study population had an arithmetic mean (± 1 standard error of the mean (s.e.m.)) of 1.57 (± 0.046), not significantly

different ($P = 0.51$) from the value of 1.6 expected for reference adult men.¹¹³

Correlation between PhIP and other dietary measures

Estimated daily intake of total PhIP explained approximately 32% of observed inter-individual variance in corresponding estimated intake of saturated fat per unit body weight (Figure 1). Similar or greater levels of positive correlation were observed between estimated total PhIP intake and energy intake ratio ($E_{\text{food}}:E_{\text{basal}}$) ($R^2 = 0.26$), total energy intake (E_{food}) ($R^2 = 0.27$) and total meat intake (g/kg day) ($R^2 = 0.68$). There were strong correlations between estimated intake of total energy and saturated fat ($R^2 = 0.84$) and between total energy and $E_{\text{food}}:E_{\text{basal}}$ ($R^2 = 0.97$).

PSA, age and family history

PSA measures were weakly positively associated with participant age and attained statistical significance for all measures < 4 ng/ml ($R^2 = 0.051$, $P = 0.00001$), but not for

Table 2 Summary of meat-specific PhIP intake for 392 study participants in an Oakland, CA study on diet and prostate screening

	Meat type						
	Chicken	Burger	Beef	Pork	Fish	Bacon	All
Mean ^a	10.3	2.5	2.4	0.3	0.4	<0.1	16.7
s.d. ^a	13.0	3.7	4.3	0.6	1.0	0.1	17
s.d.m. ^a	0.6	0.2	0.2	0.03	0.05	0.01	0.8
CVM%	6.2	7.4	9.1	8.2	15.0	30.0	5.1
%all	61.0	15.0	14.0	2.2	2.2	0.1	100.0

Abbreviations: s.d., standard deviation; s.d.m., standard deviation of the mean; CVM% = $100\% \times (\text{s.d.m.}/\text{mean})$

^aValues listed are in ng/kg day; arithmetic mean values are shown to a value of 0.1 ng/kg day. Meats listed exclude those rarely used or meats that account for relatively little HA, such as lamb and organ meats. The meat components of food mixtures were excluded from those listed individually, but are included in the 'All' category.

Table 1 Baseline characteristics of 392 participants in an Oakland, CA study on diet and prostate cancer screening

Variable ^a	Value(s)	n	Mean
Age (years)	39–50	115	47
	51–60	216	55
	61–70	61	64
	All	392	54
Weight (kg)	54–163	392	86 ^a
BMI (kg/cm ²)	<30	317	25
	≥ 30	75	34
	18–46	392	27
PSA (ng/ml)	<2	328	0.79
	2 to <4	42	3
	4 to <10	10	6
	10 to <20	7	14
	≥ 20	5	47
		PSA < 4	PSA ≥ 4 ^b
DRE	Normal ^b	233	15
	BPH ^b	108	4
	Suspicious ^b	12	3
Family history ^c	No	333	15
	Yes	37	7

Abbreviations: BMI, body mass index; BPH, benign prostatic hyperplasia; DRE, digital-rectal exam; PSA, prostate-specific antigen.

^aWeight median (interquartile range) = 84 (75–95) kg.

^bNormal = no urinary, BPH or other symptoms; suspicious = abnormal DRE result leading to medical referral. Each abnormal PSA result (≥ 4.0 ng/ml) triggered a medical follow-up referral.

^cFamily history, self-reported brother(s) and/or father diagnosed with prostate cancer. Association of positive family history with PSA ≥ 4 ng/ml; Fisher's exact test ($P = 0.007$).




Figure 1 PhIP intake estimated using individual-level doneness-preference data (Y axis, PhIP_{in}), compared with intake estimated using all individuals combined with their meat- and cooking-method-specific average levels of preferred doneness (X axis, PhIP_{inc}). The linear fit of the data shown (dashed line) has an estimated (and corresponding 95% CI) intercept of 0.58 (-1.4 – 0.23) and slope of 1.01 (0.974 – 1.045); an identity relation (solid line) is shown for comparison.

all measures ≥ 4 ng/ml ($R^2 = 0.035$, $P = 0.41$). PSA measures ≥ 20 ng/ml were observed only among older participants, aged 55–65 years. PSA measures among participants reporting vs not reporting a father and/or brother with PC were greater among those reporting such a family history ($P = 0.007$ by Wilcoxon test), particularly when the comparison was restricted to the upper quartile of PSA measures in each family history category ($P = 0.0006$ by Wilcoxon test). Positive family history also was positively associated with elevated PSA defined as ≥ 4 ng/ml ($P = 0.007$; Table 1).

PhIP and PSA

As noted, the above-estimated PhIP intake had a highly skewed distribution; so association between PhIP intake and highly elevated PSA (≥ 20 ng/ml) status was investigated using PhIP intake bin boundaries defined by the 50th, 70th and 85th percentile values of the empirical intake distribution. PhIP intake level was positively associated with highly elevated PSA status, when comparing those in the highest 15% vs the lowest 50% of PhIP consumption, with or without single-variable adjustment for father/brother family history of PC, saturated fat intake or total energy (corresponding P -values for trend: $P_{\text{trend}} = 0.01$ for saturated fat adjust-

ment, $P_{\text{trend}} \leq 0.003$ for all other tests) (Table 3). All highly elevated PhIP measures occurred in men within a fairly narrow age range (55–65 years old). In analyses limited to men aged ≥ 50 years, men in the highest 15% compared with those in the bottom 50% of estimated PhIP intake had a conditional maximum-likelihood odds ratio of 31 (95% CI: 3.1–690) for having a highly elevated PSA ($P_{\text{trend}} = 0.0002$). For this age group, adjustment for family history of PC, saturated fat intake or total energy yielded identical OR estimates and only slightly greater P_{trend} estimates (data not shown).

A similar pattern of positive association was observed between estimated PhIP intake and men with highly elevated PC risk defined as either PSA ≥ 20 ng/ml or a 'suspicious' abnormal DRE result leading to medical referral (data not shown). As expected, men with mildly elevated PSA (≥ 4.0 ng/ml) were about four times more likely to have had a suspicious DRE than did participants with PSA < 4.0 ng/ml, although this difference did not rule out chance ($P = 0.07$, Fisher's exact test). Suspicious DRE results were not obtained for any participant with a PSA measure ≥ 20 ng/ml. PSA level categorized either in quintiles or 50th, 70th, 85th percentile intervals was not associated with saturated fat intake, total energy intake or body mass index, with or without adjustment for PhIP intake ($P_{\text{trend}} > 0.10$ for each).

Table 3 Association of PhIP intake with elevated prostate-specific antigen ≥ 20 ng/ml among 392 African-American men, Oakland, CA

Adjustment or stratification ^a		PSA ≥ 20 ng/ml ^b				
Average PhIP intake ^a PR	(ng kg ⁻¹ day ⁻¹)	m	n	OR	95% CI	P _{trend}
All data						
0–50	4.8	0	196	1		
> 50–70	14.6	1	78	7.6	—	
> 70–85	25.5	1	59	10.0	—	
> 85–100	49.7	3	59	24.0	2.2–540	0.002
Adj. for FH (same as above)						
(m, n, ORs, LCLs and UCLs same as for all data)						0.003
FH–						
0–50	4.8	0	174	1		
> 50–70	14.6	1	69	7.6	—	
> 70–85	25.5	1	53	9.8	—	
> 85–100	49.7	1	52	10.0	—	0.10
FH+						
0–50	4.8	0	22	1		
> 50–70	14.6	0	9	—	—	
> 70–85	25.5	0	6	—	—	
> 85–100	49.7	2	7	18.0	1.4–540	0.002
Adj. for SatFat						
0–50:	4.8	0	196	1		
> 50–70	14.6	1	78	3.4	—	
> 70–85	25.5	1	59	4.1	—	
> 85–100	49.7	3	59	24.0	2.2–540	0.01
Adj. for kcal						
0–50	4.8	0	196	1		
> 50–70	14.6	1	78	14.	—	
> 70–85	25.5	1	59	4.3	—	
> 85–100	49.7	3	59	24.0	2.2–540	0.003

Abbreviations: CI, confidence interval (two-tailed), omitted if interval includes 1; FH, family history of prostate cancer, self-reported father or brother diagnosed; kcal, total energy intake per kg body weight; OR, maximum likelihood odds ratio estimate; PR, percentile range (PR and corresponding mean PhIP intake values pertain to all 392 participants combined); $P_{\text{trend}} = P$ -value for χ^2 test of linear, or (as indicated) adjusted linear, trend.

^aTrend analyses adjusting for Saturated fat or kcal were each performed using the adjustment variable dichotomized at its median value. All men were ≤ 70 years old.

^bm = number with PSA ≥ 20 ng/ml among total n participants included in the analysis.

Discussion

These interim data are consistent with the hypothesis that estimated dietary exposure to PhIP is related to screening indicators of PC risk such as elevated PSA levels or suspicious DRE results among African-American men. Although this conclusion remains preliminary owing to the small number of men in this prospective study to date, it is supported by the consistency of the pattern of results observed and their level of statistical significance. We observed a positive association between elevated PSA and a father/brother history of PC, which is consistent with other studies in the literature that have linked such history to elevated PC risk,^{45–48} and reflects the integrity of the study design.

A potential specific link between PhIP intake and the elevated risk of PC experienced by African-American compared with Caucasian men is suggested by relatively greater levels of PhIP and its metabolites detected in urine sampled from the two groups.¹²¹ However, urine analyses only provide a measure of PhIP exposure within 12–24 h before sampling.¹²² Such a link is supported by studies using meat- and cooking-method-specific HA concentration estimates from multi-laboratory sets of experimental cooking data.^{113–115} These studies estimated intake of PhIP and other HAs among >20 000 US participants (including >3000 African Americans)¹²³ in the nationwide, stratified, random-sample US Continuing Survey of Food Intake by Individuals (CSFII).¹²⁴ Analyzed by age, sex and race/ethnicity, these HA exposure assessments showed that African-American men consumed on average at least twice (and boys through age 15 about three times) as much PhIP (and total HA) per kg body weight per day as did corresponding US Caucasians, and that for both groups there was a small, positive correlation between estimated mean intake of PhIP and of saturated or total fat.^{113,114}

No PhIP-related PC associations were reported in a study of 317 PC cases and 480 age-matched controls that compared HA intake (primarily from cooked lamb) and PC risk in New Zealand men.⁷⁸ They did report an association between PC risk and consumption of beefsteak by the higher cooking-doneness category (OR (95% CI) = 1.8 (1.1–3.0), two-sided $P_{\text{trend}} = 0.008$). The relevance of this finding to potential associations between PC and dietary HA or PhIP intake in the US for African-American men is unknown. The incidence of PC in New Zealand is only half that of Caucasian men in the US.⁴⁴ Average meat (primarily lamb) consumption by participants (~150 g/day) was well below that of US Caucasian men (260 g/day) and even further below that of African-American men (304 g/day) (Table 10A in US Department of Agriculture (USDA)⁶⁷). Moreover, the estimated minimum PhIP intake (224 ng/day) in the *top* exposure quartile of the New Zealand men⁷⁸ was well below the estimated average PhIP intake by Caucasian men (390 ng/day) and that of African-American men (600 ng/day).¹¹³ This difference reflects dissimilar meat consumption and cooking patterns in these two groups. Assuming a linear dose-response, extrapolation from elevated OR estimates ≥ 2.2 obtained in the present study down to PhIP intakes as low as those in the New Zealand study would require that study to have detected an OR as low as ~1.2 with 95% CI at 80% statistical

power, whereas only OR values ≥ 1.6 were detectable in that study.¹²⁴ Alternatively, the lack of a PhIP-related association with PC risk in the New Zealand case-control study could reflect that men from that study (or perhaps all Caucasian men) are, for genetic and/or environmental reasons, less susceptible to the effects of this risk factor than are the African-American participants of our clinic-based study.

Ideally, a prospective study accumulates good diagnostic data and data on exposure- or treatment-related variables of interest. One key limitation of this study is that, despite ongoing work to obtain corresponding follow-up diagnostic data, a PC diagnosis is not yet available for all participants who received either positive or highly elevated PC screening results or who received a 'suspicious' DRE leading to medical referral. Compared to a case-control design, the prospective design used in this ongoing study has the advantage of being double blind, insofar as PC screening results are not known by the participant or by study investigators until after each participant has provided dietary survey data. This design eliminates potential bias in participants' self-reported cooking preferences that may be associated with knowledge of PC screening results or PC status. This is important in view of evidence that prior knowledge of cancer-related status may affect dietary recall and so induce significant differential misclassification.¹²⁵

A second key limitation was the lack of any biomarker generally considered reliable enough to assess chronic PhIP exposures. This required our study, like previous similar studies of HA-related cancer risks, to rely instead on FFQ data. Despite their tendency to slightly underestimate intakes, relative to estimates made using more accurate but also more inconvenient and expensive multiple 24 h food diaries, FFQs have been widely used as the method of choice to assess long-term nutritional exposures and to estimate specific disease risks in relation to such exposures.¹²⁶ Data from self-administered Block dietary FFQs, in particular, were shown to be effective at estimating intakes of each of four major meat-related nutrients (protein, total fat, saturated fat and cholesterol) to within <15% of corresponding estimates obtained using 24 h recall diaries among 226 men who participated in the 1997–1998 'Eating at America's Table' study.¹²⁷ A self-administered meat-specific FFQ including questions on doneness and cooking method preferences (similar to the meat-specific FFQ used in the present study) was more recently compared directly to data from 24 h recall diaries.¹¹² In that study, the FFQ method was shown to underestimate PhIP intakes, but also to classify >60% of individual intakes of each of two HAs (including PhIP) to the same or adjacent quintiles, and to correctly classify lowest vs highest quintiles for >94% of such intakes.¹¹² That study concluded that observed levels of FFQ-associated PhIP intake misclassification would tend to cause underestimation of true relative risks associated with increased PhIP intake estimated from self-administered FFQ data.¹¹² Rates of misclassification of PhIP and other dietary factors were likely reduced in the present study because our FFQs were administered in person by trained interviewers.

In conclusion, we applied methods to estimate HA concentrations in cooked meats based on individually expressed data on meat-specific intake, cooking method and doneness preference to estimate daily PhIP intake.

We observed these intake estimates to be positively associated with screening indicators of highly elevated PC risk in a prospective clinic-based PC screening study with nearly 400 African-American men in the East Bay Oakland, CA area. The observed positive association, even more significant among men 50–70 years of age, remained statistically significant after adjustments for saturated fat intake, total energy intake and self-reported father/brother history of PC. We will continue to accrue participants in this study, expand the screening indicators used to predict PC status and assess whether the observed PhIP-related association pertains to incident PC disease.

Abbreviations

HA, heterocyclic amine; PC, prostate cancer; PhIP, 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (105650-23-5).

Acknowledgements

We appreciate comments on the earlier drafts of this manuscript (Jim Felton and Mark Knize, LLNL; Jennifer Kristiansen, UCSF; an anonymous reviewer) and for interview coordination (Kristiansen). This work was performed under the auspices of the US Department of Energy by University of California, Lawrence Livermore National Laboratory under Contract W-7405-Eng-48, with support by the National Cancer Institute (NIH Grant No. P01 CA55861-01) and the Department of Defense Prostate Cancer Research Program (award no. W81XWH-05-1-0153). Views and opinions of and endorsements by the author(s) do not reflect those of the US Army or the Department of Defense.

References

- Thompson LH, Tucker JD, Stewart SA, Christensen ML, Salazar EP, Carrano AV *et al.* Genotoxicity of compounds from cooked beef in repair-deficient CHO cells versus salmonella mutagenicity. *Mutat Res* 1987; **2**: 483–487.
- Keating GA, Layton DW, Felton JS. Factors determining dietary intakes of heterocyclic amines in cooked foods. *Mutat Res* 1999; **443**: 149–156.
- Keating GA, Sinha R, Layton D, Salmon CP, Knize MG, Bogen KT *et al.* Comparison of heterocyclic amine levels in home-cooked meats with exposure indicators. *Cancer Causes Control* 2000; **11**: 731–739.
- Bogen KT. Cancer potencies of heterocyclic amines found in cooked foods. *Food Chem Toxicol* 1994; **32**: 505–515.
- Layton DW, Bogen KT, Knize MG, Hatch FT, Johnson VM, Felton J. Cancer risk assessment of heterocyclic amines in cooked foods: an analysis and implications for research. *Carcinogenesis* 1995; **16**: 39–52.
- Felton JS, Knize MG, Wood C, Wuebbles BJ, Healy SK, Stuermer DH *et al.* Isolation and characterization of new mutagens from fried ground beef. *Carcinogenesis* 1984; **5**: 95–102.
- Felton JS, Knize MG, Shen NH, Andresen BD, Bjeldanes LF, Hatch FT. Identification of the mutagens in cooked beef. *Environ Health Perspect* 1986; **67**: 17–24.
- Felton JS, Pais P, Salmon CP, Knize MG. Chemical analysis and significance of heterocyclic aromatic amines. *Z Lebensm Unters Forsch A* 1998; **207**: 434–440.
- Felton JS, Knize MG. Heterocyclic-amine mutagens/carcinogens in foods. In: Cooper CS, Grover PL (eds). *Handbook of Experimental Pharmacology*, 94/1. Springer-Verlag: Berlin, 1990, pp 471–502.
- Felton JS, Knize MG. New mutagens from cooked food. In: Pariza M, Aeschbacher H-U, Felton JS, Sato S (eds). *Mutagens and Carcinogens in the Diet*. Wiley-Liss Inc.: New York, 1990, pp 19–38.
- Sinha R, Rothman N, Brown ED, Salmon CP, Knize MG, Swanson CA *et al.* High concentrations of the carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine occur in chicken but are dependent on the cooking method. *Cancer Res* 1995; **55**: 4516–4519.
- Esumi H, Ohgaki H, Kohzen E, Takayama S, Sugimura T. Induction of lymphoma in CDF1 mice by the food mutagen, 2-amino-1-methyl-6-phenylimidazo-[4,5-*b*]pyridine. *Cancer Res* 1989; **80**: 1176–1178.
- Ochiai M, Ogawa K, Wakabayashi K, Sugimura T, Nagase S, Esumi H *et al.* Induction of intestinal adenocarcinomas by 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine in Nagase albuminemic rats. *Jap J Cancer Res (Gann)* 1991; **82**: 363–366.
- Ohgaki H, Takayama S, Sugimura T. Carcinogenicity of heterocyclic amines in cooked foods. *Mutat Res* 1991; **259**: 339–410.
- Ito N, Hasegawa R, Imaida K, Tamano S, Hagiwara A, Hirose M *et al.* Carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) in the rat. *Mutat Res* 1997; **376**: 107–114.
- Ito N, Hasegawa R, Sano M, Tamano S, Esumi H, Takayama S *et al.* A new colon and mammary carcinogen in cooked food, 2-amino-1-methyl-6-phenylimidazo-[4,5-*b*]pyridine (PhIP). *Carcinogenesis* 1991; **12**: 1503–1506.
- Hasegawa R, Sano M, Tamano S, Imaida K, Shirai T, Nagao M *et al.* Carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) in the rat. *Carcinogenesis* 1993; **14**: 2523–2557.
- Ghoshal A, Preisegger K-H, Takayama T, Thorgeirsson S, Snyderwine EG. Induction of mammary tumors in female Sprague-Dawley rats by the food-derived carcinogen 2-amino-1-methyl-6-phenylimidazo[4, 5-*b*]pyridine and effect of dietary fat. *Carcinogenesis* 1994; **15**: 2429–2433.
- Shirai T, Sano M, Tamano S, Takahashi S, Hirose M, Futakuchi M *et al.* The prostate: a target for carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) derived from cooked foods. *Cancer Res* 1997; **57**: 195–198.
- Shirai T, Cui L, Takahashi S, Futakuchi M, Asamoto M, Kato K *et al.* Carcinogenicity of 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP) in the rat prostate and induction of invasive carcinomas by subsequent treatment with testosterone propionate. *Cancer Lett* 1999; **143**: 217–221.
- Gooderham NJ, Zhu H, Lauber S, Boyce A, Creton S. Molecular and genetic toxicology of 2-amino-1-methyl-6-phenylimidazo[4,5-*b*]pyridine (PhIP). *Mutat Res* 2002; **506–507**: 91–99.
- U.S. Department of Health and Human Services (USDHHS). Final Report on Carcinogens Background Document for Selected Heterocyclic Amines: PhIP, MeIQ, and MeIQx. September 12, 2002. Prepared by the Technology Planning and Management Corp for the USDHHS National Toxicology Program, Research Triangle Park, NC. ntp.neihs.nih.gov/ntp/newhomeroc/roc11/HCApub.pdf: 2002.
- Thorgeirsson SS. Metabolic determinants in the carcinogenicity of aromatic amines. In: Greim H, Jung R, Kramer M, Marquardt H, Oesch F (eds). *Biochemical Basis of Chemical Carcinogenesis*. Raven Press: New York, NY, 1984, pp 47–56.
- Thorgeirsson SS, Glowinski IB, McManus ME. Metabolism, mutagenicity and carcinogenicity of aromatic amines. In: Hodgson E, Bend JR, Philpot RM (eds). *Reviews in Biochemical Toxicology* 5. Elsevier Biomedical: New York, Amsterdam, Oxford, 1983, pp 349–386.

- 25 Rosenkranz HS, Mermeistein R. The genotoxicity, metabolism and carcinogenicity of nitrated polycyclic aromatic hydrocarbons. *J Environ Sci Health* 1985; **C3**: 221–272.
- 26 Sato S, Negishi C, Umemoto A, Sugimura T. Metabolic aspects of pyrolysis mutagens in food. *Environ Health Perspect* 1986; **67**: 105–109.
- 27 Kato R, Yamazoe Y. Metabolic activation and covalent binding to nucleic acids of carcinogenic heterocyclic amines from cooked food and amino acid pyrolysates. *Jap J Cancer Res (Gann)* 1987; **81**: 10–14.
- 28 Snyderwine EG, Battula N. Selective mutagenic activation by cytochrome P3-450 of carcinogenic arylamines found in foods. *J Natl Cancer Inst* 1989; **81**: 223–227.
- 29 McManus ME, Burgess WM, Veronese ME, Hugget A, Quattrochi LC, Tukey RH. Metabolism of 2-acetylaminofluorene and benzo(a)pyrene and activation of food-derived heterocyclic amine mutagens by cytochromes P450. *Cancer Res* 1990; **50**: 3367–3376.
- 30 Turesky RJ, Lang NP, Butler MA, Teitel CH, Kadlubar FF. Metabolic activation of carcinogenic heterocyclic aromatic amines by human liver and colon. *Carcinogenesis* 1991; **12**: 1839–1845.
- 31 Davis CD, Schut HAJ, Adamson RH, Thorgeirsson UP, Thorgeirsson SS, Snyderwine EG. Mutagenic activation of IQ, PhIP and MeIQx by hepatic microsomes from rat, monkey and man: low mutagenic activation of MeIQx in cynomolgous monkeys *in vitro* reflects low DNA adduct levels *in vivo*. *Carcinogenesis* 1993; **14**: 61–65.
- 32 Kaderlik KR, Minchin RF, Mulder GJ, Ilett KF, Daugaard-Jenson M, Teitel CH et al. Metabolic activation pathway for the formation of DNA adducts of the carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in rat extrahepatic tissues. *Carcinogenesis* 1994; **15**: 1703–1709.
- 33 Kaderlik KR, Mulder GJ, Shaddock JG, Casciano DA, Teitel CH, Kadlubar FF. Effect of glutathione depletion and inhibition of glucuronidation and sulfation on 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) metabolism, PhIP-DNA adduct formation and unscheduled DNA synthesis in primary rat hepatocytes. *Carcinogenesis* 1994; **15**: 1711–1716.
- 34 Takahashi S, Tamano S, Hirose M, Kimoto N, Ikeda Y, Sakakibara M et al. Immunohistochemical demonstration of carcinogen-DNA adducts in tissues of rats given 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP): detection in paraffin-embedded sections and tissue distribution. *Cancer Res* 1998; **58**: 4307–4313.
- 35 Schut HAJ, Snyderwine EG. DNA adducts of heterocyclic amine food mutagens: implications for mutagenesis and carcinogenesis. *Carcinogenesis* 1999; **20**: 353–368.
- 36 Stuart GR, Holcroft J, de Boer JG, Glickman BW. Prostate mutations in rats induced by suspected human carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP). *Cancer Res* 2000; **60**: 266–268.
- 37 Williams JA, Martin FL, Muir GH, Hewer A, Grover PL, Phillips DH et al. Metabolic activation of carcinogens and expression of various cytochromes P450 in human prostate tissue. *Carcinogenesis* 2000; **21**: 1683–1689.
- 38 Lawson T, Kolar C. Human prostate epithelial cells metabolize chemicals of dietary origin to mutagens. *Cancer Lett* 2002; **175**: 141–146.
- 39 Martin FL, Cole KJ, Muir GH, Kooiman GG, Williams JA, Sherwood RA et al. Primary cultures of prostate cells and their ability to activate carcinogens. *Prostate Cancer Prostatic Dis* 2002; **5**: 96–104.
- 40 Di Paolo OA, Teitel CH, Nowell S, Coles BF, Kadlubar FF. Expression of cytochromes P450 and glutathione S-transferases in human prostate, and the potential for activation of heterocyclic amine carcinogens via acetyl-CoA-, PAPS- and ATP-dependent pathways. *Int J Cancer* 2005; **117**: 8–13.
- 41 Lauber SN, Ali S, Gooderham NJ. The cooked food derived carcinogen 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine is a potent oestrogen: a mechanistic basis for its tissue-specific carcinogenicity. *Carcinogenesis* 2004; **25**: 2509–2517.
- 42 Miller BA, Kolonel LN, Bernstein L, Young Jr JL, Swanson GM, West D et al. (eds). *Racial/Ethnic Patterns of Cancer in the United States 1988–1992*, NIH Pub. No. 96-4104. National Cancer Institute: Bethesda, MD, 1996.
- 43 Robbins AS, Whittemore AS, Van Den Eden SK. Race, prostate cancer survival, and membership in a large health maintenance organization. *J Natl Cancer Inst* 1998; **90**: 986–990.
- 44 Hsing AW, Tsao L, Devesa SS. International trends and patterns of prostate cancer incidence and mortality. *Int J Cancer* 2000; **85**: 60–67.
- 45 Schurrman AG, Zeegers PA, Goldbohm RA, van den Brandt PA. A case-control study on prostate cancer risk in relation to family history of prostate cancer. *Epidemiology* 1999; **10**: 192–195.
- 46 Hemminki K, Czene K. Age specific and attributable risks of familial prostate carcinoma from the family-cancer database. *Cancer* 2002; **95**: 1346–1353.
- 47 Zeegers MP, Jellema A, Ostrer H. Empiric risk of prostate carcinoma for relatives of patients with prostate carcinoma: a meta-analysis. *Cancer* 2003; **97**: 1894–1903.
- 48 Hemminki K, Chen B. Familial association of prostate cancer with other cancers in the Swedish Family-Center Database. *Prostate* 2005; **65**: 188–194.
- 49 Tomlins SA, Rhodes DR, Perner S, Dhanasekaran M, Mehra R, Sun X-W et al. Recurrent fusion of TMPRSS2 and ETS transcription factor genes in prostate cancer. *Science* 2005; **310**: 644–648.
- 50 Hsing AW. Hormones and prostate cancer: what's next? *Epidemiol Rev* 2001; **23**: 42–58.
- 51 Pettaway CA. Racial differences in the androgen/androgen receptor pathway in prostate cancer. *J Natl Med Assoc* 1999; **91**: 652–660.
- 52 Ross RK, Pike MC, Coetzee GA, Reichhardt JKV, Yu MC, Feigelson H et al. Androgen metabolism and prostate cancer: establishing a model of genetic susceptibility. *Cancer Res* 1998; **58**: 4497–4504.
- 53 Minami Y, Staples MP, Giles GG. The incidence of colon, breast and prostate cancer in Italian migrants to Victoria, Australia. *Eur J Cancer* 1993; **29A**: 1735–1740.
- 54 Mettlin C. Recent developments in the epidemiology of prostate cancer. *Eur J Cancer* 1997; **33**: 340–347.
- 55 Angwafo FF. Migration and prostate cancer: an international perspective. *J Natl Med Assoc* 1998; **90** (11 Suppl): S720–S723.
- 56 Whittemore AS, Kolonel LN, Wu AH, John EM, Gallagher RP, Howe GR et al. Prostate cancer in relation to diet, physical activity, and body size in Blacks, Whites, and Asians in the United States and Canada. *J Natl Cancer Inst* 1995; **87**: 652–661.
- 57 Kolonel LN, Nomura AMY, Cooney RV. Dietary fat and prostate cancer: current status. *J Natl Cancer Inst* 1999; **91**: 414–428.
- 58 Hayes RB, Ziegler RG, Gridley G, Swanson C, Greenberg RS, Swanson GM et al. Dietary factors and risks for prostate cancer among blacks and whites in the United States. *Cancer Epidemiol Biomark Prev* 1999; **8**: 25–34.
- 59 Daniels LK, Snetselaar LG, Smith BJ, Stewart RE, Robbins MEC. Problems with the assessment of dietary fat in prostate cancer studies. *Am J Epidemiol* 2004; **160**: 436–444.
- 60 Giovannucci E, Rimm EB, Stampfer MJ, Colditz GA, Willett WC. Height, body weight, and risk of prostate cancer. *Cancer Epidemiol Biomark Prev* 1997; **6**: 557–563.
- 61 Giovannucci E, Rimm EB, Wolk A, Ascherio A, Stampfer MJ, Colditz GA et al. Calcium and fructose intake in relation to risk of prostate cancer. *Cancer Res* 1998; **58**: 442–447.
- 62 Cohen JH, Kristal AR, Stanford JL. Fruit and vegetable intakes and prostate cancer. *J Natl Cancer Inst* 2000; **92**: 61–68.
- 63 Chan J, Giovannucci E. Vegetables, fruits, associated micronutrients, and risk of prostate cancer. *Epidemiol Rev* 2001; **23**: 82–86.
- 64 Chan J, Giovannucci E. Dairy products, calcium, and vitamin D, and risk of prostate cancer. *Epidemiol Rev* 2001; **23**: 87–92.
- 65 Kristal AR, Lampe JW. Brassica vegetables and prostate cancer risk: a review of the epidemiological evidence. *Nutr Cancer* 2002; **42**: 1–9.

- 66 Bodiwala D, Luscombe CJ, Liu S, Saxby M, French M, Jones PW *et al*. Cancer potencies of heterocyclic amines found in cooked foods. *Cancer Lett* 2003; **192**: 145–149.
- 67 U.S. Department of Agriculture (USDA). 1994–96 Continuing Survey of Food Intakes by Individuals and 1994–96 Diet and Health Knowledge Survey (CSFII/DHKS 1994–96 Data Set). CD-ROM. USDA Agricultural Research Service, Beltsville Human Nutrition Research Center, Food Surveys Research Group: Beltsville, MD, 1998.
- 68 Mills PK, Beeson WL, Phillips RL, Fraser GE. Cohort of diet, lifestyle and prostate cancer in Adventist men. *Cancer* 1989; **64**: 598–604.
- 69 Shiffman MH, Felton JS. Re 'Fried foods and the risk of colon cancer'. *Am J Epidemiol* 1990; **131**: 376–378.
- 70 Gerhardtsson de Verdier M, Hagman U, Peters RK, Steineck G, Övervik E. Meat, cooking methods and colorectal cancer: a case-referent study in Stockholm. *Int J Cancer* 1991; **49**: 520–525.
- 71 Talamini R, Franceschi S, La Vecchia C, Serraino D, Negri E. Diet and prostatic cancer: a case-control study in northern Italy. *Nutr Cancer* 1992; **18**: 277–286.
- 72 Lang NP, Butler MA, Massengill J, Lawson M, Stotts RC, Hauer-Hensen M *et al*. Rapid metabolic phenotypes for acetyltransferase and cytochrome P4501A2 and putative exposure to food-borne heterocyclic amines increase the risk for colorectal cancer or polyps. *Cancer Epidemiol Biomark Prev* 1994; **8**: 675–682.
- 73 De Stefani E, Fierro L, Barrios E, Ronco A. Tobacco, alcohol, diet and risk of prostate cancer. *Tumori* 1995; **81**: 315–320.
- 74 Ewings P, Bowie C. A case-control study of cancer of the prostate in Somerset and east Devon. *Br J Cancer* 1996; **74**: 661–666.
- 75 Probst-Hensch NM, Sinha R, Longnecker MP, Witte JS, Ingles SA, Frankl HD *et al*. Meat preparation and colorectal adenomas in a large sigmoidoscopy-based case-control study in California (United States). *Cancer Causes Control* 1997; **8**: 175–183.
- 76 Ward MH, Sinha R, Heinman EF, Rothman N, Markin R, Weisenburger DD *et al*. Risk of adenocarcinoma of the stomach and esophagus with meat cooking method and doneness preference. *Int J Cancer* 1997; **71**: 14–19.
- 77 Kampman E, Slattery ML, Bigler J, Leppert M, Samowitz W, Caan BJ *et al*. Meat consumption, genetic susceptibility, and colon cancer risk: a United States multicenter case-control study. *Cancer Epidemiol Biomark Prev* 1999; **8**: 15–24.
- 78 Norrish AE, Ferguson LR, Knize MG, Felton JS, Sharpe SJ, Jackson RT. Heterocyclic amine content of cooked meat and risk of prostate cancer. *J Natl Cancer Inst* 1999; **91**: 2038–2044.
- 79 Sinha R, Kulldorff M, Curtin J, Brown CC, Alavanja MC, Swanson C. Fried, well-done red meat and risk of lung cancer in women (United States). *Cancer Causes Control* 1998; **9**: 621–630.
- 80 Sinha R, Chow WH, Kulldorff M, Denobile J, Butler J, Garcia-Closas M *et al*. Well-done, grilled red meat increases the risk of colorectal adenomas. *Cancer Res* 1999; **59**: 4320–4324.
- 81 Zheng W, Gustafson DR, Sinha R, Cerhan JR, Moore D, Hong CP *et al*. Well-done meat intake and the risk of breast cancer. *J Natl Cancer Inst* 1998; **90**: 1724–1729.
- 82 Sinha R, Rothman N. Role of well-done, grilled red meat, heterocyclic amines (HCAs) in the etiology of human cancer. *Cancer Lett* 1999; **143**: 189–194.
- 83 Zheng W, Deitz AC, Campbell DR, Wen WQ, Cerhan JR, Sellers TA *et al*. N-acetyltransferase 1 genetic polymorphism, cigarette smoking, well-done meat intake, and breast cancer risk. *Cancer Epidemiol Biomark Prev* 1999; **8**: 233–239.
- 84 Sinha R. An epidemiological approach to studying heterocyclic amines. *Mutat Res* 2002; **506/507**: 197–204.
- 85 Butler LM, Sinha R, Millikan RC, Martin CF, Newman B, Gammon MD *et al*. Heterocyclic amines, meat intake, and association with colon cancer in a population-based study. *Am J Epidemiol* 2003; **157**: 434–445.
- 86 Chen K, Liang JQ, Zhang Y, Wan YZ. Meta-analysis of risk factors for colorectal cancer. *World J Gastroenterol* 2003; **9**: 1598–1600.
- 87 Murtaugh MA, Ma K, Sweeney C, Caan BJ, Slattery ML. Meat consumption patterns and preparation, genetic variants of metabolic enzymes, and their association with rectal cancer in men and women. *J Nutr* 2004; **134**: 776–784.
- 88 Chao A, Thun MJ, Connell CJ, McCullough ML, Jacobs EJ, Flanders WD *et al*. Meat consumption and risk of colorectal cancer. *JAMA* 2005; **293**: 233–234.
- 89 Rodriguez C, McCullough ML, Mondul AM, Jacobs EJ, Chao A, Patel AV *et al*. Meat consumption among black and white men and risk of prostate cancer in the Cancer Prevention Study II Nutrition Cohort. *Cancer Epidemiol Biomark Prev* 2006; **15**: 211–216.
- 90 Sadri GH, Mahjub M. Meat consumption is a risk factor for colorectal cancer: meta-analysis of case-control studies. *Pakistan J Nutr* 2006; **5**: 230–233.
- 91 Lyon JL, Mahoney AW. Fried foods and the risk of colon cancer. *Am J Epidemiol* 1988; **128**: 1000–1006.
- 92 Muscat JE, Wynder EL. The consumption of well-done red meat and the risk of colorectal cancer. *Am J Public Health* 1994; **84**: 856–858.
- 93 Augustsson K, Skog K, Jagerstad M, Dickman PW, Steineck G. Dietary heterocyclic amines and cancer of the colon, rectum, bladder, and kidney: a population based study. *Lancet* 1999; **353**: 703–707.
- 94 Delfino RJ, Sinha R, Smith C, West J, White E, Lin HJ *et al*. Breast cancer, heterocyclic aromatic amines from meat and N-acetyltransferase 2 genotype. *Carcinogenesis* 2000; **21**: 607–615.
- 95 Carter HB, Pearson JD, Metter EJ, Brant LJ, Chan DW, Andres R *et al*. Longitudinal evaluation of prostatic-specific antigen levels in men with and without prostatic disease. *J Am Med Assoc* 1992; **267**: 2215–2220.
- 96 Oesterling JE, Jacobsen SJ, Chute CG, Guess HA, Girman CJ, Panser LA *et al*. Serum prostate-specific antigen in a community-based population of healthy men. *J Am Med Assoc* 1993; **270**: 860–864.
- 97 Etzioni R, Cha R, Cowen M. Serial prostate specific antigen screening for prostate cancer: a computer model evaluates competing strategies. *J Urol* 1999; **162**: 741–748.
- 98 Abdalla I, Ray P, Vijayakumar S. Race and serum prostate-specific antigen levels: current status and future directions. *Semin Urol Oncol* 1998a; **16**: 207–213.
- 99 Abdalla I, Ray P, Ray V, Vaida F, Vijayakumar S. Comparison of serum prostate-specific antigen levels and PSA density in African American, white, and Hispanic men without prostate cancer. *Urology* 1998b; **51**: 300–305.
- 100 Vijayakumar S, Winter K, Sause W, Gallagher MJ, Michalski J, Roach M *et al*. Prostate-specific antigen levels are higher in African-American than in white patients in a multicenter registration study: results of RTOG 94-12. *Int J Radiat Oncol Biol Biophys* 1998; **40**: 17–25.
- 101 Fowler Jr JE, Bigler SA, Kilambi NK, Land SA. Relationships between prostate-specific antigen and prostate volume in black and white men with benign prostate biopsies. *Urology* 1999; **53**: 1175–1178.
- 102 Mänttinen L, Auvinen A, Stenman U-H, Tammela T, Rannikko S, Aro J *et al*. Three-year results of the Finnish Prostate Cancer Screening Trial. *J Natl Cancer Inst* 2001; **93**: 552–553.
- 103 Gerstenbluth RE, Seftel AD, Hampel N, Oefelein MG, Resnick MI. The accuracy of the increased prostate specific antigen level (greater than or equal to 20 ng/ml) in predicting prostate cancer: is biopsy always required? *J Urol* 2002; **168**: 1990–1993.
- 104 Smith RP, Malkowicz SB, Whittington R, VanArsdalen K, Tochner Z, Wein AJ. Identification of clinically significant prostate cancer by prostate-specific antigen screening. *Arch Intern Med* 2004; **164**: 1227–1230.
- 105 Vollmer RT. Predictive probability of serum prostate-specific antigen for prostate cancer: an approach using Bayes rule. *Clin Chem* 2006; **125**: 336–342.
- 106 Thompson IM, Pauler DK, Goodman PJ, Tangen CM, Lucia MS, Parnes HL *et al*. Prevalence of prostate cancer among men with

- a prostate-specific antigen level ≤ 4.0 ng per milliliter. *N Engl J Med* 2004; **350**: 2239–2246.
- 107 American Cancer Society (ACS). *California Cancer Facts and Figures, 2006*. ACS California Division and Public Health Institute California Cancer Registry: Oakland, CA, 2005.
- 108 US Census Bureau. State and County QuickFacts (<http://quickfacts.census.gov/qfd/states/06000.html>, revised 8 June 2006); and American FactFinder, 2005 American Community Survey, custom tables B12002B, B01002B, B20017B (<http://factfinder.census.gov>): 2006.
- 109 Block G, Hartman AM, Dresser CM, Carroll MD, Gannon J, Gardner L. A data-based approach to diet questionnaire design and testing. *Am J Epidemiol* 1986; **124**: 453–469.
- 110 Sinha R, Rothman N, Salmon CP, Knize MG, Brown ED, Swanson CA et al. Heterocyclic amine content in beef cooked by different methods and to varying degrees of doneness and gravy made from meat drippings. *Food Chem Toxicol* 1998; **36**: 279–287.
- 111 Sinha R, Knize MG, Salmon CP, Brown ED, Rhodes D, Felton JS et al. Heterocyclic amine content of pork products cooked by different methods and to varying degrees of doneness. *Food Chem Toxicol* 1998; **36**: 289–297.
- 112 Cantwell M, Mittl B, Curtin J, Carrol R, Potischman N, Caporaso N et al. Relative validity of a food frequency questionnaire with a meat-cooking and heterocyclic amine module. *Cancer Epidemiol Biomark Prev* 2004; **13**: 293–298.
- 113 Bogen KT, Keating GA. US dietary exposures to heterocyclic amines. *J Exposure Anal Environ Epidemiol* 2001; **11**: 155–168.
- 114 Keating GA, Bogen KT. Methods to estimate heterocyclic amine concentrations in cooked meats in the U.S. diet. *Food Chem Toxicol* 2001; **39**: 29–43.
- 115 Keating GA, Bogen KT. Estimates of heterocyclic amine intake in the U.S. population. *J Chromatogr B* 2004; **802**: 127–133.
- 116 Draper NR, Smith H. *Applied Regression Analysis*, 2nd edn. John Wiley & Sons: New York, NY, 1981.
- 117 Selvin S. *Practical Biostatistical Methods*. Duxbury Press: New York, NY, 1995.
- 118 Kendall M, Stuart A. *The Advanced Theory of Statistics*, vol. 2. MacMillan Publishing Co.: New York, NY, 1979, pp 159–160.
- 119 Breslow NE, Day NE. *Statistical Methods in Cancer Research, Volume II – The Design and Analysis of Cohort Studies*, IARC Scientific Publ. No. 82. International Agency for Cancer research (IARC): Lyon, France, 1987, pp 106–118.
- 120 Wolfram S. *Mathematica Book*, 4th edn. Cambridge University Press: Cambridge, UK, 1999.
- 121 Kidd LR, Stillwell WG, Yu MC, Wishnok JS, Skipper PL, Ross RK et al. Urinary excretion of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in White, African American, and Asian-American men in Los Angeles County. *Cancer Epidemiol Biomark Prev* 1999; **8**: 439–445.
- 122 Malfatti MA, Kulp KS, Knize MG, Davis C, Massengill JP, Williams S et al. The identification of [2-¹⁴C]2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine metabolites in humans. *Carcinogenesis* 1999; **20**: 705–713.
- 123 U.S. Department of Agriculture (USDA). 1989–91 Continuing Survey of Food Intakes by Individuals and 1989–91 Diet and Health Knowledge Survey (CSFII/DHKS 1989–91 Data Set). CD-ROM. USDA Agricultural Research Service, Beltsville Human Nutrition Research Center, Food Surveys Research Group: Beltsville, MD, 1993.
- 124 Zelterman D. *Models for Discrete Data*. Clarendon Press: Oxford, UK, 1999, pp 6–24.
- 125 Wilkens LR, Hankin JH, Yoshizawa CN, Kolonel LN, Lee J. Comparison of long-term dietary recall between cancer cases and noncases. *Am J Epidemiol* 1992; **136**: 825–835.
- 126 Stark A. An historical review of the Harvard and the National Cancer Institute Food Frequency Questionnaires: their similarities, differences, and their limitations in assessment of food intake. *Ecol Food Nutr* 2002; **41**: 35–74.
- 127 Subar AF, Thompson FE, Kipnis V, Midthune D, Hurwitz P, McNutt S et al. Comparative validation of the Block, Willett, and National Cancer Institute food frequency questionnaires – The eating at America’s table study. *Am J Epidemiol* 2001; **154**: 1089–1099.