

AD _____

Award Number: W81XWH-05-1-0045

TITLE: MT 2A Phosphorylation by PKC Mu/PKD Influences Chemosensitivity to
Cisplatin in Prostate Cancer

PRINCIPAL INVESTIGATOR: Balaji C. Kethandapatti, M.D., FRCS
Meena Jaagi
David J. Smith
Wenguang Zhang
Cheng Du
Lynette M. Smith
Anton Galich

CONTRACTING ORGANIZATION: University of Nebraska Medical Center
Omaha, NE 68198-5100

REPORT DATE: November 2006

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 01-11-2006		2. REPORT TYPE Annual		3. DATES COVERED 1 Nov 2005 – 31 Oct 20006	
4. TITLE AND SUBTITLE MT 2A Phosphorylation by PKC Mu/PKD Influences Chemosensitivity to Cisplatin in Prostate Cancer				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-05-1-0045	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Balaji C. Kethandapatti, M.D., FRCS, Meena Jaagi, David J. Smith, Wenguang Zhang, Cheng Du, Lynette M. Smith, Anton Galich Email: BalajiK@ummhc.org				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Nebraska Medical Center Omaha, NE 68198-5100				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES Original contains colored plates: ALL DTIC reproductions will be in black and white.					
14. ABSTRACT The metallothioneins (MT) are a family of small molecular weight trace metal and free radical scavenging proteins well established to play a role in resistance to chemotherapy and radiation in human cancer. MT gene expression is up regulated in response to the presence of heavy metal ions such as zinc. We have previously established a physiological in vitro cell line model of MT induction using Zn, which is significantly associated with resistance to cisplatin chemotherapy in PC. In this report, our data demonstrates that treatment with TPEN, a Zn specific chelator, significantly improves sensitivity to cisplatin in prostate cancer cells.					
15. SUBJECT TERMS Advanced prostate cancer, Chemoresistance, Metallothionein 2A, MT 2A Protein kinase C Mu, PKC Mu					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	6	19b. TELEPHONE NUMBER (include area code)

SECOND ANNUAL REPORT

Table of Contents

Title: MT 2A Phosphorylation by PKCMu/PKD Influences Chemosensitivity to Cisplatin in Prostate Cancer.

Introduction.....	4
Body.....	4
Key Research Accomplishments.....	5
Reportable Outcomes.....	6
Conclusions.....	6
References.....	6
Appendices.....	None

Introduction

Since submission of first annual report in November, 2005, the PI has moved to University of Massachusetts, Worcester, MA from University of Nebraska, Omaha, NE in March of 2006. Because we are in the process of transferring the grant to a new institution we have not accomplished our stated goals for the second year. Herein, we report on the progress made in the first three months of the second year.

Our preliminary studies had demonstrated that kinase domain of Protein kinase D1, a novel serine threonine kinase, interacts with Metallothionein 2A (MT 2A), which belongs to the family of small molecular weight proteins called metallothioneins (MT) that scavenge trace metals and free radicals and are associated with resistance to chemotherapy and radiation in human cancers¹. Our studies further demonstrated the C4-2 prostate cancer cells that express higher levels of MT compared to its parental LNCaP cells² are selectively more resistant to trace metal containing chemotherapy agent cisplatin compared to LNCaP cells. Our *in vitro* experiments also demonstrated that MT 2A was phosphorylated by PKD1. Therefore we hypothesized ““Alteration in MT 2A expression influences chemoresistance to cisplatin in prostate cancer. PKC Mu/PKD kinase activity influences sensitivity to cisplatin by MT 2A phosphorylation in prostate cancer. The expression of MT 2A is quantitatively increased in progressive human prostate cancer””.

We proposed to establish the stated hypothesis through 3 aims.

Aim 1. To determine that alteration in MT 2A expression influences resistance to cisplatin in prostate cancer.

Aim 2. Inhibition of PKC Mu/PKD kinase activity and its influences on chemoresistance in prostate cancer cells by modulating the phosphorylation of MT 2A.

Aim 3. To quantify and qualitatively evaluate MT 2A protein expression in progressive human prostate cancer.

Body

Task 1: *To determine that alteration in MT 2A expression influences resistance to cisplatin in prostate cancer.*

1a: Effect of MT 2A over expression influences resistance to cisplatin in prostate cancer

Experiment 1: Cloning and Expression of MT 2A protein in LNCaP prostate cancer and MT 2A null CHO cells. As previously stated in our first annual report we continued our efforts to develop stable LNCaP and CHO cell lines transfected with His tag labeled MT 2A cDNA. In addition, we also cloned DsRed fluorescent tagged MT 2A in LNCaP cells. We found that compared to His tag labeled MT 2A transfected cells, the DsRed tagged clones expressed high levels of exogenous MT 2A and are useful in studying subcellular localization as well (Fig 1).




Figure 1; LNCaP prostate cancer cells transfected with DsRed fused MT 2A. Figures A and B demonstrate cytoplasmic localization of MT 2A and figured C and D demonstrate nuclear localization of MT 2A follow stimulation with either PKC activator phorbol ester (Fig 1C) or PKD1 selective activator Bryostat1 (Fig 1D).

1b: Down regulation of MT2A by RNAi leads to increased sensitivity to cisplatin.

We reported in our first annual report that we were attempting to down regulate MT 2A using RNAi. Because of very low levels of endogenous MT in cells in unstimulated cells we were unable to demonstrate down regulation of MT 2A by RNAi. Therefore, we used an alternate model of MT induction by Zn followed by inhibition of MT induction by Zn specific chelator TPEN to study the effect of MT on resistance to cisplatin in prostate

cancer (Fig 2). Our results demonstrate that treatment of TPEN significantly improves sensitivity to cisplatin in prostate cancer.




Figure 2: C4-2 prostate cancer cells were plated at a concentration of 3000 cells/ well in 96 well plates, treated with equimolar concentrations (150 μ M) of zinc, followed by treatment with TPEN at 24 hours and 10 μ M of cisplatin, which we had established as the IC50 concentration

in our cell line model. Viability assays were performed at 72 hours using non radioactive cell proliferation assays (Promega, Madison, WI). Experiments were performed in triplicate, and means were compared by 2-way ANOVA models. P values less than 0.05 were considered significant.

Key Research Accomplishments

We have established a stable DsRed fused to MT 2A cell lines, which would facilitate further studies on MT 2A and prostate cancer. We also recognized that down regulation of MT 2A by RNAi may not be a feasible experimental model because of very low levels of endogenous MT in prostate cancer cells. Therefore, we used an alternate model of inhibiting Zn induced MT

expression by Zn specific chelator TPEN, which demonstrated that TPEN treatment significantly increased sensitivity to cisplatin treatment in prostate cancer cells.

Reportable Outcomes

1. Tony Mammen, Samuel Sterrett, Cheng Du and K.C. Balaji; Treatment with Zinc Specific Chelator TPEN reverse An induced Resistance to Cisplatin in Prostate Cancer; Sante Fe, New Mexico, Poster #10, 10/06

Conclusions

MT expression is associated with resistance to cisplatin chemotherapy in human prostate cancer cells.

References

1. Ebadi, M. & Iversen, P. L. Metallothionein in carcinogenesis and cancer chemotherapy. *Gen Pharmacol* 25, 1297-310. (1994).
2. Rao, P. S., Jaggi, M., Smith, D. J., Hemstreet, G. P. & Balaji, K. C. Metallothionein 2A interacts with the kinase domain of PKC μ in prostate cancer. *Biochem Biophys Res Commun* 310, 1032-8 (2003).