


Organosilicon Derivatives of Inorganic
"Benzenoid" Compounds.

by Dietmar Seyferth

Department of Chemistry
Massachusetts Institute of Technology
Cambridge 39, Massachusetts

I. Introduction.

In a previous report¹ the synthesis of thermally and oxidatively stable silicon-substituted borazines (I) was reported. The present report summarizes our



[R = (CH₃)₂R'SiCH₂-, where R' = CH₃, C₂H₅,
n-C₄H₉ and (CH₃)₃SiO]

further efforts in the borazine area and describes as well work devoted to the preparation of silicon-substituted phosphazenes and to the synthesis and characterization of perfluorovinylolithium. The latter reagent seemed of possible interest in connection with our studies of organofunctional borazines.

The work reported here was carried out principally by Drs. H. P. Kögler, G. Raab, W. Freyer and J. F. Helling and is being continued since September by Drs. H. Yamazaki and M. Takamizawa. The period 1 March 1959 - 30 November 1961 is covered in this summary.

II. Discussion.


A. Preparation of Silicon-substituted Borazines.

A compound containing a B-Si linkage (I, R = (C₆H₅)₃Si-) was prepared by the reaction of triphenylsilyllithium in THF with B-trichloro-N-trimethylborazine²:




This borazine, a white solid of m.p. 248-251°, was hydrolytically unstable:

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1962		2. REPORT TYPE		3. DATES COVERED 00-00-1962 to 00-00-1962	
4. TITLE AND SUBTITLE Organosilicon Derivatives of Inorganic				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Paul V. Galvin Library, 35 W 33rd Street, Chicago, IL, 60616				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			



Silicon-substituted borazines of higher molecule weight could be prepared by the reaction of B-chloro-B-bis(trimethylsilylmethyl)borazine with α,ω -di-Grignard reagents:



This approach to stable borazines of higher molecular weight should be applicable to other novel systems, including cross-linked polymers when B-dichloro-B-trimethylsilylmethyl-N-trimethylborazine is used in place of the monochloro compound.



Other studies showed that facile introduction of bulky trimethylsilylmethyl groups into borazines by the Grignard reaction is limited to those cases in which the groups on the nitrogen atom are fairly small. Thus various N-triphenylborazines did not react with trimethylsilylmethylmagnesium chloride or bromide, although they did react well with methylmagnesium bromide.

B. Reactions of B-Trivinyl-N-triphenylborazine.

The preparation of B-trivinyl-N-triphenylborazine has been reported and its polymerization properties have been studied.³ We have begun an investigation of the reactions of this compound with a variety of reagents with the object of preparing a wide variety of organofunctional borazines.

It might have been expected that phenyllithium would add across the double bonds in the vinylborazine, as it does with vinylsilanes and vinylgermanes. However, this was not the case; phenyllithium appeared not only to disrupt completely the borazine ring system, but also to displace the vinyl group on boron. The major boron-containing product of this reaction was triphenylborane (isolated as $(\text{C}_6\text{H}_5)_3\text{B}\cdot\text{NH}_3$).

A variety of radical reagents added to the C=C bond in B-trivinyl-N-triphenylborazine. These included bromotrichloromethane, carbon tetrabromide and benzenethiol, and borazines II, III and IV, respectively, were obtained. Dimethylphosphite and trichlorosilane reacted with the vinylborazine in the presence of peroxide initiators, but pure products have not yet been isolated. Methylchlorosilane also added to the vinylborazine when chloroplatinic acid was used as the catalyst, but again a noncrystallizable oil resulted.




C. Attempted Preparation of Silicon-substituted Phosphazenes.

It has been reported that trimeric and tetrameric phosphonitrilic chlorides can be arylated by the Grignard and by the Friedel-Crafts procedures. However, more useful procedures for obtaining alkyl or aryl derivatives of the cyclic phosphonitrilic chlorides involve using alkyl(or aryl)tetrachlorophosphoranes and dialkyl(or diaryl)trichlorophosphoranes in place of pentachlorophosphorane in the phosphazene ring closure reaction⁴:





It was decided to test this approach to organosilicon-substituted phosphazenes in this research.


The compound whose preparation was desired was:



and its synthesis was attempted using the following reactions:



Conventional methods used to prepare alkylchlorophosphines utilize the reaction of phosphorus trichloride with mild alkylating reagents such as dialkylmercury⁵ or tetraalkyllead⁶ compounds. Accordingly, bis-(trimethylsilylmethyl)mercury and tetrakis-(trimethylsilylmethyl)lead, both new compounds, were prepared by the usual Grignard procedure, and their action on phosphorus trichloride was investigated. The mercurial did not react with phosphorus trichloride either when a mixture of the reactants was refluxed in hexane solution, or in the absence of solvent at ca. 76°. In contrast, the lead compound was cleaved by phosphorus trichloride, giving crystalline, sublimable tris-(trimethylsilylmethyl)lead chloride and trimethylsilylmethyldichlorophosphine. However, the isolated yield of the phosphine was not very high, and a better synthesis was desired.



Although the Grignard procedure is not generally applicable to the preparation of pure alkylchlorophosphines because mixtures of mono-, di- and trisubstituted products result, it seemed possible that with the relatively bulky trimethylsilylmethyl group, monosubstitution could be realized in acceptable yield. Indeed, our previous work⁷ showed that complete substitution of all three chlorine atoms in phosphorus trichloride with trimethylsilylmethyl groups by the Grignard procedure requires drastic conditions. It was found in the present work that addition of one equivalent of trimethylsilylmethylmagnesium chloride to one mole of phosphorus trichloride in ether at low temperature resulted in the desired trimethylsilylmethyldichlorophosphine in ca. 40% yield. Further study of this reaction aimed at improvement of the yield was deferred until the practicality of reactions (2) and (3) could be assessed.

Attempted conversion of trimethylsilylmethyldichlorophosphine to trimethylsilylmethyltetrachlorophosphorane (reaction 2) was not successful. Even at -20° in the dark, addition of a solution of chlorine in 1,1,2,2-tetrachloroethane to trimethylsilylmethyldichlorophosphine caused cleavage of the carbon-silicon bond to form trimethylchlorosilane. This facile cleavage of the carbon-silicon bond may be due to the inductive effect of the PCl_4 group; electron withdrawal by this group would make the silicon-carbon bond more polar, hence more susceptible toward ionic fission. Investigation of the solid formed in the chlorination reaction showed it to be impure pentachlorophosphorane rather than the expected chloromethyltetrachlorophosphorane. The hydrolysis product of this solid formed a barium salt of very low carbon content, and an anilinium salt could be prepared which was shown to be $[\text{C}_6\text{H}_5\text{NH}_3][\text{H}_2\text{PO}_4]$. It is not known at what point fission of the phosphorus-carbon bond took place, nor is the fate of the methylene group of the trimethylsilylmethyldichlorophosphine known. It is, however, very likely that such cleavage took place after the cleavage of the silicon-carbon bond, since initial cleavage of trimethylsilylmethyldichlorophosphine at the phosphorus-carbon bond would have given chloromethyltrimethylsilane. This compound is stable to

silicon-carbon bond cleavage under these experimental conditions, although side chain chlorination does take place.


Since the proposed synthesis of silicon-substituted phosphazenes failed in the second of the three steps, this approach was abandoned.

B. The Synthesis of Perfluorovinyl lithium.

It would be most desirable to introduce unsaturated groups, such as the vinyl and the perfluorovinyl group, into borazines. Hopefully, functionality of this type would allow polymerization reactions to be carried out with the object of linking borazene rings. Also addition of various reagents to the unsaturated groups would permit the preparation of a large variety of borazines containing organofunctional alkyl groups linked to boron.

Active metal vinyl compounds - vinylmagnesium chloride and bromide, vinyl-lithium and vinylsodium - have been developed in recent years. More recently perfluorovinylmagnesium bromide⁸ was described. After our early attempts to prepare B-vinylated borazines, the synthesis of B-trivinyl-N-triphenylborazine was reported in the patent literature³. For this reason we initiated a study directed toward the synthesis of B-perfluorovinylborazines.

Perfluorovinylmagnesium bromide is known as a relatively unstable species that must be prepared and used in tetrahydrofuran solution⁸. It was thought that perfluorovinyl lithium might be capable of existence and that this reagent might be prepared in other, lower boiling solvents, thus extending the versatility of perfluorovinylation procedures. Our first efforts showed that perfluorovinyl lithium could not be prepared by the direct reaction between metallic lithium and bromotrifluoroethylene. Accordingly, the transmetalation procedure which had proven of exceptional utility in the synthesis of vinyl lithium⁹ and allyl lithium¹⁰, was applied in this case. Perfluorovinyltin compounds had been the subject of previous study in these Laboratories¹¹, and the action of organolithium reagents on perfluorovinyltin compounds was examined as a possible route to perfluorovinyl lithium. The reaction between triphenylperfluorovinyltin and phenyllithium, carried out in diethyl ether at room temperature, demonstrated that the exchange reaction did indeed occur, since tetraphenyltin was isolated in nearly quantitative yield. However, no evidence could be obtained for the presence of perfluorovinyl lithium in the dark brown solution which was formed. Since perfluorovinyl lithium did not appear to be stable under these conditions, it was felt that either lower reaction temperatures in the diethyl ether system or a change to a less polar solvent might favor the stability of the desired reagent. Subsequent experiments showed that perfluorovinyl lithium could be prepared as a stable entity either in diethyl ether or in pentane at -40° to -30°. The reaction of phenyllithium with phenyltriperfluorovinyltin (3:1 molar ratio) in ether at ca. -40° proceeded rapidly, giving a precipitate of tetraphenyltin and a brown solution of perfluorovinyl lithium. Exploratory experiments showed this reagent was best characterized by its reaction with triethyltin chloride:



The yield of triethylperfluorovinyltin obtained was ca. 40%, and this represents a minimum figure for the yield of perfluorovinylolithium in the transmetalation reaction. Perfluorovinylolithium also reacted normally with trimethylbromosilane and carbon dioxide.

A similar exchange reaction could be effected in pentane between n-butyllithium and n-butyiltriperfluorovinyltin. However, in this solvent exchange was not complete, and it was found best to use the lithium reagent and the tin compound in 2:1 molar ratio. When the perfluorovinylolithium preparation was carried out in -30° and triethyltin chloride used to characterize the lithium reagent formed, the work-up procedure gave triethylperfluorovinyltin in 46 - 50% yield. Since the exchange was not complete under these conditions, and since in fact a deficiency of n-butyllithium was used, the possibility existed that triethyltin chloride could undergo direct exchange with any perfluorovinyltin compounds still present in the reaction mixture. To completely exclude this possibility the reaction mixture was shaken with aqueous-alcoholic potassium fluoride solution prior to removal of solvent and distillation of the products. This treatment removed unreacted triethyltin chloride as the insoluble fluoride. Other experiments showed that the exchange reaction between triethyltin chloride and di-n-butyldiperfluorovinyltin occurred only when this mixture was heated. Thus distillation at reduced pressure of a mixture of triethyltin chloride and di-n-butyldiperfluorovinyltin produced triethylperfluorovinyltin in ca. 9% yield.

Perfluorovinylolithium was more stable in pentane than in ether solution. It appeared to be stable in pentane at 0°, but short refluxing of its pentane solution caused complete decomposition.

III. References.

1. D. Seyferth, Conference on High Temperature Polymer and Fluid Research, Materials Laboratory, WADD, Dayton, Ohio, 26-28 May 1959, p. 117.
2. D. Seyferth and H. P. Kögler, *J. Inorg. Nucl. Chem.*, **15**, 99 (1960).
3. J. Pellon, U.S. patent 2,954,366 (1960); *J. Polymer Sci.*, **55**, 153 (1961).
4. C. Haber et al., *J. Am. Chem. Soc.*, **80**, 2116 (1958).
5. G. M. Kosolapoff, "Organophosphorus Compounds", John Wiley and Sons, Inc., New York, 1950, p. 42.
6. M. S. Kharasch, E. V. Jensen and S. Weinhouse, *J. Org. Chem.*, **14**, 429 (1949).
7. D. Seyferth, *J. Am. Chem. Soc.*, **80**, 1336 (1958).
8. I. L. Knunyants, R. N. Sterlin, R. D. Yatsenko and L. N. Pinkina, *Izvest. Akad. Nauk S.S.S.R., Otdel. Khim. Nauk*, 1958, 1345.
9. D. Seyferth and M. A. Weiner, *Chem. and Ind. (London)*, 1959, 402.
10. D. Seyferth and M. A. Weiner, *J. Org. Chem.*, **24**, 1395 (1959).
11. D. Seyferth, K. Brändle and G. Raab, *Angew. Chem.*, **72**, 77 (1960); *idem.*, *J. Org. Chem.*, **26**, 2934 (1961).