

Antisense treatments for biothreat agents

Kelly L Warfield¹, Rekha G Panchal², M Javad Aman¹ & Sina Bavari^{1*}

Addresses

¹US Army Medical Research Institute of Infectious Diseases
Fort Detrick
Frederick
MD 21702
USA
Email: sina.bavari@amedd.army.mil

²Target Structure Based Drug Discovery Group
SAIC-Frederick Inc
NCI-Frederick
Frederick
MD 21702-1201
USA

*To whom correspondence should be addressed

Current Opinion in Molecular Therapeutics 2006 8(2):93-103
© The Thomson Corporation ISSN 1464-8431

Antisense oligomers (ASOs) represent a promising technology to treat viral and bacterial infections, and have already been shown to be successful against a variety of pathogens in cell culture studies and nonhuman primate models of infection. For these reasons, antisense technologies are being pursued as treatments against biothreat agents such as Ebola virus, dengue virus and Bacillus anthracis. Several generations of modified oligonucleotides have been developed to maximize nuclease resistance, target affinity, potency, cell entry, and other pharmacokinetic properties. First-generation ASOs contain phosphorothioate modifications to increase stability through nuclease resistance. Further chemical modifications in second-generation ASOs include 2'-O-methyl and 2'-O-methoxy-ethyl oligos, which increase nuclease resistance and oligo:RNA binding affinities. Third-generation ASOs contain a variety of chemical modifications that enhance stability, affinity and bioavailability. A fourth class of oligonucleotide-based compounds consists of small interfering RNAs, which have recently become widely used for gene knockdown in vitro and in vivo. This review focuses on the third-generation phosphorodiamidate morpholino oligomers, which are nonionic and contain a morpholine ring instead of a ribose, as well as phosphorodiamidate linkages in place of phosphorothioates. Multiple antisense oligomer-based therapeutics are being developed for use against biothreat agents, and antisense drugs will likely become a critical member of our arsenal in the defense against highly pathogenic, emerging or genetically engineered pathogens.

Keywords Antisense, biothreat, oligonucleotide, phosphorodiamidate morpholino, phosphorothioate, small interfering RNA

Introduction

Many viral and bacterial agents have a history of state-sponsored 'weaponization', including Marburg, Ebola, Junin, Machupo, yellow fever viruses and smallpox, as well as *Bacillus anthracis*, *Yersinia pestis* and *Francisella tularensis* [1-5]. These agents are considered serious public health threats, are handled under biosafety level (BSL)-3 or -4 conditions [6], and are listed as potential biothreat agents by

the National Institute of Allergy and Infectious Diseases (http://www3.niaid.nih.gov/Biodefense/bandc_priority.htm). Features that characterize a pathogen as a serious bioweapon threat include high morbidity and mortality rates, the potential for person-to-person or aerosol transmission, low infective dose, stability, and the feasibility of large-scale production. Unfortunately, many of the biothreat agents are also characterized by a lack of available vaccines, prophylactics or treatments, and so developing preventatives and therapeutics against these dangerous organisms has become a high public health priority in recent years [7•].

Since the first use of antisense oligomers (ASOs) for inhibiting Rous sarcoma virus replication in cell culture [8,9], there has been a keen interest in developing ASO therapeutics. ASOs function by binding complementary sequences of RNA, and inhibiting gene expression in a rapid and highly specific manner. The manner in which they inhibit expression of the target gene has made them powerful tools to dissect critical pathways, and also as potential drugs for targeting host and pathogen genes. Here, we will provide a representative, but not exhaustive, presentation of ASOs in development as therapeutics, and provide information on advances made in the ASO field for use against biothreat agents.

Antisense oligonucleotides

First-generation antisense oligomers

Unmodified oligodeoxynucleotides are highly unstable in bodily fluids because of their rapid degradation by nucleases (Table 1). To enhance oligomer stability, replacement of a nonlinking oxygen atom with sulfur within the phosphodiester linkage of each nucleotide renders first-generation antisense phosphorothioate oligomers (PTOs) inefficient substrates for both endo- and exo-nucleases (Table 1) [10]. PTOs inhibit gene expression after binding to the target mRNA, and thereby direct the mRNA for RNase H degradation as a result of the formation of the PTO:mRNA duplex. To mediate efficient suppression of fully complementary mRNAs, optimal PTOs are 15 to 21 nucleotides long; however, transient hybridization can occur between the ASO and mRNAs with short complementary sequences, and DNA:RNA heteroduplexes as short as five nucleotides can initiate RNase H1-mediated degradation of the RNA component [11-14].

Second-generation antisense oligomers

A second-generation approach to ASO design incorporates further modifications to the inter-nucleoside ribose-phosphate linkages to improve the affinity, stability and activity of these drugs (Table 1). 2'-O-(2-Methoxy)ethyl (2'MOE)-modified oligonucleotide:RNA heterodimers are not substrates for cleavage by RNase H. Therefore, mRNA cleavage is limited to a short stretch of phosphorothioate nucleotides with unmodified sugars at the center of the ASO [15-17]. PTOs with 2'MOE 'wings', consisting of five 2'MOE-modified

Report Documentation Page		Form Approved OMB No. 0704-0188
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.		
1. REPORT DATE 01 AUG 2006	2. REPORT TYPE N/A	3. DATES COVERED -
4. TITLE AND SUBTITLE Antisense treatments for biothreat agents. Current Opinion in Molecular Therapeutics 8:93 -103		5a. CONTRACT NUMBER
		5b. GRANT NUMBER
		5c. PROGRAM ELEMENT NUMBER
6. AUTHOR(S) Warfield, KL Panchal, RG Aman, MJ Bavari, S		5d. PROJECT NUMBER
		5e. TASK NUMBER
		5f. WORK UNIT NUMBER
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Unit4ee States Army Medical Research Institute of Infectious Diseases, Fort Detrick, MD		8. PERFORMING ORGANIZATION REPORT NUMBER TR-06-006
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)		10. SPONSOR/MONITOR'S ACRONYM(S)
		11. SPONSOR/MONITOR'S REPORT NUMBER(S)
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution unlimited		
13. SUPPLEMENTARY NOTES The original document contains color images.		
14. ABSTRACT Antisense oligomers (ASOs) represent a promising technology to treat viral and bacterial infections, and have already been shown to be successful against a variety of pathogens in cell culture studies and nonhuman primate models of infection. For these reasons, antisense technologies are being pursued as treatments against biothreat agents such as Ebola virus, dengue virus and Bacillus anthracis. Several generations of modified oligonucleotides have been developed to maximize nuclease resistance, target affinity, potency, cell entry, and other pharmacokinetic properties. First-generation ASOs contain phosphorothioate modifications to increase stability through nuclease resistance. Further chemical modifications in second-generation ASOs include 2'-O-methyl and 2'-O-methoxy-ethyl oligos, which increase nuclease resistance and oligo:RNA binding affinities. Third-generation ASOs contain a variety of chemical modifications that enhance stability, affinity and bioavailability. A fourth class of oligonucleotide-based compounds consists of small interfering RNAs, which have recently become widely used for gene knockdown in vitro and in vivo. This review focuses on the third-generation phosphorodiamidate morpholino oligomers, which are nonionic and contain a morpholine ring instead of a ribose, as well as phosphorodiamidate linkages in place of phosphorothioates. Multiple antisense oligomer-based therapeutics are being developed for use against biothreat agents, and antisense drugs will likely become a critical member of our arsenal in the defense against highly pathogenic, emerging or genetically engineered pathogens.		
15. SUBJECT TERMS antisense phosphorodiamidate oligomers, antiviral, antibacterial, therapeutics, review		

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 11	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Table 1. Properties of selected oligonucleotide compounds.

Antisense oligomers	Example	Chemical modifications	Net charge	Mechanism of action	Advantages	Disadvantages
Unmodified oligodeoxynucleotides	Naked DNA	None	Negative	RNase H cleavage	-	Serum half-life ~ 1 h Rapidly degraded in biological fluids by nucleases
First-generation	Phosphorothioate ODNs	Sulfur molecule replaces one of the nonbridging oxygen atoms in the phosphodiester bonds	Negative	RNase H cleavage	Serum half-life 9 to 10 h Enhanced nuclease resistance Normal Watson-Crick base pairing Activates RNase H Attractive pharmacokinetic properties Enhanced specificity of hybridization compared with unmodified ODNs	Acute toxicity resulting from activation of complement cascade Binding to proteins such as heparin-binding proteins Slightly reduced affinity toward complementary RNA molecules compared with other ODNs
Second-generation	2'-O-methyl and 2'-O-methoxy-ethyl RNA	Alkyl modifications at the 2'-position of the ribose on the nucleotide	Negative	Blocks translation (ie, steric hindrance)	Serum half-life 6 to 12 h Less toxic than phosphorothioate ODNs Slightly enhanced affinity toward complementary RNAs Can block splicing sites to increase expression of alternatively spliced protein variants	Binding to proteins such as heparin-binding proteins
Third-generation	Phosphorodiamidate morpholino oligomers	Nonionic DNA analogs with the ribose replaced with a morpholine ring and phosphorodiamidate intersubunit linkages instead of phosphodiester bonds	Neutral	Blocks translation (ie, steric hindrance)	Serum half-life 1 to 20 h Limited binding to serum proteins such as nucleic-acid-binding proteins Favorable base stacking and relatively high duplex stability Nuclease and protease resistant Limited off-target alteration of gene expression through G-quartets Limited metal chelation since there is no net charge to mediate binding No altered coagulation times Lack immune stimulation via CpG motifs of the complement cascade	Limited cellular uptake of naked PMOs Undetermined efficacious human doses against infectious diseases
RNAi	siRNAs	Double-stranded RNA	Negative	Targets RNA to RISC complex for degradation	High potent activity Chemical modifications can increase uptake Mechanisms of action are well understood	Difficult to deliver naked siRNAs into tissues May induce interferon responses Limited tolerability for subsequent chemical modifications Sensitive to nuclease degradation May require large doses for efficacy in humans

ODN oligodeoxynucleotide, **PMO** phosphorodiamidate morpholino oligomer, **RISC** RNA-induced silencing complex.

phosphorothioate-linked nucleotides at both the 5' and 3' ends, bind more strongly to RNA and deplete target mRNAs more potently than their 2' unsubstituted analogs [18-20]. Many other possible backbone modifications have been explored for RNase H-dependent ASO drugs, such as locked nucleic acid (LNA), PTO with RNase-incompetent methylphosphonate nucleotides, or 2'MOE modifications in the middle of a PTO [21-23]. Of these, 2'MOE-winged PTOs appear to be the most promising and thoroughly developed PTO-based therapeutic. In fact, the first antisense drug approved by the FDA, fomivirsen, is a 2'MOE PTO designed to treat cytomegalovirus-induced retinitis in AIDS patients [24].

Third-generation antisense oligomers

Another major class of ASO compounds, phosphorodiamidate morpholino oligonucleotides (PMOs), achieve nuclease resistance by incorporating nucleosides with a morpholino instead of ribose backbone, and also utilize uncharged internucleoside phosphorodiamidate linkages (Table 1) [25]. RNA:PMO hybrids are not generally substrates for RNase H cleavage, but suppress mRNA targets through high-affinity binding at the sites of ribosomal entry, mRNA processing, or by blocking the formation of important secondary structural elements [26-29]. Therefore, PMOs are generally designed to target important translational signals, such as the AUG start site or secondary structures immediately 5' or 3' of the AUG start site. The neutral chemistry of PMOs confers many desirable safety and pharmacokinetic properties compared with other ASOs, including avoidance of non-specific binding to host proteins and excellent resistance to nucleases and proteases (Table 1). The nonionic characteristics of PMOs have been suggested to hinder cellular uptake compared with previous-generation negatively charged ASOs. However, this aspect of PMO chemistry does not appear to affect *in vivo* efficacy and may be an artifact of cell lines, since PMOs readily enter primary cells [27]. Additionally, conjugation with specific peptide tags such as arginine-rich peptides increases PMO entry into cells [30,31••]. PMOs promise a simple method of antisense drug design, and a drug platform that is cheaper to produce than phosphorothioate-based ASO drugs [32].

RNA interference

In recent years, small interfering RNA (siRNA) has been developed as a powerful gene silencing platform both *in vitro* and *in vivo* (Table 1) [33]. The RNA interference pathway through which siRNAs are recognized in the cytoplasm, cut into short fragments, and catalytically utilized by cells to degrade other complementary transcripts in the nucleus, is well understood and described elsewhere [34-36]. Rapid tail vein injections of unmodified siRNA in mice have proved efficacious [37,38]. However, other routes of administration have not been as successful *in vivo*, seemingly as a result of rapid degradation of unmodified siRNA molecules in mammals [39]. However, introducing chemical modifications at the 3' ends of both siRNA strands confers *in vivo* stability [40••]. Ongoing clinical trials will shed light on the potential of siRNAs for use as therapeutics.

Potential of antisense oligonucleotides as therapeutics

The use of ASOs as therapeutics relies upon specific hybridization of the oligomers to the target mRNA, the capacity of the ASO:mRNA complex to inhibit gene expression, and sufficient biological stability and activity of the ASOs *in vivo*. While many of the lead candidate ASO-based therapeutics are highly specific and effective in inhibiting gene expression of the target of interest *in vitro*, there are many potential barriers toward *in vivo* efficacy for this class of therapeutics. After *in vivo* administration, the ASO must penetrate multiple barriers, gain entry to the desired area of the body, enter the target cells, and localize to the intracellular compartment necessary for inhibiting gene expression. Throughout this process, the ASO must remain intact by avoiding nuclease degradation and not be cleared from the body before gene knockdown occurs. Furthermore, especially for bacterial pathogens, the ASO may have to traverse within the pathogen to target the pathogenic mRNAs.

Among the ASOs, the pharmacokinetics of PTOs and modified PTOs in humans are fairly well understood. Intravenously administered PTO and 2'MOE-modified phosphorothioate (PS) oligonucleotides distribute primarily to highly perfused tissues (the liver, kidneys, bone marrow, skin and skeletal muscles), with only about 1% of the oligomers remaining available in the plasma [41-44]. Intravenous administration of PTO compounds may be limited by distribution mainly to highly vascularized tissues due to promiscuous interactions with serum proteins via their polyanionic backbone [45]. Encouragingly, administering PTOs subcutaneously rather than intravenously results in a much higher plasma concentration [46]. Three days after subcutaneous administration of PTOs to humans, up to 30% of PTO remains in the plasma. For this reason, subcutaneous administration may become the preferred method for delivering PTO-based drugs [46]. First- and second-generation PTOs have shown promise *in vivo* against fulminant hepatitis and hepatitis C virus infections [20,47], as well as non-liver diseases such as cancers, through the inhibition of various targets, including protein kinase C- α (aprinocarsen), Bcl-2 (oblimersen; Genta Inc), Raf kinase (ISIS-5132), Ha-ras (ISIS-2503), R1 α regulatory subunit of cAMP-dependent protein kinase type I (Gem-231) and clusterin (OGX-011; OncoGenex Technologies Inc) [48]. However, multiple reactions to PTOs during human clinical trials have been observed, including pronounced cardiovascular responses, activation of the complement cascade, thrombocytopenia, hyperglycemia and hypotension [49].

PMOs have been safely employed in more than 12 clinical trials involving over 300 individuals without adverse drug-related events occurring [49-51]. These initial clinical studies have also demonstrated pharmacokinetic elimination half-lives in humans of 1 to 20 h [49]. The potential toxicity of PMOs has been evaluated in mice, rats and nonhuman primates, and the only noted toxicity to date is the transient

appearance of basophilic granules in macrophages, Kupfer cells and other phagocytic cells in lymph nodes when PMOs are administered for at least 14 days [27]. PMO sequences have also been evaluated in Good Laboratory Practice genotoxicity batteries with no observed mutagenesis [27].

In vivo studies of siRNA biodistribution in mice show a similar profile to other classes of oligonucleotide drugs [52]. However, to date, there is only one report of a siRNA-based drug being used in humans; Sirna-027 (Sirna Therapeutics Inc/Allergan Inc) targets vascular endothelial growth factor receptor-1, a receptor involved in angiogenesis. The company aims to use this drug to treat age-related macular degeneration [53•]. Encouragingly, in six of seven patients treated with a single, intravitreal dose of Sirna-027, there was a reduction in retinal thickness [53•]. However, because

the siRNAs were directly injected into the eye, it may be difficult to extrapolate these findings to systemic siRNA treatment of humans.

Antisense treatment of viral infection

At the beginning of antisense research, the first investigators tested whether Rous-sarcoma-virus-complementary oligonucleotides could interfere with viral amplification [8,9]. Since then, gene knockdown using ASOs against a variety of agents has been demonstrated to be quite effective in squelching viral amplification (Table 2). More recently, interest in using ASO-based approaches against biothreat agents has peaked, and several new reports provide evidence that ASO therapeutic candidates will be effective for treating acute viral diseases such as Ebola and dengue (Table 2).

Table 2. Summary of studies of antisense efficacy against viral agents.

Virus	Antisense type	Model inhibition demonstrated	References
Avian myeloblastosis virus	Unmodified and modified antisense oligodeoxynucleotides	<i>In vitro</i>	[83,84]
Calicivirus	PMO	Cell culture	[85]
Coronavirus (including SARS)	PMO siRNA	Cell culture, nonhuman primates	[28,86,87•]
Encephalomyocarditis virus	Antisense oligonucleotides	Cell-free rabbit reticulocyte lysates	[88]
Ebola virus	PMO	<i>In vitro</i> , cell culture, rodents, nonhuman primates	[60,61]
Epstein-Barr virus	PTO	Cell culture	[89,90]
Equine arteritis virus	PMO	<i>In vitro</i> , cell culture	[91]
Flaviviruses (ie, dengue and West Nile viruses)	PMO	<i>In vitro</i> , cell culture	[65-68,69•]
Friend retrovirus	Oligonucleotides	Cell culture	[92]
HBV	Unmodified or chimeric DNA-RNA PTOs siRNA	<i>In vitro</i>	[93-96]
HCV	PNAs and LNAs 2'-O-methyloligoribonucleotides	<i>In vitro</i>	[97,98]
HIV	LNA/DNA chimeric oligomers 2'-O-methyloligoribonucleotides Methylphosphonate-phosphodiester hybrid oligonucleotides Phosphate-methylated DNA Unconjugated and conjugated PTOs Methylphosphonate oligodeoxynucleotides	<i>In vitro</i> , cell culture	[10,93,99-114]
HSV	Polyalkylating single-stranded oligomers Unconjugated and conjugated PTOs 2'-O-methyloligoribonucleoside methylphosphonate oligos	Cell culture	[115-119]
Infectious hematopoietic necrosis virus	PMO	Cell culture	[120]
Lentiviruses	Phosphorothioate ON	Cell culture	[121]
Mouse leukemia virus	Modified and unmodified PTOs	Cell culture and mice	[122]
Respiratory syncytial virus	Modified and unmodified PTOs	<i>In vitro</i> , cell culture, non-human primates	[123••]
Retroviruses	PTO	Cell culture	[121]
Tobacco mosaic virus	Complementary ODN	<i>In vitro</i>	[124]
Vesicular stomatitis virus	Unconjugated and conjugated PTOs Methylphosphonate oligonucleotides	Cell culture	[117,125-127]
Vesiviruses	PMO	Cell culture	[128•]

HBV hepatitis B virus, **HCV** hepatitis C virus, **HSV** herpes simplex virus, **LNA** locked nucleic acid, **ODN** oligodeoxynucleotide, **ON** oligonucleotide, **PMO** phosphorodiamidate morpholino oligomer, **PNA** peptide nucleic acid, **PTO** phosphorothioate oligomers.

Development of antisense treatments for filovirus infections

The filoviruses, Ebola (EBOV) and Marburg (MARV), are highly virulent and cause acute, severe hemorrhagic fever in humans and nonhuman primates. Although filovirus infection most often leads to death, it is not uniformly lethal in humans. Numerous studies clearly demonstrate that the survival of an infected individual or animal is critically dependent on the ability of the host's innate and adaptive immune responses to control rapid viral growth [54-57]. Thus, host- or virus-targeted therapeutics that could curb filovirus replication until generation of an effective host immune response may prove efficacious in prevention of lethal disease. MARV and EBOV have a 19-Kb, negative-sense, single-stranded RNA genome that encodes for nucleoprotein (NP), glycoprotein (GP), four viral proteins (VP), and RNA-dependent RNA polymerase (L protein) [58]. Successful transcription and replication of the viral genome depends on coordinated action of L, NP, VP30 and VP35 proteins. The VP24 and VP40 proteins coordinate budding of the virions from the cell, where the GP is incorporated into the membrane during capsid formation [58].

Currently, there are no available vaccines or therapeutics for treating EBOV infections in humans. Although many different strategies have been attempted, only the administration of a recombinant tissue Factor VIIa inhibitor is protective in 33% of treated nonhuman primates [59]. Recently, Warfield *et al* and Enterlein *et al* described successful strategies for interfering with EBOV infection *in vitro* and *in vivo* by using untagged and peptide-tagged PMOs [60,61]. Therapeutic administration of a single untagged or peptide-tagged PMO targeting the viral VP35 mRNA significantly reduced viremia and resulted in 100% survival in EBOV-infected mice [60,61]. A combination of EBOV-specific, untagged PMOs targeting VP24, VP35 and RNA polymerase L protected mice and guinea pigs in both pretreatment and post-exposure therapeutic regimens [60,61]. In a prophylactic proof-of-principal trial, the PMOs also protected 75% of rhesus macaques from lethal EBOV infection [60,61]. Screening studies for efficacious PMOs against MARV have yielded positive results in the guinea pig model and, to date, we have identified more than five compounds that resulted in high levels of survival in initial studies [Warfield KL, unpublished data].

siRNAs are also expected to provide robust treatment options for hemorrhagic fevers because siRNAs act in a self-amplifying enzymatic manner. The inhibition of filovirus amplification by siRNAs was demonstrated by targeting VP30, VP35, and the NP of MARV [62]. Recently, siRNAs targeting the EBOV L protein were shown to inhibit EBOV replication *in vitro*, as well as providing post-exposure protection to guinea pigs infected with EBOV [63]. Future studies in nonhuman primates may reveal the therapeutic potential of siRNA treatment of humans with filovirus infections.

Antisense treatment for flavivirus infections

The family *Flaviviridae* includes highly pathogenic viruses such as West Nile, yellow fever, dengue and Omsk

hemorrhagic fever. The flaviviruses are encoded by a single-stranded, positive-sense RNA genome, and are characterized by an isometric, enveloped virion. Dengue virus is the only flavivirus considered a Category A biodefense threat by the National Institute of Allergy and Infectious Diseases (http://www3.niaid.nih.gov/biodefense/bandc_priority.htm). However, it was excluded as a biowarfare threat by the American Medical Association Consensus group because it is not transmissible as a small-particle aerosol, and primary dengue infections are only rarely fatal [1]. Dengue virus causes 100 million cases of dengue fever and > 500,000 cases of hemorrhagic fever in humans each year in more than 112 countries [6,64]. Dengue-virus-infected patients present with a variety of symptoms from undifferentiated fever, hemorrhagic fever, or a 'shock-like' syndrome.


A series of peptide-conjugated (P-) PMOs have been screened and several are highly effective at inhibiting dengue viral replication by targeting RNA replication, translation of viral proteins, or both [65,66]. Among the PMOs tested is the 3'-cyclization sequence (3'CS), which is designed to bind to a perfectly conserved sequence at the 3' end of the viral genome required for circularization and replication of the dengue virus genome [67,68]. Encouragingly, the 3'CS P-PMO was able to effectively reduce viral titers to nearly undetectable levels for all four serotypes [65]. Other PMOs targeting a 5'-stem-loop (5'SL) and a 3'-stem-loop (3'SL) were also effective in inhibiting viral replication [65,66]. These studies are informative both from a therapeutic design standpoint, and in revealing portions of the flavivirus genome that regulate dengue virus replication. PMOs are also effective in viral suppression of West Nile virus, another flavivirus, by blocking viral translation and RNA replication [69].

Antisense molecules as antibacterials

Because of the emergence of antibiotic-resistant strains of bacteria, it is imperative that novel antibacterials are developed, especially ones that are effective against a broad range of bacterial agents. To date, the development of a new chemical class of antisense antibacterials is still in its infancy but has shown some promise against certain bacterial agents [70,71]. Synthetic ASOs, called peptide nucleic acids (PNA) can inhibit the expression of *Escherichia coli* bacterial genes both *in vitro* and directly in the organism [72,73]. However, the entry of PNA is inefficient because of the lipopolysaccharide layer of the outer membrane of Gram-negative bacteria [74]. Antisense PMOs have limited cellular uptake in *E. coli* but can efficiently inhibit targeted gene expression in mutant *E. coli* cells that have a leaky outer membrane [75]. The feasibility of using ASOs as antimicrobial agents against Gram-positive bacteria was recently demonstrated for the infectious agent *Mycobacterium tuberculosis*. Targeted knockdown of glutamine synthetase mRNA by using phosphorothioate-modified antisense oligodeoxyribonucleotides modestly inhibited glutamine synthetase activity, formation of the cell wall structure, and bacterial replication [76]. In search of more promising targets of *M. tuberculosis*, PTOs targeted against the 30/32-kDa mycolyl transferase complex showed increased inhibition of bacterial growth [77].

A major limitation of ASOs and their derivatives as gene-silencing agents is their poor uptake by target organisms. Studies have shown that altering the length and target position of antisense PMOs can enhance uptake and inhibit gene-specific expression in *E coli* [78]. Shorter PMOs (approximately 9 to 12 bases in length) effectively inhibit target gene expression in *E coli* culture, while longer PMOs (~ 20 bases) showed better inhibitory activity in cell-free expression systems than in culture [78]. A short, 11-base PMO targeted to an *E coli* essential gene relating to phospholipid biosynthesis (*acpP*) inhibited bacterial growth in culture and in infected mice [70]. Efficient delivery into cells can also be enhanced by conjugating synthetic oligonucleotides with peptides [31••,79,80]. Coupling peptides to PNA enhanced the entry of the antisense molecules and reduced expression of the bacterial target genes both in *E coli* [81] and *Staphylococcus aureus* [82]. Peptide-tagged PMOs can also efficiently inhibit bacterial growth in pure and infected cultures [75]. In a recent study, we observed that peptide-tagged PMOs could inhibit the outgrowth of Sterne strain of *B anthracis* (Figure 1) [Panchal RC, *et al*, unpublished data]. Studies are ongoing to determine the specificity of the target PMOs and their effect on the virulent Ames strain of *B anthracis*. While the application of antisense antibiotics to the treatment of bacterial infections looks promising, a number of issues, such as specificity, optimal size, uptake, mechanism of action, and their efficacy in animal models remain to be addressed.

Figure 1. Peptide-tagged PMOs inhibit outgrowth of *B anthracis*.



An arginine-rich peptide-tagged PMO was incubated with *B anthracis* (Sterne strain) and growth of the bacteria was assessed by light microscopy after 4 h.

Host-directed antisense-based therapeutics against biothreat viral agents

The explosion of discoveries in the area of cellular machineries and signal-transduction pathways in the past decade has had an enormous impact on medicine. Several new FDA-approved anticancer drugs (imatinib, erlotinib, gefitinib, trastuzumab and rituximab) have emerged from these studies, underscoring the importance of a deep

understanding of cellular pathways for countermeasure development. While cellular pathways involved in tumorigenesis have been extensively studied, our understanding of the host pathways in the pathogenesis of viruses is still lagging behind. Understanding the cellular machineries involved in the entry, uncoating, replication, assembly and budding of viruses, as well as delineating the signaling pathways governing the genesis of the innate immune response to pathogens or subversion of such pathways by viruses, will be key to devising such novel host-targeted therapeutics. A major challenge in studying signaling pathways, in particular in complex systems such as host-pathogen interactions, is the concurrent activation or modulation of multiple cascades and cross-talk between individual pathways. In this regard gene knockdown technologies will be invaluable as a therapeutic option as well as a tool for delineating complex cellular pathways.


PMO oligonucleotides are prime candidates for such applications due to their ability to effectively downregulate the expression of cellular genes. Our studies show that PMOs can readily enter primary human cells such as dendritic cells, monocytes and natural killer cells (Figure 2A to C). We were able to effectively block the expression of several cellular genes by using PMOs, including vacuolar protein sorting (VPS)-4 (Figure 2D). Given the demonstrated efficacy of PMOs as gene-targeted drugs in higher animals [60••], this class of compounds represents a promising approach for developing host-targeted, broad-spectrum antiviral therapeutics. One can envision a variety of cellular targets for PMO antiviral therapeutics, including virus receptors, proteolytic enzymes involved in budding and the release of virions from infected cells, cellular signaling molecules, and components of endocytic and vacuolar protein sorting machinery.

PMOs can also be used to delineate complex signaling pathways involved in host-pathogen interactions. Combined with global systems biology-based approaches, specific gene knockdowns using PMOs will allow us to closely examine complex and interconnected biological pathways and to obtain a global view of the cellular processes during host-pathogen interactions. Information extracted from such studies can be fed into bioinformatics and computational tools to create preliminary models of combined pathogen and host response systems or pathways to distill common pathogenic and protective mechanisms. These models can then be exploited to design novel therapeutic modalities for broad-spectrum antivirals.

Conclusion

Currently, multiple viral and bacterial agents represent the most likely bioterrorism or biowarfare threats by intentional transmission or natural outbreaks of disease. Furthermore, emerging and dangerous viral and bacterial infections continue to appear throughout the world, and pose legitimate worldwide health threats. While vaccination is invaluable as a defensive resource, prophylactic measures may encounter public resistance and may not meet the risk-to-

Figure 2. Entry of PMOs into human cells.



(A) HeLa cells were treated with 2 μ M fluorescein-tagged PMO, with or without peptide conjugation, and analyzed by flow cytometry after 2 h (white bars), 24 h (grey bars), or 48 h (black bars). (B) Entry of fluorescein-tagged PMOs into human dendritic cells, measured by flow cytometry as the fluorescence intensities (FL1-Height) of cells per microliter (Counts): filled trace = medium control; green trace = 2 h arginine-rich peptide-tagged PMO; red trace = 24 h arginine-rich peptide-tagged PMO; blue trace = 24 h unconjugated PMO. (C) Uptake of PMO in dendritic cell visualized by confocal microscopy: red, golgi and endoplasmic reticulum staining; green, PMO; yellow, merge (colocalization); blue, nuclear staining. (D) Western blot demonstrating that treating dendritic cells with a PMO targeted against human VPS4 gene, but not a control PMO, completely blocked VPS4A expression. Interestingly, this treatment forced the ribosomes to initiate transcription from a downstream in-frame ATG site, creating a potential dominant negative. Lower band, alternate open reading frame (Alt. ORF). Inhibition was seen as early as 24 h post-treatment and lasted for 72 h.

benefit ratio for vaccinating healthy individuals. Therefore, the most effective means of saving lives in case of an attack with a biological agent is a variety of therapeutic drugs for treating potentially exposed or ill patients in concert with prophylactic drugs and vaccines for unexposed or recently exposed populations. With the availability of rapid diagnostics and sequencing of biological agents, the causative agents in intentional or natural outbreaks of disease should be readily identified and antisense drugs could be quickly designed, if they are not already available. However, as viruses have multiple genes, a number of target sequences for each gene must be screened to identify an effective inhibitor. Therefore, the process of designing an effective antisense-based antiviral may involve the synthesis and screening of many candidates. Based on the current lack of therapeutics for most of the biothreat agents, antisense oligomer-based drugs represent one of the most efficient means of addressing emerging infectious diseases in a timely manner. The promise of specific and relatively safe therapeutics based on antisense technologies has led to recent advancements for developing treatments for diseases caused by biothreat pathogens, including Ebola and dengue viruses, as well as highly virulent bacterial agents.

Acknowledgements

The authors thank D Lane and T Kenny for excellent technical assistance, JL Smith for help preparing the manuscript, Dr G Ruthel for assistance in obtaining microscopy images, and Drs DL Swenson and CL Fuller for support and helpful discussions. A portion of the research described herein was sponsored by the Defense Threat Reduction Agency JSTO-CBD, the Medical Research and Material Command Research Plan # 03-4-7J-021, and in part with federal funds from the National Cancer Institute, National Institutes of Health, under contract N01-CO-12400, and in part by the Developmental Therapeutics Program in the Division of Cancer Treatment and Diagnosis of the National Cancer Institute. Research performed at USAMRIID was conducted in compliance with the Animal Welfare Act and other federal statutes and regulations relating to animals and experiments involving animals and adheres to principles stated in the *Guide for the Care and Use of Laboratory Animals*, National Research Council (1996). USAMRIID is fully accredited by the Association for the Assessment and Accreditation of Laboratory Animal Care International. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial

products, or organizations imply endorsement by the US Government. Opinions, interpretations, conclusions and recommendations are those of the authors and are not necessarily endorsed by the US Army.

References

- of outstanding interest
 - of special interest
1. Borio L, Inglesby T, Peters CJ, Schmaljohn AL, Hughes JM, Jahrling PB, Ksiazek T, Johnson KM, Meyerhoff A, O'Toole T, Ascher MS *et al*: **Hemorrhagic fever viruses as biological weapons: Medical and public health management.** *J Am Med Assoc* (2002) **287**(18):2391-2405.
 2. Williams P, Wallace D: *Unit 731, Japan's Secret Biological Warfare in World War II.* Free Press, New York, NY, USA (1989):38-40.
 3. Alibek K, Handelman S (Eds): *Biohazard.* Random House, New York, NY, USA (1999).
 4. Miller J, Engelberg S, Broad W (Eds): *Germs: Biological Weapons and America's Secret War.* Simon & Schuster, New York, NY, USA (2001).
 5. Center for Nonproliferation Studies: **Chemical and biological weapons: Possession and programs past and present.** *Company Web Site* (2000).
<http://cns.miis.edu/research/cbw/possess.htm>
 6. Richmond JY, McKinney RW (Eds): *Biosafety in Microbiological and Biomedical Laboratories*, 4th Edition. US Department of Health and Human Services, Centers for Disease Control and Prevention, NIH, MD, USA (1999):1-258.
 7. Burnett J, Henchal EA, Schmaljohn A, Bavari S: **The evolving field of biodefence: Therapeutic developments and diagnostics.** *Nat Rev Drug Discov* (2005) **4**(4):281-297.
 - A broad review of prospective therapeutics and therapeutics in development against biodefence agents.
 8. Stephenson ML, Zamecnik PC: **Inhibition of Rous sarcoma viral RNA translation by a specific oligodeoxyribonucleotide.** *Proc Natl Acad Sci USA* (1978) **75**(1):285-288.
 9. Zamecnik PC, Stephenson ML: **Inhibition of Rous sarcoma virus replication and cell transformation by a specific oligodeoxynucleotide.** *Proc Natl Acad Sci USA* (1978) **75**(1):280-284.
 10. Stein CA, Matsukura M, Subasinghe C, Broder S, Cohen JS: **Phosphorothioate oligodeoxynucleotides are potent sequence nonspecific inhibitors of de novo infection by HIV.** *AIDS Res Hum Retroviruses* (1989) **5**(6):639-646.
 11. Crooke S: **Basic principles of antisense technology.** In: *Antisense Drug Technology: Principles, Strategies, and Applications.* Crooke S (Ed), Marcel Dekker Inc, New York, NY, USA (2001):1-28.
 12. Dirksen ML, Crouch RJ: **Selective inhibition of RNase H by dextran.** *J Biol Chem* (1981) **256**(2):11569-11573.
 13. Monia BP, Lesnik EA, Gonzalez C, Lima WF, McGee D, Guinosso CJ, Kawasaki AM, Cook PD, Freier SM: **Evaluation of 2'-modified oligonucleotides containing 2'-deoxy gaps as antisense inhibitors of gene expression.** *J Biol Chem* (1993) **268**(19):14514-14522.
 14. McKay RA, Cummins LL, Graham MJ, Lesnik EA, Owens SR, Winniman M, Dean NM: **Enhanced activity of an antisense oligonucleotide targeting murine protein kinase C- α by the incorporation of 2'-O-propyl modifications.** *Nucleic Acids Res* (1996) **24**(3):411-417.
 15. Baker BF, Lot SS, Condon TP, Cheng-Flournoy S, Lesnik EA, Sasmor HM, Bennett CF: **2'-O-(2-Methoxy)ethyl-modified anti-intercellular adhesion molecule 1 (ICAM-1) oligonucleotides selectively increase the ICAM-1 mRNA level and inhibit formation of the ICAM-1 translation initiation complex in human umbilical vein endothelial cells.** *J Biol Chem* (1997) **272**(18):11994-12000.
 16. Monia BP, Johnston JF, Sasmor H, Cummins LL: **Nuclease resistance and antisense activity of modified oligonucleotides targeted to H-ras.** *J Biol Chem* (1996) **271**(24):14533-14540.
 17. Monia BP, Sasmor H, Johnston JF, Freier SM, Lesnik EA, Muller M, Geiger T, Altmann KH, Moser H, Fabbro D: **Sequence-specific antitumor activity of a phosphorothioate oligodeoxyribonucleotide targeted to human C-raf kinase supports an antisense mechanism of action in vivo.** *Proc Natl Acad Sci USA* (1996) **93**(26):15481-15484.
 18. Lesnik EA, Guinosso CJ, Kawasaki AM, Sasmor H, Zounes M, Cummins LL, Ecker DJ, Cook PD, Freier SM: **Oligodeoxynucleotides containing 2'-O-modified adenosine: Synthesis and effects on stability of DNA:RNA duplexes.** *Biochemistry* (1993) **32**(30):7832-7838.
 19. Freier SM, Altmann KH: **The ups and downs of nucleic acid duplex stability: Structure-stability studies on chemically-modified DNA:RNA duplexes.** *Nucleic Acids Res* (1997) **25**(22):4429-4443.
 20. Zhang H, Cook J, Nickel J, Yu R, Stecker K, Myers K, Dean NM: **Reduction of liver Fas expression by an antisense oligonucleotide protects mice from fulminant hepatitis.** *Nat Biotechnol* (2000) **18**(8):862-867.
 21. Agrawal S, Mayrand SH, Zamecnik PC, Pederson T: **Site-specific excision from RNA by RNase H and mixed-phosphate-backbone oligodeoxynucleotides.** *Proc Natl Acad Sci USA* (1990) **87**(4):1401-1405.
 22. Yamaguchi K, Papp B, Zhang D, Ali AN, Agrawal S, Byrn RA: **The multiple inhibitory mechanisms of GEM 91, a gag antisense phosphorothioate oligonucleotide, for human immunodeficiency virus type 1.** *AIDS Res Hum Retroviruses* (1997) **13**(7):545-554.
 23. Wahlestedt C, Salmi P, Good L, Kela J, Johnsson T, Hokfelt T, Broberger C, Porreca F, Lai J, Ren K, Ossipov M *et al*: **Potent and nontoxic antisense oligonucleotides containing locked nucleic acids.** *Proc Natl Acad Sci USA* (2000) **97**(10):5633-5638.
 24. Geary RS, Henry SP, Grillone LR: **Fomivirsen: Clinical pharmacology and potential drug interactions.** *Clin Pharmacokinet* (2002) **41**(4):255-260.
 25. Hudziak RM, Barofsky E, Barofsky DF, Weller DL, Huang SB, Weller DD: **Resistance of morpholino phosphorodiamidate oligomers to enzymatic degradation.** *Antisense Nucleic Acid Drug Dev* (1996) **6**(4):267-272.
 26. Giles RV, Spiller DG, Clark RE, Tidd DM: **Antisense morpholino oligonucleotide analog induces missplicing of C-myc mRNA.** *Antisense Nucleic Acid Drug Dev* (1999) **9**(2):213-220.
 27. Iversen P: **Phosphorodiamidate morpholino oligomers.** In: *Antisense Drug Technology: Principles, Strategies, and Applications.* Crooke ST (Ed), Marcel Dekker Inc, New York, NY, USA (2001):375-389.
 28. Neuman BW, Stein DA, Kroeker AD, Paulino AD, Moulton HM, Iversen PL, Buchmeier MJ: **Antisense morpholino-oligomers directed against the 5' end of the genome inhibit coronavirus proliferation and growth.** *J Virol* (2004) **78**(11):5891-5899.
 29. Nasevicius A, Ekker SC: **Effective targeted gene 'knockdown' in zebrafish.** *Nat Genet* (2000) **26**(2):216-220.
 30. Moulton HM, Hase MC, Smith KM, Iversen PL: **HIV Tat peptide enhances cellular delivery of antisense morpholino oligomers.** *Antisense Nucleic Acid Drug Dev* (2003) **13**(1):31-43.
 31. Moulton HM, Nelson MH, Hatlevig SA, Reddy MT, Iversen PL: **Cellular uptake of antisense morpholino oligomers conjugated to arginine-rich peptides.** *Bioconj Chem* (2004) **15**(2):290-299.
 - Important study showing that arginine-rich peptides enhance delivery of antisense PMOs.
 32. Summerton J, Weller D: **Morpholino antisense oligomers: Design, preparation, and properties.** *Antisense Nucleic Acid Drug Dev* (1997) **7**(3):187-195.
 33. Bertrand JR, Pottier M, Vekris A, Opolon P, Maksimenko A, Malvy C: **Comparison of antisense oligonucleotides and siRNAs in cell culture and in vivo.** *Biochem Biophys Res Commun* (2002) **296**(4):1000-1004.
 34. Elbashir SM, Harborth J, Lendeckel W, Yalcin A, Weber K, Tuschl T: **Duplexes of 21-nucleotide RNAs mediate RNA interference in cultured mammalian cells.** *Nature* (2001) **411**(6836):494-498.

35. Meister G, Landthaler M, Dorsett Y, Tuschl T: **Sequence-specific inhibition of microRNA- and siRNA-induced RNA silencing.** *RNA* (2004) **10**(3):544-550.
36. Tomari Y, Zamore PD: **Perspective: Machines for RNAi.** *Genes Dev* (2005) **19**(5):517-529.
37. McCaffrey AP, Meuse L, Pham TT, Conklin DS, Hannon GJ, Kay MA: **RNA interference in adult mice.** *Nature* (2002) **418**(6893):38-39.
38. Sato Y, Ajiki T, Inoue S, Fujishiro J, Yoshino H, Igarashi Y, Hakamata Y, Kaneko T, Murakami T, Kobayashi E: **Gene silencing in rat-liver and limb grafts by rapid injection of small interference RNA.** *Transplantation* (2005) **79**(2):240-243.
39. Urban-Klein B, Werth S, Abuharheid S, Czubyko F, Aigner A: **RNAi-mediated gene-targeting through systemic application of polyethylenimine (PEI)-complexed siRNA in vivo.** *Gene Ther* (2005) **12**(5):461-466.
40. Soutschek J, Akinc A, Bramlage B, Charisse K, Constien R, Donoghue M, Elbashir S, Geick A, Hadwiger P, Harborth J, John M *et al*: **Therapeutic silencing of an endogenous gene by systemic administration of modified siRNAs.** *Nature* (2004) **432**(7014):173-178.
- **Demonstration of gene silencing in mice using chemically modified siRNAs.**
41. Cossum PA, Sasmor H, Dellinger D, Truong L, Cummins L, Owens SR, Markham PM, Shea JP, Crooke S: **Disposition of the ¹⁴C-labeled phosphorothioate oligonucleotide ISIS 2105 after intravenous administration to rats.** *J Pharmacol Exp Ther* (1993) **267**(3):1181-1190.
42. Agrawal S, Tamsamani J, Tang JY: **Pharmacokinetics, biodistribution, and stability of oligodeoxynucleotide phosphorothioates in mice.** *Proc Natl Acad Sci USA* (1991) **88**(17):7595-7599.
43. Yu RZ, Geary RS, Monteith DK, Matson J, Truong L, Fitchett J, Levin AA: **Tissue disposition of 2'-O-(2-methoxy) ethyl modified antisense oligonucleotides in monkeys.** *J Pharm Sci* (2004) **93**(1):48-59.
44. Noll BO, McCluskie MJ, Sniatala T, Lohner A, Yuill S, Krieg AM, Schetter C, Davis HL, Uhlmann E: **Biodistribution and metabolism of immunostimulatory oligodeoxynucleotide CPG 7909 in mouse and rat tissues following subcutaneous administration.** *Biochem Pharmacol* (2005) **69**(6):981-991.
45. Krieg AM, Stein CA: **Phosphorothioate oligodeoxynucleotides: Antisense or anti-protein?** *Antisense Res Dev* (1995) **5**(4):241.
46. Nicklin PL, Bayley D, Giddings J, Craig SJ, Cummins LL, Hastewell JG, Phillips JA: **Pulmonary bioavailability of a phosphorothioate oligonucleotide (CGP 64128A): Comparison with other delivery routes.** *Pharm Res* (1998) **15**(4):583-591.
47. Soler M, McHutchison JG, Kwoh TJ, Dorr FA, Pawlotsky JM: **Virological effects of ISIS 14803, an antisense oligonucleotide inhibitor of hepatitis C virus (HCV) internal ribosome entry site (IRES), on HCV IRES in chronic hepatitis C patients and examination of the potential role of primary and secondary HCV resistance in the outcome of treatment.** *Antivir Ther* (2004) **9**(6):953-968.
48. Olie RA, Hall J, Natt F, Stahel RA, Zangemeister-Wittke U: **Analysis of ribosyl-modified, mixed backbone analogs of a bcl-2/bcl-xL antisense oligonucleotide.** *Biochim Biophys Acta* (2002) **1576**(1-2):101-109.
49. Amantana A, Iversen PL: **Pharmacokinetics and biodistribution of phosphorodiamidate morpholino antisense oligomers.** *Curr Opin Pharmacol* (2005) **5**(5):550-555.
50. Iversen PL, Arora V, Acker AJ, Mason DH, Devi GR: **Efficacy of antisense morpholino oligomer targeted to c-myc in prostate cancer xenograft murine model and a phase I safety study in humans.** *Clin Cancer Res* (2003) **9**(7):2510-2519.
51. Devi GR, Beer TM, Corless CL, Arora V, Weller DL, Iversen PL: **In vivo bioavailability and pharmacokinetics of a c-MYC antisense phosphorodiamidate morpholino oligomer, AVI-4126, in solid tumors.** *Clin Cancer Res* (2005) **11**(10):3930-3938.
52. Braasch DA, Paroo Z, Constantinescu A, Ren G, Oz OK, Mason RP, Corey DR: **Biodistribution of phosphodiester and phosphorothioate siRNA.** *Bioorg Med Chem Lett* (2004) **14**(5):1139-1143.
53. Whelan J: **First clinical data on RNAi.** *Drug Disc Today* (2005) **10**(15):1014-1015.
- **Describes first clinical trial data using siRNA to treat age-related macular degeneration.**
54. Villinger F, Rollin PE, Brar SS, Chikkala NF, Winter J, Sundstrom JB, Zaki SR, Swanepoel R, Ansari AA, Peters CJ: **Markedly elevated levels of interferon (IFN)- γ , IFN- α , interleukin (IL)-2, IL-10, and tumor necrosis factor- α associated with fatal Ebola virus infection.** *J Infect Dis* (1999) **179**(Suppl 1):S188-S191.
55. Leroy EM, Baize S, Debre P, Lansoud-Soukate J, Mavoungou E: **Early immune responses accompanying human asymptomatic Ebola infections.** *Clin Exp Immunol* (2001) **124**(3):453-460.
56. Warfield KL, Perkins JG, Swenson DL, Deal EM, Bosio CM, Aman MJ, Yokoyama WM, Young HA, Bavari S: **Role of natural killer cells in innate protection against lethal Ebola virus infection.** *J Exp Med* (2004) **200**(2):169-179.
57. Warfield KL, Olinger G, Deal EM, Swenson DL, Bailey M, Negley DL, Hart MK, Bavari S: **Induction of humoral and CD8+ T cell responses are required for protection against lethal Ebola virus infection.** *J Immunol* (2005) **175**(2):1184-1191.
58. Feldmann H, Kiley MP: **Classification, structure, and replication of filoviruses.** *Curr Top Microbiol Immunol* (1999) **235**:1-21.
59. Geisbert TW, Hensley LE, Jahrling PB, Larsen T, Geisbert JB, Paragas J, Young HA, Fredeking TM, Rote WE, Vlasuk GP: **Treatment of Ebola virus infection with a recombinant inhibitor of factor VIIa/tissue factor: A study in rhesus monkeys.** *Lancet* (2003) **362**(9400):1953-1958.
60. Warfield KL, Swenson DL, Olinger GG, Nichols DK, Pratt WD, Blouch R, Stein DA, Aman MJ, Iversen PL, Bavari S: **Gene-specific countermeasures against Ebola virus based on antisense phosphorodiamidate morpholino oligomers.** *PLoS Pathog* (2006) **2**(1):e1.
- **First demonstration of PMO-mediated protection against a virus. Survival against Ebola virus was mediated in both rodents and nonhuman primates.**
61. Enterlein S, Warfield KL, Swenson DL, Stein DA, Smith JL, Gamble CS, Kroeker AD, Iversen PL, Bavari S, Muhlberger E: **VP35 knockdown inhibits Ebola virus amplification and protects against lethal infection in mice.** *Antimicrob Agents Chemother* (2006) **50**(3):984-993.
62. Fowler T, Bamberg S, Moller P, Klenk HD, Meyer TF, Becker S, Rudel T: **Inhibition of Marburg virus protein expression and viral release by RNA interference.** *J Gen Virol* (2005) **86**(Pt 4):1181-1188.
63. Geisbert TW, Hensley LE, Kagan E, Yu EZ, Geisbert JB, Daddario-DiCaprio K, Fritz EA, Jahrling PB, McClintock K, Phelps JR *et al*: **Postexposure protection of guinea pigs against a lethal Ebola Virus challenge by RNA interference.** *J Infect Dis* (2006): in press.
64. Pinheiro FP, Corber SJ: **Global situation of dengue and dengue hemorrhagic fever and its emergence in the Americas.** *World Health Stat Q* (1997) **50**(3-4):161-168.
65. Kinney RM, Huang CY-H, Rose BC, Kroeker AD, Dreher TW, Iversen PL, Stein DA: **Inhibition of dengue virus serotypes 1 to 4 in vero cell cultures with morpholino oligomers.** *J Virol* (2005) **79**(8):5116-5128.
66. Holden KL, Stein DA, Pierson TC, Ahmed AA, Clyde K, Iversen PL, Harris E: **Inhibition of dengue virus translation and RNA synthesis by a morpholino oligomer targeted to the top of the terminal 3' stem-loop structure.** *Virology* (2006) **344**(2):439-452.
67. Hahn CS, Hahn YS, Rice CM, Lee E, Dalgarno L, Strauss EG, Strauss JH: **Conserved elements in the 3' untranslated region of flavivirus RNAs and potential cyclization sequences.** *J Mol Biol* (1987) **198**(1):33-41.
68. Khromykh AA, Meka H, Guyatt KJ, Westaway EG: **Essential role of cyclization sequences in flavivirus RNA replication.** *J Virol* (2001) **75**(14):6719-6728.
69. Deas TS, Binduga-Gajewska I, Tilgner M, Ren P, Stein DA, Moulton HM, Iversen PL, Kauffman EB, Kramer LD, Shi PY: **Inhibition of flavivirus infections by antisense oligomers specifically suppressing viral translation and RNA replication.** *J Virol* (2005) **79**(8):4599-4609.
- **Recent article examines the mechanisms of PMO-mediated inhibition of flavivirus replication.**

70. Geller BL: **Antibacterial antisense.** *Curr Opin Mol Ther* (2005) 7(2):109-113.
71. Geller BL, Deere J, Tilley L, Iversen PL: **Antisense phosphorodiamidate morpholino oligomer inhibits viability of *Escherichia coli* in pure culture and in mouse peritonitis.** *J Antimicrob Chemother* (2005) 55(6):983-988.
72. Good L, Nielsen PE: **Antisense inhibition of gene expression in bacteria by PNA targeted to mRNA.** *Nat Biotechnol* (1998) 16(4):355-358.
73. Good L, Nielsen PE: **Inhibition of translation and bacterial growth by peptide nucleic acid targeted to ribosomal RNA.** *Proc Natl Acad Sci USA* (1998) 95(5):2073-2076.
74. Good L, Sandberg R, Larsson O, Nielsen PE, Wahlestedt C: **Antisense PNA effects in *Escherichia coli* are limited by the outer-membrane LPS layer.** *Microbiology* (2000) 146(Pt 10):2665-2670.
75. Geller BL, Deere JD, Stein DA, Kroeker AD, Moulton HM, Iversen PL: **Inhibition of gene expression in *Escherichia coli* by antisense phosphorodiamidate morpholino oligomers.** *Antimicrob Agents Chemother* (2003) 47(10):3233-3239.
76. Harth G, Zamecnik PC, Tang J-Y, Tabatadze D, Horwitz MA: **Treatment of *Mycobacterium tuberculosis* with antisense oligonucleotides to glutamine synthetase mRNA inhibits glutamine synthetase activity, formation of the poly-L-glutamate/glutamine cell wall structure, and bacterial replication.** *Proc Natl Acad Sci USA* (2000) 97(1):418-423.
77. Harth G, Horwitz MA, Tabatadze D, Zamecnik PC: **Targeting the *Mycobacterium tuberculosis* 30/32-kDa mycolyl transferase complex as a therapeutic strategy against tuberculosis: Proof of principle by using antisense technology.** *Proc Natl Acad Sci USA* (2002) 99(24):15614-15619.
78. Deere J, Iversen P, Geller BL: **Antisense phosphorodiamidate morpholino oligomer length and target position effects on gene-specific inhibition in *Escherichia coli*.** *Antimicrob Agents Chemother* (2005) 49(1):249-255.
79. Gait MJ: **Peptide-mediated cellular delivery of antisense oligonucleotides and their analogues.** *Cell Mol Life Sci* (2003) 60(5):844-853.
80. Stetsenko DA, Arzumov AA, Korshun VA, Gait MJ: **[Peptide-oligonucleotide conjugates as a new generation of antisense elements].** *Mol Biol Mosk* (2000) 34(6):998-1006.
81. Good L, Awasthi SK, Dryselius R, Larsson O, Nielsen PE: **Bactericidal antisense effects of peptide-PNA conjugates.** *Nat Biotechnol* (2001) 19(4):360-364.
82. Nekhotiaeva N, Awasthi SK, Nielsen PE, Good L: **Inhibition of *Staphylococcus aureus* gene expression and growth using antisense peptide nucleic acids.** *Mol Ther* (2004) 10(4):652-659.
83. Boiziau C, Blonski C, Thuong NT, Shire D, Toulme JJ: **Inhibition of reverse transcription by unmodified and modified antisense oligodeoxynucleotides.** *Nucleic Acids Symp Ser* (1991) (24):121-125.
84. Boiziau C, Thuong NT, Toulme JJ: **Mechanisms of the inhibition of reverse transcription by antisense oligonucleotides.** *Proc Natl Acad Sci USA* (1992) 89(2):768-772.
85. Smith AW, Matson DO, Stein DA, Skilling DE, Kroeker AD, Berke T, Iversen PL: **Antisense treatment of caliciviridae: An emerging disease agent of animals and humans.** *Curr Opin Mol Ther* (2002) 4(2):177-184.
86. Li BJ, Tang Q, Cheng D, Qin C, Xie FY, Wei Q, Xu J, Liu Y, Zheng BJ, Woodle MC, Zhong N *et al*: **Using siRNA in prophylactic and therapeutic regimens against SARS coronavirus in Rhesus macaque.** *Nat Med* (2005) 11(9):944-951.
87. Neuman BW, Stein DA, Kroeker AD, Churchill MJ, Kim AM, Kuhn P, Dawson P, Moulton HM, Bestwick RK, Iversen PL, Buchmeier MJ: **Inhibition, escape, and attenuated growth of severe acute respiratory syndrome coronavirus treated with antisense morpholino oligomers.** *J Virol* (2005) 79(15):9665-9676.
88. Sankar S, Cheah KC, Porter AG: **Antisense oligonucleotide inhibition of encephalomyocarditis virus RNA translation.** *Eur J Biochem* (1989) 184(1):39-45.
89. Pagano JS, Jimenez G, Sung NS, Raab-Traub N, Lin JC: **Epstein-Barr viral latency and cell immortalization as targets for antisense oligomers.** *Ann NY Acad Sci* (1992) 660:107-116.
90. Masciarelli S, Mattioli B, Galletti R, Samoggia P, Chichiarelli S, Mearini G, Mattia E: **Antisense to Epstein Barr Virus-encoded LMP1 does not affect the transcription of viral and cellular proliferation-related genes, but induces phenotypic effects on EBV-transformed B lymphocytes.** *Oncogene* (2002) 21(26):4166-4170.
91. van den Born E, Stein DA, Iversen PL, Snijder EJ: **Antiviral activity of morpholino oligomers designed to block various aspects of *Equine arteritis virus* amplification in cell culture.** *J Gen Virol* (2005) 86(Pt 11):3081-3090.
92. Ropert C, Lavignon M, Dubernet C, Couvreur P, Malvy C: **Oligonucleotides encapsulated in pH sensitive liposomes are efficient toward Friend retrovirus.** *Biochem Biophys Res Commun* (1992) 183(2):879-885.
93. Matsukura M, Koike K, Zon G: **Antisense phosphorothioates as antivirals against human immunodeficiency virus (HIV) and hepatitis B virus (HBV).** *Toxicol Lett* (1995) 82-83:435-438.
94. Yao ZQ, Zhou YX, Feng XM, Chen ZX: **Specific inhibition of hepatitis B virus gene expression by an antisense oligonucleotide *in vitro*.** *Acta Virol* (1995) 39(4):227-230.
95. Yao Z, Zhou Y, Feng X, Chen C, Guo J: ***In vivo* inhibition of hepatitis B viral gene expression by antisense phosphorothioate oligodeoxynucleotides in athymic nude mice.** *J Viral Hepatitis* (1996) 3(1):19-22.
96. Tan TM, Zhou L, Houssais S, Seet BL, Jaenicke S, Peter F, Lim SG: **Intracellular inhibition of hepatitis B virus S gene expression by chimeric DNA-RNA phosphorothioate minimized ribozyme.** *Antisense Nucleic Acid Drug Dev* (2002) 12(4):257-264.
97. Tallet-Lopez B, Aldaz-Carroll L, Chabas S, Dausse E, Staedel C, Toulme JJ: **Antisense oligonucleotides targeted to the domain III of the hepatitis C virus IRES compete with 40S ribosomal subunit binding and prevent *in vitro* translation.** *Nucleic Acids Res* (2003) 31(2):734-742.
98. Nulf CJ, Corey D: **Intracellular inhibition of hepatitis C virus (HCV) internal ribosomal entry site (IRES)-dependent translation by peptide nucleic acids (PNAs) and locked nucleic acids (LNAs).** *Nucleic Acids Res* (2004) 32(13):3792-3798.
99. Sarin PS, Agrawal S, Civeira MP, Goodchild J, Ikeuchi T, Zamecnik PC: **Inhibition of acquired immunodeficiency syndrome virus by oligodeoxynucleoside methylphosphonates.** *Proc Natl Acad Sci USA* (1988) 85(20):7448-7451.
100. Goodchild J, Agrawal S, Civeira MP, Sarin PS, Sun D, Zamecnik PC: **Inhibition of human immunodeficiency virus replication by antisense oligodeoxynucleotides.** *Proc Natl Acad Sci USA* (1988) 85(15):5507-5511.
101. Stevenson M, Iversen PL: **Inhibition of human immunodeficiency virus type 1-mediated cytopathic effects by poly(L-lysine)-conjugated synthetic antisense oligodeoxyribonucleotides.** *J Gen Virol* (1989) 70(Pt 10):2673-2682.
102. Agrawal S, Ikeuchi T, Sun D, Sarin PS, Konopka A, Maizel J, Zamecnik PC: **Inhibition of human immunodeficiency virus in early infected and chronically infected cells by antisense oligodeoxynucleotides and their phosphorothioate analogues.** *Proc Natl Acad Sci USA* (1989) 86(20):7790-7794.
103. Matsukura M, Zon G, Shinozuka K, Robert-Guroff M, Shimada T, Stein CA, Mitsuya H, Wong-Staal F, Cohen JS, Broder S: **Regulation of viral expression of human immunodeficiency virus *in vitro* by an antisense phosphorothioate oligodeoxynucleotide against *rev* (*art/trs*) in chronically infected cells.** *Proc Natl Acad Sci USA* (1989) 86(11):4244-4248.
104. Mori K, Boiziau C, Cazenave C, Matsukura M, Subasinghe C, Cohen JS, Broder S, Toulme JJ, Stein CA: **Phosphorothioate oligodeoxynucleotides: Synthesis, physico-chemical characterization, anti-sense inhibitory properties and anti-HIV activity.** *Nucleic Acids Res* (1989) 17(20):8207-8219.

• Demonstration of PMO-mediated inhibition of the SARS coronavirus.

105. Letsinger RL, Zhang GR, Sun DK, Ikeuchi T, Sarin PS: **Cholesteryl-conjugated oligonucleotides: Synthesis, properties, and activity as inhibitors of replication of human immunodeficiency virus in cell culture.** *Proc Natl Acad Sci USA* (1989) **86**(17):6553-6556.
106. Buck HM, Koole LH, van Genderen MH, Smit L, Geelen JL, Jurriaans S, Goudsmit J: **Phosphate-methylated DNA aimed at HIV-1 RNA loops and integrated DNA inhibits viral infectivity.** *Science* (1990) **248**(4952):208-212.
107. Farooqui F, Sarin PS, Sun D, Letsinger RL: **Effect of structural variations in cholesteryl-conjugated oligonucleotides on inhibitory activity toward HIV-1.** *Bioconjug Chem* (1991) **2**(6):422-426.
108. Laurence J, Sikder SK, Kulkosky J, Miller P, Ts'o PO: **Induction of chronic human immunodeficiency virus infection is blocked *in vitro* by a methylphosphonate oligodeoxynucleoside targeted to a U3 enhancer element.** *J Virol* (1991) **65**(1):213-219.
109. Zamecnik PC, Agrawal S: **Oligodeoxynucleotide hybridization inhibition of HIV and influenza virus.** *Nucleic Acids Symp Ser* (1991) **24**:127-131.
110. Kim SG, Hatta T, Tsukahara S, Nakashima H, Yamamoto N, Shoji Y, Takai K, Takaku H: **Antiviral effect of phosphorothioate oligodeoxyribonucleotides complementary to human immunodeficiency virus.** *Bioorg Med Chem* (1995) **3**(1):49-54.
111. Kuwasaki T, Hosono K, Takai K, Ushijima K, Nakashima H, Saito T, Yamamoto N, Takaku H: **Hairpin antisense oligonucleotides containing 2'-methoxynucleosides with base-pairing in the stem region at the 3'-end: Penetration, localization, and anti-HIV activity.** *Biochem Biophys Res Commun* (1996) **228**(2):623-631.
112. Boiziau C, Tarrago-Litvak L, Sinha ND, Moreau S, Litvak S, Toulme J-J: **Antisense oligonucleotides inhibit *in vitro* cDNA synthesis by HIV-1 reverse transcriptase.** *Antisense Nucleic Acid Drug Dev* (1996) **6**(2):103-109.
113. Hamma T, Miller PS: **Interactions of hairpin oligo-2'-O-methylribonucleotides containing methylphosphonate linkages with HIV TAR RNA.** *Antisense Nucleic Acid Drug Dev* (2003) **13**(1):19-30.
114. Darfeuille F, Hansen JB, Orum H, Di Primo C, Toulme JJ: **LNA/DNA chimeric oligomers mimic RNA aptamers targeted to the TAR RNA element of HIV-1.** *Nucleic Acids Res* (2004) **32**(10):3101-3107.
115. Gao WY, Hanes RN, Vazquez-Padua MA, Stein CA, Cohen JS, Cheng YC: **Inhibition of herpes simplex virus type 2 growth by phosphorothioate oligodeoxynucleotides.** *Antimicrob Agents Chemother* (1990) **34**(5):808-812.
116. Kmetz ME, Ceruzzi M, Schwartz J: **Vmw65 phosphorothioate oligonucleotides inhibit HSV KOS replication and Vmw65 protein synthesis.** *Antiviral Res* (1991) **16**(2):173-184.
117. Leonetti JP, Machy P, Degols G, Lebleu B, Leserman L: **Antibody-targeted liposomes containing oligodeoxyribonucleotides complementary to viral RNA selectively inhibit viral replication.** *Proc Natl Acad Sci USA* (1990) **87**(7):2448-2451.
118. Kean JM, Kipp SA, Miller PS, Kulka M, Aurelian L: **Inhibition of herpes simplex virus replication by antisense oligo-2'-O-methylribonucleoside methylphosphonates.** *Biochemistry* (1995) **34**(45):14617-14620.
119. Aurelian L, Smith CC: **Herpes simplex virus type 2 growth and latency reactivation by cocultivation are inhibited with antisense oligonucleotides complementary to the translation initiation site of the large subunit of ribonucleotide reductase (RR1).** *Antisense Nucleic Acid Drug Dev* (2000) **10**(2):77-85.
120. Alonso M, Stein DA, Thomann E, Moulton HM, Leong JC, Iversen P, Mourich DV: **Inhibition of infectious haematopoietic necrosis virus in cell cultures with peptide-conjugated morpholino oligomers.** *J Fish Dis* (2005) **28**(7):399-410.
121. Archambault D, Stein CA, Cohen JS: **Phosphorothioate oligonucleotides inhibit the replication of lentiviruses and type D retroviruses, but not that of type C retroviruses.** *Arch Virol* (1994) **139**(1-2):97-109.
122. Shan YF, Gao HY, Ji H, Qiu MS, Wang DB: **Antiviral effects on mouse leukemia virus replication by oligodeoxynucleotides *in vitro* and *in vivo*.** *Chin Med J (Engl)* (1994) **107**(2):146-150.
123. Leaman DW, Longano FJ, Okicki JR, Soike KF, Torrence PF, Silverman RH, Cramer H: **Targeted therapy of respiratory syncytial virus in African green monkeys by intranasally administered 2-5A antisense.** *Virology* (2002) **292**(1):70-77.
- **First demonstration of viral knockdown in non-human primates by antisense molecules.**
124. Crum C, Johnson JD, Nelson A, Roth D: **Complementary oligodeoxynucleotide mediated inhibition of tobacco mosaic virus RNA translation *in vitro*.** *Nucleic Acids Res* (1988) **16**(10):4569-4581.
125. Degols G, Leonetti JP, Gagnor C, Lemaitre M, Lebleu B: **Antiviral activity and possible mechanisms of action of oligonucleotides-poly(L-lysine) conjugates targeted to vesicular stomatitis virus mRNA and genomic RNA.** *Nucleic Acids Res* (1989) **17**(22):9341-9350.
126. Shea RG, Marsters JC, Bischofberger N: **Synthesis, hybridization properties and antiviral activity of lipid-oligodeoxynucleotide conjugates.** *Nucleic Acids Res* (1990) **18**(13):3777-3783.
127. Levis JT, Miller PS: **Interactions of psoralen-derivatized oligodeoxyribonucleoside methylphosphonates with vesicular stomatitis virus messenger RNA.** *Antisense Res Dev* (1994) **4**(4):223-230.
128. Stein DA, Skilling DE, Iversen PL, Smith AW: **Inhibition of Vesivirus infections in mammalian tissue culture with antisense morpholino oligomers.** *Antisense Nucleic Acid Drug Dev* (2001) **11**(5):317-325.
- **First demonstration of antiviral effects of PMO in cell culture.**