

UNCLASSIFIED

E. R. D. E. 8/M/51

~~CONFIDENTIAL~~

COPY No. 18



MINISTRY OF SUPPLY

1. THIS INFORMATION IS DISCLOSED FOR OFFICIAL USE BY THE RECIPIENT GOVERNMENT. DISCLOSURE TO ANY OTHER GOVERNMENT OR RELEASE TO THE PRESS OR IN ANY OTHER WAY WOULD BE A BREACH OF THIS CONDITION.
2. THE INFORMATION SHOULD BE SAFEGUARDED UNDER RULES DESIGNED TO GIVE THE SAME STANDARD OF SECURITY AS THAT MAINTAINED BY HIS MAJESTY'S GOVERNMENT IN THE UNITED KINGDOM.
3. THE INFORMATION CONTAINED IN THIS REPORT SHOULD NOT BE CIRCULATED OUTSIDE GOVERNMENT DEPARTMENTS WITHOUT THE PRIOR PERMISSION OF THE MINISTRY OF SUPPLY.

EXPLOSIVES RESEARCH & DEVELOPMENT ESTABLISHMENT

TECHNICAL MEMORANDUM No. 8/M/51

0110
003192

Calculation of the Explosive Power of Mixtures of Chlorine Trifluoride and Ammonia (Both Liquid)

G. H. S. Young

20071109142

*Info #1
ORDG-T2182T-329*

THIS DOCUMENT IS THE PROPERTY OF H.B.M. GOVERNMENT AND ATTENTION IS CALLED TO THE PENALTIES ATTACHING TO ANY INFRINGEMENT OF THE OFFICIAL SECRETS ACTS

It is intended for the use of the recipient only, and for communication to such officers under him as may require to be acquainted with its contents in the course of their duties. The officers exercising this power of communication are responsible that such information is imparted with due caution and reserve. Any person other than the authorised holder, upon obtaining possession of this document, by finding or otherwise, should forward it together with his name and address in a closed envelope to:-
THE SECRETARY, MINISTRY OF SUPPLY, ADELPHI, LONDON, W.C. 2.

Letter postage need not be prepaid, other postage will be refunded. All persons are hereby warned that the unauthorised retention or destruction of this document is an offence against the Official Secrets Acts.

(81)
183974

UNCLASSIFIED

~~CONFIDENTIAL~~

Best Available Copy
July 1991

A.R.E.
Publishing Section

~~CONFIDENTIAL~~

OFFICER GROUP NO.1
UNITED STATES ARMY
U.S. Navy 100, Fleet Post Office
New York, N.Y.

28449

ORD-TRANT-329

4 September 1951

SUBJECT: Calculation of the Explosive Power of Mixtures of Chlorine Trifluoride and Ammonia (Both Liquid)

TO: The Chief of Ordnance
Department of the Army
Washington 25, D.C.

Attn: ORDGU-IN



350.05/901
10 SEP 1951

Englund @ DIN

003192

1. Reference - None

2. Forwarded herewith as Inclosure #1 is E.R.D.E. (Explosives Research & Development Establishment) Technical Memorandum No. 8/M/51 entitled "Calculation of the Explosive Power of Mixtures of Chlorine Trifluoride and Ammonia (Both Liquid)", dated July 1951.

3. Abstract of the Report is as follows:-

The explosive power of mixtures of Liquid NH₃ and ClF₃ within the range of ratios of 1.4 to 2.3 moles of NH₃ per mole ClF₃ are calculated. Calculations are based on the correlation of the Berthelot Product for the mixtures to its value for Picric Acid under normal condition in the explosive calorimeter. The optimum mixture of 2.8 parts by weight of ClF₃ with 1 part of NH₃ gave a calculated value of 188 which is theoretically about equivalent to Methyl Nitrate (185) and somewhat better than RDX (168). These calculations assume intimate mixture and complete reaction which would be very difficult to achieve in practice.

(Trip)
1 Incl: E.R.D.E. Technical Memo
No. 8/M/51

A.W. STODDARD
Lt. Col. Ord Corps
Technical Representative
for Artillery

Copies furnished:

w/Incl #1 - 3-OCO(ORDGU-IN); 1-R&D Bd; 2-OARMA London for ACGS-2
w/o Incl: - 1-DACS G-4 (R&D); 1-OCO(ORDTX); 1-File

PICATINNY ARSENAL LIBRARY
DOVER, N. J.

00000
M1 REC'D 22 SEP 51 OFFICE
OFFICE CHIEF DE PRODUCE

28 11 03 58.21

RECEIVED

~~CONFIDENTIAL~~

UNCLASSIFIED

E.R.D.E. 8/M/51.

~~CONFIDENTIAL~~

XR. 880/12.

MINISTRY OF SUPPLY

EXPLOSIVES RESEARCH AND DEVELOPMENT ESTABLISHMENT

TECHNICAL MEMORANDUM NO. 8/M/51.

Calculation of the Explosive Power of Mixtures of
Chlorine Trifluoride and Ammonia (Both Liquid).

by

G.H.S. Young

This memorandum does not contain classified information of
overseas origin.

Approved: L. A. Wiseman

L.A. WISEMAN
S.P.R.II.

Approved for
Circulation: C. H. Johnson

C.H. JOHNSON
C.S., E.R.D.E.

Waltham Abbey,
Essex.

~~CONFIDENTIAL~~

DISTRIBUTION STATEMENT A
Approved for Public Release
Distribution Unlimited

UNCLASSIFIED

UNCLASSIFIED

~~CONFIDENTIAL~~

DISTRIBUTION

EXTERNAL

TPA3/TIB (29)
Dr. Phillips, B.J.S.M., through TPA3/TIB.

Ministry of Supply.

Chief Scientist
C.E.A.D. (3, including Mr. Thomas, Langhurst)
C.S.A.R. (3, including Dr. Poole)
D.M.X.R.D. (2)
D.G.W.R.D.
D.W.R. (D)
P.D.S.R. (D)
Sec.O.B. (2)

Admiralty.

A.C.S.I.L.

INTERNAL

C.S.,E.R.D.E.
S.P.R.II.
S.E.I.
I.S.R.G.
Mr. N.W. Luft.
Dr. E. Roberts
Dr. G.H.S. Young
Dr. K.A. Cooper
Library (2)
Registry (2)
Information Bureau (2 + stock)

Further copies of this memorandum can be obtained from
Chief Superintendent, Explosives Research and Development
Establishment, Waltham Abbey, Essex.

~~CONFIDENTIAL~~

UNCLASSIFIED

~~CONFIDENTIAL~~


Reference: XR 880/12.

1. INTRODUCTION.

The explosive effectiveness of a given substance relative to that of the accepted standard, picric acid, as measured for example by the Trautzl Block or the Ballistic Mortar methods, is found to be fairly well correlated with the relative value of the "Bertholot Product". The latter is proportional to nQ , where Q is the heat (per gm) which would be liberated by the chemical change which actually takes place in the explosion, if it could take place in the same way at ambient temperature (300°K.), while n is the number of gram molecules of gaseous products obtained in the explosion from one gram of the substance. The Explosive Power (Pr) is therefore defined as one hundred times the ratio of the value of this Bertholot Product for the given substance under stated conditions to its value for Picric Acid under normal conditions in the explosive calorimeter.

2. SYSTEM CONSIDERED.

Assuming approximate thermodynamic equilibrium the principal reaction within the range of mixture ratios studied (1.4 to 2.3 moles. of NH_3 per mole of ClF_3) is as follows:-



As x approaches the boundary value $\frac{4}{3}$, the temperature of the adiabatic constant volume flame rises steeply (see Fig.2) and the dissociation products, atomic chlorine and atomic fluorine become appreciable in amount (3-4%), the amounts of the molecular forms, Cl_2 and F_2 , being somewhat less while the dissociation of hydrogen and nitrogen remains effectively small at the high pressures involved.

The main reaction alters in two stages as x is still further diminished; the hydrogen virtually disappears and atomic chlorine and (ultimately) fluorine appear as main products.

The necessary thermodynamic data for calculating the flame temperature and explosive power at these lower values of x can only be estimated rather imperfectly, but fortunately the optimum mixture ratio, as regards explosive power, appears to lie within the higher range of x as indicated by the graph (Fig.1). It was therefore not necessary to investigate the lower range of x in detail.

The heat of formation of gaseous ClF_3 , about which there has been some variation of opinion, was taken to be 28 k.cal. per mole (H.V. v. Wartenberg, Z.anorg.Chem., 1949, 258, 356) and the latent heat of vaporization as 5.k.cal. per mole. The remaining thermodynamic data was obtained from Pike's tables and from the compilation of the American Bureau of Standards.

The volume of the system was taken to be the sum of the volumes of the liquid components ClF_3 and NH_3 at 300°K. (assumed to be under suitable pressures).

/3.RESULTS.

~~CONFIDENTIAL~~

3. RESULTS.

The essential results of the calculations are as follows :-

Temperature °K.	x	Pr.
3870	2.24	186.55
4065	2.07	187.7
4355	1.85	187.76
4455	1.78	187.65
4645	1.65	187.3
4840	1.51	186.2
4935	1.42	184.6

The decimal places in the value of Pr are of little significance except in showing the smooth trend of the calculated values, with a definite maximum of about 188 at $x = 1.95$.

This maximum figure, 188, for a mixture of 2.8 parts by weight of ClF_3 with 1 part of NH_3 , can be compared with:

<u>Explosive</u>	<u>Pr.</u>
Picric Acid	100
T.N.T.	108
R.D.X.	168
Methyl Nitrate	185
Nitromethane	154

Thus the optimum mixture is theoretically about equivalent to methyl nitrate in explosive power and somewhat better than R.D.X.

It seems apparent from these calculations that liquid fluorine would give a higher explosive power with liquid ammonia, because the energy per unit mass is greater, and there would be no heat of formation term to be taken into account.

The calculations assume that it would be possible to get an intimate mixture, and complete reaction. In practice this would be very difficult to achieve, and explosion rather than detonation, might occur.

4. ACKNOWLEDGMENTS.

The calculations on which this note is based were carried out by Dr. B.M. Cavanagh and Miss M.A. Hooper.





FIG. 1.



~~CONFIDENTIAL~~

UNCLASSIFIED

~~CONFIDENTIAL~~

UNCLASSIFIED