

REPORT DOCUMENTATION PAGE

*Form Approved
OMB No. 0704-0188*

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to the Department of Defense, Executive Services and Communications Directorate (0704-0188). Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ORGANIZATION.

| | | | | | |
|--|-----------------------------|--|----------------------------------|---|---|
| 1. REPORT DATE (DD-MM-YYYY) 26-09-2007 | | 2. REPORT TYPE Conference Proceedings | | 3. DATES COVERED (From - To) | |
| 4. TITLE AND SUBTITLE Application of Creeping Sea-fill Methodology to the Wind Speed Over the Caspian Sea | | | | 5a. CONTRACT NUMBER | |
| | | | | 5b. GRANT NUMBER | |
| | | | | 5c. PROGRAM ELEMENT NUMBER 0602435N | |
| 6. AUTHOR(S) Murat Gundz, Birol Kara, Alan J. Wallcraft, E. Joseph Metzger | | | | 5d. PROJECT NUMBER | |
| | | | | 5e. TASK NUMBER | |
| | | | | 5f. WORK UNIT NUMBER 73-8677-06-5 | |
| 7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004 | | | | 8. PERFORMING ORGANIZATION REPORT NUMBER NRL/PP/7320-06-7035 | |
| 9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5660 | | | | 10. SPONSOR/MONITOR'S ACRONYM(S) ONR | |
| | | | | 11. SPONSOR/MONITOR'S REPORT NUMBER(S) | |
| 12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release, distribution is unlimited | | | | | |
| 13. SUPPLEMENTARY NOTES | | | | | |
| <h1>20071119059</h1> | | | | | |
| 14. ABSTRACT Numerical weather prediction (NWP) products include possible errors near the coastal regions. Effects of such errors on near surface wind speed, which is one of the critical variables for coastal applications are investigated over the Caspian Sea. For this purpose, a creeping sea-fill methodology developed by [1] was applied to the coarse resolution (1.125° x 1.125°) wind speed data obtained from European Centre for Medium-Range Weather Forecasts (ECMWF), and the results compared with the relatively fine resolution (0.25° x 0.25°) satellite based wind speed data. It is shown that the applied methodology improved the accuracy of the wind speed near the coastal regions. | | | | | |
| 15. SUBJECT TERMS Coastal Processes, Air-sea Interactions, Brackish Water | | | | | |
| 16. SECURITY CLASSIFICATION OF: | | | 17. LIMITATION OF ABSTRACT UL | 18. NUMBER OF PAGES 1 | 19a. NAME OF RESPONSIBLE PERSON Alan Wallcraft |
| a. REPORT Unclassified | b. ABSTRACT Unclassified | c. THIS PAGE Unclassified | | | 19b. TELEPHONE NUMBER (Include area code) 228-688-4813 |

38°N 48°E Caspian Sea
 Apr 9-13, 2007 in Istanbul, Turkey

APPLICATION OF CREEPING SEA-FILL METHODOLOGY TO THE WIND SPEED OVER THE CASPIAN SEA

Murat Gunduz^{1*}, Birol Kara², Alan Wallcraft² and Joe Metzger²
¹ Institute of Marine Sciences, METU, Erdemli, Turkey - gunduz@ims.metu.edu.tr
² US Naval Research Laboratory, Code 7320, NASA Stennis Space Center

Abstract

Numerical weather prediction (NWP) products include possible errors near the coastal regions. Effects of such errors on near surface wind speed, which is one of the critical variables for coastal applications are investigated over the Caspian Sea. For this purpose, a creeping sea-fill methodology developed by [1] was applied to the coarse resolution (1.125° x 1.125°) wind speed data obtained from European Centre for Medium-Range Weather Forecasts (ECMWF), and the results compared with the relatively fine resolution (0.25° x 0.25°) satellite based wind speed data. It is shown that the applied methodology improved the accuracy of the wind speed near the coastal regions.

Keywords: Coastal Processes, Air-sea Interactions, Brackish Water.

Coastal applications (e.g regional ocean models, wave models etc.) need accurate wind speed over the sea surface and proper representation of this variable is important. However, atmospheric models do not have enough resolution for inland waters like the Caspian Sea. Since many NWP models are generally designed for global or regional applications, small water bodies are not considered in great detail. Also, because of the irregularities in coastline the few grid points over inland waters can be contaminated by land. For this reason, the wind speed field requires special treatment before using it in coastal applications.

The Caspian Sea is an enclosed water body and very sensitive to the atmospheric forcing, thus improper representations of air-sea fluxes in a regional ocean model could lead to unrealistic trends of total heat, mass, and momentum. For this reason, special attention should be taken to properly represent the atmospheric forcing in this sea. The main goals of this study are (1) to show the effects of land on the surface wind speed near the coastal regions, and (2) to evaluate the success of the applied methodology for reducing the effects of land contamination.

to a 1/25° x 1/25°, cos (lat) resolution Caspian Sea Hybrid Coordinate Ocean Model (HYCOM) domain. HYCOM, described in [2], is a community ocean model (<http://oceanmodeling.rsmas.miami.edu/hycom/>). It behaves like a conventional σ (terrain-following) model in very shallow oceanic regions, like a z-level coordinate model in the mixed layer or other unstratified regions, and like an isopycnic-coordinate model in stratified regions.




Fig. 1. The land sea mask from a satellite-based product (QSCAT, right), and ECMWF interpolated to the 1/25° grid (left)

Land-sea mask values from ECMWF [2] (European Centre for Medium-Range Weather Forecasts) (Fig. 1a) and SeaWinds scatterometer on the QuikSCAT satellite (Fig. 1b) are shown. It is clearly seen that the relatively coarse resolution ECMWF land-sea mask is more contaminated by land values effecting the interior of the sea compared to QSCAT land-sea mask, which has relatively fine resolution. In this figure, a contour value of 0.8 in the land-sea masks implies that the interpolated wind speed values are approx 80% contaminated by land values on the OGCM grid.

In this study, a creeping sea-fill methodology was applied to the wind speed at 10 m obtained from ECMWF 40-year Re-Analysis (ERA40) data for June 2000. The creeping sea-fill simply means that winds are post-processed, so that only values over the ocean (or sea) are used. It replaces the value associated with each land-masked point by using only nearby sea values. For demonstration purposes, the original values were interpolated




Fig. 2. Spatial variations of wind speed from ECMWF, QSCAT and sea-fill field.

Performance of the creeping sea-fill methodology is evaluated using measurements of wind speed magnitudes at 10 m above the surface from QuikSCAT satellite. ERA40 (Figure 2a), QSCAT (Figure 2b) wind speed data and the sea-filled field (Figure 2c) are plotted to examine wind speed during June 2000. While ERA40 wind speed data shows abrupt features along the southern and western coast of the Caspian Sea, QuikSCAT data is more smooth over these regions. An improvement in the accuracy in ERA40 wind speed data is seen after applying creeping sea-fill methodology, as these fields display more or less similar features compared to QuikSCAT data.

This study shows that applying a creeping sea-fill methodology successfully reduces the error due to the land contamination associated with the coarse NWP grid, and improves the accuracy of wind speed near the coastal regions. To investigate possible effects of improved atmospheric forcing on the dynamical features of the sea, a regional ocean model should be used as a tool for a further study.

References

- 1 - Kara A.B., Wallcraft A.J. and Hurlburt H.E., 2006. Land contamination of atmospheric forcing for ocean models near land-sea boundaries. *J. Phys. Oceanogr.*, in press.
- 2 - Uppala, S.M. et al., 2005. The ERA-40 re-analysis. *Q. J. Roy. Meteor. Soc.*, 2, 131: 961-3012.
- 3 - Bleck R., 2002. An oceanic general circulation model framed in hybrid isopycnic-cartesian coordinates. *Ocean Modelling*, 4: 55-88.