

# REPORT DOCUMENTATION PAGE

*Form Approved*  
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

1. AGENCY USE ONLY <i>(Leave blank)</i>	2. REPORT DATE May, 2005	3. REPORT TYPE AND DATES COVERED Proceedings Article	
4. TITLE AND SUBTITLE NONINVASIVE WARNING INDICATOR OF THE RED ZONE OF POTENTIAL THERMAL INJURY AND PERFORMANCE IMPAIRMENT: A PILOT STUDY.		5. FUNDING NUMBERS	
6. AUTHOR(S) Miyo Yokota, Daniel Moran, Larry Berglund, Lou Stephenson, Margaret Kolka		8. PERFORMING ORGANIZATION REPORT NUMBER M05-16	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Biophysics and Biomedical Modeling Division U.S. Army Research Institute of Environmental Medicine Building 42 - Kansas Street Natick, MA 01760		10. SPONSORING / MONITORING AGENCY REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, MD 21702		11. SUPPLEMENTARY NOTES	
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited		12b. DISTRIBUTION CODE	
13. ABSTRACT <i>(Maximum 200 words)</i> The guidelines for assessing worker's level of heat strain in order to prevent thermal injury and performance impairment has been widely adapted in industries to promote workers' health and safety. Core temperature (Tcore) is a common physiological parameter used in the guidelines for heat stress; however, measuring Tcore may be invasive and impractical for real time monitoring a worker's health status. This study is a preliminary investigation of non-invasive warning indicator, Red Zone (RZ), for heat strain derived from the combination of heart rate (HR) and skin temperature (Tskin). Individual data from five datasets (Age = 22 ± 5 yrs, BMI = 23.0 ± 2.6, N = 43) were utilized to establish the RZ system. Analysis of the termination points relative to Tskin and HR were identified, along with Body Mass Index for individual differences. The data classified by RZ system were compared to their individual Tcore, using Fisher's exact test. Overall, the data points classified by the RZ system showed relatively good agreement (60-80% probability) with Tcore (p < 0.05). The RZ is a reasonable approach to warn occupational workers that they are approaching or/under thermal strain.			
14. SUBJECT TERMS Heart rate, core temperature, skin temperature, BMI, noninvasive measures, heat stress.		15. NUMBER OF PAGES 4	
17. SECURITY CLASSIFICATION OF REPORT Unclassified		16. PRICE CODE	
18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT Unlimited	

# Environmental Ergonomics XI

Proceedings of the 11<sup>th</sup> International Conference  
22-26 May, 2005 Ystad, Sweden

Ingvar Holmér, Kalev Kuklane and Chuansi Gao  
Editors



Thermal Environment Laboratory, EAT  
Department of Design Sciences  
Lund University

## Noninvasive warning indicator of the “Red Zone” of potential thermal injury and performance impairment: A pilot study.

Miyo Yokota<sup>1\*</sup>, Daniel Moran<sup>2</sup>, Larry Berglund<sup>1</sup>, Lou Stephenson<sup>1</sup>, Margaret Kolka<sup>1</sup>.

<sup>1</sup>U.S. Army Research Institute of Environmental Medicine (USARIEM), Natick, MA, USA..

<sup>2</sup>Heller Institute of Medical Research, Sheba Medical Center, Tel Hashomer, Israel.

\* corresponding author: Miyo.Yokota@na.amedd.army.mil

### INTRODUCTION

Guidelines for assessing workers' level of heat strain to prevent thermal injury and performance impairment have been widely adopted by private industries to promote workers' health and safety. Internal body temperature ( $T_{\text{core}}$ ) is the common physiological parameter for heat strain assessment as it reliably indicates impending injury (1,2). Laboratory studies as well as occupational guidelines for civilian workforces (e.g., metal and glass workers, fire fighters) and military personnel, all suggest regulating workers' operational or environment conditions when  $T_{\text{core}}$  is  $38.0^{\circ}\text{C}$  ( $100.4^{\circ}\text{F}$ ) -  $38.5^{\circ}\text{C}$  ( $101.3^{\circ}\text{F}$ ) (1,2,3). However, measuring  $T_{\text{core}}$  may be invasive and impractical for real time monitoring of worker's health status or Soldiers engaged in long hours of various multiple and unpredictable tasks in hot environments.

Skin temperature ( $T_{\text{skin}}$ ) and heart rate (HR) are non-invasive, easily measured physiological parameters that are also indicators of the thermo-vascular status of workers exposed to thermal stress. Skin blood flows and sweating are active primary control mechanisms for  $T_{\text{core}}$  (4). If the  $T_{\text{core}}$  rises above its set point ( $\sim 37^{\circ}\text{C}$ ), blood flow to the skin increases. Normal blood flow to the skin for a comfortable sedentary person is about  $6 \text{ L}\cdot\text{h}^{-1}\cdot\text{m}^{-2}$  but can increase to  $90 \text{ L}\cdot\text{h}^{-1}\cdot\text{m}^{-2}$  or higher, when a person becomes overheated (4).  $T_{\text{skin}}$  rises as a result of the warm blood flow to the skin. Under the conditions of excessive body heat,  $T_{\text{skin}}$  approaches  $T_{\text{core}}$ , and thermoregulatory insufficiency may occur due to the minimal heat transfer.

HR increases as oxygen consumption increases during work (4). The worker's sustained HR should not exceed a value of  $180 - \text{age}$ , or the limit of maximal HR ( $\text{HR}_{\text{max}}$ ) at the workplace should not exceed the  $185 - 0.65*\text{age}$  (3). Despite the documentation that these physiological non-invasive measures increase with  $T_{\text{core}}$  under excessive heat exposure (4,5), the combination of HR and skin temperature are rarely utilized for occupational performance advisories or guidance.

This paper presents a preliminary investigation of the impending heat strain warning system called Red Zone (RZ), which is based on non-invasive measures of HR and  $T_{\text{skin}}$  combined with Body Mass Index (BMI). Individual differences characterized by BMI relate to differences in physiological responses to heat exposure (6,7) and improve the accuracy of the RZ warning system for individuals.

### MATERIALS AND METHODS

Individual data from four USARIEM datasets (Age =  $25 \pm 6$  yrs, BMI =  $23.9 \pm 3.1$ , N = 19) and a heat tolerance study (Age =  $19 \pm 1$  yrs, BMI =  $24.5 \pm 3.2$ , N = 24) were used to formulate the Red Zone. Table 1 depicts the summary of five study conditions. Thigh or chest skin temperatures were examined in these data because they were less variable than  $T_{\text{skin}}$  taken from arm or calf regions.

Because  $T_{\text{skin}}$  for USARIEM study4 were not monitored,  $T_{\text{skin}}$  of these subjects were estimated by SCENARIO, a USARIEM thermoregulatory mathematical model (8). In all studies, subjects discontinued a test when  $T_{\text{core}}$  reached  $39.5^{\circ}\text{C}$ . In addition, testing was stopped when subjects reached 95% of age-estimated  $\text{HR}_{\text{max}}$  or remained at 90-94% of age-estimated  $\text{HR}_{\text{max}}$  for 5 minutes.

**Table 1.** Summary of five study conditions used in this study.

Database (reference)	N	Lab vs Field	Weather (mean)	Main activity	Clothing	Duration
USARIEM study1 (9)	5	Lab	$38^{\circ}\text{C}$ , 30%RH	walk (2 mph, 0% grade)	CP	240 min
USARIEM study2 (10)	5	Lab	$30^{\circ}\text{C}$ , 30%RH	walk ( $1.46\text{ ms}^{-1}$ , 2% grade)	CP	80 min
USARIEM study3 (11)	7	Lab	$30^{\circ}\text{C}$ , 38%RH	walk (3 mph, 2% grade)	CP	45-90 min
USARIEM study4*	2	Field	$34^{\circ}\text{C}$ , 44%RH	walk (altitude up to 350m)	BDU	all day
Heat tolerance study (5)	24	Lab	$40^{\circ}\text{C}$ , 40%RH	walk ( $5\text{ km}\cdot\text{h}^{-1}$ , 2% grade)	Shorts, T-shirt	120 min


Database: \* indicates an unpublished study

Clothing: CP = Chemical Protective Garment; BDU = Battle Dress Uniform

The final measurements of  $T_{\text{skin}}$  and HR for all tests and BMI were compared. BMI is relatively convenient to measure and is beneficial to describe associations with body fat, age, and gender. BMI, calculated by  $(\text{weight, kg})/(\text{height}^2, \text{m}^2)$ , was the only consistently measured individual parameter available between the datasets. When a BMI was unavailable (Heat tolerance study), it was randomly generated for the individual subject from the mean and standard deviation of published BMI data. Subjects were categorized into three BMI groups: a) low  $\leq 20$ ; b)  $20 < \text{medium} < 25$ ; c) high  $\geq 25$ . The RZ was also classified into three conditions: a) good/safe; b) experiencing heat/warning; and c) critical/overheated. The individual subject data labeled with these Red Zone categories based on their physiological responses were compared to their individual  $T_{\text{core}}$ , using Fisher's exact test.

## RESULTS

The termination points for combinations of  $T_{\text{skin}}$  and HR from the 5 different studies are summarized by BMI, shown in Figure 1.  $T_{\text{skin}}$  above  $38^{\circ}\text{C}$  is likely to indicate potential heat injury irrespective of HR (upper area in Figure 1). When  $T_{\text{skin}}$  and HR simultaneously occur above respective thresholds of  $36^{\circ}\text{C}$  and the subject's HR warning threshold, there is an increasing probability of impending hyperthermia, heat injury and performance decrements. When  $T_{\text{skin}}$  ranges between  $36^{\circ}\text{C}$  and  $38^{\circ}\text{C}$ , the HR warning threshold is BMI dependent and increases as BMI is lower (three dotted BMI boxes in Figure 1). When an individual's BMI is high ( $\geq 25$ ), he/she is likely to experience hyperthermia and heat stress at a lower HR than medium/light BMI individuals. Low BMI individuals ( $\leq 20$ ) are less likely to experience potential heat injury until HR limits reach to around 160-175 bpm (Figure 1).



— . . . indicates critical/warning conditions  
 ..... indicates warning conditions for addressed BMI.  
 BMI: L = low; M = medium; H = high

**Figure 1.** Red Zone model based on heart rate (HR, bpm), skin temperature ( $T_{\text{skin}}$ , °C) and three BMI categories (low, medium, high) applied for 5 studies.

Table 2 demonstrates the relationships of termination points between  $T_{\text{core}}$  classified by a standard heat guidelines ( $T_{\text{core}} = 38.5^\circ\text{C}$  as a threshold) (1,2) and  $T_{\text{skin}}$  categorized by the RZ model. The accuracy in the RZ warning that  $T_{\text{core}}$  is greater than  $38.5^\circ\text{C}$  is around 60%. Similarly,  $T_{\text{core}}$  below  $38.5^\circ\text{C}$  was classified as no warning in RZ with ~60% accuracy. False positive rate ( $T_{\text{skin}}$  was classified as a safe condition, although  $T_{\text{core}}$  was recorded above  $38.5^\circ\text{C}$ ) was around 20%. The data points classified by the three RZ categories showed relatively good agreement with  $T_{\text{core}}$  (Fisher's exact,  $p < 0.05$ ).

**Table 2.** The assessment of relationship between categorical Red Zone classifications and core temperature, obtained from 5 different studies

Core Temperature	Red Zone			Total
	Safe	Warning	Critical	
Safe ( $< 38.5^\circ\text{C}$ )	13 (59.1%)	5 (22.7%)	4 (18.2%)	22 (100.0%)
Critical ( $> 38.5^\circ\text{C}$ )	5 (23.8%)	4 (19.1%)	12 (57.1%)	21 (100.0%)
Total	18 (41.9%)	9 (20.9%)	16 (37.2%)	43 (100.0%)

## CONCLUSIONS

The RZ is an encouraging approach to warn occupational workers that they are approaching or/under excessive thermal physiological strain. This preliminary study suggests that there was a 60% – 80% probability that  $T_{\text{core}}$  is above  $38.5^\circ\text{C}$  when  $T_{\text{skin}}$  and HR of individuals fell into RZ classifications of “experiencing heat or critical conditions.” Some of the RZ false positive results

might be related to heat acclimation state because heat tolerant individuals may be able to maintain lower  $T_{skin}$  in compensable environments than heat intolerant individuals despite their high  $T_{core}$ . The RZ function, derived from non-invasive  $T_{skin}$  and HR measurements, utilizing BMI is useful to warn of likely excessive  $T_{core}$  and thermoregulatory distress as it occurs. Further field measurements including various environmental, clothing, and operational conditions will be important to refine this warning method, increase confidence intervals, and improve applications to diversified military populations.

## ACKNOWLEDGMENTS

Authors thank Mr. Bruce Cadarette, CPT Samuel Chevront, and Ms. Leslie Levine, USARIEM for assistance with this project. The investigators have adhered to the policies for protection of human subjects as prescribed in Army Regulation 70-25, and the research was conducted in adherence with the provisions of 45 CFR Part 46. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Army or the Department of Defense.

## REFERENCES

1. Amos, D., Hansen, R., Lau, Wei-Man., and Michalski, JT. Physiological and cognitive performance of Soldiers conducting routine patrol and reconnaissance operations in the tropics. *Mil. Med.* 2000; 165: 961-966.
2. Occupational Safety & Health Administration (OSHA). Heat Stress. OSHA Technical Manual (OTM). TED 1-0.15A. Section III, Chapter 4 [http://www.osha-slc.gov/dts/osta/otm/otsm\\_iii/otm\\_iii\\_4.html](http://www.osha-slc.gov/dts/osta/otm/otsm_iii/otm_iii_4.html). Accessed in July, 2003.
3. International Standard Organization (ISO) 9886. Ergonomics – Evaluation of thermal strain by physiological measurements. ISO. Geneva, Switzerland; 2004.
4. Åstrand, P, and Rodahl, K. Textbook of work physiology: Physiological bases of exercises. New York: McGraw-Hill Book Company. 3<sup>rd</sup> edition; 1986.
5. Moran, DS., Heled, Y., Still, L., Laor, A., and Shapiro, Y. Assessment of heat tolerance for post exertional heat stroke individuals. 2004; *Med. Sci. Monit.* 10:CR252-257.
6. Donoghue, AM., and Bates, GP. The risk of heat exhaustion at a deep underground metalliferous mine in relation to body-mass index and predicted  $Vo_2max$ . 2000; *Occup. Med.* 50: 259-263.
7. Wallace, R. Risk factors and mortality in relation to health illness severity. USARIEM Technical Report. T03-14. Natick, MA; 2003.
8. Kraning, KK, and Gonzalez, RR. SCENARIO.: A military/industrial heat strain model modified to account for effects of aerobic fitness and progressive dehydration. USARIEM Technical Report. T97-1. Natick, MA; 1997.
9. Levine, L, Cadarette, BS., and Kolka, MA. Endurance time in the self-contained toxic environment protective outfit (STEPO) with personal ice-cooled microclimate cooling system (PICS) in three environments. USARIEM Technical Report. T03-13. Natick, MA; 2003.
10. Chevront, SN., Kolka, MA., Cadarette, BS., Montain, SJ., and Sawka, MN. Efficacy of intermittent, regional, microclimate cooling. 2003; *J. Appl. Physiol.* 94:1841-1848.
11. Stephenson, LA., Kolka, MA. Esophageal temperature threshold for sweating decreases before ovulation in premenopausal women. 1999; *J. Appl. Physiol.* 86:22-28.