

**AFRL-ML-TY-TP-2006-4580**

**POSTPRINT**



# **SILICA-IMMOBILIZED ENZYMES FOR MULTI-STEP SYNTHESIS IN MICROFLUIDIC DEVICES**

**Heather R. Luckarift  
Air Force Research Laboratory**

**Bosung S. Ku, Jonathan S. Dordick  
Rensselaer Polytechnic Institute  
Troy, NY 12180**

**Jim C. Spain  
Georgia Institute of Technology  
Atlanta, GA 30332**

**OCTOBER 2007**

**DISTRIBUTION STATEMENT A: Approved for public release;  
distribution unlimited.**

**Published in Biotechnology and Bioengineering, Vol 98, No 3, Oct 15, 2007. Pub.  
Wiley Periodicals, Inc.**

**Air Force Research Laboratory  
Materials and Manufacturing Directorate  
Airbase Technologies Division  
139 Barnes Drive, Suite 2  
Tyndall AFB, FL 32403-5323**

**REPORT DOCUMENTATION PAGE**

*Form Approved  
OMB No. 0704-0188*

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

**PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

<b>1. REPORT DATE (DD-MM-YYYY)</b>	<b>2. REPORT TYPE</b>	<b>3. DATES COVERED (From - To)</b>
------------------------------------	-----------------------	-------------------------------------

<b>4. TITLE AND SUBTITLE</b>	<b>5a. CONTRACT NUMBER</b>
	<b>5b. GRANT NUMBER</b>
	<b>5c. PROGRAM ELEMENT NUMBER</b>

<b>6. AUTHOR(S)</b>	<b>5d. PROJECT NUMBER</b>
	<b>5e. TASK NUMBER</b>
	<b>5f. WORK UNIT NUMBER</b>

<b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b>	<b>8. PERFORMING ORGANIZATION REPORT NUMBER</b>
---	---

<b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b>	<b>10. SPONSOR/MONITOR'S ACRONYM(S)</b>
	<b>11. SPONSOR/MONITOR'S REPORT NUMBER(S)</b>

**12. DISTRIBUTION/AVAILABILITY STATEMENT**

**13. SUPPLEMENTARY NOTES**

**14. ABSTRACT**

**15. SUBJECT TERMS**

<b>16. SECURITY CLASSIFICATION OF:</b>			<b>17. LIMITATION OF ABSTRACT</b>	<b>18. NUMBER OF PAGES</b>	<b>19a. NAME OF RESPONSIBLE PERSON</b>
a. REPORT	b. ABSTRACT	c. THIS PAGE			<b>19b. TELEPHONE NUMBER (Include area code)</b>

# Silica-Immobilized Enzymes for Multi-Step Synthesis in Microfluidic Devices

Heather R. Luckarift,<sup>1</sup> Bosung S. Ku,<sup>2</sup> Jonathan S. Dordick,<sup>2</sup> Jim C. Spain<sup>3</sup>

<sup>1</sup>Air Force Research Laboratory, 139 Barnes Drive, Tyndall AFB, Florida 32403

<sup>2</sup>Department of Chemical and Biological Engineering, Rensselaer Polytechnic Institute, Troy, New York 12180; telephone: 518-276-2899; fax: 518-276-2207; e-mail: dordick@rpi.edu

<sup>3</sup>School of Civil and Environmental Engineering, 311 Ferst Drive, Georgia Institute of Technology, Atlanta, Georgia 30332-051; telephone: 404-894-0628; fax: 404-894-2278; e-mail: jspain@ce.gatech.edu

Received 5 January 2007; accepted 20 March 2007

Published online 5 April 2007 in Wiley InterScience (www.interscience.wiley.com). DOI 10.1002/bit.21447

**ABSTRACT:** The combinatorial synthesis of 2-aminophenoxazin-3-one (APO) in a microfluidic device is reported. Individual microfluidic chips containing metallic zinc, silica-immobilized hydroxylaminobenzene mutase and silica-immobilized soybean peroxidase are connected in series to create a chemo-enzymatic system for synthesis. Zinc catalyzes the initial reduction of nitrobenzene to hydroxylaminobenzene which undergoes a biocatalytic conversion to 2-aminophenol, followed by enzymatic polymerization to APO. Silica-immobilization of enzymes allows the rapid stabilization and integration of the biocatalyst within a microfluidic device with minimal preparation. The system proved suitable for synthesis of a complex natural product (APO) from a simple substrate (nitrobenzene) under continuous flow conditions.

Biotechnol. Bioeng. 2007;98: 701–705.

© 2007 Wiley Periodicals, Inc.

**KEYWORDS:** microfluidics; immobilized enzyme; aminophenoxazinone; chips; sequential catalysis

## Introduction


The use of microfluidics and 'lab-on-a-chip' devices for efficiently miniaturizing biochemical assays and chemical synthesis is now well recognized (Barry and Ivanov, 2004; Figeys and Pinto, 2000; Haswell et al., 2001; Mitchell, 2001; Polson and Hayes, 2001; Urban et al., 2006). The dramatically increased surface to volume ratios lead to rapid mass transfer and greatly reduced analysis time and sample consumption. There is increasing interest in the integration of enzymes into microfluidic systems for applications in medical diagnostics, biosensing and natural product and

organic synthesis (Hadd et al., 1997; Jones et al., 2004; Krenkova and Foret, 2004; Ku et al., 2006; Lee et al., 2003; Srinivasan et al., 2003, 2004). Such biocatalytic systems are typically enabled by covalent attachment of the biocatalysts to channel walls, physical absorption onto solid matrices, or copolymerization (Holden et al., 2004, 2005; Honda et al., 2005; Mao et al., 2002; Sakia-Kato et al., 2004). Biocatalysts encapsulated in sol-gels and hydrogels can be subject to leaching, which reduces the active enzyme concentrations (Sakai-Kato et al., 2003). We recently reported a method of enzyme immobilization in biomimetic silica that increases the mechanical stability of the immobilized enzyme and facilitates application to flow-through reaction systems (Berne et al., 2006; Luckarift et al., 2004, 2006). The silica immobilization method is simple, rapid, and applicable to a wide variety of enzymes. The strategy precludes the need to modify the surface to allow enzyme attachment, greatly reducing preparation time and enhancing the loading capacity of the reaction system.

In this study we investigated the applicability of silica-immobilized enzymes for entraining biomolecules in microfluidic devices, and in the process developed a functional chemoenzymatic microfluidic platform. Specifically, with nitrobenzene (NB) (1) as a model substrate we demonstrate a three-step continuous reaction system for the production of 2-aminophenoxazin-3-one (APO) (4) (Fig. 1). In the first step, metallic zinc reduces NB (1) to hydroxylaminobenzene (HAB) (2) (Furniss et al., 1989), which then undergoes an enzymatic intramolecular rearrangement catalyzed by HAB-mutase to form 2-aminophenol (2-AP) (3) (Davis et al., 2000; Luckarift et al., 2005; Nadeau et al., 2003). In the final stage, soybean peroxidase

Correspondence to: J.S. Dordick and J. C. Spain  
Contract grant sponsor: Air Force Office of Scientific Research  
Contract grant sponsor: NIH  
Contract grant number: GM66712

 WILEY  
InterScience®  
DISCOVER SOMETHING GREAT



**Figure 1.** Schematic of a multi-step microfluidic process for the conversion of nitrobenzene to 2-aminophenoxazin-3-one.

(SBP) catalyzes the oxidation of 2-AP to form APO (4) (Fomsgaard et al., 2004; Hishida et al., 1974; Horvath et al., 2004; Lee et al., 2003; Reihmann and Ritter, 2002; Simandi et al., 2004; Srinivasan et al., 2003; Toussaint and Lerch, 1987) APO is an intermediate in the synthesis of actinomycins; an important group of antibiotics with anti-fungal and anti-tumor properties (Barry et al., 1989; Shimizu et al., 2004; Veselkov et al., 2003).

## Materials and Methods

### Materials

Hydroxylaminobenzene was synthesized as described previously (Furniss et al., 1989). Partially purified HAB mutase A was prepared and immobilized as described previously (Luckarift et al., 2005). All other chemicals were of analytical grade and obtained from Sigma–Aldrich (St. Louis, MO).

### Fabrication of Microfluidic Chips

Standard photolithographic and molding techniques were used to fabricate a microfluidic channel (40 mm long  $\times$  1.5 mm wide  $\times$  0.1 mm deep; volume of 6  $\mu$ L) in polydimethylsiloxane as described previously (Ku et al., 2006; Srinivasan et al., 2004). The resulting chip is fabricated with three 50  $\mu$ m channels to form an exit network to retain the immobilized catalysts. Silica-immobilized biocatalysts were loaded into the channel by applying a vacuum to the product reservoir. HAB-mutase enzyme activity was determined as described previously (Davis et al., 2000). SBP enzyme activity was determined with ABTS (2, 2'-Azino-bis(3-ethylbenzthiazoline-6-sulfonic acid)) and hydrogen peroxide as described previously (Amisha Kamal and Behere, 2003).

### Chip-Based Synthesis

All substrates were dissolved in water containing NH<sub>4</sub>Cl (40 mM). Substrates were pumped at fixed flow rates and the eluate collected in an equal volume of acetonitrile for HPLC analysis. Reactants and products of all conversions

were monitored by reverse-phase HPLC on a Supelco ABZ column with an acetonitrile/water gradient as described previously (Luckarift et al., 2005).


### Synthesis of 2-Aminophenoxazin-3-One Product Standard

2-Aminophenol (100 mg in 1L of potassium phosphate buffer, pH 7.4) was incubated with 0.01 U/mL SBP enzyme and 0.6 mM H<sub>2</sub>O<sub>2</sub>. A deep red product characteristic of APO formed rapidly and precipitated from solution. The product was purified by solid phase extraction and was characterized and identified by NMR as 2-aminophenoxazin-3-one by comparison to proposed chemical shifts and previous literature reports (Gabriele et al., 2003). The yield of the reaction was approximately 83% and the product was used as a standard for future reactions. NMR spectra were recorded on a Varian Inova spectrometer equipped with a 5 mm indirect detection probe, operating at 500 MHz for <sup>1</sup>H and at 125 MHz for <sup>13</sup>C.

## Results and Discussion

### Biosynthesis of APO in a Microfluidic Chip


Silica-immobilized SBP was prepared and demonstrated good retention of enzyme activity (65–85%) relative to the soluble enzyme. The silica-immobilized SBP was packed into a microfluidic channel with an equal volume of agarose beads (1:1 vol/vol), to give a final enzyme loading of approximately 0.5 U. The presence of agarose beads prevents the silica particles from packing and reduces the void volume of the microfluidic reactor (effective volume  $\sim$ 3  $\mu$ L). 2-AP (0.5 mM) was pumped through the SBP chip at a range of flow rates and the conversion efficiency of the enzyme was determined by measuring production of APO. Hydrogen peroxide (0.5–2.5 mM in 10% DMF) was added continuously as a second reactant for the oxidation reaction. A molar ratio of 2.5:1 H<sub>2</sub>O<sub>2</sub>:2-AP was optimal for conversion of 2-AP to APO (Fig. 2). The initial reaction velocity of immobilized SBP was 83 and 98  $\mu$ M/min at 1.0 and 2.5 mM H<sub>2</sub>O<sub>2</sub>, respectively.



**Figure 2.** Effect of H<sub>2</sub>O<sub>2</sub> concentration on conversion of 2-AP to APO in a silica-immobilized SBP chip. Conversion efficiency of 2-AP catalyzed by SBP with 2.5 mM (○), 1 mM (◆), 0.5 mM (●) and 0.1 mM (■) H<sub>2</sub>O<sub>2</sub>.

### Biosynthesis of 2-Aminophenol in a Microfluidic Chip

Immobilized HAB-mutase demonstrated good retention of enzyme activity relative to soluble enzyme, with little variability among replicate preparations (45–65%). Silica-immobilized HAB-mutase was mixed with an equal volume of agarose beads and packed into the channel as described above to give a final enzyme loading of approximately 1 unit (U). HAB (1 mM) was pumped through the HAB-mutase chip at a range of flow rates and the conversion efficiency determined. Conversion efficiency decreased with increasing flow rate as expected due to the decreased residence time (Fig. 3). The maximum 2-AP product concentration obtained from 1 mM HAB was 0.61 mM (±0.09) at a flow rate of 50 μL/h. The reusability of the silica-immobilized HAB-mutase within the microfluidic chip was also determined to verify the retention of the biocatalyst within the channel during continuous flow. The



**Figure 3.** Substrate conversion efficiency as a function of flow rate. Conversion efficiency of 1 mM NB by Zn (■), 1 mM HAB by HAB-mutase (○) and 1 mM 2-AP by SBP and 1 mM H<sub>2</sub>O<sub>2</sub> (▼).


enzyme chips were washed and stored repeatedly at 4°C and retained >50% of the original enzyme activity over a number of subsequent cycles (Fig. 4).

### Biosynthesis of APO From HAB

The two immobilized-enzyme chips (HAB-mutase and SBP) were connected in series and an aqueous solution of HAB (1 mM) was introduced at a flow rate of 100 μL/h. This resulted in the production of APO (0.13 mM ±0.025) continuously for over 4 h (conversion efficiency of ~52%) (Table I). The addition of H<sub>2</sub>O<sub>2</sub> to the system diluted the substrate concentration and reduced the theoretical product concentration by 50%. The flow rate could be increased to 150 μL/h, however, with no loss in conversion efficiency.

### Chemoenzymatic Synthesis of APO

To complete the chemoenzymatic microfluidic platform, a metallic zinc chip was prepared (1 mg zinc dust, mixed 1:1 with agarose beads) and attached prior to the two enzyme reactors to demonstrate a combinatorial chemoenzymatic conversion of nitrobenzene to the resultant phenoxazinone. The zinc chip gave approximately 25% conversion of NB to HAB at low flow rates. In contrast to the chips containing biocatalysts, the metal zinc chip showed increased conversion efficiency with increasing flow rate (Fig. 3). At low flow rates, aniline was formed as a byproduct of the zinc reduction in preference to HAB. As a consequence, when the zinc chip was added in series with the two enzyme chips and NB pumped at a flow rate of 150 μL/h, the overall conversion efficiency was reduced to 19% due to the reduction in the corresponding efficiency of the zinc conversion step (Table I). Reduced conversion efficiency was also attributed to absorption and volatility



**Figure 4.** Reusability of HAB-mutase chips over several wash cycles. HAB (100 mL, 1 mM) was pumped through the HAB-mutase chips at a flow rate of 150 μL/h and the formation of 2-aminophenol was determined. The chips were washed with 20 volumes of buffer between each subsequent cycle. Data shown is the mean and standard deviation of three independent HAB-mutase chips. Chips were stored overnight at 4°C at the point indicated (\*).

**Table I.** Conversion efficiency of multi-step microfluidic synthesis.

Substrate	Chip combination	Conversion efficiency (%)	
		100 $\mu$ L/h	150 $\mu$ L/h
1 mM NB	Zinc/HAB mutase	16.7 $\pm$ 3.5	28.2 $\pm$ 2.7
1 mM HAB	HAB mutase/SBP	53.2 $\pm$ 8.8	51.7 $\pm$ 10.2
1 mM NB	Zinc/HAB mutase/SBP	ND	18.9 $\pm$ 10.14

ND, not determined.

of the substrate. In an empty chip, observed NB concentrations were reduced in the eluate relative to the starting concentration but the percentage overall loss decreased with increasing flow rate. The conversion of the chemoenzymatic system was, therefore, determined at flow rates considered optimal for the biocatalytic steps. Clearly the step involving conversion of the nitro compound to the corresponding hydroxylamino compound will require optimization for future applications. Despite the low overall yields of the chemoenzymatic steps in series, the system provides a rapid and versatile method for screening the conversion of nitroarenes.

In this work, we demonstrated the first chemoenzymatic microfluidic reactor for the synthesis of a natural product from a readily available and simple substrate, nitrobenzene. The facile silica-immobilization method provides stable heterogeneous catalysts, which can be easily incorporated into microfluidic chips. The flow-through system described could be applied to the transformation of a wide variety of nitroarene substrates into their corresponding phenoxazinone products, and provides an attractive alternative to conventional chemical synthesis. The versatility of the immobilization method makes the system amenable to a wide range of microfluidic devices incorporating biomacromolecules and coupled to chemical synthesis. Single unit reactors containing enzyme or metal catalysts can be integrated to create a variety of synthetic pathways for rapid synthesis and screening.

HRL was supported by Oak Ridge Institute for Science and Education and funding by Air Force Office of Scientific Research. JSD acknowledges funding from the National Science Foundation and NIH (GM66712). The authors acknowledge Lloyd Nadeau for helpful discussions and synthesis of HAB. NMR spectroscopy was performed by Ion Ghiviriga, University of Florida.

## References

- Amisha Kamal JK, Behere DV. 2003. Activity, stability and conformational flexibility of seed coat soybean peroxidase. *J Inorg Biochem* 94:236–242.
- Barry R, Ivanov DJ. 2004. Microfluidics in biotechnology. *Nanobiotechnology* 2(2):1–5.
- Barry CE, Nayar PG, Begley TP. 1989. Phenoxazinone Synthase: Mechanism for the formation of the phenoxazinone chromophore of actinomycin. *Biochemistry* 28:6323–6333.
- Berne C, Betancor L, Luckarift HR, Spain JC. 2006. Application of a microfluidic reactor for screening cancer prodrug activation using

- silica-immobilized nitrobenzene nitroreductase. *Biomacromolecules* 7:2631–2636.
- Davis JK, Paoli GC, He Z, Nadeau LJ, Somerville CC, Spain JC. 2000. Sequence analysis and initial characterization of two isozymes of hydroxylaminobenzene mutase from *Pseudomonas pseudoalcaligenes* JS45. *Appl Environ Micro* 66:2965–2971.
- Figey D, Pinto D. 2000. Lab-on-a-chip: A revolution in biological and medical sciences. *Anal Chem* 72(9):330A–335A.
- Fomsgaard IS, Mortensen AG, Carlsen SCK. 2004. Microbial transformation products of benzoxazoline and benzoxazinone allelochemicals—A review. *Chemosphere* 54:1025–1038.
- Furniss BS, Hannaford AJ, Smith PWG, Tatchell AR. 1989. *Vogel's Textbook of practical organic chemistry*, Longman Scientific and Technical. New York: John Wiley & Sons.
- Gabriele B, Mancuso R, Salerno G, Costa MJ. 2003. An improved procedure for the palladium-catalyzed oxidative carbonylation of  $\beta$ -amino acids to oxazolidin-2-ones. *Org Chem* 68:601–604.
- Hadd AG, Raymond DE, Halliwell JW, Jacobson SC, Ramsey JM. 1997. Microchip device for performing enzyme assays. *Anal Chem* 69:3407–3412.
- Haswell SJ, Middleton RJ, O'Sullivan B, Skelton V, Watts P, Styring P. 2001. The application of micro reactors to synthetic chemistry. *Chem Comm* 391–398.
- Hishida T, Nogami T, Shirota Y, Mikawa H. 1974. Reaction of o-Benzoquinone mono imine with o-aminophenol. *Chem Lett* 3(3):293–296.
- Holden MA, Jung S-Y, Cremer PS. 2004. Patterning enzymes inside microfluidic channels via photo-attachment chemistry. *Anal Chem* 76:1838–1843.
- Honda T, Miyazaki M, Nakamura H, Maeda H. 2005. Immobilization of enzymes on a microchannel surface through cross-linking polymerization. *Chem Comm* 5062–5064.
- Horvath T, Kaizer J, Speier G. 2004. Functional phenoxazinone synthases models. Kinetic studies on the copper-catalyzed oxygenation of 2-aminophenol. *J Mol Catalysis A Chem* 215:9–15.
- Jones F, Forrest S, Palmer J, Lu Z, Elmore J, Elmore BB. 2004. Immobilized enzyme studies in a microscale bioreactor. *Appl Biochem Biotech* 113–116:261–272.
- Krenkova J, Foret F. 2004. Immobilized microfluidic enzymatic reactors. *Electrophoresis* 25:3550–3563.
- Ku B, Cha J, Srinivasan A, Kwon SJ, Jeong J-C, Sherman DH, Dordick JS. 2006. Chip-based polyketide biosynthesis and functionalization. *Biotechnol Prog* 22:1102–1107.
- Lee M-Y, Srinivasan A, Ku B, Dordick JS. 2003. Multienzyme catalysis in microfluidic biochips. *Biotech Bioeng* 83:20–28.
- Luckarift HR, Naik RR, Stone MO, Spain JC. 2004. Enzyme immobilization in a biomimetic silica support. *Nat Biotech* 22:211–213.
- Luckarift HR, Nadeau LJ, Spain JC. 2005. Continuous synthesis of aminophenols from nitroaromatic compounds by combination of metal and biocatalyst. *Chem Comm* 383–384.
- Luckarift HR, Johnson GR, Spain JC. 2006. Silica-immobilized enzyme reactors: Application to cholinesterase-inhibition studies. *J Chrom B* 843:310–316.
- Mao H, Yang T, Cremer PS. 2002. Design and characterization of immobilized enzyme in microfluidic systems. *Anal Chem* 74:379–385.
- Mitchell P. 2001. Microfluidics-downsizing large-scale biology. *Nat Biotech* 19:717–721.
- Nadeau LJ, He Z, Spain JC. 2003. Bacterial conversion of hydroxylamino aromatic compounds by both lyase and mutase enzymes involves intramolecular transfer of hydroxyl groups. *Appl Environ Micro* 69(5):2786–2793.
- Polson NA, Hayes MA. 2001. Microfluidics controlling fluids in small places. *Anal Chem* 73(11):312A–319A.
- Reihmann MH, Ritter H. 2002. Regioselective HRP-catalyzed polymerization of 4-amino-phenol. *J Macromol Sci Pure Appl Chem* A39(12):1369–1382.
- Sakai-Kato K, Kato M, Toyooka T. 2003. Creation of an on-chip enzyme reactor by encapsulating trypsin in sol-gel on a plastic microchip. *Anal Chem* 75:388–393.

- Sakia-Kato K, Kato M, Ishihara K, Toyóoka T. 2004. An enzyme-immobilization method for integration of biofunctions on a microchip using a water-soluble amphiphilic phospholipid polymer having a reacting group. *Lab Chip* 4:4–6.
- Shimizu S, Suzuki M, Tomoda A, Arai S, Taguchi H, Hanawa T, Kamiya S. 2004. Phenoxazine compounds produced by the reactions with bovine hemoglobin show antimicrobial activity against non-tuberculosis mycobacteria. *Tohoku J Exp Med* 203(1): 47–52.
- Simandi TM, Simandi LI, Gyor M, Rockenbauer A, Gomory A. 2004. Kinetics and mechanism of the ferroxime (II)-catalysed biomimetic oxidation of 2-aminophenol by dioxygen. A functional phenoxazinone synthase model. *Dalton Trans* 7(7):1056–1060.
- Srinivasan A, Wu X, Lee M-Y, Dordick JS. 2003. Microfluidic peroxidase biochip for polyphenol synthesis. *Biotech Bioeng* 81:563–569.
- Srinivasan A, Bach H, Sherman DH, Dordick JS. 2004. Bacterial P450-catalyzed polyketide hydroxylation on a microfluidic platform. *Biotech Bioeng* 88:528–535.
- Toussaint O, Lerch K. 1987. Catalytic oxidation of 2-aminophenols and ortho hydroxylation of aromatic amines by tyrosinase. *Biochem* 26: 8567–8571.
- Urban PL, Goodall DM, Bruce NC. 2006. Enzymatic microreactors in chemical analysis and kinetic studies. *Biotech Adv* 24(1):42–57.
- Veselkov AN, Maleev VY, Glibin EN, Karawajew L, Davies DB. 2003. Structure-activity relation for synthetic phenoxazine drugs. *Eur J Biochem* 270:4200–4207.