

AD_____

Award Number: W81XWH-06-1-0092

TITLE: Synthesis of Taxol-Like Prostate Cancer Chemotherapeutic Agents

PRINCIPAL INVESTIGATOR: Hyunil Jo

CONTRACTING ORGANIZATION: University of Pennsylvania
Philadelphia, PA 19104

REPORT DATE: November 2007

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 01-11-2007		2. REPORT TYPE Annual Summary		3. DATES COVERED (From - To) 30 OCT 2006 - 29 OCT 2007	
4. TITLE AND SUBTITLE Synthesis of Taxol-Like Prostate Cancer Chemotherapeutic Agents				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-06-1-0092	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Hyunil Jo E-Mail: hyunil@sas.upenn.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Pennsylvania Philadelphia, PA 19104				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT During the study of C-O cleavage reaction toward the synthesis of Eleutherobin analogues, we have discovered a novel functionalization of α,β -epoxy ketone. This reaction seems to be very useful in preparation of vinyl functionalized allylic alcohol. Modification of hydrazine reactant is now being under investigation. On the other hand, coupling of two key fragments for the intramolecular Diels-Alder reaction failed under conventional reaction condition and the suitable reaction condition is now being studied.					
15. SUBJECT TERMS Eleutherobin, tandem Diels-Alder					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	7	19b. TELEPHONE NUMBER (include area code)

Table of Contents


Cover.....	1
SF 298.....	2
Table of Contents.....	3
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	6
Reportable Outcomes.....	6
Conclusions.....	6
References.....	7
Appendices.....	7

1. Introduction

The development of an efficient synthetic route to Eleutherobin and its analogues has been recognized as a daunting task for synthetic communities. Its scarce availability in nature and potent toxicity against cancer has attracted our attention and led us to study the synthesis of Eleutherobin utilizing a tandem Diels-Alder reaction and Grob-fragmentation strategy. During the second year of the grant, we studied the problematic C-O bond cleavage reaction using hydrazine and discovered a novel chemical transformation, which could be a powerful source of highly functionalized allylic alcohols.

2. Body


Previously, we have successfully prepared the advanced intermediate **1**, which could become a substrate for the Grob fragmentation after C-O bond cleavage. Unfortunately, cleavage of the C-O bond failed as it was extremely resistant to the SmI_2 reduction condition. Treatment of **1** to Zn-AcOH condition also failed to produce the desired Grob substrate **2**. This issue may be related with the orbital alignment of the ketyl radical anion as depicted in **Scheme 1** or an initial electron transfer process.¹⁾ Thus, successful implementation of SmI_2 reduction to triol **4** was only achieved when diol **3** was subjected to the same reaction condition.²⁾ However, further attempts to transform triol **4** to the Grob fragmentation substrate **5** were unsuccessful due to facile retro-aldol reaction. We then turned our attention to other C-O bond cleavage reactions. We believed that a Wharton rearrangement could provide a possible solution.



Scheme 1. Failure to preparation of Grob fragmentation substrate **2**


As shown in **Scheme 2**, the Wharton rearrangement can transform α , β -epoxy ketone **6** to allylic alcohol

9.




Scheme 2. Wharton Rearrangement

In our study, simple application of the Wharton rearrangement would only lead to the formation of diene alcohol that is not suitable for our purpose to retain the original carbonyl functionality. However, according to the reaction mechanism proposed by G. Stork *et al*, the intermediate of the Wharton rearrangement could be vinyl radical species from the release of nitrogen gas in vinyl diazene **8**.⁴⁾ We envisioned that this intermediate could be functionalized at the vinylic position if a suitable radical trapping agent is added. The resulting product would be a vinyl functionalized allylic alcohol which could be converted into carbonyl group again. Thus, we set to test our hypothesis with epoxy ketone **10** which can be readily prepared from verbenone.



Scheme 3. Wharton functionalization of epoxy ketone **10**


After much experimentation on the reaction condition, our hypothesis proved to be true. Vinyl sulfide **11**, vinyl halide **13** and vinyl tin species **14** could be prepared in a single pot operation in a good to moderate yield. More importantly, vinyl halide **13** and vinyl tin species **14** could be utilized in Palladium mediated C-C bond forming reactions, which further reinforces the importance and versatility of this transformation. Furthermore, formation of the 1,4 diene **15** indicates the direct C-C bond formation could be possible. Thus, this novel Wharton functionalization is extremely versatile as it provides a direct method to prepare vinyl functionalized allylic alcohols in a facile manner. However, one limitation is the formation of the normal Wharton reaction product **12**. This issue led us to investigate the source of the hydrogen atom. We ran the reaction in the CD₃OD without the trapping agent as shown in **Scheme 4**. Surprisingly, deuteriated allylic alcohol was not the predominant product which suggests hydrazine itself could be the main source of the hydrogen atom.



Scheme 4. Quenching study of Wharton reaction

Thus, it is conceivable that use of monosubstituted hydrazine would prevent the formation of the normal Wharton product. Currently, we are investigating the use of a monosilylated hydrazine.⁵⁾ If successfully implemented, this method would become very influential in the synthetic community due to its versatility and simplicity.

On the other hand, we also prepared the two key fragments for the intramolecular Diels-Alder approach as proposed in the previous annual report.



Scheme 5. Attempted preparation of Diels-Alder reaction substrate **19** or **22**

As shown in **Scheme 5**, conventional condition utilizing *n*-BuLi as a base was not successful in the preparation of the desired Diels-Alder substrate **19** as only starting materials were recovered. This suggested that the generation of alkynyl anion failed. Likewise, addition of alkyne **20** into aldehyde **21** was also unsuccessful. The only product obtained was *n*-BuLi addition to aldehyde **21**. Thus, deprotonation conditions of terminal alkyne **17** or **20** are now being studied.

3. Key Research Accomplishment

- Discovery of a Wharton functionalization
- Determination of the role of hydrazine as the main hydrogen atom source in the Wharton reaction

4. Reportable outcomes

Poster presentation

“Synthetic studies on the Taxol-like cancer chemotherapeutic agent”

Innovative Minds in Prostate Cancer Today (IMPACT) meeting, Atlanta, GA, Sep 2007

Manuscript on Wharton functionalization in preparation

5. Conclusion

During the second year of this grant, we have discovered a novel and powerful chemical transformation in studying the C-O bond cleavage reaction. This reaction would lead to a number of synthetically useful

intermediates and give further insight on the traditional Wharton reaction. Attempted assembly of the two key fragments for the intramolecular Diels-Alder reaction revealed that more extensive studies would be needed for the deprotonation reaction condition.

“So What?”

As a result of the work done over the past year, a highly useful novel transformation was discovered in the course of study on the C-O cleavage reaction. This Wharton functionalization could be very powerful tool in the synthetic community if combined with organometal catalyzed C-C bond forming reactions.

6. References

1. J. Cossy, J-L. Ranaivosata, V. Bellosta, J. Ancerewicz, R. Ferritto, and P. Vogel, “Reductive Oxa Ring Opening of 7-Oxabicyclo[2,2,1]heptan-2-ones. Synthesis of C- α -Galactosides of Carbapentopyranoses” *J. Org. Chem.* **1995**, 60, 8351-8359.
2. J. D. Winkler, K. J. Quinn, C. H. MacKinnon, S. D. Hiscock, and E. C. McLaughlin, "Tandem Diels-Alder/Fragmentation Approach to the Synthesis of Eleutherobin" *Org. Lett.* **2003**, 5, 1805-1808.
3. P. S. Wharton, and D. H. Bohlen, “Hydrazine Reduction of α,β -Epoxy Ketones to Allylic Alcohols” *J. Org. Chem.* **1961**, 26, 3615-3616.
4. G. Stork and P. G. Williard, “Five- and Six-Membered-Ring Formation from Olefinic α,β -Epoxy Ketones and Hydrazine”, *J. Am. Chem. Soc.* **1977**, 99, 7067-7068.
5. J. C. Justo de Pomar and J. A. Soderquist, “Unsymmetrical azines via triisopropylsilylhydrazine” *Tetrahedron Lett.* **2000**, 41, 3285-3289.

7. Appendices

N/A