



New Onset and Persistent Symptoms of Posttraumatic Stress Disorder Self Reported after Deployment and Combat Exposures: Prospective Population Based US Military Cohort Study

***T. C. Smith
M. A. K. Ryan, D. L. Wingard
D. J. Slymen, J. F. Sallis
D. Kritz-Silverstein
and for the Millennium Cohort Study***



Naval Health Research Center

Report No. 07-09

. Approved for public release; distribution is unlimited.

***Naval Health Research Center
140 Sylvester Road
San Diego, California 92106***

BMJ

New onset and persistent symptoms of post-traumatic stress disorder self reported after deployment and combat exposures: prospective population based US military cohort study

Tyler C Smith, Margaret A K Ryan, Deborah L Wingard, Donald J Slymen, James F Sallis, Donna Kritz-Silverstein and for the Millennium Cohort Study Team

BMJ 2008;336:366-371; originally published online 15 Jan 2008;
doi:10.1136/bmj.39430.638241.AE

Updated information and services can be found at:
<http://bmj.com/cgi/content/full/336/7640/366>

These include:

References

This article cites 29 articles, 13 of which can be accessed free at:
<http://bmj.com/cgi/content/full/336/7640/366#BIBL>

Rapid responses

You can respond to this article at:
<http://bmj.com/cgi/eletter-submit/336/7640/366>

Email alerting service

Receive free email alerts when new articles cite this article - sign up in the box at the top left of the article

Topic collections

Articles on similar topics can be found in the following collections

[Medical Consequences of Conflict](#) (484 articles)
[Anxiety disorders \(including OCD and PTSD\)](#) (228 articles)

Notes

To order reprints follow the "Request Permissions" link in the navigation box

To subscribe to *BMJ* go to:
<http://resources.bmj.com/bmj/subscribers>

New onset and persistent symptoms of post-traumatic stress disorder self reported after deployment and combat exposures: prospective population based US military cohort study

Tyler C Smith, director,^{1,2} Margaret A K Ryan, associate adjunct professor,^{1,2} Deborah L Wingard, professor and associate chief,² Donald J Slymen, professor,³ James F Sallis, professor,⁴ Donna Kritz-Silverstein, professor,⁵ for the Millennium Cohort Study Team

¹Department of Defense Center for Deployment Health Research at the Naval Health Research Center, San Diego, CA 92106, USA

²Division of Epidemiology, Department of Family and Preventive Medicine, University of California San Diego, La Jolla, CA 92093-0607

³Graduate School of Public Health, San Diego State University, San Diego, CA 92108

⁴Department of Psychology, San Diego State University, San Diego, CA 92108

⁵Division of Epidemiology, Department of Family and Preventive Medicine, University of California San Diego, La Jolla, CA 92093-0631

Correspondence to: T C Smith
tyler.c.smith@med.navy.mil

doi:10.1136/bmj.39430.638241.AE

ABSTRACT

Objective To describe new onset and persistence of self reported post-traumatic stress disorder symptoms in a large population based military cohort, many of whom were deployed in support of the wars in Iraq and Afghanistan.

Design Prospective cohort analysis.

Setting and participants Survey enrolment data from the millennium cohort (July 2001 to June 2003) obtained before the wars in Iraq and Afghanistan. Follow-up (June 2004 to February 2006) data on health outcomes collected from 50 184 participants.

Main outcome measures Self reported post-traumatic stress disorder as measured by the posttraumatic stress disorder checklist—civilian version using *Diagnostic and Statistical Manual of Mental Disorders*, fourth edition criteria.

Results More than 40% of the cohort were deployed between 2001 and 2006; between baseline and follow-up, 24% deployed for the first time in support of the wars in Iraq and Afghanistan. New incidence rates of 10-13 cases of post-traumatic stress disorder per 1000 person years occurred in the millennium cohort. New onset self reported post-traumatic stress disorder symptoms or diagnosis were identified in 7.6-8.7% of deployers who reported combat exposures, 1.4-2.1% of deployers who did not report combat exposures, and 2.3-3.0% of non-deployers. Among those with self reported symptoms of post-traumatic stress disorder at baseline, deployment did not affect persistence of symptoms.

Conclusions After adjustment for baseline characteristics, these prospective data indicate a threefold increase in new onset self reported post-traumatic stress disorder symptoms or diagnosis among deployed military personnel who reported combat exposures. The findings define the importance of post-traumatic stress disorder in this population and emphasise that specific combat exposures, rather than deployment itself, significantly affect the onset of symptoms of post-traumatic stress disorder after deployment.

INTRODUCTION

For more than five years the US military has been engaged in a combat intensive deployment in Iraq, Afghanistan, and neighbouring countries. Concern is growing among the public and veterans that post-deployment health consequences among US military personnel may be considerable and lasting. A recent report suggested that US marine and army infantry units returning from duty in Iraq and Afghanistan have higher than expected proportions of mental disorders and that as many as 10% of personnel are returning home with symptoms of post-traumatic stress disorder.¹ Combat duty in Iraq has been associated with high use of mental health services and attrition from military service and possible alterations in neural functioning after deployment.^{2,3}

Symptoms of post-traumatic stress disorder have been reported in as many as 30% of veterans after service in Vietnam and in more than 10% of US military personnel returning from the 1991 Gulf war.^{1,2,4-9} Epidemiological studies to date have largely focused on retrospective data or select groups. Prospective investigation of the causes of the disorder in large, population based military cohorts has been limited. The objective of this study was to prospectively investigate the effect of military deployment and self reported exposure to combat on new onset and persistent symptoms of post-traumatic stress disorder in a large population based US military cohort. The unique nature of the millennium cohort study allows the investigation of both personnel on active duty and Reserve/National Guard members who have remained in military service or who have separated from the military after returning from deployment.

METHODS

Study population

Between July 2001 and June 2003 the first panel of participants in the millennium cohort study, a population based US military cohort of 77 047 active military

duty and Reserve/National Guard personnel, was enrolled in a 22 year longitudinal study of health.¹⁰ The baseline enrolment ended with 36% of personnel invited consenting to participate in the 21 year study. Analyses to investigate potential reporting biases show no differences in responders' health with respect to hospital admissions and outpatient encounters in the year before enrolment (data not yet published); strong test-retest reliability¹¹; reliable vaccination reporting,^{12 13} occupation reporting,¹⁴ and deployment reporting¹⁵; and minimal differences between participants who chose web submission and those who chose paper submission.¹⁶ Between June 2004 and February 2006, 55 021 (71%) cohort members participated in the first three year follow-up questionnaire. Analyses of potential responder bias to the initial follow-up are ongoing.

To prospectively investigate new onset self reported symptoms of post-traumatic stress disorder, we removed participants deployed in support of the wars in Iraq and Afghanistan before their submission of the baseline questionnaire (n=2230) and those who were deployed while submitting the baseline questionnaire (n=621) from these analyses. Among those deployed,

we removed participants who submitted their follow-up questionnaires during their first deployment (n=1986), leaving 50 184 participants for these analyses.

We linked demographic and military personnel data to each participant; these reflected status at the time of baseline enrolment. These data included sex, birth year (categorised by groups: pre-1960, 1960-9, 1970-9, and 1980 forward), level of education (high school or less, some college or bachelor's degree, advanced degree), marital status (married, never married, divorced/other), pay grade (enlisted or officer), race/ethnicity (white non-Hispanic, black non-Hispanic, and other), service component (active duty or Reserve/National Guard), service branch (army, air force, navy/Coast Guard, or marines), and occupation (combat specialists, healthcare specialists, service supply and functional support specialists, or other). We used self reported data to supplement missing data from personnel records as needed.

Deployment data

We categorised regular active duty and Reserve/National Guard personnel who had been deployed

Table 1 | Baseline demographic and behavioural characteristics by deployment status of 50 128 millennium cohort members. Values are numbers (percentages)

Characteristic*	Deployed† (n=11 952)	Non-deployed† (n=38 176)
Sex		
Male	9727 (81.4)	26 552 (69.6)
Female	2225 (18.6)	11 624 (30.4)
Birth year		
Pre-1960	2 078 (17.4)	10 714 (28.1)
1960-9	4 943 (41.4)	15 362 (40.2)
1970-9	4 286 (35.9)	10 728 (28.1)
1980 and forward	645 (5.4)	1 372 (3.6)
Education		
High school or less diploma/equivalent	5 241 (43.9)	16 452 (43.1)
Some college or bachelor's degree	5 736 (48.0)	17 003 (44.5)
Advanced degree	975 (8.1)	4 721 (12.4)
Marital status		
Married	7 784 (65.1)	25 667 (67.2)
Never married	3 406 (28.5)	9 615 (25.2)
Divorced	762 (6.4)	2 894 (7.6)
Race/ethnicity		
White non-Hispanic	8 443 (70.6)	27 123 (71.1)
Black non-Hispanic	1 312 (11.0)	4 834 (12.7)
Other	2 197 (18.4)	6 219 (16.3)
Smoking		
Never smoker	7 126 (59.6)	22 816 (59.8)
Past smoker	2 855 (23.9)	9 768 (25.6)
Current smoker	1 971 (16.5)	5 592 (14.7)
Problem drinking		
No	9 723 (81.4)	31 327 (82.1)
Yes	2 229 (18.6)	6 849 (17.9)

*Taken at time of submission of baseline questionnaire among members who submitted questionnaires at baseline (July 2001-June 2003) and follow-up (July 2004-Jan 2006).

†Deployment in support of wars in Iraq and Afghanistan considered if full deployment occurred between submission dates of baseline and follow-up questionnaires. Cohort members who were deployed after follow-up survey are included with non-deployers in these analyses.

for one or more days in support of the wars in Iraq and Afghanistan between submission of their baseline and follow-up questionnaires as deployers. We categorised cohort members who had never been deployed or who were deployed for the first time after submission of their baseline and follow-up questionnaires as non-deployers in these analyses. The Defense Manpower Data Center, Monterey Bay, CA, provided deployment data, including entry and exit deployment dates. These data correlate strongly with millennium cohort self reported deployment status.¹⁵

Post-traumatic stress disorder assessment

Because of the complexity of diagnosing post-traumatic stress disorder and the differences in reported prevalence estimates,⁵ we used the standardised post-traumatic stress disorder checklist—civilian version, with highly sensitive and specific criteria for symptoms of post-traumatic stress disorder. The sensitive definition of symptoms of post-traumatic stress disorder used the *Diagnostic and Statistical Manual of Mental Disorders*, fourth edition (DSM-IV) criteria alone, whereas the specific definition of symptoms of post-traumatic stress disorder included the DSM-IV criteria and a requirement for a sum of 50 on a scale from 17 to 85 points for all of the questions in the checklist.¹⁷⁻²¹ The posttraumatic stress disorder checklist—civilian version is a 17 item self report measure of symptoms of post-traumatic stress disorder that requires participants to rate the severity of each symptom during the previous 30 days on a Likert-type scale ranging from 1 (not at all) to 5 (extremely). The DSM-IV criteria for post-traumatic stress disorder were met when a participant reported a moderate or higher level of at least one intrusion symptom, three avoidance symptoms, and two hyperarousal symptoms.¹⁸ The DSM-IV criteria alone are

reported to have a 100% sensitivity with a specificity of 92.0%, and the stricter criteria of at least 50 points on the checklist have shown reasonable sensitivity (0.60) and high specificity (0.99).¹⁷ The internal consistency of the standardised checklist in the millennium cohort, as measured by Cronbach's α ($\alpha=0.94$), indicated that these data were internally reliable for this population.¹¹

In addition to the assessment of self reported post-traumatic stress disorder symptoms at baseline with standard instruments, we asked participants, "Has your doctor or other health professional ever told you that you have any of the following conditions? . . . Post-traumatic stress disorder." In the follow-up questionnaire, we replaced "ever" with "in the past three years," which was used to indicate a more recent diagnosis.

We identified participants as having new onset self reported symptoms of post-traumatic stress disorder if they did not have symptoms or diagnosis at baseline but at follow-up met the criteria for the standardised checklist or reported a diagnosis by a healthcare worker of post-traumatic stress disorder within the previous three years. We identified participants as having persistent self reported symptoms of post-traumatic stress disorder if they were identified as having symptoms at baseline and at follow-up.

Other questionnaire information

We assessed cigarette smoking (never smoker, past smoker, current smoker) by using responses to the following questions: "In your lifetime, have you smoked at least 100 cigarettes (five packs)?", "In the past year have you used cigarettes?", and "Have you ever tried to quit smoking?" We assessed problem alcohol drinking (yes/no) with a standardised questionnaire.²²

Table 2 | Baseline military and occupational characteristics by deployment status, of 50 128 millennium cohort members. Values are numbers (percentages)

Characteristic*	Deployed† (n=11 952)	Non-deployed† (n=38 176)
Military rank		
Enlisted	8 802 (73.6)	27 556 (72.2)
Officer	3 150 (26.4)	10 620 (27.8)
Service component		
Reserve/National Guard	4 543 (38.0)	17 988 (47.1)
Active duty	7 409 (62.0)	20 188 (52.9)
Branch of service		
Army	5 548 (46.4)	18 438 (48.3)
Air force	4 333 (36.3)	10 608 (27.8)
Navy/Coast Guard	1 478 (12.4)	7 582 (19.9)
Marine Corps	593 (5.0)	1 548 (4.1)
Occupational category		
Other occupations	5 185 (43.4)	14 341 (37.6)
Combat specialists	2 901 (24.3)	7 114 (18.6)
Healthcare specialists	864 (7.2)	5 005 (13.1)
Service supply and functional	3 002 (25.1)	11 716 (30.7)

*Taken at time of submission of baseline questionnaire among members who submitted questionnaires at baseline (July 2001-June 2003) and follow-up (July 2004-Jan 2006).

†Deployment in support of wars in Iraq and Afghanistan considered if full deployment occurred between submission dates of baseline and follow-up questionnaires. Cohort members who were deployed after follow-up survey are included with non-deployers in these analyses.

To assess combat exposures, the follow-up survey asked: "During the past three years, have you been personally exposed to any of the following?" Answer options were "Witnessing a person's death due to war, disaster, or tragic event," "Witnessing instances of physical abuse (torture, beating, rape)," "Dead and/or decomposing bodies," "Maimed soldiers or civilians," and "Prisoners of war or refugees." We combined these with deployment experience to indicate a deployment with self reported stressful or combat exposures.

Statistical analysis

We completed descriptive and univariate analyses of population characteristics by deployment status and new onset and persistent self reported symptoms or diagnosis of post-traumatic stress disorder. We grouped deployment status as no deployment between baseline and follow-up questionnaires, deployment without self reported combat exposures between

submission of baseline and follow-up questionnaires, and deployment in support of the wars in Iraq and Afghanistan with self reported combat exposures between submission of baseline and follow-up questionnaires.

We used multivariable logistic regression to compare the adjusted odds of association between deployment in support of the wars in Iraq and Afghanistan and new onset self reported symptoms or diagnosis of post-traumatic stress disorder. We did secondary analyses of self reported persistence of post-traumatic stress disorder on a subpopulation of participants with symptoms at baseline. We used SAS software, version 9.1.3, for data management and statistical analyses.

RESULTS

Data were complete and available for these analyses for 50 128 (99.9%) of the 50 184 eligible cohort members. The mean elapsed time between baseline and

Table 3 | New onset and persistent self reported symptoms of post-traumatic stress disorder by demographic and behavioural characteristics of millennium cohort members. Values are numbers (percentages)

Characteristic*	No post-traumatic stress disorder symptoms or diagnosis at baseline		Post-traumatic stress disorder symptoms at baseline	
	New onset symptoms by specific criteria† (1347/48 447)	New onset symptoms by sensitive criteria‡ (1695/47 837)	Persisting symptoms by specific criteria† (455/995)	Persisting symptoms by sensitive criteria‡ (733/1659)
Sex				
Male	851 (2.4)	1084 (3.1)	296 (45.8)	482 (43.5)
Female	496 (3.8)	611 (4.7)	159 (45.7)	251 (45.6)
Birth year				
Pre-1960	276 (2.2)	343 (2.8)	137 (57.8)	191 (50.7)
1960-9	456 (2.3)	594 (3.1)	150 (45.7)	248 (44.1)
1970-9	512 (3.5)	627 (4.4)	138 (39.2)	237 (40.2)
1980 and forward	103 (5.4)	131 (7.0)	30 (38.5)	57 (43.5)
Education				
High school or less diploma/equivalent	859 (4.2)	1041 (5.1)	264 (42.7)	458 (43.5)
Some college or bachelor's degree	416 (1.9)	568 (2.6)	166 (49.4)	241 (44.2)
Advanced degree	72 (1.3)	86 (1.5)	25 (62.5)	34 (54.8)
Marital status				
Married	790 (2.4)	1000 (3.1)	261 (46.4)	421 (43.8)
Never married	420 (3.4)	532 (4.3)	141 (43.3)	233 (44.1)
Divorced	137 (3.9)	163 (4.8)	53 (50.0)	79 (46.8)
Race/ethnicity				
White non-Hispanic	889 (2.6)	1134 (3.3)	310 (46.6)	505 (44.1)
Black non-Hispanic	207 (3.5)	247 (4.2)	73 (47.4)	112 (46.5)
Other	251 (3.1)	314 (3.9)	72 (40.9)	116 (42.3)
Smoking at baseline				
Never smoker	654 (2.2)	831 (2.9)	185 (43.7)	294 (41.6)
Past smoker	356 (2.9)	463 (3.9)	136 (46.0)	220 (45.5)
Current smoker	337 (4.7)	401 (5.8)	134 (48.6)	219 (46.7)
Problem drinking at baseline				
No	1008 (2.5)	1278 (3.2)	289 (45.0)	459 (42.3)
Yes	339 (4.0)	417 (5.0)	166 (47.0)	274 (47.7)

*Taken at time of baseline submission of questionnaire among members who submitted questionnaires at baseline (July 2001-June 2003) and follow-up (July 2004-Jan 2006).

†Self reported post-traumatic stress disorder symptoms based on posttraumatic stress disorder patient checklist—civilian version with DSM-IV criteria or diagnosis of post-traumatic stress disorder within previous three years.

‡Self reported post-traumatic stress disorder symptoms based on posttraumatic stress disorder patient checklist—civilian version with DSM-IV criteria and a sum of 50 points out of 85 points possible or diagnosis of post-traumatic stress disorder within previous three years.

submission of follow-up questionnaire was 2.7 years (SD 0.5 years; median 2.8 years). Deployed cohort members were proportionately more likely than non-deployed cohort members to be male, born between 1970 and 1979, less educated, on active duty, and combat specialists (table 1 and table 2).

When we applied the specific criteria for post-traumatic stress disorder, 1681 participants had self reported symptoms of post-traumatic stress disorder or a previous diagnosis at baseline. When we applied the sensitive criteria for post-traumatic stress disorder, 2291 participants had self reported symptoms of post-traumatic stress disorder or a previous diagnosis at baseline. This left 48 447 or 47 837 cohort members respectively available for analyses of new onset self reported symptoms or diagnosis of post-traumatic stress disorder (table 3, table 4).

Tables 3 and 4 present analyses of new onset self reported symptoms or diagnosis of post-traumatic stress disorder. We found new onset symptoms defined by the specific criteria of DSM-IV with a sum of 50 points on the checklist in 4.3% of deployed cohort members and 2.3% of non-deployed cohort members.

We found new onset symptoms defined by the specific criteria of DSM-IV with a sum of 50 points on the checklist in 7.6% of cohort members who were deployed and self reported combat exposures, 1.4% of cohort members who were deployed and did not self report combat exposures, and 2.3% of non-deployed cohort members. Applying the more sensitive DSM-IV criteria alone, we identified new onset self reported symptoms of post-traumatic stress disorder in 8.7%, 2.1%, and 3.0% of these categories of cohort members. These data correspond to new onset self reported symptoms or diagnosis of post-traumatic stress disorder in up to 21 per 1000 non-combat deployers and up to 87 per 1000 combat deployed military personnel. New onset self reported symptoms of post-traumatic stress disorder were proportionately higher among cohort members who were female, younger, high school or less educated, never married or divorced, black non-Hispanic, enlisted, Reserve/National Guard members, army personnel, healthcare specialists, and service supply and functional specialists, and in those who self reported being a current smoker or problem drinker at baseline.

Table 4 | New onset and persistent self reported post-traumatic stress disorder symptoms by military and occupational characteristics of millennium cohort members. Values are numbers (percentages)

Characteristic*	No post-traumatic stress disorder symptoms or diagnosis at baseline		Post-traumatic stress disorder symptoms at baseline	
	New onset symptoms by specific criteria† (1347/48 447)	New onset symptoms by sensitive criteria‡ (1695/47 837)	Persisting symptoms by specific criteria† (455/995)	Persisting symptoms by sensitive criteria‡ (733/1659)
Deployment§				
Not deployed	849 (2.3)	1106 (3.0)	391 (47.6)	614 (45.9)
Deployed without combat exposures	89 (1.4)	128 (2.1)	17 (26.2)	30 (22.4)
Deployed with combat exposures	409 (7.6)	461 (8.7)	47 (43.5)	89 (47.9)
Military rank				
Enlisted	1162 (3.3)	1451 (4.2)	403 (45.0)	656 (43.9)
Officer	185 (1.4)	244 (1.8)	52 (52.0)	77 (46.4)
Service component				
Reserve/National Guard	630 (2.9)	784 (3.7)	218 (49.1)	338 (45.6)
Active duty	717 (2.7)	911 (3.5)	237 (43.0)	395 (43.1)
Branch of service				
Army	906 (3.9)	1103 (4.9)	280 (45.0)	459 (45.8)
Air force	184 (1.3)	267 (1.8)	82 (48.0)	114 (37.9)
Navy/Coast Guard	195 (2.2)	250 (2.9)	67 (42.7)	118 (42.6)
Marine Corps	62 (3.0)	75 (3.7)	26 (57.8)	42 (53.2)
Occupational category				
Other occupations	517 (2.7)	683 (3.7)	175 (42.8)	288 (42.4)
Combat specialists	237 (2.4)	290 (3.0)	70 (45.5)	18 (45.7)
Healthcare specialists	161 (2.9)	193 (3.5)	57 (56.4)	86 (50.0)
Service supply and functional	432 (3.1)	529 (3.8)	153 (46.2)	41 (43.8)

*Taken at time of baseline submission of questionnaire among members who submitted questionnaires at baseline (July 2001-June 2003) and follow-up (July 2004-Jan 2006).

†Self reported post-traumatic stress disorder symptoms based on posttraumatic stress disorder patient checklist—civilian version with DSM-IV criteria or diagnosis of post-traumatic stress disorder within previous three years.

‡Self reported post-traumatic stress disorder symptoms based on posttraumatic stress disorder patient checklist—civilian version with DSM-IV criteria and a sum of 50 points out of 85 points possible or diagnosis of post-traumatic stress disorder within previous three years.

§Deployment in support of wars in Iraq and Afghanistan considered if full deployment occurred between submission dates of baseline and follow-up questionnaires. Cohort members who were deployed after follow-up survey are included with non-deployers in these analyses.

Tables 3 and 4 also present persistent self reported symptoms of post-traumatic stress disorder among cohort members with symptoms at baseline (995 on the basis of the specific criteria and 1659 on the basis of the sensitive criteria). We removed members with a self reported diagnosis of post-traumatic stress disorder at

baseline but no current symptoms at baseline from these analyses. We found persistent self reported symptoms of post-traumatic stress disorder defined by the specific criteria in 43.5% of cohort members who were deployed to the wars in Iraq and Afghanistan and reported combat exposures, 26.2% of cohort members

Table 5 | Adjusted odds of new onset post-traumatic stress disorder*, stratified by US army and air force, among millennium cohort members without baseline post-traumatic stress disorder symptoms

Characteristic†	Army cohort members (n=22 959)		Air force cohort members (n=14 608)	
	Percentage	Odds ratio (95% CI)	Percentage	Odds ratio (95% CI)
Deployment‡				
Not deployed	3.0	1.00	1.2	1.00
Deployed without combat exposures	2.6	0.87 (0.64 to 1.18)	0.7	0.56 (0.35 to 0.89)
Deployed with combat exposures	9.3	3.59 (3.08 to 4.17)	3.5	3.38 (2.29 to 4.98)
Sex				
Male	3.6	1.00	1.0	1.00
Female	4.9	1.70 (1.44 to 2.00)	2.0	2.00 (1.41 to 2.83)
Birth year				
Pre-1960	3.1	1.00	1.2	1.00
1960-9	3.5	0.90 (0.74 to 1.10)	1.0	0.84 (0.56 to 1.24)
1970-9	4.9	1.09 (0.88 to 1.36)	1.6	1.40 (0.89 to 2.21)
1980 and forward	6.0	1.17 (0.83 to 1.65)	2.5	1.72 (0.70 to 4.26)
Education				
High school or less diploma/equivalent	5.0	1.00	2.2	1.00
Some college or bachelor's degree	2.8	0.83 (0.69 to 1.00)	1.2	0.75 (0.50 to 1.14)
Advanced degree	1.9	0.80 (0.54 to 1.19)	0.6	0.82 (0.35 to 1.92)
Marital status				
Married	3.7	1.00	1.1	1.00
Never married	4.4	0.80 (0.67 to 0.96)	1.3	0.79 (0.51 to 1.23)
Divorced	4.7	1.09 (0.85 to 1.40)	2.7	1.85 (1.25 to 2.74)
Race/ethnicity				
White non-Hispanic	4.0	1.00	1.3	1.00
Black non-Hispanic	4.5	1.14 (0.93 to 1.38)	1.2	0.86 (0.51 to 1.47)
Other	3.5	1.37 (1.13 to 1.65)	1.4	1.06 (0.63 to 1.76)
Smoking				
Never smoker	3.2	1.00	1.1	1.00
Past smoker	4.1	1.21 (1.03 to 1.44)	1.3	1.05 (0.73 to 1.50)
Current smoker	6.4	1.69 (1.42 to 2.01)	2.1	1.40 (0.94 to 2.07)
Problem drinking				
No	3.6	1.00	1.2	1.00
Yes	5.4	1.47 (1.25 to 1.73)	1.9	1.69 (1.17 to 2.43)
Military rank				
Enlisted	4.8	2.20 (1.70 to 2.86)	1.5	2.31 (1.24 to 4.30)
Officer	1.9	1.00	0.6	1.00
Service component				
Reserve/National Guard	3.8	1.00	1.5	1.00
Active duty	4.1	0.88 (0.76 to 1.03)	1.1	0.79 (0.56 to 1.13)
Occupational category				
Other occupations	4.4	1.00	1.3	1.00
Combat specialists	3.3	0.91 (0.75 to 1.11)	0.7	0.71 (0.41 to 1.25)
Healthcare specialists	3.7	1.02 (0.81 to 1.29)	1.3	0.78 (0.46 to 1.33)
Service supply and functional	4.1	0.95 (0.80 to 1.12)	1.4	0.93 (0.65 to 1.33)

*Self reported post-traumatic stress disorder symptoms based on posttraumatic stress disorder patient checklist—civilian version with DSM-IV criteria and a sum of 50 points out of 85 points possible or diagnosis of post-traumatic stress disorder within previous three years.

†Characteristic taken at time of submission of baseline questionnaire among members who submitted questionnaires at baseline (July 2001-June 2003) and follow-up (July 2004-Jan 2006).

‡Deployment in support of wars in Iraq and Afghanistan considered if full deployment occurred between submission dates of baseline and follow-up questionnaires. Cohort members who were deployed after follow-up survey are included with non-deployers in these analyses.

who were deployed and did not report combat exposures, and 47.6% of non-deployed cohort members. Applying the more sensitive criteria alone, we identified persistent self reported symptoms of post-traumatic stress disorder in 47.9%, 22.4%, and 45.9% of these groups. Persistent self reported symptoms were proportionately higher among older, higher educated,

and divorced personnel; among officers, Reserve/National Guard members, marines, and healthcare specialists; and in those who self reported being a smoker or problem drinker at baseline.

Using a variance inflation level of 4, we found no demographic, behavioural, or military variables that exhibited multicollinearity. Tables 5 and 6 present

Table 6 | Adjusted odds of new onset post-traumatic stress disorder*, stratified by US Navy/Coast Guard and Marine Corps, among millennium cohort members without baseline post-traumatic stress disorder symptoms

Characteristic†	Navy and Coast Guard cohort members (n=8655)		Marine cohort members (n=2077)	
	Percentage	Odds ratio (95% CI)	Percentage	Odds ratio (95% CI)
Deployment‡				
Not deployed	2.2	1.00	2.3	1.00
Deployed without combat exposures	1.4	0.60 (0.35 to 1.02)	2.8	1.42 (0.57 to 3.51)
Deployed with combat exposures	5.2	2.48 (1.48 to 4.14)	5.7	2.78 (1.52 to 5.07)
Sex				
Male	1.8	1.00	2.8	1.00
Female	3.4	1.73 (1.25 to 2.38)	4.4	1.92 (0.94 to 3.94)
Birth year				
Pre-1960	2.0	1.00	1.2	1.00
1960-9	2.1	0.99 (0.66 to 1.48)	1.4	0.71 (0.18 to 2.79)
1970-9	2.4	0.97 (0.60 to 1.56)	3.9	1.51 (0.40 to 5.70)
1980 and forward	5.2	1.73 (0.79 to 3.78)	6.8	2.00 (0.42 to 9.56)
Education				
High school or less diploma/equivalent	2.7	1.00	3.8	1.00
Some college or bachelor's degree	1.6	0.89 (0.57 to 1.39)	1.4	0.70 (0.22 to 2.20)
Advanced degree	1.6	1.31 (0.60 to 2.88)	0.7	0.57 (0.05 to 6.17)
Marital status				
Married	1.9	1.00	2.0	1.00
Never married	2.7	1.14 (0.81 to 1.61)	4.3	1.09 (0.56 to 2.10)
Divorced	4.9	2.53 (1.31 to 4.91)	5.7	2.75 (1.06 to 7.14)
Race/ethnicity				
White non-Hispanic	2.0	1.00	2.9	1.00
Black non-Hispanic	3.5	1.54 (1.02 to 2.34)	1.6	0.70 (0.21 to 2.36)
Other	2.6	1.22 (0.80 to 1.88)	4.3	1.53 (0.78 to 3.01)
Smoking				
Never smoker	1.8	1.00	2.1	1.00
Past smoker	2.5	1.33 (0.94 to 1.88)	3.6	1.55 (0.83 to 2.88)
Current smoker	3.4	1.59 (1.08 to 2.34)	5.7	1.84 (0.94 to 3.59)
Problem drinking				
No	2.0	1.00	2.4	1.00
Yes	3.1	1.59 (1.08 to 2.34)	4.7	1.73 (1.00 to 2.99)
Military rank				
Enlisted	2.7	2.14 (1.16 to 3.94)	3.7	1.92 (0.52 to 7.13)
Officer	1.3	1.00	1.1	1.00
Service component				
Reserve/National Guard	2.1	1.00	2.5	1.00
Active duty	2.3	1.24 (0.86 to 1.77)	3.1	1.24 (0.62 to 2.47)
Occupational category				
Other occupations	2.0	1.00	2.8	1.00
Combat specialists	1.7	1.14 (0.72 to 1.79)	3.7	1.63 (0.86 to 3.06)
Healthcare specialists	3.1	1.49 (0.96 to 2.32)	—	—
Service supply and functional	2.6	1.23 (0.85 to 1.79)	2.7	1.02 (0.52 to 1.99)

*Self reported post-traumatic stress disorder symptoms based on posttraumatic stress disorder patient checklist—civilian version with DSM-IV criteria and a sum of 50 points out of 85 points possible or diagnosis of post-traumatic stress disorder within previous 3 years.

†Taken at time of baseline questionnaire submission among members who submitted questionnaires at baseline (July 2001-June 2003) and follow-up (July 2004-Jan 2006).

‡Deployment in support of wars in Iraq and Afghanistan considered if full deployment occurred between submission dates of baseline and follow-up questionnaires. Cohort members who were deployed after follow-up survey are included with non-deployers in these analyses.

logistic regression results for new onset self reported symptoms or diagnosis of post-traumatic stress disorder based on the more specific criteria, stratified by branch of service. Analyses included adjustment for baseline characteristics including sex, age, education, marital status, race/ethnicity, rank, service component, occupation, cigarette smoking, and problem alcohol drinking. Among all of the services, deployed personnel who reported combat exposures had significantly higher odds of post-deployment self reported post-traumatic stress disorder symptoms than did those who were not deployed (army odds ratio 3.59, 95% confidence interval 3.08 to 4.17; air force odds ratio 3.38, 2.29 to 4.98; navy/Coast Guard odds ratio 2.48, 1.48 to 4.14; Marine Corps odds ratio 2.78, 1.52 to 5.07). Being female, divorced, or enlisted and reporting problem alcohol drinking were also associated with increased odds of new onset self reported symptoms or diagnosis in at least three of the four service branches. Among the air force members, deployers who did not report combat exposures had significantly lower odds of post-deployment self reported symptoms or diagnosis of post-traumatic stress disorder (odds ratio 0.56, 0.36 to 0.90) than did non-deployers. We found no statistical difference between deployed army, navy/Coast Guard, and Marine Corps cohort members who did not report combat exposures when compared with non-deployers.

DISCUSSION

In this study, we document an increased risk of new onset self reported symptoms of post-traumatic stress disorder among cohort members who were female, divorced, or enlisted, and in those who self reported being a current smoker or problem drinker at baseline. Furthermore, we report overall new incidence rates of 10-13 cases per 1000 person years and a threefold increase in new onset self reported symptoms of post-traumatic stress disorder in deployed personnel who reported combat exposures compared with non-deployers. The complexity and diversity of combat deployments and associated morbidity is a cause for much concern among veterans and the public about the health of military personnel returning from deployments to Iraq and Afghanistan.²³⁻²⁵ The unpredictability and intensity of urban combat, constant risk of roadside bombs, multiple and prolonged tours, and complex problems of differentiating enemies from allies can leave many troops with high stress levels and possible lasting health consequences. Millennium cohort study data allow a unique opportunity to observe new onset and persistent self reported symptoms of post-traumatic stress disorder in a large population based military cohort.

Limitations and strengths

Limitations to these analyses should be noted. The millennium cohort, by design, oversampled female, previously deployed, and Reserve/National Guard personnel and may not be representative of the military population in general or of all deployers. However,

approximately 40% of the cohort were deployed in support of the wars in Iraq and Afghanistan between 2001 and 2006; 24% were deployed for the first time between baseline and follow-up. Approximately 30% of the US military on rosters when the cohort was established were deployed during the same period (data not shown). This indicates a robust cohort for investigating deployment related concerns.

The millennium cohort study was able to track and obtain a follow-up questionnaire from more than 70% of those who submitted a baseline questionnaire; however, response biases may exist. Although participants self selected in accepting the invitation to become part of the cohort, and investigation of possible biases in follow-up participants continues, reports of millennium cohort baseline data suggest a representative sample of military personnel measured by demographic and health characteristics and reliable health and exposure reporting.^{10-16,26} Even so, assessment of exposure and outcomes through self report is challenging and represents an estimate of the true prevalence of exposures and disease. Studies have shown that increased reporting of exposure and symptoms occurs among personnel who seek medical compensation,²⁷ as well as among those with symptoms of post-traumatic stress disorder.²⁸ If the onset of symptoms in this cohort caused an increase in reporting of exposure, or if significant exposures caused an artificial increase in reporting of symptoms, the results of this study would overestimate the true burden of the disease. Furthermore, we categorised these data into deployment with and without combat on the basis of a set of five questions that may not truly indicate that a person specifically saw combat. This may in turn have biased these findings towards the null, and only true isolation of exposure and disease outcome would quantify the true effect. Compensation has not been offered to millennium cohort participants, which should reduce potential bias associated with compensation. Personnel may conceivably participate, however, and self report conditions and exposures likely to result in potential future healthcare benefits. More research into millennium cohort participants presenting for Veterans Administration care will help us to understand this potential reporting bias. Also, we could not discern those exposures that are unique to combat from those that might be encountered through other occupational or non-occupational activities. Furthermore, although these data have shown good reliability in test-retest analyses and strong internal consistency,¹¹ self reported data in general have inherent limitations that may lead to information bias in these analyses.

Lastly, the use of a standardised instrument for self reported data along with no specific date of trauma as a surrogate for post-traumatic stress disorder diagnosis is imperfect. However, the posttraumatic stress disorder checklist—civilian version using the DSM-IV criteria along with a sum of 50 points has been found to correlate well with a physician's assessment of post-traumatic stress disorder symptoms,¹⁷ and it is internally valid in millennium cohort members.¹¹

WHAT IS ALREADY KNOWN ON THIS TOPIC

Post-traumatic stress disorder is associated with comorbidities and independently with military combat deployments

No published data describe the incidence of post-traumatic stress disorder or the persistence of symptoms in a large population based cohort

WHAT THIS STUDY ADDS

Female, divorced, or enlisted personnel and those who self reported being a current smoker or problem drinker at baseline had an increased risk of new onset self reported symptoms of post-traumatic stress disorder

Deployed personnel who reported combat exposures had a threefold increase in new onset self reported symptoms of post-traumatic stress disorder compared with non-deployers

Unique strengths of these analyses should also be considered. This study is the first large, population based, prospective analysis of new onset self reported symptoms of post-traumatic stress disorder after combat deployment. Self reported data offer information not accessible elsewhere. Furthermore, post-traumatic stress disorder is often under-reported in electronic healthcare databases among populations who do not readily present for care for mental disorders.²⁹ The use of a standardised instrument allows for comparison with other populations, such as the general US population and other military populations.

Implications

Studies have estimated that as many as 30% of Vietnam war veterans developed post-traumatic stress disorder at some point after the war; between 9% and 15% had the disorder by the end of the 1980s.^{4 30 31} Among 1991 Gulf war veterans, as many as 10% were reported to have post-traumatic stress disorder symptoms years after returning from deployment.^{7 9} Although not all post-traumatic stress disorder is war related, trends of persistent symptoms of post-traumatic stress disorder in veterans suggest what may be expected after the current combat deployments in Iraq and Afghanistan. Early identification of personnel with symptoms of post-traumatic stress disorder may lead to a smaller burden of the disorder in the years to come if appropriate and timely treatments are provided. In this study, the prevalence of new onset self reported symptoms or diagnosis of post-traumatic stress disorder varied from 1% to 10% from baseline to follow-up.

Differences in the amount of self reported post-traumatic stress disorder in returning US military found in these analyses compared with past reports are likely to be due to variations in the populations studied, time periods between assessment, and prospective methods.¹⁻³² We used the millennium cohort to prospectively investigate new onset and persistent self reported symptoms of post-traumatic stress disorder by differentiating those with self reported symptoms at baseline. The average time between baseline and follow-up was nearly three years, whereas the average time between the end of first deployment and follow-up

was just over one year. A recent study reported that compared with 11.8% of US army personnel reporting symptoms of post-traumatic stress disorder soon after combat deployment, 16.7% reported symptoms six months post-deployment.³² This difference was larger among Reserve army personnel, 12.7% of whom reported symptoms after the end of deployment compared with 24.5% reporting symptoms six months post-deployment. In addition to the populations studied and the time periods between assessments, differences between these results and the results from our study may be due to the ability to consider those with and without symptoms or diagnosis of post-traumatic stress disorder at baseline. Nearly half of those with symptoms at baseline reported symptoms at follow-up, which would have increased the burden of post-deployment post-traumatic stress disorder symptoms if reported in the aggregate. In a recent report,¹ the authors found a pre-deployment prevalence of symptoms of post-traumatic stress disorder of 5-9%, compared with the millennium cohort baseline prevalence of 2-4%. The millennium cohort allowed us to investigate a large, population based sample of US military members, which differs in composition from those in specific combat units who are involved with combat operations on a daily basis. Future longitudinal investigation of the millennium cohort will yield information that may be more temporally comparable to studies of symptoms in Vietnam and 1991 Gulf war veterans, conducted many years after deployment.

Overall, the rate of new onset self reported symptoms of post-traumatic stress disorder in deployed personnel was 4.3%, a rate that is nearly twice that seen in non-deployed personnel (2.3%). A better comparison group than non-deployers, however, might be deployers without combat exposures. The new onset of as many as 87 cases of self reported symptoms of post-traumatic stress disorder per 1000 combat deployers, compared with 21 cases per 1000 non-combat deployers, suggests that up to 76% of new onset self reported symptoms or diagnosis of post-traumatic stress disorder may be attributed to combat exposures in deployers. This report documents a threefold adjusted increase in risk of new onset self reported symptoms of post-traumatic stress disorder among those who were deployed and who also report combat exposures compared with non-deployed cohort members. The increase in risk would be much larger if the comparison was with non-combat deployers, suggesting that deployment in itself may not lead to the onset of post-traumatic stress disorder symptoms but rather that exposures during deployment contribute to the onset of symptoms.

We found persistent self reported symptoms of post-traumatic stress disorder at follow-up in 40-50% of the 2.4% of millennium cohort members who had symptoms at baseline.²⁶ This implies resiliency or recovery among more than half of the population between baseline and follow-up. Even so, this underscores a considerable burden of persistent symptoms of post-traumatic stress disorder among those with baseline self

reported symptoms and suggests that resolution of post-traumatic stress disorder may not be expected for many years. Interestingly, cohort members who were older, higher educated, officers, and marines were more likely to experience persisting symptoms at follow-up. This contrasts with the lower odds of new onset symptoms in these personnel. This may reflect subgroups of the population who are concerned about the perceived stigma often associated with post-traumatic stress disorder. These groups may be less likely to report new onset symptoms; however, once they accept the diagnosis, they may be more willing to report problems they are experiencing. As found with new onset self reported symptoms of post-traumatic stress disorder, personnel with persisting symptoms were more likely to be divorced, Reserve/National Guard members, or healthcare specialists and to have reported problem alcohol drinking and cigarette smoking at baseline.

These prospective data show a threefold statistically significant increase in new onset self reported symptoms or diagnosis of post-traumatic stress disorder among recently deployed military personnel who report combat exposures. Compared with non-deployed personnel, we found no significant difference in new onset self reported symptoms or diagnosis of post-traumatic stress disorder among deployers who did not report combat exposures. Although the overall prevalence of post-traumatic stress disorder in the military is not high, a substantial number of cases can be expected on the basis of the number of service members deployed and exposed to combat over time in the wars in Iraq and Afghanistan. The risk of post-traumatic stress disorder conferred by combat exposure may not be preventable, but subsets of this population may exist who are even more vulnerable or, conversely, who retain a certain level of resilience to combat exposure. Future research should include efforts to better understand the resiliency and vulnerability to symptoms of post-traumatic stress disorder among subpopulations of combat deployers. Different types of research are also needed to investigate development of better validated and standardised screening and diagnostic approaches for post-traumatic stress disorder; population based controlled clinical trials on treatment strategies; and evaluations of how increased awareness, improved access to care, or both can reduce morbidity. Continued longitudinal investigation of post-traumatic stress disorder in the millennium cohort will yield even greater insight into the temporal onset of symptoms, the persistence or recurrence of symptoms, and recovery from this disorder.

We are indebted to all millennium cohort study participants. We thank Scott L Seggerman and Greg D Boyd from the Management Information Division, Defense Manpower Data Center, Seaside, CA. We also thank Lacy Farnell, Isabel Jacobson, Cynthia Leard, Travis Leleu, Robert Reed, Steven Spiegel, Karl Welch, and James Whitmer from the Department of Defense Center for Deployment Health Research; Michelle Stoia, from the Naval Health Research Center, San Diego, CA; and Karl E Friedl and professionals at the US Army Medical Research and Materiel Command, especially those from the Military Operational Medicine Research Program, Fort Detrick,

MD. We appreciate the support of the Henry M Jackson Foundation for the Advancement of Military Medicine, Rockville, MD.

In addition to the authors, the Millennium Cohort Study Team includes Paul J Amoroso, Madigan Army Medical Center, Tacoma, WA; Edward J Boyko, Seattle Epidemiologic Research and Information Center, Department of Veterans Affairs Puget Sound Health Care System, Seattle, WA; Gary D Gackstetter and Tomoko I Hooper, Department of Preventive Medicine and Biometrics, Uniformed Services University of the Health Sciences, Bethesda, MD; Gregory C Gray, College of Public Health, University of Iowa, Iowa City, IA; James R Riddle and Timothy S Wells, Air Force Research Laboratory, Wright-Patterson Air Force Base, OH; and Besa Smith, Department of Defense Center for Deployment Health Research at the Naval Health Research Center, San Diego, CA.

Contributors: TCS, MAKR, DLW, DJS, JFS, and DKS were involved in study design and concept. TCS and MAKR were involved in acquisition of data. TCS, MAKR, DLW, DJS, JFS, and DKS were responsible for analysis and interpretation of data. TCS, MAKR, DLW, DJS, JFS, and DKS drafted the manuscript. TCS, MAKR, DLW, DJS, JFS, and DKS did the statistical analysis. TCS, MAKR, DLW, DJS, JFS, and DKS critically revised the manuscript and approved the final version. TCS is the guarantor.

Funding: This represents Naval Health Research Center report 07-09, supported by the Department of Defense, under work unit No 60002. Funding sources did not play any other role. The views expressed in this article are those of the authors and do not reflect the official policy or position of the US Department of the Navy; US Department of the Army; US Department of the Air Force; US Department of Defense; US Department of Veterans Affairs; US Government; University of California, San Diego; or San Diego State University.

Competing interests: None declared.

Ethical approval: This research has been conducted in compliance with all applicable federal regulations governing the protection of human subjects in research (Protocol NHRC.2000.007).

Provenance and peer review: Not commissioned; externally peer reviewed.

- 1 Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL. Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *N Engl J Med* 2004;351:13-22.
- 2 Hoge CW, Auchterlonie JL, Milliken CS. Mental health problems, use of mental health services, and attrition from military service after returning from deployment to Iraq or Afghanistan. *JAMA* 2006;295:1023-32.
- 3 Vasterling JJ, Proctor SP, Amoroso P, Kane R, Heeren T, White RF. Neuropsychological outcomes of Army personnel following deployment to the Iraq war. *JAMA* 2006;296:519-29.
- 4 Dohrenwend BP, Turner JB, Turse NA, Adams BG, Koenen KC, Marshall R. The psychological risks of Vietnam for U.S. veterans: a revisit with new data and methods. *Science* 2006;313:979-82.
- 5 Thompson WW, Gottesman, II, Zalewski C. Reconciling disparate prevalence rates of PTSD in large samples of US male Vietnam veterans and their controls. *BMC Psychiatry* 2006;6:19.
- 6 Kang HK, Hyams KC. Mental health care needs among recent war veterans. *N Engl J Med* 2005;352:1289.
- 7 Kang HK, Natelson BH, Mahan CM, Lee KY, Murphy FM. Post-traumatic stress disorder and chronic fatigue syndrome-like illness among Gulf War veterans: a population-based survey of 30,000 veterans. *Am J Epidemiol* 2003;157:141-8.
- 8 Engel CJ, Engel A, Campbell S, McFall M, Russo J, Katon W. Posttraumatic stress disorder symptoms and precombat sexual and physical abuse in Desert Storm veterans. *J Nerv Ment Dis* 1993;181:683-8.
- 9 Stretch RH, Marlowe DH, Wright KM, Bliese PD, Knudson KH, Hoover CH. Post-traumatic stress disorder symptoms among Gulf war veterans. *Mil Med* 1996;161:407-10.
- 10 Ryan MA, Smith TC, Smith B, Amoroso P, Boyko EJ, Gray GC, et al. Millennium cohort: enrollment begins a 21-year contribution to understanding the impact of military service. *J Clin Epidemiol* 2007;60:181-91.
- 11 Smith TC, Smith B, Jacobson IG, Corbeil TE, Ryan MA. Reliability of standard health assessment instruments in a large, population-based cohort study. *Ann Epidemiol* 2007;17:271-84.
- 12 Smith B, Leard CA, Smith TC, Reed RJ, Ryan MA. Anthrax vaccination in the millennium cohort: validation and measures of health. *Am J Prev Med* 2007;32:347-53.
- 13 Leardmann CA, Smith B, Smith TC, Wells TS, Ryan MA. Smallpox vaccination: comparison of self-reported and electronic vaccine records in the millennium cohort study. *Hum Vaccin* 2007;3:245-51.
- 14 Smith TC, Jacobson IG, Smith B, Hooper TI, Ryan MA, Team FT. The occupational role of women in military service: validation of

- occupation and prevalence of exposures in the millennium cohort study. *Int J Environ Health Res* 2007;17:271-84.
- 15 Smith B, Wingard DL, Ryan MAK, Macera CA, Patterson TL, Slymen DJ. US military deployment during 2001-2006: comparison of subjective and objective data sources in a large prospective health study. *Ann Epidemiol* 2007;17:976-82.
 - 16 Smith B, Smith TC, Gray GC, Ryan MA. When epidemiology meets the internet: web-based surveys in the millennium cohort study. *Am J Epidemiol* 2007;166:1345-54.
 - 17 Brewin CR. Systematic review of screening instruments for adults at risk of PTSD. *J Trauma Stress* 2005;18:53-62.
 - 18 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*, 4th ed. Washington, DC: American Psychiatric Association, 1994.
 - 19 Weathers FW, Litz BT, Herman DS, Huska JA, Keane TM. The PTSD checklist (PCL): reliability, validity, and diagnostic utility. Annual Meeting of International Society for Traumatic Stress Studies, October 1993, San Antonio, Texas.
 - 20 Blanchard EB, Jones-Alexander J, Buckley TC, Forneris CA. Psychometric properties of the PTSD checklist (PCL). *Behav Res Ther* 1996;34:669-73.
 - 21 Wright KM, Huffman AH, Adler AB, Castro CA. Psychological screening program overview. *Mil Med* 2002;167:853-61.
 - 22 Ewing JA. Detecting alcoholism: the CAGE questionnaire. *JAMA* 1984;252:1905-7.
 - 23 Jones E, Hodgins-Vermaas R, McCartney H, Everitt B, Beech C, Poynter D, et al. Post-combat syndromes from the Boer war to the Gulf war: a cluster analysis of their nature and attribution. *BMJ* 2002;324:321-4.
 - 24 Jones E, Vermaas RH, McCartney H, Beech C, Palmer I, Hyams K, et al. Flashbacks and post-traumatic stress disorder: the genesis of a 20th-century diagnosis. *Br J Psychiatry* 2003;182:158-63.
 - 25 Hyams KC, Wignall FS, Roswell R. War syndromes and their evaluation: from the U.S. civil war to the Persian Gulf war. *Ann Intern Med* 1996;125:398-405.
 - 26 Riddle JR, Smith TC, Smith B, Corbeil TE, Engel CC, Wells TS, et al. Millennium cohort: the 2001-2003 baseline prevalence of mental disorders in the U.S. military. *J Clin Epidemiol* 2007;60:192-201.
 - 27 Frueh BC, Elhai JD, Grubaugh AL, Monnier J, Kashdan TB, Sauvageot JA, et al. Documented combat exposure of US veterans seeking treatment for combat-related post-traumatic stress disorder. *Br J Psychiatry* 2005;186:467-72.
 - 28 Koenen KC, Stellman SD, Dohrenwend BP, Sommer JF Jr, Stellman JM. The consistency of combat exposure reporting and course of PTSD in Vietnam war veterans. *J Trauma Stress* 2007;20:3-13.
 - 29 Magruder KM, Frueh BC, Knapp RG, Davis L, Hamner MB, Martin RH, et al. Prevalence of posttraumatic stress disorder in Veterans Affairs primary care clinics. *Gen Hosp Psychiatry* 2005;27:169-79.
 - 30 Kulka RA, Schlenger WE, Fairbank JA, Hough RL, Jordan BK, Marmar CR, et al. Trauma and the Vietnam war generation: report of findings from the national Vietnam veterans readjustment study. New York: Brunner/Mazel, 1990.
 - 31 Eisen SA, Griffith KH, Xian H, Scherrer JF, Fischer ID, Chantarujikapong S, et al. Lifetime and 12-month prevalence of psychiatric disorders in 8,169 male Vietnam war era veterans. *Mil Med* 2004;169:896-902.
 - 32 Milliken CS, Auchterlonie JL, Hoge CW. Longitudinal assessment of mental health problems among active and reserve component soldiers returning from the Iraq war. *JAMA* 2007;298:2141-8.

Accepted: 25 November 2007

REPORT DOCUMENTATION PAGE

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB Control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. Report Date (DD MM YY) 20 02 07		2. Report Type New		3. DATES COVERED (from - to)	
4. TITLE AND SUBTITLE New Onset and Persistent Symptoms of Post-traumatic Stress Disorder Self Reported After Deployment and Combat Exposure: Prospective Population Based US Military Cohort Study				5a. Contract Number: 5b. Grant Number: 5c. Program Element: 5d. Project Number: 5e. Task Number: 5f. Work Unit Number: 60002	
6. AUTHORS Tyler C. Smith, Margaret A.K. Ryan, Deborah L. Wingard, Donald J. Slymen, James F. Sallis, and Donna Kritz-Silverstein				9. PERFORMING ORGANIZATION REPORT NUMBER Report No. 07-09	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Health Research Center P.O. Box 85122 San Diego, CA 92186-5122					
8. SPONSORING/MONITORING AGENCY NAMES(S) AND ADDRESS(ES) Commanding Officer Commander Naval Medical Research Center Navy Medical Support Command 503 Robert Grant Ave P.O. Box 240 Silver Spring, MD 20910-7500 Jacksonville, FL 332212 0140				10. Sponsor/Monitor's Acronyms(s) NMRC/NMSC	
				11. Sponsor/Monitor's Report Number(s)	
12 DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited.					
13. SUPPLEMENTARY NOTES Published in: <u>British Medical Journal</u> , 2008, 336, 366-71					
14. ABSTRACT (maximum 200 words) Context There has been much concern over the health impact of military deployment to Iraq and Afghanistan. Posttraumatic stress disorder (PTSD) is reported to be prevalent in combat-exposed veterans. Objective To describe new onset and persistence of PTSD symptoms in a large population-based military cohort, many of whom deployed in support of the Global War on Terrorism (GWOT). Design, Setting, and Participants Millennium Cohort Study baseline enrollment data (July 2001 to June 2003) were obtained before the current GWOT. Follow-up (June 2004 to January 2006) data on health outcomes were collected from over 50 000 participants. Main Outcome Measures PTSD as measured by the PTSD Checklist-Civilian version (PCL-C) with the <i>Diagnostic and Statistical Manual of Mental Disorders</i>, 4th ed. (DSM-IV) criteria. Results Over 40% of the Cohort was deployed between 2001 and 2006, with 24% deployed for the first time in support of the GWOT between baseline and follow-up questionnaire submission. New-onset PTSD symptoms or diagnosis was identified in 7.6% to 8.7% of deployers reporting combat exposures, 1.4% to 2.1% of deployers not reporting combat exposures, and 2.3% to 3.0% of nondeployers. Among those with baseline PTSD symptoms, persistent symptoms were found in 47.9% of deployers reporting combat exposures and 45.9% of nondeployers. Temporal assessment of new-onset PTSD symptoms after deployment with combat exposures suggests a bimodal distribution of when symptoms may be most prevalent. Conclusions After adjusting for baseline characteristics, these prospective data indicate a threefold increase in new-onset PTSD symptoms or diagnosis among deployed military personnel reporting combat exposures. Findings define the importance of PTSD in this population and emphasize that specific combat exposures, rather than deployment itself, significantly affect the onset of PTSD symptoms postdeployment.					
15. SUBJECT TERMS combat disorders, stress disorders, posttraumatic, cohort studies					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UNCL	18. NUMBER OF PAGES 12	19a. NAME OF RESPONSIBLE PERSON Commanding Officer
a. REPORT UNCL	b. ABSTRACT UNCL	b. THIS PAGE UNCL			19b. TELEPHONE NUMBER (INCLUDING AREA CODE) COMM/DSN: (619) 553-8429