

AD _____

Award Number: **W81XWH-08-1-0062**

TITLE:

Merlin, the Hippo Pathway, and Tumor Suppression in Meningiomas

PRINCIPAL INVESTIGATOR:

Anita Lal, Ph.D.

CONTRACTING ORGANIZATION:

Regents of the University of California
San Francisco, CA 94143-0962

REPORT DATE:

February 2009

TYPE OF REPORT:

Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 07-02-2009		2. REPORT TYPE Final		3. DATES COVERED (From - To) 08 JAN 2008 - 7 JAN 2009	
4. TITLE AND SUBTITLE Merlin, the Hippo Pathway, and Tumor Suppression in Meningiomas				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-08-1-0062	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Anita Lal; Gilson S Baia				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Regents of the University of California c/o Office of Sponsored Research 3333 California Street, Ste. 315 San Francisco, CA 94143-0962				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The goal of this proposal was to determine whether merlin exerts its tumor suppressive effects in meningiomas by signaling through the Hippo pathway. Using paired meningioma cell lines, where the only difference is expression of merlin, we have shown that loss of merlin is associated with a clear increase in the protein levels of YAP, a transcriptional co-activator and downstream effector of the Hippo pathway. In addition to this increase, merlin loss was associated with nuclear localization of YAP. Using RNA interference and flow cytometry, we have shown that depletion of YAP results in a reversal of the enhanced proliferation phenotype caused by the loss of merlin. Thus, Merlin signals through YAP, a core component of the Hippo pathway, to regulate cell proliferation in meningiomas.					
15. SUBJECT TERMS Merlin, meningioma, Hippo pathway, YAP, RNA interference, flow cytometry					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 18	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	4
Body.....	4
Key Research Accomplishments.....	5
Reportable Outcomes.....	5
Conclusion.....	5
References.....	6
Appendices.....	10

INTRODUCTION

Intracranial meningiomas, often multicentric, occur in ~50% of NF2 patients and are associated with an increased risk of mortality. Mutations in the NF2 gene are detected in up to 60% of sporadic meningiomas and mice with a conditional knockout of NF2 in the leptomeninges develop meningiomas. Thus, it is clear that meningiomas are a serious clinical problem for NF2 patients and that loss of NF2 is critical for the development of meningiomas. However, the mechanism(s) by which the protein product of the NF2 gene, merlin, functions as a tumor suppressor in meningiomas is unclear. In *Drosophila*, it has been shown that merlin signals via the Hippo pathway to regulate growth and apoptosis. Components and signaling mechanism(s) of the Hippo pathway are conserved in vertebrates and several orthologues of this pathway have been implicated in the genesis of mammalian cancers. The aim of this proposal was to link the hippo signaling pathway to loss of merlin and its tumor suppressive functions in human meningiomas.

BODY

Using funding from a New Investigator Award and either RNA interference or retroviral mediated gene transfer, we have generated three paired meningioma and arachnoidal cell lines where the only difference is expression of merlin. These cell lines and the phenotypes generated by merlin loss in these cell lines are described in greater detail in the manuscript (recently published in Neoplasia) included in Appendix 1, and in the annual and final reports associated with proposal NF050061. In this proposal, these cell lines have been used to show a link between merlin loss and signaling through the hippo signaling pathway. Experimental details and the key findings from these experiments are also described in detail in the manuscript in Appendix 1.

Outlined below is a summary of the research accomplishments associated with each task outlined in the approved Statement of Work. Tables and Figures related to the text are included in the Supporting Data Section. We refer to Appendix 1 when referring to a figure in the included manuscript. For example, (Figure 1 in Appendix 1) would imply Figure 1 in the first attached manuscript in Appendix 1, while (Figure 1) refers to the Supporting Data Section.

Task 1. Characterize the effect of merlin expression on activation of components of the hippo pathway in meningioma cell lines (Months 1-5)

We firstly assessed protein levels and phosphorylation status of Mst1, Mst2, Lats1, Lats2 and YAP in meningioma cell lines. Protein levels of Mst2 (antibody from cell signaling #3952) were unaffected by merlin loss in meningioma cells (Figure 1). Similar results were obtained for Lats1 (antibody from cell signaling #9351S), Phospho-Lats1 (antibody from cell signaling #9157), while several attempts at optimizing western blot conditions for Mst1 (antibody from cell signaling), Phospho-Mst1/2 (antibody from cell signaling #3681S), Lats2 (antibody from Novus, #NB 200-199) did not result in clean, definite western blot results. In contrast, YAP protein levels were clearly induced by merlin loss (Figure 5A in Appendix 1) in meningioma cell lines. Phospho-YAP levels were also induced by merlin loss (Figure 2). However, given that total YAP levels are altered, the extent to which phosphorylation of YAP contributes to YAP function in meningiomas is unclear. Previous studies have reported that YAP translocates between the nucleus and the cytoplasm (1) and that it would need to be localized in the nucleus in order to perform its transcriptional coactivator functions. Therefore, we also performed immunocytochemistry for YAP in the paired cell lines and found that YAP was translocated to the nucleus in merlin-deficient cells (Figure 5B in Appendix 1).

We used quantitative PCR to assess the transcript levels of downstream target genes and found that, among the genes analyzed, Cyclin E1 transcript levels were induced in merlin-deficient cells (Figure 6 in Appendix 1). In addition, we showed by western blot that Cyclin E1 protein levels were also induced (Figure 6 in Appendix 1). Cyclin E1 is a downstream target of the Hippo pathway in *Drosophila* (2, 3) and our results suggest that merlin signals through YAP to Cyclin E1 to effect cell proliferation.

LAL, ANITA

Task 2. Evaluate the functional effects of depleting and overexpressing YAP (orthologue of Yorkie) in meningioma cell lines (Months 5-12)

We generated a YAP siRNA construct in the pSUPERneo retroviral vector instead of the pSUPERpuro retroviral construct since we had serious toxicity issues while using puromycin to determine kill curves in our meningioma cell lines and would be unable to use puromycin to generate stables in meningioma cell lines. Since our stable cell lines were already resistant to neomycin owing to the expression of the NF2 siRNA, we performed transient transfections with the YAPsiRNA construct. We optimized transfection conditions such as DNA concentration, time post-transfection to harvest, cell densities to obtain optimal depletion of YAP levels. Merlin loss induces cell proliferation, and this can be measured as an increase in the percentage of cells in the S-phase of the cell cycle using BrdU incorporation and flow cytometry. We used a similar methodology to assess the functional effects of depletion of YAP on the paired merlin-positive and merlin-negative cell lines. We found that depletion of YAP reversed the growth phenotype associated with merlin loss. While merlin loss enhanced S-phase entry, reduction of YAP caused ~50% decrease in the percentage of cells in the S-phase (Figure 7 in Appendix 1). These experiments were repeated three times with the same result and transfections of the YAP siRNA in AC1 cells also resulted in a similar trend. Thus, these results clearly demonstrate that YAP was

We subcloned the coding region of YAP in pBABEhygro and generated stable cell lines in merlin positive cell lines, AC1 and MENII-1/SF6717 using retroviral mediated gene transfer. However, several attempts at confirming YAP over-expression by western blot analysis were negative (Figure 3), indicating a problem with either the construct or the retroviral infection. Given that the experiments using RNA interference gave us clear results and confirmed a functional link between merlin and YAP, the hippo signaling pathway effector, we did not pursue this line of investigation any further.

KEY RESEARCH ACCOMPLISHMENTS

- 1) We have shown that protein levels of YAP are upregulated in response to merlin loss using three paired meningioma and arachnoidal cell lines where the only difference is expression of merlin
- 2) We have also shown that loss of merlin is associated with nuclear localization of YAP
- 3) We have shown that merlin loss is associated with an increase in transcript and protein levels of cyclin E1.
- 4) We have used RNA interference to deplete YAP levels and reverse the enhanced proliferation phenotype caused by merlin loss.
- 5) We have evidence that links the tumor suppressive mechanism of merlin to the hippo signaling pathway in human meningiomas.

REPORTABLE OUTCOMES

- 1) We are using the results of this research as preliminary data to apply for an investigator initiated neurofibromatosis grant in 2009.
- 2) We have successfully published a research publication in Neoplasia using the data described in this proposal.

CONCLUSION

The aim of this proposal was to link the hippo signaling pathway to loss of merlin and its tumor suppressive functions in human meningiomas. We have shown that in three independent paired meningioma and arachnoidal cell lines, where the only difference is merlin expression, loss of merlin is associated with an

LAL, ANITA

increase in protein levels of YAP and a concurrent nuclear localization of the induced YAP. In addition, we have shown that merlin loss is associated with an increase in transcript and protein levels of Cyclin E1 while levels of Cyclin D1 were unaffected. These results suggest that merlin signals through YAP which in turn effects Cyclin E1 transcription and cell proliferation. Using a shRNA specific for YAP, we have shown that depletion of YAP eliminates the enhanced proliferation induced in merlin-deficient cells, suggesting that merlin regulates cell proliferation by signaling through YAP. Thus, we have linked merlin to YAP, the downstream effector of the Hippo pathway, and have shown that merlin function is dependent on YAP induction.

REFERENCES

1. Hao Y, Chun A, Cheung K, Rashidi B, Yang X. Tumor suppressor LATS1 is a negative regulator of oncogene YAP. J Biol Chem 2008;283(9):5496-509.
2. Saucedo LJ, Edgar BA. Filling out the Hippo pathway. Nat Rev Mol Cell Biol 2007;8(8):613-21.
3. Pan D. Hippo signaling in organ size control. Genes Dev 2007;21(8):886-97.

BIBLIOGRAPHY

PUBLICATIONS

Striedinger KS, VandenBerg SR, Baia GS, McDermott MW, Gutmann DH and Lal A: The neurofibromatosis 2 tumor suppressor gene product, merlin, regulates meningioma cell growth by signaling through YAP. Neoplasia 10:1204-12, 2008

PERSONNEL RECEIVING PAY FROM THIS RESEARCH EFFORT

Anita Lal, Ph.D.
Gilson S Baia, MS




Figure 1. MST2 protein levels was unaffected in merlin-deficient cells. Western blot analysis of cell lysates derived from AC1, SF6717, and KT21MG1 stable cells were used to show that the phosphorylation status of MST2 was unchanged in the absence of merlin. Levels of α -tubulin were determined in the same samples as a loading control.




Figure 2. Total YAP and Phospho YAP expression was increased in merlin-deficient cells. Total YAP and Phospho YAP levels were measured by western blot analysis in cell lysates from SF6717 and KT21MG1 stable cells. Total YAP and Phospho YAP levels were induced by merlin loss.



Figure 3. Total YAP expression was negative in SF6717 stable clones. Western blot analysis of cell lysates derived from SF6717 stable cell lines did not confirm expression of YAP. Given that our experiments using shRNA specific for YAP were reproducible, definitive and conclusive, we did not pursue the overexpressing YAP line of investigation.

The Neurofibromatosis 2 Tumor Suppressor Gene Product, Merlin, Regulates Human Meningioma Cell Growth by Signaling through YAP¹

Katherine Striedinger*, Scott R. VandenBerg*, Gilson S. Baia*, Michael W. McDermott*, David H. Gutmann[†] and Anita Lal*

*Brain Tumor Research Center, Department of Neurological Surgery, University of California, San Francisco, CA 94143, USA; [†]Department of Neurology, Washington University School of Medicine, St. Louis, MO 63110, USA

Abstract

Neurofibromatosis type 2 (NF2) is an autosomal dominant disorder characterized by the occurrence of schwannomas and meningiomas. Several studies have examined the ability of the *NF2* gene product, merlin, to function as a tumor suppressor in diverse cell types; however, little is known about merlin growth regulation in meningiomas. In *Drosophila*, merlin controls cell proliferation and apoptosis by signaling through the Hippo pathway to inhibit the function of the transcriptional coactivator Yorkie. The Hippo pathway is conserved in mammals. On the basis of these observations, we developed human meningioma cell lines matched for merlin expression to evaluate merlin growth regulation and investigate the relationship between *NF2* status and Yes-associated protein (YAP), the mammalian homolog of Yorkie. *NF2* loss in meningioma cells was associated with loss of contact-dependent growth inhibition, enhanced anchorage-independent growth and increased cell proliferation due to increased S-phase entry. In addition, merlin loss in both meningioma cell lines and primary tumors resulted in increased YAP expression and nuclear localization. Finally, siRNA-mediated reduction of YAP in *NF2*-deficient meningioma cells rescued the effects of merlin loss on cell proliferation and S-phase entry. Collectively, these results represent the first demonstration that merlin regulates cell growth in human cancer cells by suppressing YAP.

Neoplasia (2008) 10, 1204–1212

Introduction

Neurofibromatosis type 2 (NF2) is a cancer predisposition syndrome phenotypically characterized by the occurrence of multiple nervous system tumors. The two most common tumors in this inherited syndrome are schwannomas and meningiomas [1]. Whereas meningiomas from individuals with NF2 exhibit biallelic inactivation of the *NF2* gene, loss of *NF2* expression is also detected in as many as 60% of sporadic meningiomas [2]. Similarly, genetically engineered mice with leptomeningeal *NF2* inactivation also develop meningiomas [3,4]. These findings strongly implicate the *NF2* gene in the pathogenesis of meningiomas; however, the molecular mechanism by which *NF2* regulates cell growth relevant to meningioma tumorigenesis remains unsolved.

Merlin (or schwannomin), the product of the *NF2* gene, is a member of the protein 4.1 family that links the actin cytoskeleton to plasma membrane proteins [5]. Although few studies have examined merlin loss in meningioma cells, loss of merlin in fibroblasts and Schwann cells results in loss of contact-dependent inhibition of proliferation, enhanced growth in soft agar and tumor formation in mice

[6,7]. In these cell types, merlin has been implicated in epidermal growth factor receptor [8], β_1 -integrin [9], and CD44 [7] function as well as Ras [10], Rac1 [11,12], phosphatidylinositol 3-kinase [13], mitogen-activated protein kinase [14], and signal transducer and activator of transcription [15] intracellular signaling. It is not known whether any of these growth control pathways are deregulated in *NF2*-deficient meningioma tumors.

Negative regulation of growth by merlin is conserved in *Drosophila*, where it acts upstream of the Hippo signaling pathway to coordinately

Address all correspondence to: Anita Lal, Brain Tumor Research Center, Department of Neurological Surgery, Box 0520, University of California, San Francisco, CA-94143. E-mail: anita.lal@ucsf.edu

¹Grant support: Department of Defense New Investigator Award (W81XWH-06-1-0221) and Concept Award (NF073066) to A. Lal and Department of Defense Award to D. Gutmann.

Received 29 May 2008; Revised 4 August 2008; Accepted 5 August 2008

Copyright © 2008 Neoplasia Press, Inc. All rights reserved 1522-8002/08/\$25.00
DOI 10.1593/neo.08642

regulate cell proliferation and apoptosis [16,17]. Mutations in merlin or other components of the Hippo pathway such as the serine/threonine kinases, Hippo and Warts, the adaptor molecule, Salvador, or Mats, results in activation of the transcriptional coactivator, Yorkie. Yorkie regulates expression of downstream target genes including *cyclin E* and *DIAP1* (*Drosophila* inhibitor of apoptosis protein 1) causing increased growth, delayed cell cycle exit, inhibition of apoptosis, and enhanced cell survival [16,17].

Individual components of the Hippo pathway are highly conserved in mammals, where they also regulate cell proliferation and apoptosis (Figure 1) [18,19]. Mice and humans have two Warts orthologs, Lats1 and Lats2. Mice deficient for Lats1 develop soft-tissue sarcomas and ovarian tumors [20]. The human ortholog of Salvador, hWW45, is mutated in cancer cell lines [21]. The two mammalian Hippo homologues, Mst1 and Mst2, promote apoptosis and regulate cell cycle exit [22]. Vertebrate Mst2 can rescue the lethality and overgrowth phenotypes of Hippo mutants in *Drosophila* [23]. Similar to their *Drosophila* counterparts, human Mst2 phosphorylates and activates both Lats1 and Lats2 [24]. The Yes-associated protein (YAP), the mammalian ortholog of Yorkie, is the primary effector of the mammalian Hippo pathway. Similar to the function of Yorkie in *Drosophila*, YAP causes aberrant tissue expansion in mice and induces epithelial transformation in mammary cells [25,26].

Given the conservation of components and mechanisms that operate downstream of merlin between *Drosophila* and mammals, we tested the functional relationship among merlin, Hippo pathway regulation, and growth suppression in human meningioma tumors. We developed nonneoplastic and neoplastic meningeal cell lines that mimic gain or loss of *NF2* expression and used these matched lines

to examine merlin regulation of YAP. We found that absence of merlin results in loss of contact-dependent inhibition of growth and promotes anchorage-independent growth. Merlin loss enhances cell proliferation by increasing entry into the S-phase and cyclin E1 expression. Inactivation of merlin results in increased YAP expression and nuclear accumulation of YAP in these meningioma cell lines and in primary human meningioma tumors. Finally, we show that YAP suppression reverses the proliferation effects associated with merlin loss in meningiomas. Collectively, these data demonstrate that merlin regulates cell growth in a YAP-dependent manner in meningiomas and suggests that YAP is a compelling target for therapeutic inhibition of human meningioma tumor growth.

Materials and Methods

Tumor Samples, Cell Lines, and Culture

All human meningioma tumor samples were collected by the Neurological Surgery Tissue Bank using protocols approved by the University of California, San Francisco Committee on Human Research. Human meningioma cell lines used were KT21MG1 [27] and MENII-1. MENII-1 cells were isolated from a surgically resected grade II meningioma and were immortalized by the expression of telomerase and the human papillomavirus *E6/E7* genes as described earlier [28]. Human arachnoidal cells (AC1) were cultured by plating small fragments of surgically resected spinal arachnoid tissue on scores scratched on the bottom of six-well tissue culture plates. Within 7 days, cells with characteristic arachnoidal morphology growing as a monolayer of polygonal cells with large cytoplasmic arcs migrate out of these scores. The arachnoidal origins of these cells were verified by positive staining for vimentin and desmoplakin. Primary cultures of arachnoidal cells were immortalized by stable transfection with the human papillomavirus *E6/E7* oncogenes and telomerase as described earlier [28]. All cell lines were maintained in Dulbecco's modified Eagle's medium supplemented with 10% fetal bovine serum and appropriate antibiotic selection markers.

Expression Constructs and Antibodies

The pSUPER.retro.neo-*NF2*-siRNA construct was generated by digesting the pSUPER.retro.neo vector (Oligoengine) with *Bgl*II and *Hind*III and ligating the annealed oligos (5'-gatccccGCAGCAAGCACAATACCATttcaagagaATGGTATTGTGCTTGCTGCTttttta and 5'-agcttaaaaaGCAGCAAGCACAATACCATtctcttgaaATGGTATTGTGCTTGCTGCTGCGggg) that contain a 19-nucleotide *NF2* target sequence (in caps) using the strategy described earlier [29]. The pSUPER.retro.neo-YAP-siRNA construct was also generated in pSUPER.retro.neo using the strategy described above and a previously described YAP target sequence (CCAGAGAATCAGTCAGAGA) [30]. A nonspecific mammalian scramble sequence (Oligoengine, Seattle, WA) in pSUPER.retro.neo (pSUPER.retro.neo-Control) was used as a control for the development of stable cell lines. Wild type *NF2*, S518A *NF2*, and S518D *NF2* mutant constructs in pUHD10.3 have been previously described [7,31]. These three constructs were subcloned into pBABE-Hygro using standard techniques. Merlin polyclonal (A19, #sc-331) and monoclonal (B12, #sc-55575) antibodies and the cyclin E1 monoclonal antibody (13A3, #sc-56310) were from Santa Cruz Biotechnology, Santa Cruz, CA. The YAP monoclonal antibody (#4912) was from Cell Signaling (Danvers, MA), the cyclin D1 monoclonal antibody




Figure 1. Schematic of the mammalian Hippo signaling pathway. The Hippo pathway is an evolutionary conserved cellular pathway that coordinately regulates cell proliferation and apoptosis. Merlin has been proposed to interact with unknown membrane proteins and transduce a signal that stimulates the phosphorylation of LATS1/2 by the serine/threonine kinases MST1/2 that interact with hWW45. LATS1/2 inhibits the transcriptional coactivator YAP resulting in suppressed expression of downstream target genes such as cyclins that are involved in cell growth and proliferation.




Figure 2. *In vitro* model system of merlin expression in human meningeal cells. Endogenous merlin was silenced in AC1 and MENII-1 cells using *NF2* specific siRNA. In parallel, merlin isoform 1 was exogenously expressed in KT21MG1 cells using retroviral mediated gene transfer. (A) *NF2* transcript levels were measured by quantitative PCR and showed a 5.6-fold reduction in MENII-1-*NF2*-siRNA cells compared with MENII-1-Control cells. Asterisk denotes statistical significance ($P < .05$). (B) Western blot analysis of cell lysates derived from AC1, MENII-1, and KT21MG1 stable cell populations was used to confirm loss or gain of merlin. Whereas merlin expression was observed in AC1-Control and MENII-1-Control cells, *NF2* siRNA abolished expression of merlin in AC1-*NF2*-siRNA and MENII-1-*NF2*-siRNA cells. In parallel, KT21MG1-Control cells lacked merlin, whereas KT21MG1-*NF2* cells expressed wild type merlin. Levels of α -tubulin were determined in the same samples as a loading control. Immunoblot of one representative experiment of three with similar results is shown. (C) Immunofluorescence using the A19 polyclonal antibody against merlin revealed the presence of cytoplasmic staining in AC1-Control and MENII-1-Control cells and its absence in AC1-*NF2*-siRNA and MENII-1-*NF2*-siRNA cells. In contrast, KT21MG1-Control cells had no staining, whereas KT21MG1-*NF2* cells had cytoplasmic staining. Merlin immunolabeling is shown in green, and nuclear DAPI counterstaining is shown in blue.

(clone DCS-6) was from BD Pharmingen (Franklin Lakes, NJ), and the α -tubulin (#CP06) antibody was from Calbiochem (San Diego, CA).

Retroviral Infection and Selection of Stable Cell Populations

To stably suppress *NF2* in MENII-1 and AC1 cell lines, retroviral supernatants were generated by transfecting Phoenix A packaging cells with pSUPER.retro.neo-*NF2*-siRNA or pSUPER.retro.neo-Control using Lipofectamine 2000 Plus Reagent (Invitrogen, Carlsbad, CA). The 48-hour posttransfection supernatant was harvested, filtered, and used to infect MENII-1 and AC1 cell lines in the presence of 8 μ g/ml polybrene. Stable cell populations were selected using 500 μ g/ml G418. In parallel, stable cell populations expressing wild type or mutant *NF2* were generated by transfecting Phoenix A cells with the particular pBABE-Hygro construct and infecting KT21MG1 cells with the 48-hour posttransfection supernatant. Stable cell populations were selected using 200 μ g/ml hygromycin. Empty pBABE-Hygro vector was used as a negative control.

Transient Suppression of YAP

Merlin-positive and -negative MENII-1 stable cell populations generated above were plated at 80% confluency in 100-mm dishes and transfected with 10 μ g of pSUPER.retro.neo-YAP siRNA using Lipofectamine 2000 Plus Reagent (Invitrogen). Empty pSUPER.retro.neo vector was used as a negative control. The 72-hour post-

transfection cells were subjected to flow cytometry to quantify bromodeoxyuridine (BrdU) uptake (described below).

Quantitative Polymerase Chain Reaction

Quantitative polymerase chain reaction (PCR) was performed in three independent experiments using cDNA templates with the I-cycler machine (Bio-Rad, Hercules, CA) and SYBR Green I (Molecular Probes, Eugene, OR) using PCR conditions and data analysis as described earlier [32]. Primers specific for GAPDH and actin were used to verify the integrity of the cDNA and to normalize cDNA yields. The primers used were: GAPDH, 5'-GGAAGCTTGTCATCAATGGAA and 5'-AAATGAGCCCCAGCCTTCTC; Actin, 5'-CAGGAGGAGCAATGATCTTG and 5'-ACTCTTC-CAGCCTTCCTTCC; *NF2*, 5'-ACCGTTGCCTCCTGACATAC and 5'-TCGGAGTTCTCATTGTGCAG; YAP, 5'-GCAGTTGG-GAGCTGTTTCTC and 5'-GCCATGTTGTTGTCTGATCG; cyclin E1, 5'-CCATCCTTCTCCACCAAAGA and 5'-TTTGA-TGCCATCCACAGAAA; cyclin D1, 5'-TGTTTGCAAGCAG-GACTTTG and 5'-CCTTCCGGTGTGAAACATCT.

Western Blot Analysis

Total cell lysates were prepared either in buffer A (50 mM Tris-HCl, pH 7.5; 1 mM EDTA pH 8.0, 1% Triton) for merlin detection or in 1 $\times$ SDS buffer following manufacturer's instructions (Cell

Signaling) for YAP and cyclin protein detection. Protein (50–200 μ g) was resolved by electrophoresis for each sample and was transferred to a polyvinylidene difluoride membrane. Membranes were blocked in 5% low-fat dry milk in Tris-buffered saline–Tween 20 and incubated overnight at 4°C with either merlin A19 or B12 antibodies, or the YAP, cyclin E1, or cyclin D1 antibody, or α -tubulin. Incubation with horseradish peroxidase–conjugated goat antirabbit or antimouse immunoglobulin (Jackson ImmunoResearch, West Grove, PA) was performed for 1 hour at room temperature. Bound antibody was visualized by chemiluminescence using the SuperSignal West Pico substrate (Pierce Chemical Co., Rockford, IL). The molecular weights were determined with the use of prestained protein ladders (BioRad, Hercules, CA, and Invitrogen). Films were scanned and exported as TIFF files.

Immunofluorescence Microscopy

Indirect immunofluorescence for merlin (A19 antibody or B12 antibody) and YAP was performed as described earlier [28]. Briefly, cells were fixed, permeabilized, blocked, and sequentially incubated with primary and secondary (Alexa 488 goat antirabbit IgG or Alexa 546 goat antimouse IgG) antibodies. Cells were mounted in DAPI mounting media, examined, and photographed with a microscope (Zeiss, Thornwood, NY).

Immunohistochemistry

Immunohistochemical staining was performed on 5- μ m formalin-fixed, paraffin-embedded sections from meningioma tissue (8 primary tumors) and from a meningioma tissue microarray (29 primary tumors) for merlin (B12 antibody) and YAP as described earlier [33]. Slides were reviewed in consultation with a neuropathologist (S.R.V.) and included 33 WHO grade I and 4 WHO grade III meningiomas.

Growth Curves and Soft Agar Assay

Merlin-positive and -negative stable cell populations (20,000 cells) were plated in 24-well plates, and cells from three wells were counted at 3, 6, 12, 18, 25, and 30 days. To assess colony growth in soft agar, 50,000 cells were plated in Dulbecco's modified Eagle's medium in 0.4% low melting temperature agarose upon a layer of 0.8% agarose. After 8 weeks, colonies were stained with 0.005% crystal violet, and colonies larger than 100 μ m in diameter were scored by counting under a microscope.

Flow Cytometry

Cells (70–80% confluent) were incubated with 1 mM BrdU for 3 hours at 37°C and processed using the fluorescein isothiocyanate BrdU Flow Kit (BD Biosciences, San Jose, CA) following manufacturer's instructions. Briefly, 1×10^6 trypsinized cells were fixed, permeabilized, and digested with DNase. Cells were then stained with fluorescein isothiocyanate–conjugated anti-BrdU and 7-aminocincomycin (7-AAD). Flow cytometry was performed on a Becton Dickinson FACSCalibur machine. For each experiment, 10,000 events were counted. Data acquisition was performed with the CellQuest software (BD Biosciences), and data were analyzed using Flow Jo v8.5.3.

Statistics

All data are expressed as mean $\pm$ SEM. GraphPad Prism version 4 was used for statistical analysis, consisting of unpaired *t* test and significant differences with a *P* < .05.

Results

Establishment of Matched NF2-Expressing and NF2-Deficient Human Arachnoidal and Meningioma Cell Lines

To develop a meningioma-specific NF2 *in vitro* model system, we determined the expression levels of endogenous merlin in normal human arachnoidal and meningioma cell lines (data not shown). On the basis of these results, we selected one arachnoidal cell line (AC1), one merlin-positive grade II meningioma cell line (MENII-1), and one merlin-negative grade III meningioma cell line (KT21MG1) for further analysis. Paired cell lines were generated by either suppressing NF2 expression in AC1 and MENII-1 cells using RNA interference (siRNA) or by overexpressing the full-length human NF2 cDNA (isoform 1, lacking exon 16 sequences) in KT21MG1 cells after retroviral-mediated gene transfer. Stable cell populations expressing either NF2 siRNA (AC1-NF2-siRNA or MENII-1-NF2-siRNA) or a nonspecific target siRNA (AC1-Control or MENII-1-Control) were




Figure 3. Suppression of merlin causes loss of contact-dependent inhibition of growth and promotes anchorage-independent growth. (A) Growth curves in the presence and absence of merlin expression. Cultures were subconfluent during the first 6 days. MENII-1-NF2-siRNA cells (dotted lines) continued growing after confluent conditions and have less contact-dependent inhibition of growth compared with MENII-1-Control cells (solid lines). Each line corresponds to representative cultures. (B) NF2 suppression promotes anchorage-independent growth. Marked increased in colony formation in soft agar was observed in cells without merlin expression as MENII-1-NF2-siRNA and KT21MG1-Control cells compared with MENII-1-Control and KT21MG1-NF2, respectively. Representative images of the colonies formed (upper panel) and the mean number of colonies per well (lower panel) are shown. Error bars equal $\pm$ SE of three independent experiments. Asterisks denote statistical significance (*P* < .05).




Figure 4. Merlin loss enhances S-phase entry. AC1 and meningioma cell lines (MENII-1 and KT21MG1) were labeled with BrdU and 7-AAD to assess the cell cycle distribution of individual cells by flow cytometry. (A) Representative flow cytometric histograms indicate an increase in the percent of BrdU-positive cells in AC1-*NF2*-siRNA, MENII-1-*NF2*-siRNA, and KT21MG1-Control cells compared with AC1-Control, MENII-1-Control, and KT21MG1-*NF2* cells, respectively. (B) Bar graphs depict the percentage of cells in the S-phase of the cell cycle (BrdU-positive cells) averaged from three independent experiments. Error bars correspond to $\pm$ SE. Asterisks denote statistical significance using unpaired *t* test ($P < .05$). (C) Table shows the mean of the percentage of cells $\pm$ SE in each phase of the cell cycle from three independent experiments.

selected. *NF2* transcript levels were 5.6-fold lower in MENII-1-*NF2*-siRNA cells when compared with MENII-1-Control cells using quantitative PCR (Figure 2A). Similarly, merlin protein expression was undetectable in AC1-*NF2*-siRNA and MENII-1-*NF2*-siRNA cells by Western blot analysis and immunofluorescence (Figure 2, B and C). In parallel, stable cell populations expressing either exogenous merlin (KT21MG1-*NF2*) or empty vector (KT21MG1-Control) were selected. Expression of merlin in KT21MG1-*NF2* cells was confirmed by Western blot and immunofluorescence (Figure 2, B and C). In total, we have generated three separate human meningeal and meningioma cell lines that differ only in their expression of merlin for subsequent study.

Merlin Loss Increases S-Phase Entry, Cell Proliferation, and Promotes Anchorage-Independent Growth

Next, we assessed the effect of merlin loss on the growth properties of these human meningioma cell lines. MENII-1-*NF2*-siRNA cells exhibited a more pronounced loss of contact-dependent inhibition of growth compared with MENII-1-Control cells (Figure 3A). Merlin loss also promoted colony formation in soft agar. In these experiments, MENII-1-*NF2*-siRNA cells formed a greater number of colonies (10.6 ± 3.2) larger than 100 μ m in diameter compared with MENII-1-Control cells (0.4 ± 0.4 ; $P = .01$; Figure 3B). Conversely, merlin expression in KT21MG1 cells significantly decreased the formation of colonies (32 ± 6.3) compared with merlin-negative KT21MG1 cells (361 ± 4.9 ; $P \leq .0001$; Figure 3B).

To determine the effect of merlin on cell cycle progression, we measured BrdU incorporation and total DNA content by flow cytometry. Loss of merlin in AC1 and MENII-1 cells resulted in a significant increase in the percentage of BrdU-positive cells, indicated by an increase in S-phase entry (Figure 4). This is consistent with the observed increase in proliferation in *NF2*-deficient cells. Conversely, expression of exogenous merlin in KT21MG1 cells induced G₀/G₁ arrest and a concomitant decrease in the S-phase cell population (Figure 4). These results demonstrate that merlin functions as a negative growth regulator for both nonneoplastic leptomeningeal cells and meningioma cells and establishes this system as a tractable experimental platform for examining growth regulatory pathways.

Merlin Loss Is Associated with an Increase in YAP Protein Expression

In *Drosophila*, merlin controls cell proliferation and apoptosis by signaling through the Hippo pathway and its effector protein Yorkie, the ortholog of YAP [16,18]. To determine whether merlin might regulate meningioma cell growth by modulating Hippo pathway signaling, we investigated whether changes in merlin expression were associated with altered levels of YAP using our human *NF2* meningioma model system. Transcript levels of YAP were unaffected by merlin loss in MENII-1 cells (data not shown); however, YAP protein expression was elevated in arachnoidal and meningioma cells lacking merlin expression and decreased in KT21MG1 cells expressing wild type merlin (Figure 5A). These results suggest that merlin regulates YAP expression at the translational or posttranslational level.




Figure 5. Protein levels of YAP are up-regulated and localized to the nucleus in *NF2*-deficient cells. (A) Total cell lysates were subjected to Western blot using a YAP- or merlin-specific antibody. Increased YAP protein expression was observed when *NF2* was suppressed in AC1 and MENII-1 cells compared with controls. Conversely, exogenous expression of merlin decreased YAP in KT21MG1 cells compared with controls. Expression of a nonphosphorylated, active merlin (S518A *NF2*) was also associated with lower levels of YAP compared with the expression of pseudophosphorylated inactive merlin (S518D *NF2*). Levels of α -tubulin were determined in the same samples as loading control. Results were reproduced in three independent experiments. (B) YAP was translocated to the nucleus in merlin-deficient cells. Immunofluorescence staining was used to show that YAP was localized to the nucleus in AC1-*NF2*-siRNA, MENII-1-*NF2*-siRNA, and KT21MG1-Control cells. In contrast, YAP was primarily cytoplasmic in AC1-Control, MENII-1-Control, and KT21MG1-*NF2* cells. Merlin immunolabeling is shown in red; YAP staining is shown in green; and nuclear DAPI counterstaining is shown in blue. (C) *In situ* immunostaining of merlin and YAP in serial sections of primary human meningioma tumors. We surveyed 37 primary meningiomas by immunohistochemistry. YAP expression was minimal to absent in 95% of merlin-positive meningiomas (a representative tumor is shown here as *Meningioma 1*). In contrast, YAP was expressed and localized to the nucleus in 92% of merlin-negative meningiomas (a representative tumor is shown here as *Meningioma 2*). Arrow depicts an example of YAP nuclear localization. Insets show images at higher magnification.

Merlin becomes active after dephosphorylation of the conserved C-terminal serine 518 (S518) residue [31,34]. Previous studies have shown that merlin mutants in which this phosphorylatable residue is changed to alanine (S518A) are constitutively nonphosphorylated and active, whereas those containing aspartic acid (S518D) are pseudophosphorylated and nonfunctional as negative growth regulators [31,34]. Using these mutants, we assessed the effect of merlin phosphorylation status on YAP protein levels in KT21MG1 human meningioma cells. YAP protein levels were markedly up-regulated in KT21MG1 cells expressing the S518D inactive mutant compared with those with active merlin S518A expression (Figure 5A), suggesting that active and functional merlin was required to inhibit YAP.

YAP has been reported to shuttle between the cytoplasm and the nucleus where it can function as a transcriptional coactivator [35]. To determine whether merlin regulates the subcellular localization of YAP, we used fluorescence immunocytochemistry to examine *NF2*-deficient and *NF2*-expressing AC1 and MENII-1 cells. YAP was preferentially expressed in the cytoplasm of AC1-Control and MENII-1-Control cells. In contrast, YAP was localized in the nucleus in AC1-*NF2*-siRNA and MENII-1-*NF2*-siRNA cells (Figure 5B). Collectively, these data indicate that merlin regulates YAP protein expression and YAP nuclear localization.

To assess whether a similar association exists between merlin and YAP in human meningioma tumors, we surveyed 37 sporadic primary meningioma tumors by immunohistochemistry. After immunostaining with a merlin-specific monoclonal antibody, meningiomas were classified as either merlin-positive, if tumors exhibited any positive immunoreactivity, or merlin-negative, if tumors had no immunoreactivity. We then assessed protein levels of YAP in adjacent serial sections. 13 (92%) of 14 merlin-negative meningiomas exhibited strong nuclear YAP immunoreactivity. In contrast, 22 (95%) of 23 merlin-positive meningiomas had weak to no YAP immunoreactivity (Figure 5C). These results further support the *in vitro* results demonstrating that merlin regulates YAP protein levels *in vivo*.

Merlin Loss Is Associated with an Increase in Protein Levels of Cyclin E1

Previous studies have identified cyclin E1 as a transcriptional target of the Hippo pathway in *Drosophila* [36–38] and an essential regulator of progression from G₁ to S cell cycle progression in mammalian cells [39,40]. Also, merlin has been reported to inhibit cell proliferation by repressing cyclin D1 in human mesothelioma cells [41]. To determine whether merlin regulates cyclin E1 and/or cyclin D1 levels in human meningioma cells, we measured cyclin E1 and D1 RNA and protein expression by quantitative PCR and Western blot analysis, respectively. Cyclin E1 transcript levels were at least 2.5-fold higher in MENII-1-*NF2*-siRNA cells compared with MENII-1-Control cells, whereas cyclin D1 transcript levels were the same in MENII-1-*NF2*-siRNA and MENII-1-Control cells (Figure 6A). In addition, cyclin E1 protein levels were elevated in MENII-1-*NF2*-siRNA and AC1-*NF2*-siRNA cells compared with MENII-1-Control and AC1-Control cells, respectively (Figure 6B). Conversely, exogenous expression of merlin in *NF2*-deficient KT21MG1 cells resulted in decreased cyclin E1 protein levels. In contrast, cyclin D1 protein levels were unaffected by the absence or presence of merlin in both MENII-1 and KT21MG1 cells (Figure 6C). These data suggest that merlin likely regulates cell

growth by modulating cyclin E1 expression at the transcriptional level in meningiomas.

YAP Reduction Reverses the Growth Phenotype Associated with Merlin Loss

Lastly, to determine whether YAP expression is necessary for the enhanced S-phase entry induced by merlin loss in meningioma cells, we transiently depleted YAP using RNA interference in MENII-1-Control and MENII-1-*NF2*-siRNA cells (Figure 7). Reduced YAP expression in *NF2*-deficient MENII-1 meningioma cells caused a ~50% decrease in the percentage of cells in S-phase (9 ± 1.2) compared with mock-transfected cells (20 ± 0.2 ; $P = .001$; Figure 7, B and C). In contrast, YAP siRNA treatment had a minor effect on MENII-1-Control cells (~20% reduction; $P = .08$; Figure 7, B and C). Collectively, these experiments demonstrate that YAP transduces the merlin growth regulatory signal in meningiomas and that the Hippo growth control pathway is responsible for merlin tumor suppressor function in this tumor type.

Discussion

The Hippo signaling pathway is emerging as an evolutionarily conserved mechanism that controls organ size and growth relevant to tumorigenesis [18,37]. In this study, we investigated the functional association between *NF2* gene expression and the downstream effector of the Hippo pathway in human meningiomas. We provide several lines of converging and complementary evidence that merlin functions through YAP in meningiomas. First, using paired meningioma cell lines differing only in *NF2* expression, we show that

YAP expression is increased in a dose-response manner upon merlin loss. This regulation occurs at the translational or posttranslational level and results in YAP nuclear localization. Second, the relationship between merlin expression and YAP nuclear localization is also observed in human surgical meningioma specimens *in vivo*. These observations are consistent with findings made in other solid cancers in which Hippo signaling is deregulated [42]. Third, suppressed YAP expression in merlin-deficient meningioma cells attenuates the cell growth and S-phase cell cycle progression associated with merlin loss. To the best of our knowledge, these results represent the first demonstration that merlin regulates cell growth in human meningioma cells by suppressing YAP, the main Hippo pathway effector protein.

YAP has previously been implicated in other human cancers, including pancreatic ductal adenocarcinoma, and has been shown to function as an oncogene that induces epithelial transformation of mouse mammary cells [26,43]. Nuclear localization of YAP is dependent on the phosphorylation status of a conserved Ser residue and is necessary for its cotranscriptional activator function [35,44]. YAP associates with multiple transcription factors in the nucleus such as p73 and TEAD/TEF [44,45]. ErbB4 receptors have been reported to recruit YAP and relocate to the nucleus to regulate transcription [46,47]. In contrast, LATS1 inactivates YAP oncogenic function by sequestering YAP in the cytoplasm and, consequently, suppressing its transcriptional regulation of cellular genes [35]. Our findings are in accordance with these and suggest that the tumorigenic behavior of meningioma cells is driven in part by YAP nuclear localization and the transcription of genes involved in increased proliferation.




Figure 6. Cyclin E1 and cyclin D1 expression was increased in merlin-deficient cells. (A) Transcript levels of cyclin E1 and cyclin D1 were measured in MENII-1 cells using quantitative PCR. At least a 2.5-fold increase in the transcript levels of cyclin E1 was seen in MENII-1-*NF2*-siRNA cells compared with MENII-1-Control cells, whereas transcript levels of cyclin D1 were unchanged. Asterisk denotes statistical significance ($P < .05$). (B) Western blot analysis of cell lysates derived from AC1, MENII-1, and KT21MG1 stable cells was used to show that protein levels of cyclin E1 were increased in the absence of merlin. Levels of α -tubulin were determined in the same samples as a loading control. (C) Western blot analysis of cell lysates derived from MENII-1 and KT21MG1 stable cells was used to show that protein levels of cyclin D1 were unchanged in the absence of merlin. Levels of α -tubulin were determined in the same samples as a loading control.




Figure 7. Down-regulation of YAP decreased proliferation in merlin-deficient cells. MENII-1-Control and MENII-1-NF2-siRNA cells were transiently transfected with YAP-specific siRNA or empty vector. (A) YAP transcript levels were measured in MENII-1-Control cells using quantitative PCR. The YAP-specific siRNA caused a 50% reduction in YAP transcript levels compared with controls in MENII-1 cells. (B, C) BrdU incorporation and 7-AAD staining were measured by flow cytometry. (B) One representative experiment shows the distribution of cells in G₀-G₁, S, and G₂ phases of the cell cycle. The table below shows the mean of the percentage of cells ±SE in each phase of the cell cycle from three independent experiments. (C) Bar graphs depict the percentage of cells in the S-phase of the cell cycle (BrdU-positive cells) averaged from three independent experiments. Error bars correspond to ±SE. Suppression of YAP decreased the percentage of cells in S-phase in MENII-1-NF2-siRNA cells to levels similar to MENII-1-Control cells.

We demonstrate that merlin controls the cell cycle in meningiomas. Merlin suppression results in increased S-phase entry, and merlin expression resulted in G₀/G₁ arrest. Similar results were obtained in mesotheliomas [41] and schwannomas [48]. These results provide strong evidence that merlin functions as a tumor suppressor by controlling the G₀/G₁ to S-phase checkpoint of the cell cycle. Cyclin E1 is thought to be essential for this cell cycle transition in humans [39,40] and has been identified as a downstream target of Yorkie in *Drosophila* [36–38]. In contrast, cyclin E1 was not in-

duced after overexpression of YAP in mammary epithelial cells [26]. Instead, it has been shown that cyclin D1 is regulated by YAP in mouse intestine [25] and by merlin in human mesotheliomas [41]. Our experiments show that merlin regulates cyclin E1 but not cyclin D1 in human meningiomas. These differences most likely represent cell type-specific differences in gene expression. Further studies are needed to determine whether merlin regulates cyclin E1 by signaling through YAP.

The fact that NF2 patients develop only certain CNS tumors also emphasizes the cell type-specific tumorigenic effects of merlin. Merlin loss results specifically in aberrant growth of Schwann and meningeal cell types. However, not all meningiomas have NF2 loss, suggesting that other mechanisms that do not involve the Hippo pathway result in meningioma formation. Alternatively, it remains to be elucidated whether other components of the Hippo pathway, downstream of merlin, are possibly deregulated in these tumors resulting in the same phenotype as merlin loss.

Treatment strategies for NF2 and sporadic meningioma patients are restricted to traditional forms of cancer therapy such as surgery and radiation therapy. These options are sometimes insufficient because of the location of these tumors, the recurrence despite therapies and the occurrence of multiple tumors. Although receptor tyrosine kinases inhibitors are emerging as pathway targets in other CNS tumors such as glioblastomas [49], targeted therapies have not yet been proposed for meningioma patients. This is mainly because of a lack of knowledge regarding relevant signaling pathways. Our results argue that YAP is an attractive candidate as a key mediator of NF2 growth regulation and tumorigenesis in meningioma, making it a potential target for the development of therapies for NF2 and meningioma patients.

Acknowledgments

The authors thank the Neurological Surgery Tissue Bank at the University of California, San Francisco, CA, for the primary meningioma tumor samples. The authors also thank Jason Pomerantz for fruitful discussions and critical review of the manuscript.

References

- [1] Ferner RE (2007). Neurofibromatosis 1 and neurofibromatosis 2: a twenty first century perspective. *Lancet Neurol* 6, 340–351.
- [2] Rutledge MH, Sarrazin J, Rangaratnam S, Phelan CM, Twist E, Merel P, Delattre O, Thomas G, Nordenskjold M, Collins VP, et al. (1994). Evidence for the complete inactivation of the NF2 gene in the majority of sporadic meningiomas. *Nat Genet* 6, 180–184.
- [3] Kalamirides M, Niwa-Kawakita M, Leblais H, Abramowski V, Perricaudet M, Janin A, Thomas G, Gutmann DH, and Giovannini M (2002). NF2 gene inactivation in arachnoid cells is rate-limiting for meningioma development in the mouse. *Genes Dev* 16, 1060–1065.
- [4] Kalamirides M, Stemmer-Rachamimov AO, Takahashi M, Han ZY, Chareyre F, Niwa-Kawakita M, Black PM, Carroll RS, and Giovannini M (2008). Natural history of meningioma development in mice reveals: a synergy of NF2 and p16 (Ink4a) mutations. *Brain Pathol* 18, 62–70.
- [5] Gautreau A, Louvard D, and Arpin M (2002). ERM proteins and NF2 tumor suppressor: the yin and yang of cortical actin organization and cell growth signaling. *Curr Opin Cell Biol* 14, 104–109.
- [6] Johnson KC, Kissil JL, Fry JL, and Jacks T (2002). Cellular transformation by a FERM domain mutant of the NF2 tumor suppressor gene. *Oncogene* 21, 5990–5997.
- [7] Morrison H, Sherman LS, Legg J, Banine F, Isacke C, Haipiek CA, Gutmann DH, Ponta H, and Herrlich P (2001). The NF2 tumor suppressor gene product, merlin, mediates contact inhibition of growth through interactions with CD44. *Genes Dev* 15, 968–980.

- [8] Curto M, Cole BK, Lallemand D, Liu CH, and McClatchey AI (2007). Contact-dependent inhibition of EGFR signaling by Nf2/Merlin. *J Cell Biol* **177**, 893–903.
- [9] Fernandez-Valle C, Tang Y, Ricard J, Rodenas-Ruano A, Taylor A, Hackler E, Biggerstaff J, and Iacovelli J (2002). Paxillin binds schwannomin and regulates its density-dependent localization and effect on cell morphology. *Nat Genet* **31**, 354–362.
- [10] Tikoo A, Varga M, Ramesh V, Gusella J, and Maruta H (1994). An anti-Ras function of neurofibromatosis type 2 gene product (NF2/Merlin). *J Biol Chem* **269**, 23387–23390.
- [11] Shaw RJ, Paez JG, Curto M, Yaktine A, Pruitt WM, Saotome I, O'Bryan JP, Gupta V, Ratner N, Der CJ, et al. (2001). The Nf2 tumor suppressor, merlin, functions in Rac-dependent signaling. *Dev Cell* **1**, 63–72.
- [12] Morrison H, Sperka T, Manent J, Giovannini M, Ponta H, and Herrlich P (2007). Merlin/neurofibromatosis type 2 suppresses growth by inhibiting the activation of Ras and Rac. *Cancer Res* **67**, 520–527.
- [13] Rong R, Tang X, Gutmann DH, and Ye K (2004). Neurofibromatosis 2 (NF2) tumor suppressor merlin inhibits phosphatidylinositol 3-kinase through binding to PIKE-L. *Proc Natl Acad Sci USA* **101**, 18200–18205.
- [14] Chadee DN, Xu D, Hung G, Andalibi A, Lim DJ, Luo Z, Gutmann DH, and Kyriakis JM (2006). Mixed-lineage kinase 3 regulates B-Raf through maintenance of the B-Raf/Raf-1 complex and inhibition by the NF2 tumor suppressor protein. *Proc Natl Acad Sci USA* **103**, 4463–4468.
- [15] Scoles DR, Nguyen VD, Qin Y, Sun CX, Morrison H, Gutmann DH, and Pulst SM (2002). Neurofibromatosis 2 (NF2) tumor suppressor schwannomin and its interacting protein HRS regulate STAT signaling. *Hum Mol Genet* **11**, 3179–3189.
- [16] Hamaratoglu F, Willecke M, Kango-Singh M, Nolo R, Hyun E, Tao C, Jafar-Nejad H, and Halder G (2006). The tumour-suppressor genes *NF2/Merlin* and *Expanded* act through Hippo signalling to regulate cell proliferation and apoptosis. *Nat Cell Biol* **8**, 27–36.
- [17] Harvey K and Tapon N (2007). The Salvador-Warts-Hippo pathway—an emerging tumour-suppressor network. *Nat Rev Cancer* **7**, 182–191.
- [18] Dong J, Feldmann G, Huang J, Wu S, Zhang N, Comerford SA, Gayyed ME, Anders RA, Maitra A, and Pan D (2007). Elucidation of a universal size-control mechanism in *Drosophila* and mammals. *Cell* **130**, 1120–1133.
- [19] Zeng Q and Hong W (2008). The emerging role of the hippo pathway in cell contact inhibition, organ size control, and cancer development in mammals. *Cancer Cell* **13**, 188–192.
- [20] Turenchalk GS, St John MA, Tao W, and Xu T (1999). The role of lats in cell cycle regulation and tumorigenesis. *Biochim Biophys Acta* **1424**, M9–M16.
- [21] Tapon N, Harvey KF, Bell DW, Wahrer DC, Schiripo TA, Haber DA, and Hariharan IK (2002). Salvador promotes both cell cycle exit and apoptosis in *Drosophila* and is mutated in human cancer cell lines. *Cell* **110**, 467–478.
- [22] Dan I, Watanabe NM, and Kusumi A (2001). The Ste20 group kinases as regulators of MAP kinase cascades. *Trends Cell Biol* **11**, 220–230.
- [23] Wu S, Huang J, Dong J, and Pan D (2003). Hippo encodes a Ste-20 family protein kinase that restricts cell proliferation and promotes apoptosis in conjunction with salvador and warts. *Cell* **114**, 445–456.
- [24] Chan EH, Nousiainen M, Chalamalasetty RB, Schafer A, Nigg EA, and Sillje HH (2005). The Ste20-like kinase Mst2 activates the human large tumor suppressor kinase Lats1. *Oncogene* **24**, 2076–2086.
- [25] Camargo FD, Gokhale S, Johnnidis JB, Fu D, Bell GW, Jaenisch R, and Brummelkamp TR (2007). YAP1 increases organ size and expands undifferentiated progenitor cells. *Curr Biol* **17**, 2054–2060.
- [26] Overholtzer M, Zhang J, Smolen GA, Muir B, Li W, Sgroi DC, Deng CX, Brugge JS, and Haber DA (2006). Transforming properties of YAP, a candidate oncogene on the chromosome 11q22 amplicon. *Proc Natl Acad Sci USA* **103**, 12405–12410.
- [27] Tanaka K, Sato C, Maeda Y, Koike M, Matsutani M, Yamada K, and Miyaki M (1989). Establishment of a human malignant meningioma cell line with amplified *c-myc* oncogene. *Cancer* **64**, 2243–2249.
- [28] Baia GS, Slocum AL, Hyer JD, Misra A, Sehati N, Vandenberg SR, Feuerstein BG, Deen DF, McDermott MW, and Lal A (2006). A genetic strategy to overcome the senescence of primary meningioma cell cultures. *J Neurooncol* **78**, 113–121.
- [29] Brummelkamp TR, Bernards R, and Agami R (2002). Stable suppression of tumorigenicity by virus-mediated RNA interference. *Cancer Cell* **2**, 243–247.
- [30] Levy D, Adamovich Y, Reuven N, and Shaul Y (2007). The Yes-associated protein 1 stabilizes p73 by preventing Itch-mediated ubiquitination of p73. *Cell Death Differ* **14**, 743–751.
- [31] Surace EI, Haipke CA, and Gutmann DH (2004). Effect of merlin phosphorylation on neurofibromatosis 2 (NF2) gene function. *Oncogene* **23**, 580–587.
- [32] Cuevas IC, Slocum AL, Jun P, Costello JF, Bollen AW, Riggins GJ, McDermott MW, and Lal A (2005). Meningioma transcript profiles reveal deregulated notch signaling pathway. *Cancer Res* **65**, 5070–5075.
- [33] Yoo H, Baia GS, Smith JS, McDermott MW, Bollen AW, Vandenberg SR, Lamborn KR, and Lal A (2007). Expression of the hypoxia marker carbonic anhydrase 9 is associated with anaplastic phenotypes in meningiomas. *Clin Cancer Res* **13**, 68–75.
- [34] Rong R, Surace EI, Haipke CA, Gutmann DH, and Ye K (2004). Serine 518 phosphorylation modulates merlin intramolecular association and binding to critical effectors important for NF2 growth suppression. *Oncogene* **23**, 8447–8454.
- [35] Hao Y, Chun A, Cheung K, Rashidi B, and Yang X (2008). Tumor suppressor LATS1 is a negative regulator of oncogene YAP. *J Biol Chem* **283**, 5496–5509.
- [36] Pan D (2007). Hippo signaling in organ size control. *Genes Dev* **21**, 886–897.
- [37] Edgar BA (2006). From cell structure to transcription: Hippo forges a new path. *Cell* **124**, 267–273.
- [38] Saucedo LJ and Edgar BA (2007). Filling out the Hippo pathway. *Nat Rev Mol Cell Biol* **8**, 613–621.
- [39] Ohtsubo M, Theodoras AM, Schumacher J, Roberts JM, and Pagano M (1995). Human cyclin E, a nuclear protein essential for the G₁-to-S phase transition. *Mol Cell Biol* **15**, 2612–2624.
- [40] Ohtsubo M and Roberts JM (1993). Cyclin-dependent regulation of G₁ in mammalian fibroblasts. *Science* **259**, 1908–1912.
- [41] Xiao GH, Gallagher R, Shetler J, Skele K, Altomare DA, Pestell RG, Jhanwar S, and Testa JR (2005). The NF2 tumor suppressor gene product, merlin, inhibits cell proliferation and cell cycle progression by repressing cyclin D1 expression. *Mol Cell Biol* **25**, 2384–2394.
- [42] Zhao B, Wei X, Li W, Udan RS, Yang Q, Kim J, Xie J, Ikenoue T, Yu J, Li L, et al. (2007). Inactivation of YAP oncoprotein by the Hippo pathway is involved in cell contact inhibition and tissue growth control. *Genes Dev* **21**, 2747–2761.
- [43] Guo J, Kleeff J, Zhao Y, Li J, Giese T, Esposito I, Buchler MW, Korc M, and Friess H (2006). Yes-associated protein (YAP65) in relation to Smad7 expression in human pancreatic ductal adenocarcinoma. *Int J Mol Med* **17**, 761–767.
- [44] Basu S, Totty NF, Irwin MS, Sudol M, and Downward J (2003). Akt phosphorylates the Yes-associated protein, YAP, to induce interaction with 14-3-3 and attenuation of p73-mediated apoptosis. *Mol Cell* **11**, 11–23.
- [45] Vassilev A, Kaneko KJ, Shu H, Zhao Y, and DePamphilis ML (2001). TEAD/TEF transcription factors utilize the activation domain of YAP65, a Src/Yes-associated protein localized in the cytoplasm. *Genes Dev* **15**, 1229–1241.
- [46] Omerovic J, Puggioni EM, Napoletano S, Visco V, Fraioli R, Frati L, Gulino A, and Alimandi M (2004). Ligand-regulated association of ErbB-4 to the transcriptional co-activator YAP65 controls transcription at the nuclear level. *Exp Cell Res* **294**, 469–479.
- [47] Komuro A, Nagai M, Navin NE, and Sudol M (2003). WW domain-containing protein YAP associates with ErbB-4 and acts as a co-transcriptional activator for the carboxyl-terminal fragment of ErbB-4 that translocates to the nucleus. *J Biol Chem* **278**, 33334–33341.
- [48] Schulze KM, Hanemann CO, Muller HW, and Hanenberg H (2002). Transduction of wild-type merlin into human schwannoma cells decreases schwannoma cell growth and induces apoptosis. *Hum Mol Genet* **11**, 69–76.
- [49] Stommel JM, Kimmelman AC, Ying H, Nabioullin R, Ponugoti AH, Wiedemeyer R, Stegh AH, Bradner JE, Ligon KL, Brennan C, et al. (2007). Coactivation of receptor tyrosine kinases affects the response of tumor cells to targeted therapies. *Science* **318**, 287–290.