

AD _____

Award Number: W81XWH-07-1-0131

TITLE: TAF1, From a General Transcription Factor to Modulator of
Androgen Receptor in Prostate Cancer

PRINCIPAL INVESTIGATOR: Peyman Tavassoli M.D., Paul Rennie Ph.D.

CONTRACTING ORGANIZATION: University of British Columbia
Vancouver, BC, Canada V6T 1Z3

REPORT DATE: February 2009

TYPE OF REPORT: Annual Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				<i>Form Approved</i> OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 14-02-2009		2. REPORT TYPE Annual Summary		3. DATES COVERED 15 JAN 2008 - 14 JAN 2009	
4. TITLE AND SUBTITLE TAF1, From a General Transcription Factor to Modulator of Androgen Receptor in Prostate Cancer				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-07-1-0131	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Peyman Tavassoli M.D., Paul Rennie Ph.D. Email: peyman.tavassoli@vch.ca				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of British Columbia Vancouver, BC, Canada V6T 1Z3				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The androgen receptor (AR) is a ligand-activated transcription factor that is essential for development and progression of prostate cancer. Using the N-terminus of AR as bait in the repressed transactivator yeast two-hybrid system, TATA binding protein-associated factor 1 (TAF1) was identified. TAF1, a multifunctional protein that contains acetylation, ubiquitin activating and kinase domains, can interact with several proteins to promote or suppress gene transcription. In the present study, using GST pull-down assay we confirmed that TAF1 binds to NTD of AR through the E1/E2 and HAT domains. We demonstrated by co-immunoprecipitation and ChIP assays that TAF1 and AR bind in the nucleus and associate with an androgen response element (ARE) at the proximal promoter of the prostate specific antigen (PSA) gene in the presence of androgen. In addition, we found that TAF1 acts as a coactivator and enhances AR transcriptional activity through its NTK and E1/E2 domains without influencing the general transcriptional activity of a non-androgen responsive promoter. By using in-cell and in-tube ubiquitination assays, our finding also suggested that TAF1 could ubiquitinate AR.					
15. SUBJECT TERMS Androgen receptor, Prostate cancer, transcription					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 16	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	4
Hypothesis.....	4
Specific Aims.....	4
Progress and Results.....	4
Summary of last year report.....	4
Specific Aim 2.....	5
TAF1 interacts with the N-terminus of AR mainly through its HAT and E1/E2 domains	
Specific Aim 3.....	6
TAF1 and AR phosphorylation.....	6
Androgen receptor is ubiquitinated by TAF1.....	6
TAF1 ubiquitinates AR <i>in vitro</i>	6
Key Research Accomplishments.....	7
Reportable Outcomes.....	7
 Conclusion.....	 7
Figure 1.....	10
Figure 2.....	11
Figure 3.....	12
Appendix.....	13-14
Figure I.....	13
Figure II.....	14
 References.....	 15-16

INTRODUCTION

Prostate cancer is the most commonly diagnosed non-skin cancer in men and one of the leading causes of cancer death. Androgen withdrawal therapies are still the most effective treatment for advanced disease, although it eventually progresses to the lethal castration-resistant stage. Evidence suggests that most cells in castration-resistant tumors retain androgen receptor (AR) that maintains its transcriptional activity through alternative pathways (1-7). Using the N-terminal domain of AR (NTD) as bait in the repressed transactivator yeast two-hybrid system (RTA), TATA binding protein associated factor 1 (TAF1) was detected (8). TAF1 is a multifunctional protein that contains acetylation (HAT), ubiquitin activating/conjugating (E1/E2) and bipartite, kinase domains consisting of N- and C-terminal kinases (NTK and CTK, respectively). TAF1 is capable of autophosphorylation as well as specific phosphorylation of TFIIF (9), p53 (10), and Mdm2 proto-oncogene (11). TAF1 binds and modulates transcriptional activity of proteins, such as c-Jun (12), the Mdm2 (13), and cyclin D1 (14) that they can also influence AR activity and hence prostate cancer progression (15-17). The focus of this study is to identify if and how TAF1 modulates AR transcription, and to determine the role of TAF1 in prostate cancer progression. In particular, this research proposal will test the hypothesis that TAF1 directly modulates AR activity, and aids in the development and maintenance of castration-resistant prostate cancer. We are testing this hypothesis with the following four specific aims:

- Specific Aim 1: To determine whether the TAF1/AR interaction specifically modulates AR transactivation
- Specific Aim 2: To map functional interaction site(s) of AR and TAF1
- Specific Aim 3: To determine the mechanisms through which TAF1 coactivates AR
- Specific Aim 4: To assess the expression profile of TAF1 versus AR in prostate tumors:
 - a) To characterize TAF1 and AR expression at mRNA and protein levels in the *Pten*^{-/-} prostate-specific knock out mouse model.
 - b) To determine the expression level of TAF1 and AR in malignant versus benign human prostate tissues.

PROGRESS AND RESULTS

A. Summary of last year report (specific aim 1)

We have demonstrated that TAF1 interacted with AR within nucleus of LNCaP cells, a prostate cancer cell line that expresses AR. Using ChIP assay, we also showed that TAF1 associated with AR at the prostate specific antigen (PSA) promoter in the presence of hormone. To assess if TAF1 modulates AR transcriptional activity, transactivation assays in prostate cancer cells were performed. TAF1 was either over-expressed (pHA-TAF1, kindly given by Dr. Tjian and Dr. Wong) or knocked down (siRNA duplexes; AAGACCCAAACAACCCCGCAT-3' and 5'-AACTACGACTACGCTCCACCA-3') in prostate cancer cells and luciferase assays were performed. The results indicated that the expression level of TAF1 is directly correlated to AR activity only in the presence of hormone, when the receptor has been activated (Fig. I, appendix).

Since TAF1 is a member of the general transcription machinery complex it is expected that it influences on promoters of other genes. In our transactivation assays, we also found that the non-androgenic reporter (thymidine kinase-renilla (pRLtk-renilla)) is also modulated by TAF1 in a similar manner as androgen responsive reporters (Fig. IIB, appendix). To differentiate the effect of TAF1 on AR from its general effects on transcription and to determine which TAF1 domains are specifically involved in AR activation, we cloned various domains of TAF1 and repeated the luciferase assay in LNCaP cells. In contrast to full-length TAF1, TAF1 domains had no effect on the generic renilla construct with pRL-tk promoter, implying that general transcription is not affected. By comparison, while HAT and CTK domains had no significant effect on AR activity, NTK and the E1/E2 domains of TAF1 did enhance AR activity in a ligand dependent manner. NTK significantly enhanced AR transactivation by 2.4-fold, which is almost as much as the full-length TAF1. However, E1/E2 domain had even greater effect, enhancing AR activity over 3.4-fold (Fig. II, appendix).

In 2008, the specific aims 2 and 3 were tested and the results are discussed as follows:

B. Specific Aim 2: To map functional interaction site(s) of AR and TAF1

TAF1 interacts with the N-terminus of AR mainly through its HAT and E1/E2 domains

To confirm the AR/TAF1 interaction and to determine the domains involved, GST pull-down assays were performed using GST-fusion protein with AR/NTD₁₋₅₅₉, DBD₅₄₁₋₆₆₅, or LBD₆₄₉₋₉₁₉ (Fig. 1). Purified GST-AR domains were analyzed by SDS-PAGE followed by Coomassie Blue staining (Fig. 1A). Equimolar amounts of non-degraded protein were determined for each of the GST protein products (*) and used in pull-down assays to assess relative TAF1 binding to AR domains, as described before (8). Radiolabeled TAF1 and TAF1 truncated domains were generated *in vitro* and verified by SDS-PAGE followed by autoradiography (Fig. 1B). The presence of multiple bands for radiolabeled proteins is due to several ATG sites within hTAF1 cDNA. Radiolabeled TAF1 (Fig. 1B, lane 1) was allowed to interact with GST protein or GST-AR fragments bound to agarose beads. As shown in Fig. 1C (top row), TAF1 did not interact with GST alone (lane 2), whereas it did bind to all domains of AR (lane 3-5) with the strongest interaction with the NTD of AR (lane 3).

GST pull-down experiments were repeated with radiolabeled TAF1 domains (NTK, HAT, E1/E2, and CTK proteins, Fig. 1B) to identify domains essential for interaction between TAF1 and AR (Fig. 1C, rows 2-4). All domains except the NTK of TAF1 bind to GST fusion AR proteins with different affinities. We performed the above pull-down assay with two more independent experiments and calculated an averaged percentage of total input bound to the GST-AR domains ($[\text{CNT}^* \text{ mm}^2] \text{ bound} / [\text{CNT}^* \text{ mm}^2] \text{ input}$), as described before (8). A summary of the quantified GST pull-down data (percentage of the input bound) for these TAF1 truncations and the full-length protein is shown in Fig. 1D. Similar to the full-length protein, HAT and E1/E2 domains interact most strongly with the NTD of AR, but the HAT domain binds this region with 1.5-times higher affinity. The CTK domain that was originally pulled out with the RTA assays with N-terminus of AR as a bait, also strongly interacts with DBD and LBD domains of AR. Taken together, these data demonstrate that TAF1 interacts directly with AR and mapping the interaction domains of TAF1 and AR suggests that the HAT, E1/E2 and CTK

domains of TAF1 are all involved in binding to AR. The pattern of binding of HAT and E1/E2 domains is similar to that seen with full length TAF1.

C. Specific Aim 3: To determine the mechanisms through which TAF1 coactivates AR

TAF1 and AR phosphorylation

To explore the molecular mechanism by which TAF1 enhances AR transcriptional activity, we tested the effect of apigenin, a TAF1 protein kinase inhibitor (10, 11), on AR transcriptional activity. Using MTS assay, we first found that apigenin at 5 μ M did not influence LNCaP cell viability (data not shown). This concentration was used to treat LNCaP cells transfected with the PSA-Luc reporter and luciferase assays were performed. As shown in Fig. 2A, AR activity is suppressed more than 4-fold in the presence of apigenin. To show whether TAF1 can phosphorylate AR, *in vitro* kinase activity was performed. We were able to show that TAF1 autophosphorylates itself and the DNA binding domain of AR can be phosphorylated with ERK/MAPK, consistent with others (9, 18, 19) (Fig. 2B). However, we could not detect AR phosphorylation by TAF1 (data not shown).

Androgen receptor is ubiquitinated by TAF1

Since NTK does not bind to AR and the CTK and HAT domains do not enhance AR transcriptional activity, we focused on the E1/E2 domain, which binds to AR and has the most profound effect on its transactivation. Ubiquitination is a posttranslational modification that mediates the covalent conjugation of ubiquitin to protein substrates. The functional role of ubiquitination was originally considered to be targeting proteins to the proteasome for degradation. However, it is now known that ubiquitination regulates many other processes in the cell, including membrane trafficking, DNA repair, and transcription (20). AR is also a direct target for mono and poly-ubiquitination (16, 21). To address whether TAF1 can ubiquitinate AR, LNCaP cells were co-transfected with pHis₆-Ubiquitin and either pHA-TAF1 or empty vector. Cells were then treated with 10% FBS RPMI followed by 6h treatment with vehicle or MG132, a proteasome inhibitor. To show the His-ubiquitin conjugated status of AR in the presence and absence of MG132, after saving 5% input (Fig. 3A), His-conjugated proteins were purified followed by a Western blot with AR and His antibodies. Fig. 3B shows that in the absence of MG132, there is no His-conjugated AR when TAF1 is not overexpressed (lane 1 & 5). Whereas, ubiquitinated AR is seen once TAF1 is overexpressed. (lane 2, arrow). As expected, in the presence of proteasome inhibitor, the total amount of conjugated AR is increased irrespective of TAF1 status level (lane 3, 4, 7 & 8). However, there is a drastic increase in the level of ubiquitinated AR once TAF1 is overexpressed (lane 4 & 8 versus 3 & 7). The results indicate that TAF1 facilitates AR ubiquitination in a spectrum from mono to poly-ubiquitin form.

TAF1 ubiquitinates AR *in vitro*

To confirm the ubiquitination of AR by TAF1 and to assess whether TAF1 is able to directly ubiquitinate AR, an in-tube ubiquitination assay was performed (22). As an abundant source of TAF1, HeLa cells were subjected to pull-down with anti-TAF1 antibodies and immobilized on beads. Protein extracts from LNCaP cells were incubated with radiolabeled ubiquitin, 10 nM ATP and TAF1 immobilized on agarose beads or beads alone at room temperature. After an hour, AR was immunoprecipitated and subjected to Western blot analysis. Fig. 3B shows that

AR can be ubiquitinated *in vitro* (out of cells) (lane 2) and that the enriched amount TAF1 enhances the amount of both mono and poly ubiquitinated forms AR (lane 3).

KEY RESEARCH ACCOMPLISHMENTS

1. TAF1 interacts with AR within prostate cancer cells and associates on the promoter of the PSA gene.
2. TAF1 binds to the N-terminal domain of AR. This interaction is mainly through HAT and E1/E2 domains of TAF1
3. The NTK and E1/E2 domains of TAF1 specifically enhance AR transcriptional activity and that mechanism is different from the role of TAF1 on general transcription machinery.
4. TAF1 can ubiquitinate AR inside a prostate cancer cell line and in test tubes.

REPORTABLE OUTCOME

1. A summary of the above data has been presented in the following meetings:
2. Poster presentation: The Endocrine Society's meeting 89th Annual Meeting, Toronto, Canada, June 2007
3. Oral presentation: The 6th Annual Canadian Prostate Cancer-BioNet Meeting, Montreal, Canada, April 22-24, 2007
4. Poster presentation: The Endocrine Society's meeting 90th Annual Meeting, San Francisco, CA, June 2008
5. Oral presentation: The urologic science meeting, Vancouver, Canada, June 12, 2008
6. Poster presentation: the 4th PacRim Breast and Prostate Cancer, Whistler, Canada, August 2008
7. Manuscript in preparation for the Molecular Endocrinology journal:
Tavassoli P., Wafa L. A., Cheng H., Zoubeidi A., Fazli L., Gleave M., Rennie P. S.
TATA binding protein-associated factor 1 (TAF1) can bind and differentially enhance androgen receptor transcriptional activity via its ubiquitin activating and conjugating domain

CONCLUSION

The molecular mechanisms responsible for the development of CR prostate cancer are largely unknown, but typically they do not appear to involve the loss of AR expression (23). The up-regulation of AR-target genes and over-expression of AR at the protein and mRNA levels support the notion that AR activity is altered in castration-resistant states (24-27). There are a variety of molecular alterations that could lead to continued or amplified AR signaling following surgical or medical castration. Recent evidence suggests that AR-specific gene regulation may

occur through interactions with unique coregulatory proteins. Since the N-terminus of AR (AR-NTD) is the least conserved, protein interactions in this region may dictate receptor-specific coregulation capacity. Using the RTA yeast two-hybrid system TAF1 was identified as a novel AR-NTD-interacting protein (8) and this interaction was confirmed with the full-length TAF1 using GST pull-down assays (Fig. 1C). Mapping of the TAF1 and AR interacting domains showed that HAT and E1/E2 domains bind strongly to AR-NTD mimicking the full length TAF1. The CTK domain that was originally isolated by the RTA system interacts with all AR domains but most strongly with the AR-DBD. In contrast, NTK does not have affinity for any AR domains, further indicating the specificity of these interactions (Fig. 1C & D). It has been reported by others that the N-terminus of TAF1 binds to the concave surface of TATA binding protein (TBP) and consequently inhibits TBP/ TATA box contact, thus repressing transcription (12, 28). However, binding of activators, such as c-Jun with the N-terminus of TAF1 releases this inhibition resulting in transcription initiation (29). Accordingly, the ability of TAF1 to interact with AR through multiple domains other than NTK suggests that TAF1 may play a role in modulating AR and one can speculate that upon interaction with AR, NTK release from the concave surface of TBP will initiate transcription. This hypothesis is supported by the fact that upon overexpression of TAF1 in both PC3 and LNCaP cells in the presence of nuclear AR (hormone induced activation), AR activity is increased (Fig. IA&B in appendix), whereas siRNA knock down of TAF1 suppresses AR activity (Fig. IC&D in appendix).

Co-immunoprecipitation assays of the nuclear extracts of LNCaP cells demonstrate that the interaction between AR and TAF1 is enhanced in the presence of hormone (R1881) (data was shown in last year report). Since TAF1 is a component of the general transcription machinery within the TFIID complex and directly associates with AR to modulate AR activity, we explored whether TAF1 co-localizes with AR at the PSA promoter. Using ChIP assays with LNCaP cells, we found that TAF1 is associated with an ARE in the proximal promoter of the PSA gene (data was shown in last year report). About 18% of genes of a hamster cell line are TAF1-dependent genes (30, 31). The essential nature of TAF1 can be attributed to its broad requirement during RNA-pol-II-dependent transcription (32). In our studies, we also found that TAF1 can modulate the transcription of non-androgen responsive reporters (Fig. IIB, appendix). To show that there is a specific effect of TAF1 on AR, various domains of TAF1 were tested for their effect on androgen and non-androgen responsive reporters in transient transfection assays (Fig. II, appendix). Although TAF1 is a component of the transcriptional machinery, it appears to be a specific coactivator of AR, enhancing AR transcription through its NTK and E1/E2 domains (Fig. IIA, appendix). In contrast, unlike the full-length TAF1, these truncated proteins did not modify generic promoters such as ptk-Renilla (Fig. IIB, appendix).

The AR is a substrate for phosphorylation, acetylation and ubiquitination (16, 21, 33-35). This makes AR as a potential substrate for TAF1 that possesses all the above enzymatic activities. We did not pursue to determine whether TAF1 can acetylate AR because no significant changes on AR activity were seen in transfection assays when the HAT domain was overexpressed (Fig. IIA). However, we have demonstrated that apigenin, a TAF1 kinase inhibitor (10, 11) can suppress AR transcriptional activity by 4-fold (Fig. 2A). To further explore the TAF1 kinase activity on AR, *in vitro* kinase assays were performed. Although we could show that TAF1 can autophosphorylates itself (Fig. 2B), but we were unable to show AR phosphorylation by TAF1. This could be due to inadequacy of cell-free assays, which are not representative of cell biology.

It has been reported that TAF1 can mono-ubiquitinate histone 1(H1) and poly-ubiquitinate p53 (10, 32). Since E1.E2 has the most profound effect on AR activity, we also sought to determine whether ubiquitination of AR can be increased as a consequence of TAF1 overexpression. Interestingly, in the presence of proteasome inhibitor (MG132) and overexpression of ubiquitin protein, full length-TAF1 enhances the total amount of ubiquitinated AR (Fig. 3A, lane 4 & 8). This was further validated using an *in vitro* ubiquitination assay (Fig. 3B, lane 3).

Taken together, we have demonstrated that TAF1 binds to the N-terminus of AR mainly through its HAT and E1/E2 domains and associates with AR in the nucleus and at the PSA promoter. In addition, we showed that TAF1 through its NTK and E1/E2 domains can affect AR transactivation without influencing the general transcription machinery. Finally, we have demonstrated that TAF1 is able to ubiquitinate AR both *in vivo* and *in vitro*. Currently, we are assessing the level of TAF1 expression in human prostate cancer samples using our gleason grade and neo-adjuvant tissue microarray (aim 4).





Figure 1. TAF1 binds AR through HAT and E1/E2 domains *in vitro*.

A. GST-fused AR domains (N-terminus =NTD, DNA binding domain =DBD, ligand binding domain =LBD) were expressed in *E. Coli* BL21 and purified using glutathione beads. Fusion protein-bound bead volumes were titrated, eluted with sample buffer and analyzed by SDS-PAGE followed by Coomassie Brilliant Blue staining. The eluent in each case was run alongside known amounts of BSA (ranging from 250 to 1000 ng) to generate a standard curve for protein concentrations. Equimolar amounts of non-degraded proteins (*) were used in GST pull-down assays. **B.** [35 S] Radiolabeled TAF1 and its domains (N-terminal kinase (NTK), Histone acetylation (HAT), Ubiquitin activating conjugating (E1/E2), and C-terminal kinase (CTK) were generated using *in vitro* Transcription/Translation kit. **C.** GST pull-down assay. Equivolume of [35 S] labeled TAF1, [35 S] labeled NTK, [35 S] labeled HAT, [35 S] labeled E1/E2, [35 S] labeled CTK were incubated with GST-AR fragments bound to agarose beads. GST alone coupled to agarose beads was used as negative control. **D.** Dried gels were also analyzed using a Phosphorimaging screen. Quantity One software was used to obtain data (counts/mm²) for radiolabelled protein bands. All pull-downs were done in triplicate, averaged, and normalized as a function of the percentage input bound.

A



B




Fig. 2: A) LNCaP cells were co-transfected with 0.7 μ g/well pPSA-Luc and 0.1 μ g/well pRLtk-renilla. Transfected cells were grown in the presence of 1 nM R1881 and either 5 μ M apigenin or vehicle for 24 h before harvesting for luciferase assay. Luciferase units were normalized to protein levels and fold-induction was calculated. All values are given as the mean ($\pm$ SEM) of triplicate readings. Graphs are representative of the 3 independent experiments. * indicates a $p < 0.05$ compared to empty vector control. B) HeLa cells were subjected to pulldown with normal IgG (lane 1) or anti-TAF1 (lane 2) antibodies immobilized on beads and incubated in kinase buffer (1M Hepes pH 7, 1M $MgCl_2$, 1M $MnCl_2$, 100 mM Na_3VO_4 , 10X microcystin) with the presence of [γ^{32} P]-ATP at 30°C for 20 min. Mock enzyme (lane 3) or ERK (lane 4) were incubated in kinase buffer with the presence of [γ^{32} P]-ATP and GST fusion of DBD of AR at 30°C for 20 min.




Figure 3. TAF1 ubiquitinates AR. LNCaP cells were transfected with 2 µg pHis₆-ubiquitin and either 6 µg of pHA-TAF1 or empty vector. Cells were then treated with 5% FBS RPMI followed by ±10 µM MG132 for 6h. After harvesting and lysing the cells in RIPA buffer, 5% of cell lysate was used as an input (A) and the remainder was mixed with 50 µl Ni²⁺-NTA-agarose beads. The mixture was rotated at 4°C for 3 h and then affinity pulled down followed by Western blot analysis for AR and His (B). The input was blotted for HA, AR and β-actin. C) TAF1 was immunoprecipitated from nuclear extracts of Hela cells. LNCaP cell lysate was incubated with 1 mM ATP, ubiquitin and either TAF1 immobilized on agarose beads (lane 3) or beads alone (lane 2) in 20 µl 0.4 M HEMG. After 1 h incubation at room temperature, AR was immunoprecipitated followed by a Western blot for AR. Lane 1 = Input (5%).

APPENDIX




Figure I. TAF1 modulates AR transactivation.

PC3 cells were co-transfected with 1.5 µg/well full length AR (pAR₆) and 0.2 µg/well pARR3-tk-Luc reporter, 0.1 µg/well pRLtk-renilla, and increasing amounts of pHA-TAF1 (**A**) or increasing amounts of TAF1 RNAi duplexes (**C**). LNCaP cells were co-transfected with 1 µg/well pPSA-Luc, 0.1 µg/well pRLtk-renilla and 1 µg/well pCS2+HA-hTAF1 (**B**) or 10 nM TAF1 RNAi duplexes (**D**). Transfected cells were growing in the presence or absence of 1 nM R1881 for 24 h (A, B) or 48 h (C & D) before harvesting for luciferase assay and Western blot analysis. Luciferase units (RLU) are expressed relative to protein values for each sample. All luciferase values are given as the mean (± SEM) of triplicate readings. Graphs are representative of the 3 independent experiments. * indicates a $p < 0.05$ compared to empty vector control. ** indicates a $p < 0.05$ compared to *.




Figure II. N-terminal kinase and ubiquitin activating/conjugating domains of TAF differentially enhance AR transactivation. LNCaP cells were co-transfected with either pHA-TAF1 or one of its four domains (pV5-NTK, pV5-HAT, V5-E1/E2, or pV5-CTK) (1 ug/well) and the pPSA-Luc and pRLtk-renilla. Transfected cells were growing in the presence or absence of 1 nM R1881 for 24 h before harvesting for luciferase assay. A) luciferase units were normalized to protein. B) fold-induction of renilla units were plotted against empty vevtor, TAF1 or its domains. * indicates a $p < 0.05$ compared to empty vector control. C) Western blot analysis for AR, β -actin, empty vector (EV) (1), TAF1 (2), NTK (3), HAT (4), E1/E2 (5) and CTK (6).

REFERENCES

1. **Blaszczyk N, Masri BA, Mawji NR, Ueda T, McAlinden G, Duncan CP, Bruchovsky N, Schweikert HU, Schnabel D, Jones EC, Sadar MD** 2004 Osteoblast-derived factors induce androgen-independent proliferation and expression of prostate-specific antigen in human prostate cancer cells. *Clin Cancer Res* 10:1860-1869
2. **Culig Z, Steiner H, Bartsch G, Hobisch A** 2005 Interleukin-6 regulation of prostate cancer cell growth. *J Cell Biochem* 95:497-505
3. **Degeorges A, Tatoud R, Fauvel-Lafeve F, Podgorniak MP, Millot G, de Cremoux P, Calvo F** 1996 Stromal cells from human benign prostate hyperplasia produce a growth-inhibitory factor for LNCaP prostate cancer cells, identified as interleukin-6. *Int J Cancer* 68:207-214
4. **Lu Y, Zhang J, Dai J, Dehne LA, Mizokami A, Yao Z, Keller ET** 2004 Osteoblasts induce prostate cancer proliferation and PSA expression through interleukin-6-mediated activation of the androgen receptor. *Clin Exp Metastasis* 21:399-408
5. **Wang MH, Abreu-Delgado Y, Young CY** 2003 Effects of vitamin C on androgen receptor mediated actions in human prostate adenocarcinoma cell line LAPC-4. *Urology* 62:167-171
6. **Nishimura K, Nonomura N, Satoh E, Harada Y, Nakayama M, Tokizane T, Fukui T, Ono Y, Inoue H, Shin M, Tsujimoto Y, Takayama H, Aozasa K, Okuyama A** 2001 Potential mechanism for the effects of dexamethasone on growth of androgen-independent prostate cancer. *J Natl Cancer Inst* 93:1739-1746
7. **Ueda T, Mawji NR, Bruchovsky N, Sadar MD** 2002 Ligand-independent activation of the androgen receptor by interleukin-6 and the role of steroid receptor coactivator-1 in prostate cancer cells. *J Biol Chem* 277:38087-38094
8. **Wafa LA, Cheng H, Rao MA, Nelson CC, Cox M, Hirst M, Sadowski I, Rennie PS** 2003 Isolation and identification of L-dopa decarboxylase as a protein that binds to and enhances transcriptional activity of the androgen receptor using the repressed transactivator yeast two-hybrid system. *Biochem J* 375:373-383
9. **Dikstein R, Ruppert S, Tjian R** 1996 TAFII250 is a bipartite protein kinase that phosphorylates the base transcription factor RAP74. *Cell* 84:781-790
10. **Li HH, Li AG, Sheppard HM, Liu X** 2004 Phosphorylation on Thr-55 by TAF1 mediates degradation of p53: a role for TAF1 in cell G1 progression. *Mol Cell* 13:867-878
11. **Allende-Vega N, McKenzie L, Meek D** 2008 Transcription factor TAFII250 phosphorylates the acidic domain of Mdm2 through recruitment of protein kinase CK2. *Mol Cell Biochem* 316:99-106
12. **Lively TN, Ferguson HA, Galasinski SK, Seto AG, Goodrich JA** 2001 c-Jun binds the N terminus of human TAF(II)250 to derepress RNA polymerase II transcription in vitro. *J Biol Chem* 276:25582-25588
13. **Leveillard T, Wasylyk B** 1997 The MDM2 C-terminal region binds to TAFII250 and is required for MDM2 regulation of the cyclin A promoter. *J Biol Chem* 272:30651-30661
14. **Siegert JL, Rushton JJ, Sellers WR, Kaelin WG, Jr., Robbins PD** 2000 Cyclin D1 suppresses retinoblastoma protein-mediated inhibition of TAFII250 kinase activity. *Oncogene* 19:5703-5711
15. **Cai C, Hsieh CL, Shemshedini L** 2007 c-Jun has multiple enhancing activities in the novel cross talk between the androgen receptor and Ets variant gene 1 in prostate cancer. *Mol Cancer Res* 5:725-735
16. **Gaughan L, Logan IR, Neal DE, Robson CN** 2005 Regulation of androgen receptor and histone deacetylase 1 by Mdm2-mediated ubiquitylation. *Nucleic Acids Res* 33:13-26
17. **Narayanan BA, Narayanan NK, Davis L, Nargi D** 2006 RNA interference-mediated cyclooxygenase-2 inhibition prevents prostate cancer cell growth and induces differentiation: modulation of neuronal protein synaptophysin, cyclin D1, and androgen receptor. *Mol Cancer Ther* 5:1117-1125

18. **Azzam DG, Tay JW, Greeve MA, Harvey JM, Bentel JM** 2008 ERK/MAPK regulation of the androgen responsiveness of breast cancer cells. *Adv Exp Med Biol* 617:429-435
19. **Gioeli D, Black BE, Gordon V, Spencer A, Kesler CT, Eblen ST, Paschal BM, Weber MJ** 2006 Stress kinase signaling regulates androgen receptor phosphorylation, transcription, and localization. *Mol Endocrinol* 20:503-515
20. **Miranda M, Sorkin A** 2007 Regulation of receptors and transporters by ubiquitination: new insights into surprisingly similar mechanisms. *Mol Interv* 7:157-167
21. **Burgdorf S, Leister P, Scheidtmann KH** 2004 TSG101 interacts with apoptosis-antagonizing transcription factor and enhances androgen receptor-mediated transcription by promoting its monoubiquitination. *J Biol Chem* 279:17524-17534
22. **Belz T, Pham AD, Beisel C, Anders N, Bogin J, Kwozynski S, Sauer F** 2002 In vitro assays to study protein ubiquitination in transcription. *Methods* 26:233-244
23. **Taplin ME, Balk SP** 2004 Androgen receptor: a key molecule in the progression of prostate cancer to hormone independence. *J Cell Biochem* 91:483-490.
24. **Chen CD, Welsbie DS, Tran C, Baek SH, Chen R, Vessella R, Rosenfeld MG, Sawyers CL** 2004 Molecular determinants of resistance to antiandrogen therapy. *Nat Med* 10:33-39
25. **Rennie PS, Nelson CC** 1998 Epigenetic mechanisms for progression of prostate cancer. *Cancer Metastasis Rev* 17:401-409
26. **Sirotnak FM, She Y, Khokhar NZ, Hayes P, Gerald W, Scher HI** 2004 Microarray analysis of prostate cancer progression to reduced androgen dependence: studies in unique models contrasts early and late molecular events. *Mol Carcinog* 41:150-163
27. **Tilley WD, Lim-Tio SS, Horsfall DJ, Aspinall JO, Marshall VR, Skinner JM** 1994 Detection of discrete androgen receptor epitopes in prostate cancer by immunostaining: measurement by color video image analysis. *Cancer Res* 54:4096-4102
28. **Kokubo T, Swanson MJ, Nishikawa JI, Hinnebusch AG, Nakatani Y** 1998 The yeast TAF145 inhibitory domain and TFIIA competitively bind to TATA-binding protein. *Mol Cell Biol* 18:1003-1012
29. **Lively TN, Nguyen TN, Galasinski SK, Goodrich JA** 2004 The basic leucine zipper domain of c-Jun functions in transcriptional activation through interaction with the N terminus of human TATA-binding protein-associated factor-1 (human TAF(II)250). *J Biol Chem* 279:26257-26265
30. **Irvin JD, Pugh BF** 2006 Genome-wide transcriptional dependence on TAF1 functional domains. *J Biol Chem* 281:6404-6412
31. **O'Brien T, Tjian R** 2000 Different functional domains of TAFII250 modulate expression of distinct subsets of mammalian genes. *Proc Natl Acad Sci U S A* 97:2456-2461
32. **Wassarman DA, Sauer F** 2001 TAF(II)250: a transcription toolbox. *Journal of cell science* 114:2895-2902
33. **Fu M, Wang C, Reutens AT, Wang J, Angeletti RH, Siconolfi-Baez L, Ogryzko V, Avantiaggiati ML, Pestell RG** 2000 p300 and p300/cAMP-response element-binding protein-associated factor acetylate the androgen receptor at sites governing hormone-dependent transactivation. *J Biol Chem* 275:20853-20860
34. **Zhou ZX, Kemppainen JA, Wilson EM** 1995 Identification of three proline-directed phosphorylation sites in the human androgen receptor. *Mol Endocrinol* 9:605-615
35. **Zhu Z, Becklin RR, Desiderio DM, Dalton JT** 2001 Identification of a novel phosphorylation site in human androgen receptor by mass spectrometry. *Biochem Biophys Res Commun* 284:836-844