

SCIENTIFIC NOTE

HABITATS AND DISTRIBUTION OF *ANOPHELES SINENSIS* AND ASSOCIATED *ANOPHELES* HYRCANUS GROUP IN JAPAN

LEOPOLDO M. RUEDA,¹ MASASHIRO IWAKAMI,^{2,6} MONICA O'GUINN,^{2,6} MOTYOYOSHI MOGI,³
BRIAN F. PRENDERGAST,^{4,7} ICHIRO MIYAGI,³ TAKAKO TOMA,³ JAMES E. PECOR¹ AND
RICHARD C. WILKERSON¹

ABSTRACT. Mosquito collections were carried out in August 2002 and July 2003 in Japan. *Anopheles sinensis* of the Hyrcanus Group, Myzomyia Series of *Anopheles*, was the most common species encountered. The distribution and habitats of 5 *Anopheles* Hyrcanus Group species are included. Eight species belonging to *Aedes*, *Culex*, and *Uranotaenia* were found associated with *An. sinensis* in rice paddies and a variety of other larval habitats.

KEY WORDS Mosquitoes, *Anopheles sinensis*, Hyrcanus Group, *Aedes*, *Culex*, *Uranotaenia*, Japan, Culicidae

The *Anopheles* Hyrcanus Group consists of several species that are vectors of malaria and other mosquito-borne diseases in the Oriental and Palearctic regions. The group, as currently defined, includes about three quarters of the species that comprise the Myzorhynchus Series of genus *Anopheles* subgenus *Anopheles*. The group has about 30 species, with the addition of 2 newly described species (Li et al. 2005, Rueda 2005) in the Oriental and Palearctic regions.

Anopheles hyrcanus (Pallas 1771) was initially recorded (as *Culex*) from the southern shores of the Caspian Sea. It was shown to have a wide distribution from Europe to east and southeastern Asia, including islands in the Indian and Pacific Oceans. It became apparent that the taxon included a number of morphologically distinct forms and that some were involved in the transmission of malarial and filarial parasites in the Oriental and Eastern Palearctic regions. Subsequently, about 30 species have been described and named (Ramsdale 2001, Harbach 2004, Rueda 2005). Harbach (2004) listed 27 species in the Hyrcanus Group, with 6 species placed in the *Lesteri* Subgroup, 4 in the *Nigerrimus*

Subgroup, and 17 in the unassigned subgroup. He omitted *An. yatsushiroensis* Miyazaki and *An. albotaeniatus* (Theobald) from his Hyrcanus Group list, with the latter being transferred to the *Albotaeniatus* Group and the former currently a synonym of *An. pullus* M. Yamada (Shin and Hong 2001, Hwang et al. 2004). Twenty-seven species of the Hyrcanus Group are found in the Oriental and Eastern Palearctic regions, and only 3 species have a western Palearctic distribution (from western China and south of 50° N) (Ramsdale 2001). In this paper, we treated *An. yatsushiroensis* as a valid species because the genetic comparison to type specimens from the type locality of *An. yatsushiroensis*, Yatsushiro City, Japan, is still needed to rule out the possibility that this is a valid species.

In Japan, there are only 5 known species of the Hyrcanus Group: *An. sinensis* Wiedemann, *An. sineroides* Yamada, *An. engarensis* Kanda and Oguma, *An. lesteri* Baisas and Hu, and *An. yatsushiroensis* (Tanaka et al. 1979). The type locality of each species is listed in Table 1.

Anopheles sinensis is the most common anopheline species in Japan, including Ryukyu Islands (Tanaka et al. 1979). It has long been suspected that it is the most important vector of malaria in Japan, including Okinawa and Hokkaido. Even though indigenous malaria has disappeared, this vector remains abundant throughout Japan. It is a known vector of malaria in Korea and China, and it has a wide distribution in Asia (Harrison and Scanlon 1975, Tanaka et al. 1979, Rueda et al. 2005a). *Anopheles lesteri* is a very important vector of malaria in China. Based on a combination of published and newly generated rDNA ITS2 sequences, Wilkerson et al. (2003) found that *An. lesteri* from the Philippines (type locality of true *lesteri*) and Korea and so called *An. anthropophagus* from China are synonymous, with *An. anthropophagus* as the junior synonym. To clarify and stabilize the taxon, Rueda et al. (2005b) designated and described the

¹ Walter Reed Biosystematics Unit, Department of Entomology, Walter Reed Army Institute of Research, 503 Robert Grant Avenue, Silver Spring, MD 20910.

² Center for Human Health and Preventive Medicine-Pacific, Entomology Program, CP Zama Building 715, Room 242, Tokyo APO, AP 96343.

³ Department of Pathology and Biodefense, Faculty of Medicine, Saga University, Nabeshima 5-1-1, Saga 849-8501, Japan.

⁴ United States Marine Corps, 3rd Medical Battalion, Unit 38447, Okinawa FPO, AP 96604-8447.

⁵ University of Ryukyus, Health Science School, 207 Uehara Nishihara, Okinawa 903-0215, Japan.

⁶ Present address: MCMR-UIV, Building 1425, Room A, Fort Detrick, MD 21702.

⁷ Present address: Navy Environmental and Preventive Medicine Unit 2, 1887 Powhatan Street, Norfolk, VA 23511.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2005		2. REPORT TYPE		3. DATES COVERED 00-00-2005 to 00-00-2005	
4. TITLE AND SUBTITLE Habitats and Distribution of Anopheles Sinensis and Associated Anopheles Hyrcanus Group in Japan				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Walter Reed Army Institute of Research, Department of Entomology, Silver Spring, MD, 20910				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT see report					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			
			Same as Report (SAR)	6	

Table 1. Type locality of *Anopheles* Hyrcanus Group species in Japan.

Species	Type locality	Reference
<i>An. sinensis</i>	China: Guangzhou	Wiedemann 1828
<i>An. engarensis</i>	Japan: Hokkaido (Engaru)	Kanda and Oguma 1978
<i>An. yatsushiroensis</i>	Japan: Kyushu (Yatsushiro)	Miyazaki 1951
<i>An. sineroides</i>	Japan: Hokkaido (Bibai)	Yamada 1924
<i>An. lesteri</i>	Philippines: Manila (Sta. Mesa)	Baisas and Hu 1936
	Laguna (Calauan)—neotype	Rueda et al. 2005b

neotype and alloneotype of *An. lesteri*. This species was suspected as an important vector of indigenous malaria in Japan, but particularly in Hokkaido, where it commonly occurs in great numbers. It is also common in the Ryukyus Islands and has been found more frequently in coastal regions than inland in Honshu and Kyushu (Tanaka et al. 1979). Shin and Hong (2001) and Hwang et al. (2004) considered *An. yatsushiroensis* as a synonym of *An. pullus* based on molecular and morphological data of Korean specimens. We did not collect *An. pullus* during our collections in 2002 and 2003 in Japan,

and no report indicates the existence of *An. pullus* in that country. However, because the type locality of *An. yatsushiroensis* is in Japan, it is necessary to do a genetic comparison of *An. pullus* from Korea with the topotypic specimens from Japan to resolve definitively if the two are synonyms or not. *Anopheles yatsushiroensis* is not known as a vector of indigenous malaria in Japan. *Anopheles pullus* is considered a potential vector of *vivax* malaria in the Korean Peninsula. The other Hyrcanus Group species, *An. sineroides* and *An. engarensis*, are not known vectors of indigenous malaria in Japan.




Fig. 1. Map of Japan showing the distribution of *Anopheles* Hyrcanus Group (based on observed and published specimens). (A) Hokkaido district. (B–F) Honshu. (G) Shikoku district. (H) Kyushu district. (I) Amami Guntō. (J) Okinawa Guntō. (K) Miyako and Yaeyama Guntō.

Table 2. Distribution of *Anopheles Hyrcanus* Group species in Japan (based on observed and published specimens).

Location	<i>An. engarensis</i>	<i>An. lesteri</i>	<i>An. sineroides</i>	<i>An. sinensis</i>	<i>An. yatsu- shiroensis</i>
Amami Oshima Is.: Akagina				6 ¹	
Amami Oshima Is.: Koniya				1	
Amami Oshima Is.: Nishinakama		6		6	
Amami Oshima Is.: Yuwan		6, 7		6	
Hokkaido: Bibai			6	7	
Hokkaido: Engaru	6, 7	7	6, 7		
Hokkaido: Hakodate		1			
Hokkaido: Nagayama				7	
Hokkaido: Rubeshibe	6, 7		6	7	
Hokkaido: Sakura-oka			7	7	
Hokkaido: Sapporo	6				
Honshu: Aomori Pref., Sukayu			6, 7		
Honshu: Chiba Pref., Imba-Numa					2, 5
Honshu: Chiba Pref., Shirahama				1	
Honshu: Chiba Pref., Tateyama			1		
Honshu: Chiba Pref., Teganuma				6	
Honshu: Ishikawa Pref., Kamiosaki				1	
Honshu: Ishikawa Pref., Wajima				1	
Honshu: Kanagawa Pref., Atsugi			6, 7		
Honshu: Kanagawa Pref., Ayase			6		
Honshu: Kanagawa Pref., Hiratsuka				7	
Honshu: Kanagawa Pref., Nakatsu Valley				6	
Honshu: Kanagawa Pref., Sagami-hara			6, 7	6	
Honshu: Kanagawa Pref., Toya			6, 7		
Honshu: Kanagawa Pref., Yamato			6, 7	6	
Honshu: Kanagawa Pref., Zama			6	6	
Honshu: Kyoto Pref., Hikone					5
Honshu: Kyoto Pref., Ogura					4, 5
Honshu: Nagano Pref., Norikura-kogen				6	
Honshu: Okoyama Pref., Kurashiki				6	
Honshu: Okoyama Pref., Utoma		6		6	
Honshu: Shizouka Pref., Gotenba			1	1	
Honshu: Shizouka Pref., Misakubo				6	
Honshu: Shizouka Pref., Shimoda			1		
Honshu: Tokyo Pref., Hachioji			6	6	
Honshu: Tokyo Pref., Kamiosaki				1	
Honshu: Tokyo Pref., Seroganeshizenen				1	
Honshu: Tokyo Pref., Shakuji				1	
Honshu: Tokyo Pref., Tachikawa				6	
Honshu: Tokyo Pref., Ueno			6	6	
Irimote Is.: Itokawa-rindo				6	
Irimote Is.: Mt. Bama		6			
Irimote Is.: Ohara				6	
Irimote Is.: Uehara		6			
Ishigaki Is., Inoda				6	
Ishigaki Is.: Miyara Riv.		6			
Ishigaki Is.: Mt. Maeshi				6	
Ishigaki Is.: Mt. Omoto		6		6	
Ishigaki Is.: Nagura				6	
Ishigaki Is.: near Kabira				6	
Ishigaki Is.: Yashigawa		6			
Ishigaki Is.: Yoshiwara				6	
Kyushu: Fukuoka Pref., Yanagawa					3
Kyushu: Kagoshima Pref., Kagoshima				6, 7	
Kyushu: Kagoshima Pref., Shiroyama				6	
Kyushu: Kagoshima Pref., Taniyama				6	
Kyushu: Kumamoto Pref., Minamata					3
Kyushu: Kumamoto Pref., Nishisato				7	
Kyushu: Kumamoto Pref., Tamana				6, 7	
Kyushu: Kumamoto Pref., Ueki				7	
Kyushu: Kumamoto Pref., Uto				6	
Kyushu: Kumamoto Pref., Yatsushiro				6, 7	3

Table 2. Continued.

Location	<i>An. engarensis</i>	<i>An. lesteri</i>	<i>An. sineroides</i>	<i>An. sinensis</i>	<i>An. yatsushiroensis</i>
Kyushu: Nagasaki Pref., Aino				6	
Kyushu: Nagasaki Pref., Fukushima Is.				7	
Kyushu: Nagasaki Pref., Isahaya				6	
Kyushu: Saga Pref., Ahikari				7	
Kyushu: Saga Pref., Saga and south of Saga				7	3
Kyushu: Saga Pref., Tosu				6	
Kyushu: Wakayama, Kushimoto				1	
Okinawa Is.: Ginowan		6			
Okinawa Is.: Igei				7	
Okinawa Is.: Kena Dam				7	
Okinawa Is.: Kin				7	
Okinawa Is.: Ojina		6, 7			
Okinawa Is.: Toguchi				6	
Okinawa Is.: Yona				6	
Shikoku: Kochi, Ashizuri-misaki				6	
Tsushima Is.: Asamo				6	
Tsushima Is.: Kuwa				6	
Tsushima Is.: Nain				6	
Yonaguni Is.: near Sonai		6		6	

¹ 1 = Kurihara (1999); 2 = Miyake (1950); 3 = Miyazaki (1951); 4 = Otsuru and Konoe (1951); 5 = Otsuru and Ohmori (1960); 6 = Tanaka et al. (1979); 7 = observed specimens (museum and field collected).

We conducted 40 mosquito collections in Japan in August 2002 and July 2003, resulting in 223 individually reared pinned adults, 138 individually reared adults preserved in 100% ethyl alcohol for molecular tests, and 351 with exuviae of larvae and pupae and whole larvae. In addition, 3 type localities in Japan were visited to collect mosquitoes in 2003, namely Yatsushiro for *An. yatsushiroensis*, Bibai for *An. sineroides*, and Engaru for *An. engarensis*. In 2002, only the type locality in Yatsushiro was visited for collection. We initially targeted larval collections from those habitats where mosquitoes in the genus *Anopheles* were likely to be found. Larvae and pupae from various habitats were collected using dippers and individually reared to the adult stage (Walter Reed Biosystematics Unit 2001). Larval and pupal skins were preserved in 80% ethyl alcohol and slide mounted using standard protocols. Emerged adults were

pinned, provided with appropriate collection data, and identified. Some adults and whole larvae were separately preserved in 100% ethyl alcohol for molecular analysis. Adult specimens and associated larval/pupal exuviae were identified using information from Tanaka et al. (1979). Voucher specimens are deposited in the National Museum of Natural History, Smithsonian Institution, Washington, DC.

Specimens of *Anopheles* Hyrcanus Group species deposited at various museums were examined, including those in the Entomology Museum of the National Institute of Infectious Diseases, Tokyo, Japan; University of Hokkaido, Insect Museum, Sapporo, Japan; and the National Museum of Natural History, Smithsonian Museum, Washington, DC. To further confirm the identification of 2 *Anopheles* species (i.e., *An. sinensis* and *An. sineroides*), DNA was isolated from individual adult mosquitoes by

Table 3. Habitats of *Anopheles* Hyrcanus Group species in Japan and other Asian countries.

Species	Larval habitat ¹	Locality	Reference
<i>An. sinensis</i>	RP, SP ID, MA, SW	Japan Thailand	Tanaka et al. 1979; ² Harrison and Scanlon et al. 1975
<i>An. engarensis</i>	CR, GP, PO, RP, GP	Japan	Tanaka et al. 1979
<i>An. yatsushiroensis</i>	CR, ID, RP	Japan	Tanaka et al. 1979
<i>An. sineroides</i>	GP, PO, RD, SM RH	Japan Korea	Tanaka et al. 1979; ³ Tanaka et al. 1979
<i>An. lesteri</i> ³	GP, IW, MA, PO, RP, SM HD	Japan Philippines	Tanaka et al. 1979; ² Rueda et al. 2005b ³

¹ CR, creek; GP, ground pool; HD, hill ditch; ID, irrigation ditch; IW, impounded water; MA, marsh; PO, pond; RD, road and other ditches; RH, rock hole; RP, rice paddy; SM, stream margin; SP, stream pool; SW, swamp.

² Observed museum and field-collected specimens.

³ Prefer cooler and shaded areas unlike *An. sinensis*, and more common in coastal areas than inland (Tanaka et al. 1979).

phenol-chloroform extraction as described in Wilkerson et al. (1993). Direct sequencing of ribosomal DNA ITS2 was carried out, and the results were compared with those of Wilkerson et al. (2003).

Figure 1 and Table 2 show the distribution of 5 *Anopheles* Hyrcanus Group species, based on our field-collected data, museum specimen data, and published information (Miyake 1950, Miyazaki 1951, Otsuru and Konoe 1951, Otsuru and Ohmori 1960, Tanaka et al. 1979, Kurihara 1999). *Anopheles sinensis* occurs almost all throughout Japan (Honshu, Shikoku, and Kyushu districts and Rukyu Archipelago). *Anopheles lesteri* is found from the southern areas (Ryukyu Archipelago) to the northern areas (e.g., Hokkaido). *Anopheles yatsushiroensis* is known to occur primarily in the Honshu and Kyushu districts. *Anopheles engarensis* is confined to Hokkaido, whereas *An. sineroides* occurs in Hokkaido and other areas of Japan.

Table 3 shows the habitats of *An. sinensis*, *An. sineroides*, *An. engarensis*, *An. lesteri*, and *An. yatsushiroensis* at various areas of Japan based on our field-collected data, museum specimen data, and published information. The larvae of all species, except *An. sineroides*, are found in rice paddies. During our 2003 survey, we found that the mean water pH (6.8), conductivity (0.17 μ S), and salinity (0.07 ppm) of the larval/pupal habitats of *An. sinensis* and associated species in rice paddies were greater compared with the irrigation ditches (6.5, 0.07, and 0.03, respectively). The mean water temperature (27.5°C) of these rice paddy habitats was lower compared with the irrigation ditches (30.5°C). We found the larvae of *An. sinensis* associated in the rice paddies with *Culex* (*Culex*) *tritaeniorhynchus* Giles at Saga (Saga Prefecture) and Yatsushiro (Kumamoto Prefecture), in the rice paddies with *Cx.* (*Cux.*) *bitaeniorhynchus* Giles at Nishisato (Kumamoto Prefecture), and in the taro or "rotas" irrigated fields with *Cx.* (*Lophoceraomyia*) *infantulus* Edwards, *Cx.* (*Cux.*) *tritaneorhynchus*, *Cx.* (*Lutzia*) *fuscanus* Wiedemann, and *Uranotaenia* (*Uranotaenia*) *macfarlanei* Edwards at Kin, Okinawa. We also collected from other larval habitats where *Anopheles* species were absent and found the following species: *Aedes* (*Aedimorphus*) *vexans nipponii* (Theobald) in rice paddies at Kin, Okinawa; *Cx.* (*Cux.*) *tritaneorhynchus* and *Cx.* (*Eumelanomyia*) *hayashii ryukyuanus* Tanaka, Mizusawa, and Saugstad in creeks or ditches at Camp Gonsalves, Okinawa; *Cx.* (*Eumelanomyia*) *hayashii* Yamada in rice paddies at Asahikawa (Hokkaido); *Cx.* (*Cux.*) *pipiens pallens* Coquillett and *Cx.* (*Lop.*) *infantulus* in stream margins at Sapporo (Hokkaido); and *Cx.* (*Cux.*) *mimeticus* Noe in rice paddies at Futoni and Bibai (Hokkaido).

This paper presents the habitats and distribution of the 5 species of Hyrcanus Group of *Anopheles* in Japan based on specimens collected in 2002 and 2003, museum specimens, and available references. Our work provides readily available information to

preventive medicine and public health personnel who deal with malaria vector problems if sporadic malaria cases appear in the future.

We thank Masaaki Suwa and Kazunori Yoshizawa of Hokkaido University, Sapporo, for allowing L. M. Rueda to examine mosquito specimens; Takeshi Kurihara, Hiromu Kurahashi, Kyoko Sawabe, and Akihiko Shinohara for their kindness, hospitality, and help to L. M. Rueda when he examined specimens in the Entomology Museum, National Institute of Infectious Diseases, Tokyo; Noboru Minakawa, Saga University, Saga, for help with collecting specimens at Saga; Cong Li for DNA analysis; and Tom Anderson for preparing the map. This research was performed under a Memorandum of Understanding between the Walter Reed Army Institute of Research and the Smithsonian Institution, with institutional support provided by both organizations. The opinions and assertions contained herein are those of the authors and are not to be construed as official or reflecting the views of the Department of the Army or the Department of Defense.

REFERENCES CITED

- Baisas FE, Hu SMK. 1936. *Anopheles hyrcanus* var. *sinensis* of the Philippines and certain parts of China, with some comments on *Anopheles hyrcanus* var. *nigerrimus* of the Philippines. Manila, Philippines: *Monthly Bull Bureau of Health* 16:205-241.
- Harbach RE. 2004. The classification of genus *Anopheles* (Diptera: Culicidae): a working hypothesis of phylogenetic relationships. *Bull Entomol Res* 94:537-553.
- Harrison BA, Scanlon JE. 1975. Medical entomology studies—II. The subgenus *Anopheles* in Thailand (Diptera: Culicidae). *Contrib Am Entomol Inst (Ann Arbor)* 12:1-307.
- Hwang UW, Yong TS, Ree HI. 2004. Molecular evidence for synonymy of *Anopheles yatsushiroensis* and *An. pullus*. *J Am Mosq Control Assoc* 20:99-104.
- Kanda T, Oguma M. 1978. *Anopheles engarensis*, new species related to *sinensis* from Hokkaido, Island, Japan. *Mosq Syst* 10:45-52.
- Kurihara T. 1999. Family Culicidae (Insecta: Diptera)/mosquitoes. *Database Rec Entomol Collect* 1:1-34.
- Li C, Lee JS, Groebner JL, Kim HC, Klein TA, O'Guinn ML, Wilkerson RC. 2005. A newly recognized species in the *Anopheles* Hyrcanus Group and molecular identification of related species from the Republic of South Korea (Diptera: Culicidae). *Zootaxa* 939:1-8.
- Miyake M. 1950. Study on the anopheline mosquitoes in Kyushu. *Hukuoka Acta Med* 41:918-927.
- Miyazaki I. 1951. On a new anopheline mosquito *Anopheles yatsushiroensis* n. sp. found in Kyushu, with some remarks on two related species of the genus. *Kyushu Mem Med Sci* 2:195-206.
- Otsuru M, Konoe I. 1951. On the distribution of the *Anopheles* called "Yatsushiro type" in Ogura area, the suburbs of Kyoto city. *Med Biol* 19:194-196.
- Otsuru M, Ohmori Y. 1960. Malaria studies in Japan after World War II, part II. The search for *Anopheles sinensis* sibling species group. *Japan J Exp Med* 30:33-65.
- Pallas PS. 1771. *Reise durch verschiedene provinzen des*

- russischen reichs. Vol. 1. Graz, Austria: Akademische Druck-u Verlagsanstalt.
- Ramsdale CD. 2001. Internal taxonomy of the Hyrcanus group of *Anopheles* (Diptera: Culicidae) and its bearing on the incrimination of vectors of malaria in the west of the Palearctic Region. *Eur Mosq Bull* 10:1–8.
- Rueda LM. 2005. Two new species of *Anopheles* (*Anopheles*) Hyrcanus Group (Diptera: Culicidae) from the Republic of South Korea. *Zootaxa* 941:1–26.
- Rueda LM, Ma Y, Song GH, Gao Q. 2005a. Notes on the distribution of *Anopheles* (*Anopheles*) *sinensis* Wiedemann (Diptera: Culicidae) in China and the status of some *Anopheles* Hyrcanus Group type specimens from China. *Proc Entomol Soc Wash* 107:235–238.
- Rueda LM, Wilkerson RC, Li C. 2005b. *Anopheles* (*Anopheles*) *lesteri* Baisas and Hu (Diptera: Culicidae): neotype designation and description. *Proc Entomol Soc Wash* 107:604–622.
- Shin EH, Hong HK. 2001. A new synonym of *Anopheles* (*Anopheles*) *pullus* Yamada, 1937: *A. (A.) yatsushiroensis* Miyazaki, 1951. *Korean J Entomol* 31:1–5.
- Tanaka K, Mizusawa K, Saugstad E. 1979. A revision of the adult and larval mosquitoes of Japan (including the Ryukyu Archipelago and the Ogasawara Islands) and Korea (Diptera: Culicidae). *Contrib Am Entomol Inst (Ann Arbor)* 16:1–987.
- Walter Reed Biosystematics Unit. 2001. *Laboratory and field protocols*. Washington, DC: Walter Reed Biosystematics Unit, Smithsonian Institution.
- Wiedemann CRG. 1828. *Aussereuropäische zweiflügelige Insekten*. Vol. 1. Hamm, Germany: Erster Theil Schulz.
- Wilkerson RC, Li C, Rueda LM, Kim HC, Klein TA, Song GH, Strickman D. 2003. Molecular confirmation of *Anopheles* (*Anopheles*) *lesteri* from the Republic of South Korea and its genetic identity with *An. (Ano.) anthropophagus* from China (Diptera: Culicidae). *Zootaxa* 378:1–14.
- Yamada S. 1924. A revision of the adult anopheline mosquitoes of Japan: systematic descriptions, their habits and their relations to human diseases, together with an account of three new species. *Sci Rep Gov Inst Infect Dis Tokyo* 3:215–241.