

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2008		2. REPORT TYPE		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE Development of Turbulent Biological Closure Parameterizations				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Massachusetts Dartmouth,School for Marine Science and Technology (SMAST),706 South Rodney French Blvd,New Bedford,MA,02744				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Development of Turbulent Biological Closure Parameterizations

Louis Goodman

School for Marine Science and Technology (SMAST)

University of Massachusetts Dartmouth

706 South Rodney French Blvd

New Bedford, MA 02744

phone: (508) 910-6375 fax: (508) 910-6376 email: lgoodman@umassd.edu

Grant Number: N000140810631

<http://www.smast.umassd.edu/Turbulence/>

LONG-TERM GOAL

The long-term goal of this project is to understand and represent the dynamical role of planktonic fluctuations induced by physical turbulence. Additionally, this work is intended to be the basis of an efficient representation and methodology which will allow their inclusion in real ocean mean field dynamical models.

OBJECTIVES

- I. Extend Goodman and Robinson (2007) statistically based theory to include a greater variety of fundamental biophysical dynamical processes;
- II. Development of turbulent subgrid scale parameterizations for non linear biophysical interaction;
- III. Application of the PDF theory and the parameterized Advective Diffusive Reaction (ADR) model to oceanographically relevant processes.


APPROACH AND CONTEXT

A nonlinear model for biological and physical dynamical interactions in a laminar upwelling flow field Robinson(1999) has been extended to turbulent flow (Goodman and Robinson, 2007). Lagrangian and Eulerian probability density functions have been derived and applied to obtain the statistics of the biological state variables and mean dynamical effects of fluctuation correlations. This approach has been shown to lead to closed form solutions for the ensemble averaged biological state variables; no additional assumption is required to close the biodynamical interaction terms. The nonlinear uptake of Nutrient (N) by Phytoplankton (P) in a linear strain upwelling field and random walk turbulence has been explored in detail. It has been shown that the mean P , N fields can be obtained in terms of α , the ratio of mean advection to rate of uptake of nutrients, and Pe , (Peclet number), the ratio of time scale of turbulent diffusion to that of mean advection. The mean net primary production was found to be equal in the laminar limit of $Pe = \infty$ and strong turbulence limit of $Pe = 0$. For the advection parameter $\alpha = .1$, characteristic of a typical ocean near surface layer, mean net primary production was found to exhibit a maximum at some intermediate value of Pe . The mean total production $\langle PN \rangle$ is composed of two terms, $\langle P \rangle \langle N \rangle$ and $\langle P'N' \rangle$ which are opposite in sign resulting in $\langle PN \rangle$ being much smaller than each of the two contributing terms.

WORK COMPLETED

Many fundamental biological dynamical processes in the sea are strongly non-linear. Thus, physical turbulence in the sea not only enhances advective transport processes by eddy diffusivity but also importantly induces turbulent fluctuations in biological state variables which can correlate in the mean and effect mean biological dynamics. Goodman and Robinson (Proc. Roy. Soc A, 2008, 484, 555-572; doi: 10.1098/rspa.2007.0251) have formulated a theory of turbulent biological-physical interactions in terms of probability density functions (pdfs) and applied it to the simple example of nutrient (N) and seed phytoplankton (P) upwelling into a surface turbulent layer. The mean uptake $\langle PN \rangle$ is the sum of the product of the means $\langle P \rangle \langle N \rangle$ and the often neglected correlation fluctuations $\langle P'N' \rangle$. $\langle PN \rangle$ and $\langle P \rangle \langle N \rangle$ are positive, but $\langle P'N' \rangle$ is found to be negative and to significantly reduce $\langle PN \rangle$. We have parameterized $\langle P'N' \rangle$ by $\langle P \rangle \langle N \rangle$ with a proportionality constant a function of two non-dimensional parameters: i) the ratio of the biological uptake time scale to the upwelling time, α , and ii) the ratio of the upwelling time scale to the turbulent diffusive time, Pe , (Peclet number).

In the figure below for the steady state the integrated phytoplankton in the near surface layer is plotted versus Pe for different values of α . The integrated phytoplankton density is nearly independent of Pe . However note that there is a maximum at $Pe \approx 5$ for $\alpha = .1$. An ADR (advective-diffusive-reactive) model has been formulated for the mean fields $\langle P \rangle$, $\langle N \rangle$ and solved analytically for the intense turbulent limit of small Peclet number. It is found that for $Pe = 0$, the fully turbulent case, $\langle P \rangle$ is constant. In the table below are shown results comparing the parameter $C = \langle P \rangle_{Pe=0}$ for the pdf and ADR theories. Agreement of $\langle P \rangle$, $\langle N \rangle$ values in the mixed layer between the two theories is excellent. The ADR results provide useful insight into coupled turbulent physical-biological dynamical processes.



Table

	α	.01	.05	.1	.25	.5
PDF Theory	C	.96	.80	.63	.35	.15
ADR Model	C	.96	.81	.64	.29	.19

IMPACT/APPLICATIONS

The often neglected turbulence induced correlated fluctuations of biological and biophysical fields should be able to be explicitly included in biodynamical processes. This in turn will allow the inclusion of important biological and biophysical processes in interdisciplinary predictive environmental models. It is expected that this work will contribute significantly to the new subfield of biogeophysical fluid dynamics.

REFERENCES

Robinson, A.R. 1999 On the theory of Advective effects on biological dynamics in the sea, II Localization, Light limitation, and nutrient saturation, *Proc R. Soc. London A.*, **455**, 1813-1828.

Goodman and Robinson, *Proc R Soc A* (2008) 469, 555-572.