

DTIC COPY

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-01-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY) 26-06-2009		2. REPORT TYPE REPRINT		3. DATES COVERED (From - To)	
4. TITLE AND SUBTITLE Nonlinear interaction of radiation belt electrons with electromagnetic ion cyclotron waves			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER 62601F		
6. AUTHORS J. M. Albert and J. Bortnik*			5d. PROJECT NUMBER 1010		
			5e. TASK NUMBER RS		
			5f. WORK UNIT NUMBER A1		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory /RVBXT 29 Randolph Road Hanscom AFB, MA 01731-3010			8. PERFORMING ORGANIZATION REPORT NUMBER AFRL-RV-HA-TR-2009-1092		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S) AFRL/RVBXR		
			11. SPONSOR/MONITOR'S REPORT		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for Public Release; distribution unlimited.			20091019023		
13. SUPPLEMENTARY NOTES Reprinted from <i>Geophysical Research Letters</i> , Vol. 36, L12110, doi:10.1029/2009GL038904, 2009 ©2009, American Geophysical Union *Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, CA					
14. ABSTRACT The rapid loss of radiation belt electrons in the main phase of geomagnetic storm is believed to be aided by EMIC waves, and is usually analyzed with quasi-linear theory. However, even moderate EMIC wave intensities easily cause resonant electrons to respond nonlinearly, with drastically different results. We map out the region of nonlinear behavior with a single parameter, and show that both the direction and magnitude of scattering can be estimated by analytical expressions. The nonlinear interactions typically lead to advection toward large pitch angles, rather than diffusion toward the loss cone. This is expected to reduce the overall loss rate and greatly affect the distribution of trapped electrons.					
15. SUBJECT TERMS Radiation belt electrons Electromagnetic ion cyclotron waves EMIC waves Cyclotron resonant interaction					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			Jay M. Albert
UNCL	UNCL	UNCL	UNL		19b. TELEPHONE NUMBER (Include area code)

Click
Here
for
Full
Article

Nonlinear interaction of radiation belt electrons with electromagnetic ion cyclotron waves

J. M. Albert¹ and J. Bortnik²

Received 27 April 2009; revised 15 May 2009; accepted 26 May 2009; published 26 June 2009.

[1] The rapid loss of radiation belt electrons in the main phase of geomagnetic storms is believed to be aided by EMIC waves, and is usually analyzed with quasi-linear theory. However, even moderate EMIC wave intensities easily cause resonant electrons to respond nonlinearly, with drastically different results. We map out the region of nonlinear behavior with a single parameter, and show that both the direction and magnitude of scattering can be estimated by analytical expressions. The nonlinear interactions typically lead to advection toward large pitch angles, rather than diffusion toward the loss cone. This is expected to reduce the overall loss rate and greatly affect the distribution of trapped electrons. **Citation:** Albert, J. M., and J. Bortnik (2009), Nonlinear interaction of radiation belt electrons with electromagnetic ion cyclotron waves, *Geophys. Res. Lett.*, 36, L12110, doi:10.1029/2009GL038904.

1. Introduction

[2] Strong electromagnetic ion cyclotron (EMIC) waves, with typical intensities of ~ 1 – 10 nT, are commonly present in the outer radiation belts [Meredith *et al.*, 2003; Fraser *et al.*, 2006]. They are excited by unstable distributions of ring current ions, and can scatter both ring current and radiation belt ions into the loss cone. They may also strongly scatter radiation belt electrons, especially near the plasmapause and in high density plumes [Thorne *et al.*, 2005, 2006; Shprits *et al.*, 2008]. Such scattering has mostly been modeled as quasi-linear diffusion [Lyons, 1974; Summers and Thorne, 2003; Gamayunov and Khazanov, 2007; Jordanova *et al.*, 1996, 2008; Miyoshi *et al.*, 2008], which indicates that EMIC waves with magnetic field amplitudes of 1 nT, in conjunction with whistler mode hiss and chorus, give decay time scales for 1 MeV electrons of less than one day [Albert, 2003; Li *et al.*, 2007]. Similar calculations with stronger EMIC waves can reach the strong diffusion limit, where particles scatter into the loss cone faster than they can be removed by the atmosphere [Shprits *et al.*, 2009]. However, such strong amplitudes raise doubts about the validity of the quasi-linear framework, and a nonlinear treatment seems called for, as noted by Millan and Thorne [2007]. Such analysis for test particles has previously been developed and applied to large amplitude whistler mode chorus [e.g., Nunn, 1974; Albert, 2002; Omura *et al.*, 2008; Bortnik *et al.*, 2008]. Here, we use a similar approach to demonstrate

the potentially nonlinear character of electron interactions with EMIC waves.

2. Inhomogeneity Parameter

[3] For field-aligned EMIC waves, the cyclotron resonance condition for electrons is

$$\omega - kv_{\parallel} = -\Omega_e/\gamma \quad (1)$$

where $\Omega_e = eB/mc$, m is the electron rest mass, and e is the electron charge (in absolute value). Resonances involving multiples of the cyclotron frequency do not play a role for field-aligned waves. Assuming the wave is generated near the equator and propagates away from it [e.g., Fraser *et al.*, 1996], the electron must be “co-streaming,” i.e., directed toward high latitude. (This is opposite to the usual requirement for resonance with field-aligned whistler mode waves.)

[4] The dynamics of resonant wave-particle interactions have been treated by many authors [e.g., Nunn, 1974; Inan *et al.*, 1978]. Albert [1993, 2000] derived a “ $1\frac{1}{2}$ dimensional” Hamiltonian for the normalized first adiabatic invariant $I = \omega(p_{\perp}/mc)^2/2\Omega_e$, the wave-particle phase ξ , and distance s along a field line, of the form

$$K(I, \xi, s) = K_0(I, s) + K_1(I, s) \sin \xi, \quad (2)$$

where K_0 describes adiabatic motion along a field line and K_1 captures the effect of the resonant wave. With field-aligned wave propagation, they take the form

$$K_0 = \frac{kc}{\omega} (I - I_0) + \sqrt{(I - I_0)^2 - 1 - 2\Omega_e I/\omega}, \quad (3)$$

$$K_1 = \frac{eB_w/mc}{kc} \tan \alpha,$$

where α is the local particle pitch angle and I_0 is a constant.

[5] For fixed s the Hamiltonian is analogous to that of a plane pendulum, whose phase portrait features two classes of periodic trajectories, with a separatrix between them. The frequency of small amplitude oscillations is $(K_1 \partial^2 K_0 / \partial I^2)^{1/2}$, and the island width is $[4K_1 / (\partial^2 K_0 / \partial I^2)]^{1/2}$. Albert [1993] considered the timescales for particle motion within a frozen phase portrait and for motion of the island itself, finding that the particle behavior was determined by their ratio,

$$R = \frac{\partial^2 K_0 / \partial I^2}{K_1 (\partial^2 K_0 / \partial s \partial I)}. \quad (4)$$

For $|R| \gg 1$, “linear” motion is expected, with an effectively random value of the phase at resonance, leading

¹Space Vehicles Directorate, Air Force Research Laboratory, Hanscom Air Force Base, Bedford, Massachusetts, USA.

²Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, California, USA.

DTIC COPY




Figure 1. Inhomogeneity parameter R versus equatorial pitch angle and energy, for a model of 2 nT EMIC waves in a high density plume at $L = 4$. The full equation (5) was used for the left-hand plot, and the approximate equation (7) was used on the right. The triangles indicate the particles simulated numerically. Nonlinear interactions are expected for $R < 1$.

to diffusion in I , while $|R| \ll 1$ leads to essentially nonlinear motion, with phase bunching (with or without phase trapping), and deterministic changes in I . R represents a competition between the strength of the wave, in K_1 , and the inhomogeneity of the background magnetic field (and plasma properties), which enters through the derivative with respect to s (effectively, latitude λ).

[6] *Omura et al.* [2008] recently presented a related analysis for field-aligned whistler-mode chorus waves, working directly with the resonance-averaged equations of motion, and obtained a related nonlinear oscillator equation with the same properties. Repeating their derivation for arbitrary (but field-aligned) waves yields an “inhomogeneity parameter” which coincides exactly with R , and can be written explicitly as

$$R = \frac{-B}{B_w} \frac{\mu}{\mu^2 - 1} \frac{c}{v_\perp} \frac{c}{\omega} \left\{ \left[\mu \frac{\gamma}{2} \frac{\omega}{\Omega_e} \frac{v_\perp^2}{c^2} - \frac{v_\parallel}{c} \right] \frac{1}{\Omega_e} \frac{\partial \Omega_e}{\partial s} - \mu \frac{\gamma}{2} \frac{\omega}{\Omega_e} \frac{v_\perp^2}{c^2} \frac{1}{\mu^2} \frac{\partial (\mu^2)}{\partial s} \right\}, \quad (5)$$

evaluated at resonance. Here μ is the refractive index, kc/ω .

[7] For parallel-propagating EMIC waves, μ^2 is given by the Stix L parameter [Stix, 1962], which is conveniently written as

$$\mu^2 = 1 - \frac{\omega_{pe}^2}{\omega^2} \left\{ \frac{\omega}{\omega + \Omega_e} + \sum_i \frac{\eta_i Z_i^2}{\beta_i} \frac{1}{M} \frac{\omega}{\omega - \Omega_i} \right\}, \quad (6)$$

where M is the electron-to-proton mass ratio m/m_p , i indexes the ion species, $\beta_i = m_i/m_p$, $\eta_i = n_i/n_e$, and Z_i is the charge of ion i . Quasineutrality requires $\sum_i Z_i \eta_i = 1$. EMIC waves only propagate where $\mu^2 > 0$, which occurs in frequency bands bounded below by cutoffs and above by resonances at the ion cyclotron frequencies.

[8] We take the ions to be H^+ , He^+ , and O^+ , so that $\beta_H = 1$, $\beta_{He} = 4$, $\beta_O = 16$, with all $Z_i = 1$. Typical storm-time values of the ion concentrations are $\eta_H = 0.77$, $\eta_{He} = 0.20$, $\eta_O = 0.03$ [Jordanova et al., 2008]. The value of μ in equation (6) will be dominated by the term corresponding to

the appropriate ion band. Making this approximation, using a small-latitude approximation of the dipole magnetic field, and ignoring density variation in equation (5) yield the estimate

$$R \approx \frac{-9\lambda}{2L} \frac{mc}{eB_w} \frac{c}{R_e} \frac{p}{mc} \frac{\cos^2 \alpha}{\sin \alpha} \frac{1}{1 - \omega/\Omega_i}. \quad (7)$$

From now on, it will be understood that “ R ” refers to $|R|$.

[9] Figure 1 shows R as a function of energy and equatorial pitch angle α_0 for field-aligned, 2 nT EMIC waves in the helium band, evaluated in a high density plume at $L = 4$, with $\omega_{pe}/\Omega_e = 15$. The frequency $\omega/2\pi$ is taken to be about 1.84 Hz, so that $\omega/\Omega_i = 0.96$ at the equator. Such a large ratio is necessary to bring the minimum resonant energy down to about 1 MeV [Summers and Thorne, 2003; Albert, 2003]. Equations (5) and (7) give very similar results for R . Near the loss cone, the interactions are expected to be linear (that is, diffusive, as discussed below). However, for a wide range of pitch angle and energy values, the interactions are expected to be nonlinear, so that quasi-linear diffusion will not be a valid description.

3. Equations of Motion and Analytical Estimates

[10] To test these ideas, we follow particles numerically, using the gyroresonance-averaged equations of motion following Chang and Inan [1983]. These equations take the form

$$\begin{aligned} \frac{dp_\parallel}{dt} &= \frac{eB_w}{c} \frac{p_\perp}{m\gamma} \sin \phi - \frac{p_\perp^2}{2m\gamma B} \frac{\partial B}{\partial s} \\ \frac{dp_\perp}{dt} &= \frac{eB_w}{c} \left(\frac{\omega}{k} - \frac{p_\parallel}{m\gamma} \right) \sin \phi + \frac{p_\perp p_\parallel}{2m\gamma B} \frac{\partial B}{\partial s} \\ \frac{d\phi}{dt} &= \left(\frac{kp_\parallel}{m\gamma} - \omega - \frac{\Omega_e}{\gamma} \right) + \frac{eB_w}{c} \left(\frac{\omega}{k} - \frac{p_\parallel}{m\gamma} \right) \frac{\cos \phi}{p_\perp} \\ \frac{ds}{dt} &= \frac{p_\parallel}{m\gamma} \end{aligned} \quad (8)$$




Figure 2. Trajectories of 2 MeV electrons with $\alpha_0 = 15^\circ$, interacting with 2 nT EMIC waves in a high density plume. Even for these large amplitude waves, the interaction is fairly linear, leading to nearly symmetric scattering in α_0 and in E .

In a dipole field, s is easily transformed to latitude λ . Both the wave frequency ω and magnetic field strength B_w , as well as the background plasma density, are treated as constant. The particles were not allowed to reach the location of the crossover frequency, where the Stix parameter $D = 0$, and the wave can switch from left hand to right hand circular polarization [Stix, 1962]. The wave-particle phase ϕ differs from ξ in equation (2) only by sign and phase conventions.

[11] The Hamiltonian analysis of Albert [1993] yields analytical estimates of the net, nonadiabatic change in I due to either linear or nonlinear resonant interactions. These are:

$$\begin{aligned} \Delta I_{lin} &= -K_1 \sqrt{\frac{2\pi}{|\partial^2 K_0 / \partial s \partial I|}} \cos \tilde{\xi}, \quad R > 1, \\ \Delta I_{NL} &= \frac{8}{\pi} \sqrt{\left| \frac{K_1}{\partial^2 K_0 / \partial I^2} \right|}, \quad R < 1. \end{aligned} \quad (9)$$

Here $\tilde{\xi}$ is related to the value of ξ evaluated right at resonance, and is effectively random. Thus in the linear regime, I suffers a random walk with zero mean, or diffuses, while in the nonlinear regime the resonant value of ξ is well defined and I experiences systematic drift, or advection. Corresponding changes in energy and α_0 are [Albert, 2000]

$$\Delta \alpha_0 = \frac{\gamma \sin^2 \alpha_0 + \Omega_{eq} / \omega}{(p/mc)^2 \sin \alpha_0 \cos \alpha_0} \Delta I, \quad \frac{\Delta E}{mc^2} = -\Delta I, \quad (10)$$

where Ω_{eq} is the equatorial value of Ω_e . Since $\omega \ll \Omega_e$, pitch angle changes will be much larger than normalized energy changes.

4. Results

[12] First, the trajectories of 24 particles interacting with a small amplitude wave, $B_w = 5$ pT, were calculated according to equation (8). The particles were started at the equator with $E = 2$ MeV, $\alpha_0 = 15^\circ$, and a uniform spread in initial phase, ϕ_0 . For this arrangement, R at the expected resonance location is nearly 1000. After passing through the

resonance, both α_0 and E are sharply scattered. The distributions of $\Delta \alpha_0$ and ΔE , though not shown, are symmetric about zero and depend sinusoidally on the initial phases, which should be similar to the phases just before the resonant interactions. Furthermore, the scattering amplitudes agree quite well with the prediction ΔI_{lin} .

[13] Figure 2 shows a similar simulation with $B_w = 2$ nT, which gives $R = 2.4$. This distribution of changes is also fairly sinusoidal, though not perfectly so, and the amplitudes of the changes are well predicted by ΔI_{lin} , shown as the double-arrowed line. The maximum change in $\Delta \alpha_0$ is about 10° , which is nearly enough to move the particles into the loss cone in one bounce. The pitch angle diffusion coefficient, $\langle (\Delta \alpha_0)^2 \rangle / \tau_b \approx 3 \times 10^3 \text{ day}^{-1}$, is in reasonable agreement with calculations based on broadband quasi-linear theory applied to similar models of EMIC waves at $L = 4.5$ [Li et al., 2007; Albert, 2008]. The bottom two panels show the expected, nearly sinusoidal dependence on initial phase, as well as mean changes (dashed lines) and mean changes plus or minus one standard deviation (dotted lines).

[14] In Figure 3, the initial pitch angle of the 2 MeV particles is changed to $\alpha_0 = 45^\circ$, for which $R = 0.54$, a moderately nonlinear value. In fact, the distribution of changes is very different from above, with most particles tracing very similar paths. This is due to the underlying "bunching" of wave-particle phases as the particles encounter and circumnavigate the separatrix between trapped and untrapped trajectories (see, e.g., Nunn [1974], Albert [1993, 2000], Omura et al. [2008] for details). Some paths are exceptions, reflecting the fact that R is not very much less than 1. Nevertheless, the predicted changes based on ΔI_{NL} , indicated by the plus symbols, give good estimates of the maximum, and most common, values. Repeated interactions would cause the particle distributions to drift, not diffuse, in pitch angle and energy; the cumulative changes would grow proportionally to time, not as $t^{1/2}$. Also note that the typical change in α_0 is positive, away from the loss cone. This is expected to cause lower phase space density at the relevant pitch angles, and lower overall loss rates, than estimates based on pure diffusion.




Figure 3. Trajectories of 2 MeV electrons with $\alpha_0 = 45^\circ$, interacting with 2 nT EMIC waves in a high density plume. For these particles, the interaction is nonlinear, leading to phase bunching with systematic increases in α_0 and decreases in E .

[15] None of the particles in Figure 3 exhibit phase trapping, whereby particles cross the separatrix and spend many phase oscillations within it. This is not surprising since the particles are moving away from the equator, as required for resonance with the EMIC waves, and so experience increasing magnetic field inhomogeneity. Even if $R < 1$ at resonance, R will be increasing, which is not conducive to trapping [Albert, 2000]. On the other hand, particles may start out inside the separatrix. Figure 4 shows 2 MeV particles starting with $\alpha_0 = 55^\circ$ at the equator. The linear resonance condition, equation (1), is satisfied at $\lambda = 1.7^\circ$, where $R = 0.2$. About half the particles behave nonlinearly in accordance with ΔI_{NL} , as shown by the plus marks. However, almost half the particles experience even larger, and oppositely signed changes, which develop over a sustained interval of nonadiabatic motion, as is characteristic of phase trapping. Evidently the initial conditions are near enough to resonance that some particles complete several phase oscillations before becoming nonresonant. Detailed examination of the trajectories verifies that for

the trapped particles, ϕ oscillates in a limited range starting at $t = 0$. Physically, this would occur if $B_{\parallel}$, which has been held constant, grew rapidly to “envelop” regions of phase space containing nearly resonant particles.

5. Summary

[16] We have demonstrated that the cyclotron resonant interaction of relativistic electrons with field-aligned EMIC waves can be highly nonlinear. Regions of expected nonlinear behavior were mapped out with the parameter R , using a realistic magnetic field amplitude of 2 nT. Moreover, we have shown that previously developed theoretical expressions for the expected pitch angle and energy changes give good estimates of the behavior seen in numerical simulation of the particle motion. Phase bunching without trapping, the more commonly expected version of nonlinear behavior, leads to rapid pitch angle increases, away from the loss cone, with small changes in energy. In combination with diffusion, this could lead to major, complex effects on




Figure 4. Trajectories of 2 MeV electrons with $\alpha_0 = 55^\circ$, interacting with 2 nT EMIC waves in a high density plume. For these particles, the interaction is nonlinear and some particles are initially phase trapped, giving both systematic increases and sustained decreases in α_0 , and oppositely directed changes in E .

both the loss rate and the trapped particle distribution. Since EMIC waves are believed to play a major role in the loss of radiation belt electrons during storms, such behavior should be accounted for in future models. One approach would be to use the analytical expressions in a combined diffusion-advection formulation.

[17] **Acknowledgments.** This work was supported by the Space Vehicles Directorate of the Air Force Research Laboratory. J.B. gratefully acknowledges support from NASA grant NNX08A1135G.

References

- Albert, J. M. (1993), Cyclotron resonance in an inhomogeneous magnetic field, *Phys. Fluids B*, **5**, 2744.
- Albert, J. M. (2000), Gyroresonant interactions of radiation belt particles with a monochromatic electromagnetic wave, *J. Geophys. Res.*, **105**, 21,191, doi:10.1029/2000JA000008.
- Albert, J. M. (2002), Nonlinear interaction of outer zone electrons with VLF waves, *Geophys. Res. Lett.*, **29**(8), 1275, doi:10.1029/2001GL013941.
- Albert, J. M. (2003), Evaluation of quasi-linear diffusion coefficients for EMIC waves in a multispecies plasma, *J. Geophys. Res.*, **108**(A6), 1249, doi:10.1029/2002JA009792.
- Albert, J. M. (2008), Efficient approximations of quasi-linear diffusion coefficients in the radiation belts, *J. Geophys. Res.*, **113**, A06208, doi:10.1029/2007JA012936.
- Bortnik, J., R. M. Thorne, and U. S. Inan (2008), Nonlinear interaction of energetic electrons with large amplitude chorus, *Geophys. Res. Lett.*, **35**, L21102, doi:10.1029/2008GL035500.
- Chang, H. C., and U. S. Inan (1983), Quasi-relativistic electron precipitation due to interactions with coherent VLF waves in the magnetosphere, *J. Geophys. Res.*, **88**, 318.
- Fraser, B. J., H. J. Singer, W. J. Hughes, J. R. Wygant, R. R. Anderson, and Y. D. Hu (1996), CRRES Poynting vector observations of electromagnetic ion cyclotron waves near the plasmapause, *J. Geophys. Res.*, **101**, 15,331.
- Fraser, B. J., T. M. Loto'ainu, and H. J. Singer (2006), Electromagnetic ion cyclotron waves in the magnetosphere, in *Magnetospheric ULF Waves: Synthesis and New Directions*, *Geophys. Monogr. Ser.*, vol. 169, edited by K. Takahashi et al., p. 195, AGU, Washington, D. C.
- Gamayunov, K. V., and G. V. Khazanov (2007), Effect of oblique electromagnetic ion cyclotron waves on relativistic electron scattering: Combined Release and Radiation Effects Satellite (CRRES)-based calculation, *J. Geophys. Res.*, **112**, A07220, doi:10.1029/2007JA012300.
- Inan, U. S., T. F. Bell, and R. A. Helliwell (1978), Nonlinear pitch angle scattering of energetic electrons by coherent VLF waves in the magnetosphere, *J. Geophys. Res.*, **83**, 3235.
- Jordanova, V. K., J. U. Kozyra, and A. F. Nagy (1996), Effects of heavy ions on the quasi-linear diffusion coefficients from resonant interactions with electromagnetic ion cyclotron waves, *J. Geophys. Res.*, **101**, 19,771.
- Jordanova, V. K., J. Albert, and Y. Miyoshi (2008), Relativistic electron precipitation by EMIC waves from self-consistent global simulations, *J. Geophys. Res.*, **113**, A00A10, doi:10.1029/2008JA013239.
- Li, W., Y. Y. Shprits, and R. M. Thorne (2007), Dynamic evolution of energetic outer zone electrons due to wave-particle interactions during storms, *J. Geophys. Res.*, **112**, A10220, doi:10.1029/2007JA012368.
- Lyons, L. R. (1974), Pitch angle and energy diffusion coefficients from resonant interactions with ion-cyclotron and whistler waves, *J. Plasma Phys.*, **12**, 417.
- Meredith, N. P., R. M. Thorne, R. B. Horne, D. Summers, B. J. Fraser, and R. R. Anderson (2003), Statistical analysis of relativistic electron energies for cyclotron resonance with EMIC waves observed on CRRES, *J. Geophys. Res.*, **108**(A6), 1250, doi:10.1029/2002JA009700.
- Millan, R. M., and R. M. Thorne (2007), Review of radiation belt relativistic electron loss, *J. Atmos. Sol. Terr. Phys.*, **69**, 362.
- Miyoshi, Y., K. Sakaguchi, K. Shiokawa, D. Evans, J. Albert, M. Connors, and V. Jordanova (2008), Precipitation of radiation belt electrons by EMIC waves, observed from ground and space, *Geophys. Res. Lett.*, **35**, L23101, doi:10.1029/2008GL035727.
- Nunn, D. (1974), A self-consistent theory of triggered VLF emissions, *Planet. Space Sci.*, **22**, 349.
- Omura, Y., Y. Katoh, and D. Summers (2008), Theory and simulation of the generation of whistler-mode chorus, *J. Geophys. Res.*, **113**, A04223, doi:10.1029/2007JA012622.
- Shprits, Y. Y., D. A. Subbotin, N. P. Meredith, and S. R. Elkington (2008), Review of modeling losses and sources of relativistic electrons in the outer radiation belt II: Local acceleration and loss, *J. Atmos. Sol. Terr. Phys.*, **70**, 1694, doi:10.1016/j.jastp.2008.06.014.
- Shprits, Y. Y., L. Chen, and R. M. Thorne (2009), Simulations of pitch angle scattering of relativistic electrons with MLT-dependent diffusion coefficients, *J. Geophys. Res.*, **114**, A03219, doi:10.1029/2008JA013695.
- Stix, T. H. (1962), *The Theory of Plasma Waves*, McGraw-Hill, New York.
- Summers, D., and R. M. Thorne (2003), Relativistic electron pitch-angle scattering by electromagnetic ion cyclotron waves during geomagnetic storms, *J. Geophys. Res.*, **108**(A4), 1143, doi:10.1029/2002JA009489.
- Thorne, R. M., R. B. Horne, S. A. Glauert, N. P. Meredith, Y. Y. Shprits, D. Summers, and R. R. Anderson (2005), Dynamic evolution of energetic outer zone electrons due to wave-particle interactions during storms, in *Inner Magnetosphere Interactions: New Perspectives from Imaging*, *Geophys. Monogr. Ser.*, vol. 159, edited by J. L. Burch, M. Schulz, and H. Spence, p. 101, AGU, Washington, D. C.
- Thorne, R. M., R. B. Horne, V. K. Jordanova, J. Bortnik, and S. A. Glauert (2006), Interaction of EMIC waves with thermal plasma and radiation belt particles, in *Magnetospheric ULF Waves: Synthesis and New Directions*, *Geophys. Monogr. Ser.*, vol. 169, edited by K. Takahashi et al., p. 213, AGU, Washington, D. C.

J. M. Albert, Air Force Research Laboratory, 29 Randolph Road, Hanscom AFB, MA 01731-3010, USA. (jay.albert@hanscom.af.mil)

J. Bortnik, Department of Atmospheric and Oceanic Sciences, University of California, Los Angeles, CA 90095-1565, USA. (jbortnik@gmail.com)