

AD_____

AWARD NUMBER: W81XWH-08-2-0653

TITLE: Validation of the Military Acute Concussion Evaluation (MACE) for In-Theater
Evaluation of Combat-Related Traumatic Brain Injury

PRINCIPAL INVESTIGATOR: Michael McCrea, Ph.D.
COL Michael Jaffee, M.D.
Kathy Helmick
Guskiewicz, Ph.D.
Selina Doncevic

CONTRACTING ORGANIZATION: Waukesha Memorial Hospital
Waukesha, WI 53188

REPORT DATE: October 2009

TYPE OF REPORT: Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE 1 October 2009		2. REPORT TYPE Annual		3. DATES COVERED 15 Sep 2008 – 14 Sep 2009	
4. TITLE AND SUBTITLE Validation of the Military Acute Concussion Evaluation (MACE) for In-Theater Evaluation of Combat-Related Traumatic Brain Injury				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-08-2-0653	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Michael McCrea, Ph.D.; COL Michael Jaffee, M.D., Kathy Helmick; Kevin Guskiewicz, Ph.D.; Selina Doncevic E-Mail: michael.mccrea@phci.org				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Waukesha Memorial Hospital Waukesha, WI 53188				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT BACKGROUND: Traumatic brain injury (TBI) has been widely considered the “signature injury” among United States military personnel involved in combat in Iraq and Afghanistan. The Military Acute Concussion Evaluation (MACE) was designed by DVBIC and civilian brain injury experts to assess the mechanism of injury, acute characteristics and cognitive deficits in military personnel with suspected MTBI in an austere environment. PURPOSE AND SCOPE: This study investigates the clinical and operational utility of the MACE in military operational settings. Through a systematic, retrospective review of MACE data, the specific aims of this study are: Epidemiological: To advance our understanding of the acute injury characteristics of MTBI in the current military operational setting (e.g., mechanisms of injury, influence of personal protective equipment (PPE), clinical indicators, severity range). Clinical: To determine the clinical utility of the MACE in assessing the acute signs and symptoms of MTBI, measuring the acute cognitive effects, and objectively tracking recovery; To assess the unique contribution of the MACE in clinical decision-making and modulating risk around fitness to return to duty after MTBI. Operational: To assess the practical feasibility and user-friendliness of the MACE for medical and other personnel in the military operational setting. PROGRESS: Significant delays were initially confronted in gaining necessary IRB approval for this retrospective study. These issues have now been resolved and data analysis is in process. No formal results or conclusions are available at the time of this abstract submission.					
15. SUBJECT TERMS Traumatic brain injury; concussion; MACE					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 10	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

Table of Contents

	<u>Page</u>
Introduction.....	4
Body.....	5
Key Research Accomplishments.....	6
Reportable Outcomes.....	6
Conclusion.....	6
References.....	8
Appendices.....	10

INTRODUCTION

Traumatic brain injury (TBI) has been widely considered the “signature injury” among United States military personnel involved in combat in Iraq and Afghanistan. In previous wars such as Desert Storm, approximately 20% of military personnel treated for wounds had primary or concurrent head injuries (Carey, 1991, 1996; Leadham, Newland, & Blood, 1993). Due to several factors, however, the rate of traumatic brain injury in Operation Enduring Freedom (OEF) and Operation Iraqi Freedom (OIF) is thought to be significantly higher than any previous war (Warden, 2006). In brief, advances in protective armor (e.g., helmets and Kevlar vests) and medical triage have saved lives of military personnel that likely would have died from the same serious injuries in previous wars. Additionally, the frequency of explosive or blast attacks in Iraq and Afghanistan is significantly higher than in past military conflicts that create a new set of concerns about the risks and dynamics of closed head injury (Scott, Belanger, Vanderploeg, Massengale, & Scholten, 2006; Taber, Warden, & Hurley, 2006).

Data from the Defense and Veterans Brain Injury Center (DVBIC) headquarters at Walter Reed Army Medical Center (WRAMC) show that, among OIF and OEF veterans, 30 percent of battle injured OIF/OEF veterans were found to have traumatic brain injury, with an even greater percentage meeting TBI criteria when their mechanism of injury was blast related. As in the civilian setting, the overwhelming majority of TBI's (> 85%) in the current military conflict are categorized as MTBI based on acute injury characteristics and accepted injury definition criteria. The prevalence of MTBI in the austere environment is considered very high, a precise estimate being difficult to establish due to the fact that milder injuries may go untreated or unreported, just as in the civilian sector. Additionally, an estimated 10-20 percent of combat veterans meet the criteria for MTBI on post-deployment screening (Zoroya, 2006).

Unfortunately, MTBI presents a unique set of challenges in terms of injury detection, diagnosis, assessment and management due to the more subtle nature of injury characteristics in the absence of classic indicators (e.g., unconsciousness, amnesia, focal neurological deficit, positive neuroimaging findings). Military personnel have explicitly requested a clinical tool to assist in the acute triage of TBI that is appropriate for the frontline military operational setting.

As a result, assessment, management, and rehabilitation of deployment-related TBI has garnered increasing attention from the medical community (both military and civilian), multiple government agencies, patient advocacy groups, and the media. The Department of Defense (DoD) assembled the Defense and Veterans Brain Injury Center (DVBIC) Working Group on the Acute Management of Mild Traumatic Brain Injury (MTBI) in Military Operational Settings, which included representation from neuropsychology and generated the first Clinical Practice Guideline (CPG) in December 2006 (DVBIC, 2006). The Working Group's CPG focused primarily on standardized algorithms for the operational assessment and management of MTBI *in-theater*, but also generated recommendations for pre-deployment baseline cognitive testing and military educational initiatives around MTBI.

The Military Acute Concussion Evaluation (MACE) (see Appendix A) was designed by DVBIC and civilian brain injury experts specifically for the purposes of assessing and documenting the mechanism of injury, acute characteristics and cognitive deficits in military personnel with suspected MTBI in an austere environment. The MACE was developed by a team of military and civilian TBI experts and first distributed for clinical use by military personnel in August 2006. The instrument is currently the only standardized and most widely used method for evaluation of acute MTBI in military operational settings. Embedded in the MACE is the Standardized Assessment of Concussion (SAC), a brief cognitive screening tool with demonstrated reliability, validity, sensitivity, and specificity in assessing the acute cognitive effects of sport-related MTBI (DVBIC, 2006; McCrea et al., 2003; McCrea, Kelly, Randolph, Cisler, & Berger, 2002).

Although the MACE and SAC have sound basis extrapolated from the sport concussion literature, *neither has been formally validated for the unique purpose of evaluating military-related MTBI, particularly in combat theater.* The current study represents the first formal investigation of the clinical and operational utility of the MACE in military operational settings.

BODY

This study is designed to focus on investigating the clinical and operational utility of the MACE in military operational settings. It is hypothesized that the MACE is a valid and reliable tool that has significant clinical utility in the acute triage of TBI in an austere environment. In keeping with the parameters of the TBI Concept Award with respect to prospective study of human subjects, a retrospective review of MACE data already collected on deployed MTBI patients since August 2006 will be executed through access to existing DoD databases. Through a systematic review of MACE data, the specific aims of this study are:

Epidemiological: To advance our understanding of the acute injury characteristics of MTBI in the current military operational setting (e.g., mechanisms of injury, influence of personal protective equipment (PPE), clinical indicators, severity range)

Specifically, the epidemiological objectives are:

- To document the frequency of specific acute injury characteristics (e.g., loss of consciousness, posttraumatic amnesia, specific symptoms) as markers of traumatic brain injury in this setting.
- To document known characteristics causing traumatic brain injury in the current setting (e.g., blast vs. blunt trauma, acceleration/deceleration, etc.)
- To document the distribution of injury severity gradient (mild, moderate, severe) in the current setting.
- To document other vital statistics relevant to traumatic brain injury (e.g., was protective helmet or other equipment worn at time of injury, etc.).
- Other objectives as identified during study

Clinical: To determine the clinical utility of the MACE in assessing the acute signs and symptoms of MTBI, measuring the acute cognitive effects, and objectively tracking recovery;
To assess the unique contribution of the MACE in clinical decision-making and modulating risk around fitness to return to duty after MTBI

Specifically, the clinical objectives are:

- To analyze MACE scores from earliest post injury assessment point to final assessment point to establish sensitivity/specificity of MACE score in detecting cognitive abnormalities after TBI, and plotting MACE score recovery curves as done in previous studies using the SAC.
- When possible, compare post-injury MACE scores to pre-injury baseline score to determine sensitive and specificity of MACE change scores as a marker of cognitive dysfunction after TBI (and track recovery back to baseline MACE score).

- When no preinjury baseline MACE score is available, analyze the distribution of postinjury MACE scores to help determine evidence-based cutoff scores for determining cognitive dysfunction that minimize the risk of Type I or Type II errors in clinical decision making based on MACE scores.
- To analyze MACE data and determine what symptoms are most common after TBI in this setting, both acutely and persistently
- Other objectives as identified during study

Operational: To assess the practical feasibility and user-friendliness of the MACE for medical and other personnel in the military operational setting

Specifically, the operational objectives are:

- To analyze clinician feedback on the clinical utility of the MACE as a tool to assist them in making a more accurate assessment of TBI effects and recovery in the current setting.
- To assess recommendations from end users of the MACE that could enhance the instrument, either clinically or operationally
- Other objectives as identified during study

Our hypothesis tested by this study is that the MACE is a reliable, valid, sensitive and specific tool to assess traumatic brain injury that is a valuable resource to users of the tool in the current setting.

KEY RESEARCH ACCOMPLISHMENTS

To date, the following project-related tasks have been accomplished:

1. Secured approval from relevant Institutional Review Boards (IRB) for completion of this retrospective data study.
2. Refined the plan for data extraction for specific DoD and DVBIC databases that contain the information relevant to the specific aims of this study.
3. Secured part-time, contracted biostatistician to assist in data extraction, management and analysis.
4. Determined the planned sequence of tasks to effectively capture relevant data for analysis.
5. Developed a detailed plan for statistical analysis of data in accordance with the study's specific aims.
6. Presented oral and poster presentations at the 2009 Congressional Directed Medical Research Program Forum, providing an overview of the study significance, specific aims, design and expected impact on military and civilian populations.

REPORTABLE OUTCOMES

Data aggregation and analysis is underway and no reportable findings or results are available at the time of this annual report.

CONCLUSION

We anticipate significant progress ahead in completing this study according to the intended specific aims and statement of work.

As it relates to military application, this study is predicted to directly address valid criticisms currently being voiced by military and civilian clinicians as to the existing gap in established validity of the Military Acute Concussion Evaluation (MACE) and Standardized Assessment of Concussion (SAC) as the main methods used to evaluate mild traumatic brain injury (MTBI) in Operation Enduring Freedom (OEF) and Operation Iraqi Freedom (OIF).

The findings from this study are also expected to inform future prospective investigations of traumatic brain injury that utilize the MACE and similar methods, particularly for assessment of military-related MTBI. Because there is currently no widely accepted standard for objectively evaluating MTBI in either general research or clinical environments, findings from this study are predicted to have more global, ground-breaking, and translational implications for establishment of a standardized clinical instrument and research tool to evaluate MTBI in settings of mass casualty, breaches to homeland security, and other trauma settings.

REFERENCES

- APA. (1994). *Diagnostic and Statistical Manual of Mental Disorders Fourth Edition*. Washington, DC: American Psychiatric Association.
- Barr, W. B. (2001). Methodologic Issues in Neuropsychological Testing. *J Athl Train*, 36(3), 297-302.
- Belanger, H. G., Curtiss, G., Demery, J. A., Lebowitz, B. K., & Vanderploeg, R. D. (2005). Factors moderating neuropsychological outcomes following mild traumatic brain injury: a meta-analysis. *J Int Neuropsychol Soc*, 11(3), 215-227.
- Borg, J., Holm, L., Peloso, P. M., Cassidy, J. D., Carroll, L. J., von Holst, H., Paniak, C., & Yates, D. (2004). Non-surgical intervention and cost for mild traumatic brain injury: results of the WHO Collaborating Centre Task Force on Mild Traumatic Brain Injury. *J Rehabil Med*(43 Suppl), 76-83.
- Brolinson, P. G., Manoogian, S., McNeely, D., Goforth, M., Greenwald, R., & Duma, S. (2006). Analysis of linear head accelerations from collegiate football impacts. *Curr Sports Med Rep*, 5(1), 23-28.
- Carey, M. E. (1991). Analysis of wounds incurred by U.S. Army Seventh Corps personnel in Corps hospitals during Operation Desert Storm, February 20 to March 10, 1991. *Journal of Trauma*, 40(3), S165-S169.
- Carey, M. E. (1996). Analysis of wounds incurred by U.S. Army Seventh Corps personnel treated in Corps hospitals during Operation Desert Storm, February 20 to March 10, 1991. *J Trauma*, 40(3 Suppl), S165-169.
- DVBIC. (2006). *Defense and Veterans Brain Injury Center Working Group on the Acute Management of Mild Traumatic Brain Injury in Military Operational Settings*, Washington, DC.
- Guskiewicz, K. M., Bruce, S. L., Cantu, R. C., Ferrara, M. S., Kelly, J. P., McCrea, M., Putukian, M., & McLeod, T. C. (2004). Recommendations on management of sport-related concussion: summary of the National Athletic Trainers' Association position statement. *Neurosurgery*, 55(4), 891-895; discussion 896.
- Hoge, C. W., Auchterlonie, J. L., & Milliken, C. S. (2006). Mental health problems, use of mental health services, and attrition from military service after returning from deployment to Iraq or Afghanistan. *Jama*, 295(9), 1023-1032.
- Hoge, C. W., Castro, C. A., Messer, S. C., McGurk, D., Cotting, D. I., & Koffman, R. L. (2004). Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *N Engl J Med*, 351(1), 13-22.
- Holm, L., Cassidy, J. D., Carroll, L. J., & Borg, J. (2005). Summary of the WHO Collaborating Centre for Neurotrauma Task Force on Mild Traumatic Brain Injury. *J Rehabil Med*, 37(3), 137-141.
- Iverson, G. L. (2005). Outcome from mild traumatic brain injury. *Curr Opin Psychiatry*, 18(3), 301-317.
- Iverson, G. L., Zasler, N. D., & Lange, R. T. (2006). Post-Concussive Disorder. In N. D. Zasler & D. I. Katz & R. D. Zafonte (Eds.), *Brain Injury Medicine: Principles and Practice* (pp. 373-405). New York: Demos Medical Publishing.
- Jennett, B., & Teasdale, G. (1981). *Management of Head Injuries*. Philadelphia, PA: FA Davis.
- Kashluba, S., Paniak, C., Blake, T., Reynolds, S., Toller-Lobe, G., & Nagy, J. (2004). A longitudinal, controlled study of patient complaints following treated mild traumatic brain injury. *Arch Clin Neuropsychol*, 19(6), 805-816.
- Kay, T., Harrington, D. E., Adams, R. E., Anderson, T. W., Berrol, S., Cicerone, K., Dahlberg, C., Gerber, D., Goka, R. S., Harley, J. P., Hilt, J., Horn, L. J., Lehmkuhl, D., & Malec, J. (1993). Definition of mild traumatic brain injury: Report from the Mild Traumatic Brain Injury Committee of the Head Injury Interdisciplinary Special Interest Group of the

- American Congress of Rehabilitation Medicine. *Journal of Head Trauma Rehabilitation*, 8(3), 86-87.
- Leadham, C. S., Newland, C., & Blood, C. G. (1993). A descriptive analysis of wounds among U.S. Marines treated at second echelon facilities in the Kuwaiti theater of operation. *Military Medicine*, 158(8), 508-512.
- Luis, C. A., Vanderploeg, R. D., & Curtiss, G. (2003). Predictors of postconcussion symptom complex in community dwelling male veterans. *J Int Neuropsychol Soc*, 9(7), 1001-1015.
- McCrea, M. (2007). *Mild Traumatic Brain Injury and Post-Concussion Syndrome: The New Evidence Base for Diagnosis and Treatment*. New York: Oxford Press.
- McCrea, M., Barr, W. B., Guskiewicz, K., Randolph, C., Marshall, S. W., Cantu, R., Onate, J. A., & Kelly, J. P. (2005). Standard regression-based methods for measuring recovery after sport-related concussion. *J Int Neuropsychol Soc*, 11(1), 58-69.
- McCrea, M., Guskiewicz, K. M., Marshall, S. W., Barr, W., Randolph, C., Cantu, R. C., Onate, J. A., Yang, J., & Kelly, J. P. (2003). Acute effects and recovery time following concussion in collegiate football players: the NCAA Concussion Study. *Jama*, 290(19), 2556-2563.
- McCrea, M., Hammeke, T., Olsen, G., Leo, P., & Guskiewicz, K. (2004). Unreported concussion in high school football players: implications for prevention. *Clin J Sport Med*, 14(1), 13-17.
- McCrea, M., Kelly, J. P., Randolph, C., Cisler, R., & Berger, L. (2002). Immediate neurocognitive effects of concussion. *Neurosurgery*, 50(5), 1032-1040; discussion 1040-1032.
- Mittenberg, W., Tremont, G., Zielinski, R. E., Fichera, S., & Rayls, K. R. (1996). Cognitive-behavioral prevention of postconcussion syndrome. *Arch Clin Neuropsychol*, 11(2), 139-145.
- Ponsford, J., Willmott, C., Rothwell, A., Cameron, P., Ayton, G., Nelms, R., Curran, C., & Ng, K. (2001). Impact of early intervention on outcome after mild traumatic brain injury in children. *Pediatrics*, 108(6), 1297-1303.
- Randolph, C., McCrea, M., & Barr, W. B. (2005). Is neuropsychological testing useful in the management of sport-related concussion? *J Athl Train*, 40(3), 139-152.
- Report to Congress on Mild Traumatic Brain Injury in the United States: Steps to Prevent a Serious Public Health Problem*. (2003). Atlanta, GA: National Center for Injury Prevention and Control, Centers for Disease Control and Injury Prevention.
- Scott, S. G., Belanger, H. G., Vanderploeg, R. D., Massengale, J., & Scholten, J. (2006). Mechanism-of-injury approach to evaluating patients with blast-related polytrauma. *J Am Osteopath Assoc*, 106(5), 265-270.
- Taber, K. H., Warden, D. L., & Hurley, R. A. (2006). Blast-related traumatic brain injury: what is known? *J Neuropsychiatry Clin Neurosci*, 18(2), 141-145.
- Warden, D. (2006). Military TBI during the Iraq and Afghanistan wars. *J Head Trauma Rehabil*, 21(5), 398-402.
- Warden, D. L., Ryan, L. M., Helmick, K. M., Schwab, K., French, L. M., Lu, W., Lux, W. E., Ecklund, J., & Ling, G. (2005). War neurotrauma: The Defense and Veterans Brain Injury Center (DVBIC) experience at Walter Reed Army Medical Center (WRAMC) (abstract). *Journal of Neurotrauma*, 22(10), 1178.
- WHO. (1992). *International Statistical Classification of Diseases and Related Health Problems - 10th Edition*. Geneva, Switzerland: World Health Organization.
- Zhang, L., Yang, K. H., & King, A. I. (2004). A proposed injury threshold for mild traumatic brain injury. *J Biomech Eng*, 126(2), 226-236.
- Zoroya, G. (2006, August 8, 2006). Center for war-related brain injuries faces budget cut. *USA Today*.

APPENDICES

Appendix A: Poster presentation at CDMRP Forum 2009 – Kansas City, MO, 9/1/09



VALIDATION OF THE MILITARY ACUTE CONCUSSION EVALUATION (MACE) FOR IN-THEATER EVALUATION OF COMBAT-RELATED TRAUMATIC BRAIN INJURY

Michael A. McCrea, PhD^{1,2}, COL Michael Jaffee, MD³, Katherine Helmick, MS, CNRN, CRNP⁴,
Kevin Guskiewicz, PhD, ATC⁴, Selina Donoiovic, MSN, RN⁴

¹ProHealth Care Neuroscience Center, ²Department of Neurology, Medical College of Wisconsin, ³Defense and Veterans Brain Injury Center, ⁴University of North Carolina at Chapel Hill

BACKGROUND

Traumatic brain injury (TBI) has been widely considered the "signature injury" among United States military personnel involved in combat in Iraq and Afghanistan. Due to several factors, however, the rate of traumatic brain injury in Operation Enduring Freedom (OEF) and Operation Iraqi Freedom (OIF) is thought to be significantly higher than any previous war.

As in the civilian setting, the overwhelming majority of TBIs (> 80%) in the current military conflict are categorized as MTBI based on acute injury characteristics and accepted injury definition criteria. Unfortunately, MTBI presents a unique set of challenges in terms of injury detection, diagnosis, assessment and management due to the more subtle nature of injury characteristics in the absence of classic indicators (e.g., unconsciousness, amnesia, focal neurological deficit, positive neuroimaging findings). Military personnel have explicitly requested a clinical tool to assist in the acute triage of MTBI that is appropriate for the frontline military operational setting.

The Military Acute Concussion Evaluation (MACE) was designed by the Defense and Veterans Brain Injury Center (DVBIC) and civilian brain injury experts specifically for the purposes of assessing and documenting the mechanism of injury, acute characteristics and cognitive deficits in military personnel with suspected MTBI in an austere environment. The MACE was developed by a team of military and civilian TBI experts and first distributed for clinical use by military personnel in August 2006. The instrument is currently the only standardized and most widely used method for evaluation of acute MTBI in military operational settings.

This study was designed to investigate the clinical and operational utility of the MACE in military operational settings. In keeping with the parameters of the TBI Concept Award with respect to prospective study of human subjects, a retrospective review of MACE data already collected on deployed MTBI patients since August 2006 is being executed through access to existing DoD databases.

OBJECTIVES

Through a systematic review of MACE data, the specific aims of this study are:

- **Epidemiological:** To advance our understanding of the acute injury characteristics of MTBI in the current military operational setting (e.g., mechanisms of injury, influence of personal protective equipment (PPE), clinical indicators, severity range)
- **Clinical:** To determine the clinical utility of the MACE in assessing the acute signs and symptoms of MTBI, measuring the acute cognitive effects, and objectively tracking recovery. To assess the unique contribution of the MACE in clinical decision-making and modulating risk around fitness to return to duty after MTBI
- **Operational:** To assess the practical feasibility and user-friendliness of the MACE for medical and other personnel in the military operational setting

METHODS

This is a retrospective data analysis study. Per the eligibility requirements for the Concept Award, this study does not involve prospective collection of data from human subjects. Retrospective data is being extracted from several existing DoD and DVBIC databases. Data from all-severity TBI will initially be collected and stratified across mild, moderate and severe based on established TBI classification systems.

Existing data from injuries evaluated using the MACE between 8/1/06 and 7/31/08 is being systematically analyzed in accordance with the study's specific epidemiological, clinical and operational aims. The target population for this study is military personnel who have sustained a traumatic brain injury in the deployed setting while serving in OEF or OIF. Traumatic brain injury will be defined based on the United States Department of Defense definition.

RESULTS & IMPACT OF RESEARCH

Data analysis is in progress; no formal results or conclusions are available at this time.

As it relates to military application, this study is predicted to directly address valid criticisms currently being voiced by military and civilian clinicians as to the existing gap in established validity of the MACE as the main method used to evaluate mild traumatic brain injury (MTBI) in OEF and OIF.

The findings from this study are also expected to inform future prospective investigations of traumatic brain injury that utilize the MACE and similar methods, particularly for assessment of military-related MTBI.

Because there is currently no widely accepted standard for objectively evaluating MTBI in either general research or clinical environments, findings from this study are predicted to have more global, ground breaking, and translational implications for establishment of a standardized clinical instrument and research tool to evaluate MTBI in settings of mass casualty, breaches to homeland security, and other trauma settings.

*Study funded through the Congressionally-Directed Medical Research Program (CDMRP), U.S. Army Medical Research and Materiel Command (DMRMC), Award Number W81XWH-06-2-0033.

MILITARY ACUTE CONCUSSION EVALUATION (FORM A)

*Adapted from the MACE form in use, which cognitive screening applies from abbreviated assessment of concussion (AAC)