



A Prospective Study of Depression Following Combat Deployment in Support of the Wars in Iraq and Afghanistan

T. S. Wells

C. A. LeardMann, S. O. Fortuna

B. Smith, T. C. Smith

M.A.K. Ryan, E.J. Boyko

D. Blazer



Naval Health Research Center

Report No. 08-05

. Approved for Public Release; Distribution Unlimited.

*Naval Health Research Center
140 Sylvester Road
San Diego, California 92106*

A Prospective Study of Depression Following Combat Deployment in Support of the Wars in Iraq and Afghanistan

Timothy S. Wells, DVM, PhD, MPH, Cynthia A. LeardMann, MPH, Sarah O. Fortuna, MD, Besa Smith, PhD, MPH, Tyler C. Smith, PhD, MS, Margaret A. K. Ryan, MD, MPH, Edward J. Boyko, MD, MPH, and Dan Blazer, MD, PhD, for the Millennium Cohort Study Team

Depression is one of the most prevalent and costly of all public health problems.^{1,2} Estimates of the 12-month and lifetime prevalence for *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM-IV)*³ major depressive disorders is 5.3% to 6.6% and 13.2% to 16.2%, respectively.^{4,5} Much of the burden associated with depression results from comorbidities and reduced ability to maintain employment. Estimates predict that by 2020 depression will be second only to ischemic heart disease as a cause of disability.⁶ Even those with minor depression have higher levels of impairment based on measures of functional status and social health, and incur higher rates of medical services utilization.⁷

Female gender and heredity are reported risk factors for depression.^{4,8} Women are twice as likely to be diagnosed with depression as men, which may be the result of differences in biological vulnerabilities and environmental experiences.⁹ Individuals experiencing threats to personal safety, such as living in unsafe neighborhoods, and individuals living close to the World Trade Center following the events of September 11, 2001, were found to be at increased risk for depression.^{10,11}

Deployments pose unique and often stressful situations to military personnel. Although the association between deployment and post-traumatic stress disorder (PTSD) has been investigated,^{12–20} little is known regarding the risk for depression. One cross-sectional survey that utilized previously studied UK veterans at high risk for mental disorders found depression more common than posttraumatic stress disorder.²¹ Another cross-sectional study utilizing National Comorbidity Survey data found an increased risk for major depressive disorders among male participants with combat experience.²² These findings are supported by other population-based studies that observed an increased risk for mental disorders among 1991 Gulf War veterans,^{23,24} and 1991 Gulf War or

Objective. We investigated relations between deployment and new-onset depression among US service members recently deployed to the wars in Iraq and Afghanistan.

Methods. We included 40219 Millennium Cohort Study participants who completed baseline and follow-up questionnaires and met inclusion criteria. Participants were identified with depression if they met the Primary Care Evaluation of Mental Disorders Patient Health Questionnaire criteria for depression at follow-up, but not at baseline.

Results. Deployed men and women with combat exposures had the highest onset of depression, followed by those not deployed and those deployed without combat exposures. Combat-deployed men and women were at increased risk for new-onset depression compared with nondeployed men and women (men: adjusted odds ratio [AOR]=1.32; 95% confidence interval [CI]=1.13, 1.54; women: AOR=2.13; 95% CI=1.70, 2.65). Conversely, deployment without combat exposures led to decreased risk for new-onset depression compared with those who did not deploy (men: AOR=0.66; 95% CI=0.53, 0.83; women: AOR=0.65; 95% CI=0.47, 0.89).

Conclusions. Deployment with combat exposures is a risk factor for new-onset depression among US service members. Post-deployment screening may be beneficial for US service members exposed to combat. (*Am J Public Health.* 2010;100:90–99. doi:10.2105/AJPH.2008.155432)

Bosnia veterans²⁵; however, none were of longitudinal design.

More recently, members of the US Army and US Marine Corps were at increased risk for depression upon return from combat duties in Afghanistan and Iraq, compared with soldiers who completed questionnaires within 1 week prior to deployment.¹³ However, questionnaires were administered anonymously, making it difficult to identify new-onset depression upon return from deployment. Similarly, a cross-sectional survey of Canadian Forces identified significant associations between depression and exposure to combat or atrocities.²⁶ Finally, depression following deployment may be comorbid with PTSD²⁷ and other mental disorders.²⁵

Per Department of Defense policy, all members are screened for mental disorders before deployment.^{28,29} Screening includes a review of medical records and an interview with a credentialed provider. Diagnosis or symptoms

of depression and PTSD must be assessed for deployment suitability. Disqualifying mental health conditions include, among others, bipolar disorder and psychotic disorders, and disqualifying medications include, but are not limited to, lithium, antipsychotics, and anticonvulsants. Also, those with changes in mental disorder treatment with less than 3 months of stability, or other concerns regarding fitness for duty while deployed, are disqualified.

The Millennium Cohort is the largest-ever population-based prospective study of a military population designed to improve the scientific knowledge of the long-term health effects associated with military service.³⁰ Between 2001 and 2008, 3 panels of US service members were enrolled into the study. Baseline enrollment for the 3 panels ended with 31% of those invited consenting to participate in the 21-year study. The first enrollment cycle was conducted between July 2001 and June 2003, with

TABLE 1—Demographic Characteristics of Participants (N=40219), by Deployment Status: Millennium Cohort Study, 2001–2006

Baseline Characteristic	Men (n = 30 041)			Women (n = 10 178)		
	Not Deployed (n = 22 126), No. (%)	Deployed Without Combat Exposures (n = 3940), No. (%)	Deployed With Combat Exposures (n = 3975), No. (%)	Not Deployed (n = 8543), No. (%)	Deployed Without Combat Exposures (n = 891), No. (%)	Deployed With Combat Exposures (n = 744), No. (%)
Birth year						
Pre-1960	6 599 (29.8)	854 (21.7)	634 (16.0)	2073 (24.3)	100 (11.2)	95 (12.8)
1960–1969	9 371 (42.4)	1813 (46.0)	1663 (41.8)	3147 (36.8)	323 (36.3)	249 (33.5)
1970–1979	5 687 (25.7)	1156 (29.3)	1494 (37.6)	2785 (32.6)	370 (41.5)	321 (43.2)
1980–present	469 (2.1)	117 (3.0)	184 (4.6)	538 (6.3)	98 (11.0)	79 (10.6)
Education						
High school or less	9 342 (42.2)	1309 (33.2)	2025 (50.9)	3532 (41.3)	372 (41.8)	388 (52.2)
Some college	5 597 (25.3)	1517 (38.5)	801 (20.2)	2132 (24.9)	308 (34.6)	134 (18.0)
College degree	7 187 (32.5)	1114 (28.3)	1149 (28.9)	2880 (33.7)	211 (23.7)	222 (29.8)
Marital status						
Never married	4 505 (20.4)	834 (21.2)	1066 (26.8)	3024 (35.4)	386 (43.3)	355 (47.7)
Married	16 546 (74.8)	2897 (73.5)	2727 (68.6)	4460 (52.2)	405 (45.5)	294 (39.5)
Divorced	1 075 (4.9)	209 (5.3)	182 (4.6)	1059 (12.4)	100 (11.2)	95 (12.8)
Race/ethnicity						
Non-Hispanic White	16 344 (73.9)	3021 (76.7)	2754 (69.3)	5391 (63.1)	544 (61.1)	429 (57.7)
Non-Hispanic Black	2 094 (9.5)	348 (8.8)	353 (8.9)	1736 (20.3)	193 (21.7)	145 (19.5)
Other	3 688 (16.7)	571 (14.5)	868 (21.8)	1416 (16.6)	154 (17.3)	170 (22.9)
Smoking						
Never smoker	13 156 (59.5)	2427 (61.6)	2333 (58.7)	5587 (65.4)	581 (65.2)	469 (63.0)
Past smoker	5 710 (25.8)	911 (23.1)	915 (23.0)	1861 (21.8)	174 (19.5)	152 (20.4)
Current smoker	3 260 (14.7)	602 (15.3)	727 (18.3)	1095 (12.8)	136 (15.3)	123 (16.5)
CAGE/alcohol^a						
No	17 884 (80.8)	3197 (81.1)	3203 (80.6)	7595 (88.9)	787 (88.3)	651 (87.5)
Yes	4 242 (19.2)	743 (18.9)	772 (19.4)	948 (11.1)	104 (11.7)	93 (12.5)
Baseline PTSD^b						
No	21 762 (98.4)	3903 (99.1)	3905 (98.2)	8380 (98.1)	884 (99.2)	732 (98.4)
Yes	364 (1.7)	37 (0.9)	70 (1.8)	163 (1.9)	7 (0.8)	12 (1.6)
Military rank						
Enlisted	15 674 (70.8)	2899 (73.6)	2805 (70.6)	5977 (70.0)	715 (80.3)	536 (72.0)
Officer	6 452 (29.2)	1041 (26.4)	1170 (29.4)	2566 (30.0)	176 (19.8)	208 (28.0)
Service component						
Reserve/National Guard	10 057 (45.5)	1561 (39.6)	1456 (36.6)	4401 (51.5)	366 (41.1)	351 (47.2)
Active duty	12 069 (54.6)	2379 (60.4)	2519 (63.4)	4142 (48.5)	525 (58.9)	393 (52.8)
Branch of service						
US Army	10 321 (46.7)	948 (24.1)	2539 (63.9)	4262 (49.9)	342 (38.4)	516 (69.4)
US Air Force	6 113 (27.6)	2125 (53.9)	849 (21.4)	2475 (29.0)	399 (44.8)	153 (20.6)
US Navy/Coast Guard	4 597 (20.8)	720 (18.3)	284 (7.1)	1625 (19.0)	139 (15.6)	65 (8.7)
US Marine Corps	1 095 (5.0)	147 (3.7)	303 (7.6)	181 (2.1)	11 (1.2)	10 (1.3)
Occupational category						
Combat specialists	5 496 (24.8)	927 (23.5)	1301 (32.7)	491 (5.8)	83 (9.3)	65 (8.7)
Health care specialists	1 705 (7.7)	120 (3.1)	311 (7.8)	2198 (25.7)	68 (7.6)	183 (24.6)
Service supply and functional	5 524 (25.0)	828 (21.0)	838 (21.1)	3706 (43.4)	418 (46.9)	261 (35.1)
Other occupations	9 401 (42.5)	2065 (52.4)	1525 (38.4)	2148 (25.1)	322 (36.1)	235 (31.6)

Continued

TABLE 1—Continued

Cumulative length of deployments ^c						
0 d	22 126 (100.0)	...	...	8543 (100.0)	...	...
1-180 d	...	2694 (68.4)	1670 (42.0)	...	567 (63.6)	345 (46.4)
181-270 d	...	757 (19.2)	932 (23.5)	...	165 (18.5)	157 (21.1)
≥271 d	...	489 (12.4)	1373 (34.5)	...	159 (17.9)	242 (32.5)

Note. PTSD = posttraumatic stress disorder. Deployed defined as at least 1 deployment in support of the wars in Iraq and Afghanistan between the baseline and follow-up questionnaire. Percentages are rounded and may not sum to 100.

^aAt baseline, participant self-reported ever feeling at least 1 of the following: (1) a need to cut back on drinking; (2) annoyed at anyone who suggested cutting back on drinking; (3) a need for an “eye-opener,” or early morning drink; and (4) guilty about drinking.

^bSelf-reported a PTSD diagnosis or screened positive for PTSD symptoms at baseline.

^cNumber of total days deployed in support of the wars in Iraq and Afghanistan between the baseline and follow-up questionnaire.

more than 77 000 individuals completing either a postal or Internet questionnaire. Between 2004 and 2006, approximately 71% of the first panel members completed a follow-up questionnaire. Investigations of potential reporting biases showed no differential in health care utilization in the year prior to enrollment³¹; strong test–retest reliability³²; reliable reporting of vaccinations,^{33,34} occupations,³⁵ and deployments³⁶; and minimal differences between participants choosing submission by Internet survey or paper submission.³⁷ Analyses of potential responder bias in ongoing follow-ups continue.

The Millennium Cohort questionnaire includes, among other questions, mental health measures based on 2 standardized scoring instruments: the Primary Care Evaluation of Mental Disorders Patient Health Questionnaire (PHQ)³⁸ and the Medical Outcomes Study Short Form 36-Item Health Survey for Veterans.³⁹ We used Millennium Cohort Study data to investigate the association of new-onset depression and deployment in support of the wars in Iraq and Afghanistan.

METHODS

The Millennium Cohort Study was launched in 2001 to collect and evaluate population-based data on behavioral and occupational risks related to military service that may be associated with poor health.^{30,40} The first panel of invited Millennium Cohort Study participants was randomly selected from all US military personnel serving in October 2000. To ensure sufficient power to detect differences in smaller subgroups, those with previous deployment to Kosovo, Bosnia, or Southwest Asia; Reserve and National Guard members; and women were

oversampled. Through a modified Dillman approach, 77 047 consenting participants enrolled in the first panel of the Millennium Cohort Study.⁴¹ The methodology of the study has been described elsewhere in detail.³⁰ Of the 77 047 enrolled in this first panel, 55 021 (71.4%) completed the follow-up questionnaire between June 2004 and February 2006. Those completing the follow-up questionnaire were slightly more likely to be born in 1969 or earlier, married, more highly educated, officers, and non-Hispanic White compared with participants who completed the baseline survey.

The population for this study included participants in the first panel of Millennium Cohort Study who had complete demographic, behavioral, and military-specific data and who completed the baseline and follow-up questionnaires. We excluded participants from the final sample if, at baseline, they reported ever having a diagnosis for depression, met the criteria for mild or other depression, or reported taking medicine for anxiety, stress, or depression. We also excluded those completing the baseline questionnaire after deployment, completing either questionnaire while on deployment, or not answering questions on depression diagnosis or symptoms or use of medication for anxiety, stress, or depression.

The Defense Manpower Data Center provided demographic and military-specific data from electronic personnel files, including gender, birth date, highest education level, marital status, race/ethnicity, deployment experience, pay grade, service component (active duty or Reserve/National Guard), service branch (US Army, US Air Force, US Navy, US Coast Guard, or US Marine Corps), and military occupations. Race/ethnicity was classified by the Defense Manpower Data

Center, and was incorporated into analyses because of differences in prevalence and incidence for depression by race/ethnicity.^{42,43}

Depression Data

The PHQ, embedded in the Millennium Cohort questionnaire, provides a psychosocial assessment based on scores of several health concepts.^{38,44,45} For this study, we used the PHQ to assess depression. Sensitivity and specificity have been reported as high for the PHQ-defined major depressive disorder (sensitivity=0.93; specificity=0.89).⁴⁶ Major depressive disorder, as measured by 9 items from the PHQ, corresponds to the depression diagnosis from the *DSM-IV*.⁴⁷ Participants used a 4-point Likert scale to rate the severity of each depressive symptom from “not at all” to “nearly every day” during the previous 2 weeks prior to questionnaire completion. For this investigation, we defined participants as having new-onset depression at follow-up if they met the following 2 criteria: (1) endorsed having a depressed mood or anhedonia and (2) responded “more than half the days” or “nearly every day” to at least 5 of the 9 items, where suicidal ideation was counted if present at all.⁴⁴

Deployment Status

We considered participants who had completed a deployment in support of the wars in Iraq and Afghanistan between the baseline and follow-up questionnaires to be deployed. We further categorized deployment status on the basis of self-reported exposure to combat at follow-up. We classified participants as deployed with combat exposure if they reported at least one combat experience in the past 3 years, including witnessing death, trauma,

TABLE 2—Percentage of New-Onset Depression Among Male Participants, by Deployment Status: Millennium Cohort Study, 2001–2006

Baseline Characteristics	Not Deployed, No. (%)	Deployed Without Combat Exposures, No. (%)	Deployed With Combat Exposures, No. (%)
Total	872 (3.9)	92 (2.3)	225 (5.7)
Birth year			
Pre-1960	219 (3.3)	22 (2.6)	26 (4.1)
1960–1969	346 (3.7)	38 (2.1)	85 (5.1)
1970–1979	280 (4.9)	26 (2.3)	90 (6.0)
1980–present	27 (5.8)	6 (5.1)	24 (13.0)
Education			
High school or less	471 (5.0)	48 (3.7)	165 (8.2)
Some college	212 (3.8)	26 (1.7)	38 (4.7)
College degree	189 (2.6)	18 (1.6)	22 (1.9)
Marital status			
Never married	211 (4.7)	22 (2.6)	76 (7.1)
Married	610 (3.7)	63 (2.2)	136 (5.0)
Divorced	51 (4.7)	7 (3.4)	13 (7.1)
Race/ethnicity			
Non-Hispanic White	671 (4.1)	71 (2.4)	167 (6.1)
Non-Hispanic Black	85 (4.1)	6 (1.7)	22 (6.2)
Other	116 (3.2)	15 (2.6)	36 (4.2)
Smoking			
Never smoker	422 (3.2)	50 (2.1)	107 (4.6)
Past smoker	247 (4.3)	22 (2.4)	51 (5.6)
Current smoker	203 (6.2)	20 (3.3)	67 (9.2)
CAGE/alcohol ^a			
No	655 (3.7)	73 (2.3)	176 (5.5)
Yes	217 (5.1)	19 (2.6)	49 (6.4)
Baseline PTSD ^b			
No	809 (3.7)	87 (2.2)	209 (5.4)
Yes	63 (17.3)	5 (13.5)	16 (22.9)
Military rank			
Enlisted	730 (4.7)	73 (2.5)	199 (7.1)
Officer	142 (2.2)	19 (1.8)	26 (2.2)
Service component			
Reserve/National Guard	404 (4.0)	34 (2.2)	94 (6.5)
Active duty	468 (3.9)	58 (2.4)	131 (5.2)
Branch of service			
US Army	485 (4.7)	31 (3.3)	170 (6.7)
US Air Force	185 (3.0)	35 (1.7)	32 (3.8)
US Navy/Coast Guard	160 (3.5)	21 (2.9)	11 (3.9)
US Marine Corps	42 (3.8)	5 (3.4)	12 (4.0)
Occupational category			
Combat specialists	188 (3.4)	15 (1.6)	55 (4.2)
Health care specialists	81 (4.8)	4 (3.3)	18 (5.8)
Service supply and functional	233 (4.2)	24 (2.9)	50 (6.0)
Other occupations	370 (3.9)	49 (2.4)	102 (6.7)

Continued

prisoners of war, or refugees. We identified those who deployed and did not report any of these experiences in the past 3 years as deployed without combat exposure.

Posttraumatic Stress Disorder

The Millennium Cohort questionnaire includes the PTSD Checklist–Civilian Version, a 17-item self-report measure of PTSD symptoms.^{48,49} Participants used a 5-point Likert scale (from 1=not at all to 5=extremely) to rate the severity of each intrusion, avoidance, and hyperarousal symptom during the past 30 days. Following standard procedures, we scored participants as having PTSD symptoms at baseline if they reported a moderate or higher level of at least 1 intrusion symptom, 3 avoidance symptoms, and 2 hyperarousal symptoms at baseline.³ We defined participants as having a PTSD diagnosis at baseline if they reported ever having a diagnosis of PTSD from a health care professional.

Statistical Analysis

We completed a descriptive investigation to examine deployment status; demographic, behavioral, and occupational characteristics; and baseline PTSD symptoms or diagnosis with new-onset depression, stratified by gender. To calculate associations between deployment status and new-onset depression, we used multivariable logistic regression, adjusting for demographic, behavioral, and occupational variables, as well as baseline PTSD symptoms or diagnosis. We defined alcohol misuse by positive endorsement of 1 or more CAGE (cut down, annoyed, guilt, eye-opener) questions,⁵⁰ which is noted as “CAGE/alcohol” hereafter. Because of known associations between smoking and depression,^{51,52} and smoking and deployment,⁵³ there were concerns these relations may interact. We investigated this by entering a first-order multiplicative interaction term of deployment by smoking status into the regression model. We performed regression diagnostics, including examining covariates for multicollinearity and goodness-of-fit tests. We conducted all data analyses with SAS version 9.1.3 (SAS Institute Inc, Cary, NC).

RESULTS

There were 55 021 Millennium Cohort participants who completed the 2001–2003 and

TABLE 2—Continued

Cumulative length of deployments ^c			
0 d	872 (3.9)	...	...
1–180 d	...	57 (2.1)	96 (5.8)
181–270 d	...	23 (3.0)	53 (5.7)
≥271 d	...	12 (2.5)	76 (5.5)

Note. PTSD = posttraumatic stress disorder. Deployed defined as at least 1 deployment in support of the wars in Iraq and Afghanistan between the baseline questionnaire and follow-up questionnaire.

^aAt baseline, participant self-reported ever feeling at least 1 of the following: (1) a need to cut back on drinking; (2) annoyed at anyone who suggested cutting back on drinking; (3) a need for an “eye-opener,” or early morning drink; and (4) guilty about drinking.

^bSelf-reported a PTSD diagnosis or screened positive for PTSD symptoms at baseline.

^cNumber of total days deployed in support of the wars in Iraq and Afghanistan between the baseline and follow-up questionnaire.

2004–2006 questionnaires. We excluded individuals with indicators of depression at baseline (n=6537), who deployed prior to completing the 2001–2003 questionnaire (n=1853), who completed a questionnaire

while deployed (n=2553), or who did not have complete data (n=3859), leaving 40 219 Cohort members for inclusion in this study. We conducted separate analyses for the 30 041 men and 10 178 women who met study criteria.

We categorized men and women as not deployed (22 126 men and 8543 women), deployed without combat exposures (3940 men and 891 women), or deployed with combat exposures (3975 men and 744 women). Study participants had, on average, completed the baseline survey 404.1 days prior to deployment (range=1 to 1402 days), and had completed the follow-up survey 349.4 days, on average, after returning from deployment (range=1 to 1366 days).

Compared with nondeployers, deployed men and women were proportionately more likely to be younger and serving on active duty (Table 1). Men and women deployed without combat exposures were proportionately more likely to have some college education, be in the US Air Force, and have occupations listed as “other” (i.e., all other occupations excluding combat, health care, service, supply, and functional specialists) compared with the other groups. Conversely, men and women deployed with combat exposures were proportionately more likely to have a high school education or less, be serving in the US Army, and to be combat specialists, compared with nondeployed men and women. Deployed women were more likely to have never been married compared with nondeployed women. In addition, deployed men and women who reported combat exposures were proportionately more likely to have been deployed 181 days or more compared with those deployed without combat exposures.

Men and women deployed with combat exposures had the highest occurrence of new-onset depression, (5.7% and 15.7%, respectively), followed by those not deployed (3.9% and 7.7%, respectively), whereas participants deployed without combat exposures exhibited the lowest occurrence (2.3% and 5.1%, respectively; Tables 2 and 3). This trend remained consistent when we assessed demographic, behavioral, and military-related characteristics for new-onset depression (Tables 2 and 3). Across all 3 groups, men with the highest percentages of new-onset depression were proportionately more likely to be born in 1980 or later, less educated, other than married, current smokers, enlisted, and serving in the US Marine Corps or US Army. Furthermore, they were more likely to positively endorse CAGE/alcohol questions and

TABLE 3—Percentage of New-Onset Depression Among Female Participants, by Deployment Status: Millennium Cohort Study, 2001–2006

Baseline Characteristics	Not Deployed, No. (%)	Deployed Without Combat Exposures, No. (%)	Deployed With Combat Exposures, No. (%)
Total	654 (7.7)	45 (5.1)	117 (15.7)
Birth year			
Pre-1960	125 (6.0)	3 (3.0)	19 (20.0)
1960–1969	212 (6.7)	13 (4.0)	32 (12.9)
1970–1979	258 (9.3)	24 (6.5)	52 (16.2)
1980–present	59 (11.0)	5 (5.1)	14 (17.7)
Education			
High school or less	326 (9.2)	25 (6.7)	73 (18.8)
Some college	161 (7.6)	14 (4.6)	18 (13.4)
College degree	167 (5.8)	6 (2.8)	26 (11.7)
Marital status			
Never married	234 (7.7)	18 (4.7)	49 (13.8)
Married	327 (7.3)	22 (5.4)	54 (18.4)
Divorced	93 (8.8)	5 (5.0)	14 (14.7)
Race/ethnicity			
Non-Hispanic White	449 (8.3)	30 (5.5)	74 (17.3)
Non-Hispanic Black	108 (6.2)	9 (4.7)	26 (17.9)
Other	97 (6.9)	6 (3.9)	17 (10.0)
Smoking			
Never smoker	363 (6.5)	24 (4.1)	68 (14.5)
Past smoker	175 (9.4)	10 (5.8)	25 (16.5)
Current smoker	116 (10.6)	11 (8.1)	24 (19.5)

Continued

TABLE 3—Continued

CAGE/alcohol ^a			
No	560 (7.4)	38 (4.8)	95 (14.6)
Yes	94 (9.9)	7 (6.7)	22 (23.7)
Baseline PTSD ^b			
No	615 (7.3)	44 (5.0)	114 (15.6)
Yes	39 (23.9)	1 (14.3)	3 (25.0)
Military rank			
Enlisted	517 (8.7)	43 (6.0)	98 (18.3)
Officer	137 (5.3)	2 (1.1)	19 (9.1)
Service component			
Reserve/National Guard	312 (7.1)	20 (5.5)	54 (15.4)
Active duty	342 (8.3)	25 (4.8)	63 (16.0)
Branch of service			
US Army	361 (8.5)	23 (6.7)	89 (17.3)
US Air Force	147 (5.9)	16 (4.0)	16 (10.5)
US Navy/Coast Guard	133 (8.2)	6 (4.3)	12 (18.5)
US Marine Corps	13 (7.2)	0 (0.0)	0 (0.0)
Occupational category			
Combat specialists	37 (7.5)	3 (3.6)	5 (7.7)
Health care specialists	158 (7.2)	5 (7.4)	18 (9.8)
Service supply and functional	281 (7.6)	18 (4.3)	59 (22.6)
Other occupations	178 (8.3)	19 (5.9)	35 (14.9)
Cumulative length of deployments ^c			
0 d	654 (7.7)	...	...
1-180 d	...	26 (4.6)	45 (13.0)
181-270 d	...	7 (4.2)	23 (14.7)
≥ 271 d	...	12 (7.6)	49 (20.3)

Note. PTSD = posttraumatic stress disorder. Deployed defined as at least 1 deployment in support of the wars in Iraq and Afghanistan between the baseline questionnaire and follow-up questionnaire.

^aAt baseline, participant self-reported ever feeling at least 1 of the following: (1) a need to cut back on drinking; (2) annoyed at anyone who suggested cutting back on drinking; (3) a need for an "eye-opener," or early morning drink; and (4) guilty about drinking.

^bSelf-reported a PTSD diagnosis or screened positive for PTSD symptoms at baseline.

^cNumber of total days deployed in support of the wars in Iraq and Afghanistan between the baseline and follow-up questionnaire.

have PTSD symptoms or diagnosis at baseline. Similar to men, women with the highest odds of new-onset depression across all 3 deployment categories included those who were more likely to be current smokers, positively endorse CAGE/alcohol questions, have PTSD symptoms or diagnosis at baseline, be enlisted, and have a high school education or less (Table 3).

After we adjusted for deployment status, birth year, education, marital status, race/ethnicity, smoking status, positive CAGE/alcohol question responses, baseline PTSD symptoms or diagnosis, rank, service component, service branch, and occupation, both men and women

deployed with combat exposure were significantly more likely to develop new-onset depression than were those not deployed (odds ratio [OR]=1.32; 95% confidence interval [CI]=1.13, 1.54; and OR=2.13; 95% CI=1.70, 2.65, respectively; Table 4). In contrast, men and women deployed without combat exposures were less likely to develop new-onset depression than were those who did not deploy (OR=0.66; 95% CI=0.53, 0.83; and OR=0.65; 95% CI=0.47, 0.89, respectively). In the adjusted model, men and women who were past or current smokers, had PTSD symptoms or diagnosis at baseline, positively endorsed CAGE/alcohol questions, or were

enlisted had higher odds for new-onset depression. Additionally, women with new-onset depression were more likely to be born in 1970 or later, married or divorced, non-Hispanic White, active duty, and in the US Army or US Navy/Coast Guard. Men who were born in 1980 or later, served in the US Army, and worked in health care, service supply, or functional occupations had higher odds for new-onset depression.

No interaction was observed between smoking status and deployment status, nor was there interaction between baseline PTSD symptoms or diagnosis and deployment as related to depression. Additionally, length of deployment was collinear with deployment status, and excluded from the main analysis. However, to assess the significance of deployment length, we conducted a subanalysis incorporating deployment length with data of those deployed to examine the association between deployment length and new-onset depression. In this analysis, deployment length was not statistically significant.

DISCUSSION

We used Millennium Cohort Study data to identify more than 30 000 male and 10 000 female US service members who were free of depression at baseline. To our knowledge, this is the first population-based longitudinal study with baseline data prior to deployment to report a temporal association between combat exposure among deployed US service members and new-onset depression compared with nondeployed peers. However, finding an increased risk for a mental disorder associated with combat exposure is not unique, as Smith et al. observed similar findings for PTSD with Millennium Cohort data.¹⁴

Other research, though cross-sectional in design, has found similar depression risk with exposure to combat or witnessing atrocities among members of the Canadian military.²⁶ Although PTSD was associated with increased length of deployment among Vietnam veterans⁵⁴ and UK veterans serving in Afghanistan or Iraq,⁵⁵ this trend was not found for depression in this study. Additionally, a large historical prospective study found an inverse association between mental health hospitalizations and US servicewomen who were members of the US

TABLE 4—Multivariable Logistic Regression Adjusted Odds Ratio (AOR) of Depression, by Gender: Millennium Cohort Study, 2001–2006

Baseline Characteristics	Men, AOR (95% CI)	Women, AOR (95% CI)
2001–2006 deployment status ^a		
No deployment (Ref)	1.00	1.00
Deployed without combat exposures	0.66 (0.53, 0.83)	0.65 (0.47, 0.89)
Deployed with combat exposures	1.32 (1.13, 1.54)	2.13 (1.70, 2.65)
Birth year		
Pre-1960 (Ref)	1.00	1.00
1960–1969	1.06 (0.90, 1.24)	1.00 (0.80, 1.24)
1970–1979	1.20 (0.99, 1.45)	1.39 (1.10, 1.76)
1980–present	1.56 (1.10, 2.21)	1.61 (1.13, 2.28)
Education		
High school or less (Ref)	1.00	1.00
Some college	0.90 (0.75, 1.07)	1.03 (0.84, 1.27)
College degree	0.85 (0.68, 1.07)	1.10 (0.85, 1.43)
Marital status		
Never married (Ref)	1.00	1.00
Married	0.99 (0.84, 1.17)	1.25 (1.05, 1.49)
Divorced	1.22 (0.92, 1.62)	1.49 (1.16, 1.92)
Race/ethnicity		
Non-Hispanic White (Ref)	1.00	1.00
Non-Hispanic Black	0.88 (0.72, 1.08)	0.73 (0.59, 0.90)
Other	0.84 (0.71, 1.01)	0.75 (0.60, 0.93)
Smoking		
Never smoker (Ref)	1.00	1.00
Past smoker	1.18 (1.02, 1.36)	1.30 (1.09, 1.55)
Current smoker	1.52 (1.31, 1.77)	1.35 (1.10, 1.66)
CAGE/alcohol ^b		
No (Ref)	1.00	1.00
Yes	1.19 (1.04, 1.37)	1.27 (1.03, 1.57)
Baseline PTSD ^c		
No (Ref)	1.00	1.00
Yes	4.29 (3.34, 5.50)	2.98 (2.07, 4.28)
Military rank		
Enlisted	1.67 (1.31, 2.11)	1.65 (1.26, 2.16)
Officer (Ref)	1.00	1.00
Service component		
Reserve/National Guard (Ref)	1.00	1.00
Active duty	1.01 (0.88, 1.15)	1.21 (1.03, 1.42)
Branch of service		
US Army	1.52 (1.27, 1.82)	1.64 (1.33, 2.01)
US Air Force (Ref)	1.00	1.00
US Navy/Coast Guard	1.13 (0.91, 1.40)	1.55 (1.21, 2.00)
US Marine Corps	1.11 (0.81, 1.52)	0.88 (0.48, 1.60)
Occupational category		
Combat specialists (Ref)	1.00	1.00
Health care specialists	1.52 (1.20, 1.93)	1.09 (0.77, 1.55)

Continued

Navy or US Marine Corps serving in combat support occupations.⁵⁶ These contrasting observations may be caused by underlying differences in PTSD and depression risk factors, or because the Vietnam study was conducted on a relatively small sample many years after the war's end, making it difficult to rule out misclassification of exposure or outcome.

In this study, deployed men and women without combat exposures were significantly less likely to develop depression than were nondeployed men and women. This finding may suggest that individuals who are healthier and at less risk for depression are more likely to deploy and, thus, mitigate stress-related effects of deployment^{57–60} among men and women who do not face combat exposures. In other words, deployment with low probability of exposure to combat (or the effects of combat, such as being in the line of fire or handling casualties, prisoners, or treating the wounded) may not be significantly stressful to the majority of US service members.

Military hardiness, the context-specific adaptation of psychological hardiness,⁶¹ has been observed to moderate the impact of deployment stressors on depression.⁶² Military hardiness may also partially explain the observation that male combat specialists were at lower risk for depression than either health care specialists or men whose duties include service supply and functional support, after adjustment for a number of variables, including combat exposure. It is possible that combat specialists are more mentally prepared for deployment-related stresses either through training or selection factors for their positions. Medical personnel have significant exposure to the after effects of the trauma of combat but may not be as well-prepared to cope with the effects of this exposure because of different training or constitutional make-up. Another possibility is that the range and type of combat stressors medical personnel experience are fundamentally different in unmeasured ways than those experienced by combat specialists. Men working in service supply and functional-support occupations conduct convoy and other duties that randomly and frequently put them in dangerous situations, and similarly may not have the advantages of combat specialist training or constitution.

The reporting of baseline PTSD symptoms or diagnosis was associated with new-onset

TABLE 4—Continued

Service supply and functional	1.20 (1.01, 1.43)	1.09 (0.78, 1.54)
Other occupations	1.09 (0.93, 1.28)	1.04 (0.74, 1.47)

Notes. CI = confidence interval; PTSD = posttraumatic stress disorder. Odds ratios for men and women are adjusted for all variables listed.

^aDeployment defined as at least 1 deployment in support of the wars in Iraq and Afghanistan between the baseline questionnaire and follow-up questionnaire. Combat exposure defined as reporting at least 1 combat-like exposure on the follow-up questionnaire.

^bAt baseline, participant self-reported ever feeling at least 1 of the following: (1) a need to cut back on drinking; (2) annoyed at anyone who suggested cutting back on drinking; (3) a need for an “eye-opener,” or early morning drink; and (4) guilty about drinking.

^cSelf-reported a PTSD diagnosis or screened positive for PTSD symptoms at baseline.

depression in adjusted models for both female and male US service members. Finding comorbid PTSD and depression following exposure to traumatic events has received significant attention in the scientific literature.^{63–68} However, the causes of this comorbidity remain poorly understood.

One study found that trauma variables known to be related to PTSD development, such as type and horror of the trauma, were also related to the occurrence of comorbid mental disorders, including depression, generalized anxiety disorder, and agoraphobia.⁶⁹ Similarly, depression was observed to develop with or without comorbid PTSD following exposure to a traumatic event.^{66,70} Some believe that PTSD and depression comorbidity can be largely explained by common genetic influences.^{71,72} Further research to unravel this comorbidity is paramount as comorbid PTSD and depression have been reported to increase dysfunction,^{63,64} health care utilization,⁶⁵ and the risk for suicide.^{65,68} Data from the Millennium Cohort Study have contributed to the understanding of PTSD among US military populations.^{14,15,73} Following the publication of this article and as additional cohort data become available, subsequent analyses will investigate PTSD and depression comorbidity.

Other risk factors for depression observed in this study were consistent with previous reports that used data from military and civilian populations.^{4,5,73–75} In adjusted models, male and female US service members were at increased risk for depression if they were younger, past or current smokers, positively endorsed CAGE/alcohol questions, and had served in the US Army. Additionally, women who were married or divorced, non-Hispanic White, active duty, and served in the US Navy/Coast Guard

were at increased risk of depression. The increased risk of depression among married women compared with never-married women may be related to postpartum depression,⁷⁶ separation anxiety associated with leaving family members while deployed, or a lack of social support in the deployed setting. Although few studies have examined increased separation anxiety or increased lack of social support among deployed married women, animal studies provide evidence that separation may induce depression,^{77,78} and studies have found an association between sense of belonging and depression.^{79,80} It is also plausible that differences exist in role expectations and family or social support for married male and female deployed service members. For example, married women may be expected to resume routine family care while attempting to cope with the events of the deployment, more so than their male counterparts.

Limitations to these analyses should be noted. A population accounting for 31% of invited cohort members as well as self-report of exposures and symptoms has the potential to introduce bias. However, multiple investigations of possible reporting and selection biases in baseline Millennium Cohort data suggest reliable reporting, minimal response bias, and a representative sample of military personnel, including deployers.^{14,15,30–35,37,73,81} Use of the PHQ along with the *DSM-IV* criteria has shown to correlate well with a physician's assessment of depression symptoms,³² and the PHQ is internally valid in Millennium Cohort members.³² However, the use of a standardized instrument for self-reported data as a surrogate for depression diagnosis is imperfect. Finally, the definition of combat exposure in this study was quite broad, encompassing a number of stressful

events that often accompany combat, but do not necessarily indicate actual engagement in combat operations (i.e., exchanging fire with an enemy).

Despite limitations, these analyses offer the first large, prospective epidemiological investigation of new-onset depression symptoms in combat-deployed military men and women. The large sample of both men and women along with adjustment for multiple potential confounding variables allowed for a robust investigation of the association between deployment and new-onset depression. Depression is often underreported in electronic health care databases among populations that do not readily access care for mental disorders, making it often advantageous to use self-reported measures.

In summary, male and female US service members who deployed and reported combat exposures were at increased risk for depression compared with nondeployed service members, after adjustment for baseline PTSD symptoms or diagnosis and other potentially confounding variables. Conversely, men and women who deployed and did not report combat exposures were at lower risk for depression than nondeployed men and women. These findings support hypotheses that stress associated with combat may lead to depression, and that stress related to deployment, in the absence of combat exposure, may be mitigated by selective deployment of service members who are at decreased risk for the development of depression in comparison with nondeployed men and women. ■

About the Authors

Timothy S. Wells and Sarah O. Fortuna are with the US Air Force Research Laboratory, Wright-Patterson AFB, OH. Cynthia A. LeardMann, Besa Smith, Tyler C. Smith, and Margaret A.K. Ryan are with the Department of Defense Center for Deployment Health Research, Naval Health Research Center, San Diego, CA. Edward J. Boyko is with the Seattle Epidemiologic Research and Information Center, Veterans Affairs Puget Sound Health Care System, Seattle, WA. Dan Blazer is with Duke University Medical Center, Durham, NC.

Correspondence should be sent to Timothy S. Wells, 711th HPW/RHPA, Building 824 Room 206, 2800 Q St, Wright-Patterson AFB, OH 45433-7947 (e-mail: timothy.wells@wpafb.af.mil). Reprints can be ordered at <http://www.ajph.org> by clicking the “Reprints/Eprints” link.

This article was accepted on April 12, 2009.

Contributors

T. S. Wells led the design, analyses, and article development. C. A. LeardMann contributed to the design, conducted

most of the analyses, and assisted with article writing. S.O. Fortuna assisted with design, made recommendations during the analyses, and wrote the introduction of the article. B. Smith, T. C. Smith, M.A.K. Ryan, E.J. Boyko, and D. Blazer made significant contributions to the design, helped to guide analyses, and provided critical review of the article.

Acknowledgments

This work represents report 08-06, supported by the Department of Defense, under work unit no. 60002. The views expressed in this article are those of the authors and do not reflect the official policy or position of the Department of the Navy, Department of the Army, Department of the Air Force, Department of Defense, Department of Veterans Affairs, or the US Government.

In addition to the authors, the Millennium Cohort Study Team includes Lacy Farnell, Gia Gumbs, Isabel Jacobson, Molly Kelton, Travis Leleu, Jamie McGrew, Katherine Snell, Steven Spiegel, Kari Welch, Martin White, James Whitmer, and Charlene Wong from the Department of Defense Center for Deployment Health Research, Naval Health Research Center, San Diego, CA; Paul J. Amoroso, from the Madigan Army Medical Center, Tacoma, WA; Gary D. Gackstetter, from the Department of Preventive Medicine and Biometrics, Uniformed Services University of the Health Sciences, Bethesda, MD, and Analytic Services Inc, Arlington, VA; Gregory C. Gray, from the College of Public Health, University of Iowa, Iowa City; Tomoko I. Hooper, from the Department of Preventive Medicine and Biometrics, Uniformed Services University of the Health Sciences; and James R. Riddle, from the US Air Force Research Laboratory, Wright-Patterson Air Force Base, OH. We are indebted to the Millennium Cohort Study participants, without whom these analyses would not be possible. We thank Scott L. Seggerman and Greg D. Boyd from the Management Information Division, Defense Manpower Data Center, Seaside, CA. Additionally, we thank Michelle Stoia from the Naval Health Research Center. We also thank the professionals from the US Army Medical Research and Materiel Command, especially those from the Military Operational Medicine Research Program, Fort Detrick, MD. We appreciate the support of the staff of the Henry M. Jackson Foundation for the Advancement of Military Medicine, Rockville, MD.

Human Participant Protection

This research has been conducted in compliance with all applicable federal regulations governing the protection of human subjects in research, and was approved by the institutional review board, Naval Health Research Center, San Diego, CA (Protocol NHRC.2000.007).

References

1. Stoudemire A, Frank R, Hedemark N, Kamlet M, Blazer D. The economic burden of depression. *Gen Hosp Psychiatry*. 1986;8(6):387–394.
2. Greenberg PE, Kessler RC, Birnbaum HG, et al. The economic burden of depression in the United States: how did it change between 1990 and 2000? *J Clin Psychiatry*. 2003;64(12):1465–1475.
3. *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition*. Washington, DC: American Psychiatric Association; 1994.
4. Hasin DS, Goodwin RD, Stinson FS, Grant BF. Epidemiology of major depressive disorder: results from the National Epidemiologic Survey on Alcoholism and Related Conditions. *Arch Gen Psychiatry*. 2005;62(10):1097–1106.
5. Kessler RC, Berglund P, Demler O, et al. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003;289(23):3095–3105.
6. Murray CJ, Lopez AD. Alternative projections of mortality and disability by cause 1990–2020: Global Burden of Disease Study. *Lancet*. 1997;349(9064):1498–1504.
7. Wagner HR, Burns BJ, Broadhead WE, Yarnall KS, Sigmon A, Gaynes BN. Minor depression in family practice: functional morbidity, co-morbidity, service utilization and outcomes. *Psychol Med*. 2000;30(6):1377–1390.
8. Ebmeier KP, Donaghey C, Steele JD. Recent developments and current controversies in depression. *Lancet*. 2006;367(9505):153–167.
9. Kessler RC. Epidemiology of women and depression. *J Affect Disord*. 2003;74(1):5–13.
10. Cutrona CE, Wallace G, Wesner KA. Neighborhood characteristics and depression: an examination of stress processes. *Curr Dir Psychol Sci*. 2006;15(4):188–192.
11. Galea S, Ahern J, Resnick H, et al. Psychological sequelae of the September 11 terrorist attacks in New York City. *N Engl J Med*. 2002;346(13):982–987.
12. Hoge CW, Auchterlonie JL, Milliken CS. Mental health problems, use of mental health services, and attrition from military service after returning from deployment to Iraq or Afghanistan. *JAMA*. 2006;295(9):1023–1032.
13. Hoge CW, Castro CA, Messer SC, McGurk D, Cotting DI, Koffman RL. Combat duty in Iraq and Afghanistan, mental health problems, and barriers to care. *N Engl J Med*. 2004;351(1):13–22.
14. Smith TC, Ryan MA, Wingard DL, Slymen DJ, Sallis JF, Kritz-Silverstein D. New onset and persistent symptoms of post-traumatic stress disorder self reported after deployment and combat exposures: prospective population based US military cohort study. *BMJ*. 2008;336(7640):366–371.
15. Smith TC, Wingard DL, Ryan MA, Kritz-Silverstein D, Slymen DJ, Sallis JF. Prior assault and posttraumatic stress disorder after combat deployment. *Epidemiology*. 2008;19(3):505–512.
16. Milliken CS, Auchterlonie JL, Hoge CW. Longitudinal assessment of mental health problems among active and reserve component soldiers returning from the Iraq war. *JAMA*. 2007;298(18):2141–2148.
17. Ismail K, Kent K, Brugha T, et al. The mental health of UK Gulf war veterans: phase 2 of a two phase cohort study. *BMJ*. 2002;325(7364):576.
18. Unwin C, Blatchley N, Coker W, et al. Health of UK servicemen who served in Persian Gulf War. *Lancet*. 1999;353(9148):169–178.
19. Orcutt HK, Erickson DJ, Wolfe J. A prospective analysis of trauma exposure: the mediating role of PTSD symptomatology. *J Trauma Stress*. 2002;15(3):259–266.
20. Orcutt HK, Erickson DJ, Wolfe J. The course of PTSD symptoms among Gulf War veterans: a growth mixture modeling approach. *J Trauma Stress*. 2004;17(3):195–202.
21. Iversen A, Dyson C, Smith N, et al. 'Goodbye and good luck': the mental health needs and treatment experiences of British ex-service personnel. *Br J Psychiatry*. 2005;186:480–486.
22. Prigerson HG, Maciejewski PK, Rosenheck RA. Population attributable fractions of psychiatric disorders and behavioral outcomes associated with combat exposure among US men. *Am J Public Health*. 2002;92(1):59–63.
23. The Iowa Persian Gulf Study Group. Self-reported illness and health status among Gulf War veterans. *JAMA*. 1997;277(3):238–245.
24. Dlugosz LJ, Hocter WJ, Kaiser KS, et al. Risk factors for mental disorder hospitalization after the Persian Gulf War: U.S. Armed Forces, June 1, 1991–September 30, 1993. *J Clin Epidemiol*. 1999;52(12):1267–1278.
25. David AS, Farrin L, Hull L, Unwin C, Wessely S, Wykes T. Cognitive functioning and disturbances of mood in UK veterans of the Persian Gulf War: a comparative study. *Psychol Med*. 2002;32(8):1357–1370.
26. Sareen J, Cox BJ, Afifi TO, et al. Combat and peacekeeping operations in relation to prevalence of mental disorders and perceived need for mental health care: findings from a large representative sample of military personnel. *Arch Gen Psychiatry*. 2007;64(7):843–852.
27. Erickson DJ, Wolfe J, King DW, King LA, Sharkansky EJ. Posttraumatic stress disorder and depression symptomatology in a sample of Gulf War veterans: a prospective analysis. *J Consult Clin Psychol*. 2001;69(1):41–49.
28. Individual protection and individual unit deployment policy, Oct 2001 (with updates through Jan 2009) [policy letter]. MacDill AFB, FL: US Department of Defense, Commander, Central Command; 2009.
29. Policy guidance for deployment-limiting psychiatric conditions and medications, Nov 2006 [policy letter]. Washington, DC: US Department of Defense, Assistant Secretary of Defense for Health Affairs; 2006.
30. Ryan MA, Smith TC, Smith B, et al. Millennium Cohort: enrollment begins a 21-year contribution to understanding the impact of military service. *J Clin Epidemiol*. 2007;60(2):181–191.
31. Wells TS, Jacobson IG, Smith TC, et al. Prior health care utilization as a potential determinant of enrollment in a 21-year prospective study, the Millennium Cohort Study. *Eur J Epidemiol*. 2008;23(2):79–87.
32. Smith TC, Smith B, Jacobson IG, Corbeil TE, Ryan MA. Reliability of standard health assessment instruments in a large, population-based cohort study. *Ann Epidemiol*. 2007;17(7):525–532.
33. Smith B, Leard CA, Smith TC, Reed RJ, Ryan MA. Anthrax vaccination in the Millennium Cohort: validation and measures of health. *Am J Prev Med*. 2007;32(4):347–353.
34. LeardMann CA, Smith B, Smith TC, Wells TS, Ryan MA. Smallpox vaccination: comparison of self-reported and electronic vaccine records in the Millennium Cohort Study. *Hum Vaccin*. 2007;3(6):245–251.
35. Smith TC, Jacobson IG, Smith B, Hooper TI, Ryan MA, For The Millennium Cohort Study Team. The occupational role of women in military service: validation of occupation and prevalence of exposures in the Millennium Cohort Study. *Int J Environ Health Res*. 2007;17(4):271–284.

36. Smith B, Wingard DL, Ryan MA, Macera CA, Patterson TL, Slymen DJ. US Military deployment during 2001-2006: comparison of subjective and objective data sources in a large prospective health study. *Ann Epidemiol.* 2007;17(12):976-982.
37. Smith B, Smith TC, Gray GC, Ryan MA. When epidemiology meets the Internet: Web-based surveys in the Millennium Cohort Study. *Am J Epidemiol.* 2007;166(11):1345-1354.
38. Spitzer RL, Williams JB, Kroenke K, et al. Utility of a new procedure for diagnosing mental disorders in primary care. The PRIME-MD 1000 study. *JAMA.* 1994;272(22):1749-1756.
39. Ware JE Jr, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care.* 1992;30(6):473-483.
40. Gray GC, Chesbrough KB, Ryan M, et al. The Millennium Cohort Study: a 21-year prospective cohort study of 140,000 military personnel. *Mil Med.* 2002;167(6):483-488.
41. Dillman DA. *Mail and Telephone Surveys: The Total Design Method.* Vol xvi. New York, NY: Wiley; 1978.
42. Sclar DA, Robison LM, Skaer TL. Ethnicity/race and the diagnosis of depression and use of antidepressants by adults in the United States. *Int Clin Psychopharmacol.* 2008;23(2):106-109.
43. Riolo SA, Nguyen TA, Greden JF, King CA. Prevalence of depression by race/ethnicity: findings from the National Health and Nutrition Examination Survey III. *Am J Public Health.* 2005;95(6):998-1000.
44. Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire. *JAMA.* 1999;282(18):1737-1744.
45. Spitzer RL, Williams JB, Kroenke K, Hornyak R, McMurray J. Validity and utility of the PRIME-MD patient health questionnaire in assessment of 3000 obstetric-gynecologic patients: the PRIME-MD Patient Health Questionnaire Obstetrics-Gynecology Study. *Am J Obstet Gynecol.* 2000;183(3):759-769.
46. Fann JR, Bombardier CH, Dikmen S, et al. Validity of the Patient Health Questionnaire-9 in assessing depression following traumatic brain injury. *J Head Trauma Rehabil.* 2005;20(6):501-511.
47. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16(9):606-613.
48. Blanchard EB, Jones-Alexander J, Buckley TC, Forneris CA. Psychometric properties of the PTSD Checklist (PCL). *Behav Res Ther.* 1996;34(8):669-673.
49. Weathers FW, Litz BT, Herman DS, Huska JA, Keane TM. The PTSD Checklist (PCL): reliability, validity, and diagnostic utility. Paper presented at: Annual Meeting of International Society for Traumatic Stress Studies; October 1993; San Antonio, TX. Available at: http://www.pdhealth.mil/library/downloads/PCL_sychometrics.doc. Accessed October 9, 2009.
50. Bush B, Shaw S, Cleary P, Delbanco TL, Aronson MD. Screening for alcohol abuse using the CAGE questionnaire. *Am J Med.* 1987;82(2):231-235.
51. Wiesbeck GA, Kuhl HC, Yaldizli O, Wurst FM. Tobacco smoking and depression—results from the WHO/ISBRA study. *Neuropsychobiology.* 2008;57(1-2):26-31.
52. Martini S, Wagner FA, Anthony JC. The association of tobacco smoking and depression in adolescence: evidence from the United States. *Subst Use Misuse.* 2002;37(14):1853-1867.
53. Smith B, Ryan MA, Wingard DL, Patterson TL, Slymen DJ, Macera CA. Cigarette smoking and military deployment: a prospective evaluation. *Am J Prev Med.* 2008;35(6):539-546.
54. Dohrenwend BP, Turner JB, Turse NA, Adams BG, Koenen KC, Marshall R. The psychological risks of Vietnam for U.S. veterans: a revisit with new data and methods. *Science.* 2006;313(5789):979-982.
55. Rona RJ, Fear NT, Hull L, et al. Mental health consequences of overstretch in the UK armed forces: first phase of a cohort study. *BMJ.* 2007;335(7620):603.
56. Lindstrom KE, Smith TC, Wells TS, et al. The mental health of U.S. military women in combat support occupations. *J Womens Health (Larchmt).* 2006;15(2):162-172.
57. Gaylord KM. The psychosocial effects of combat: the frequently unseen injury. *Crit Care Nurs Clin North Am.* 2006;18(3):349-357.
58. Adler AB, Huffman AH, Bliese PD, Castro CA. The impact of deployment length and experience on the well-being of male and female soldiers. *J Occup Health Psychol.* 2005;10(2):121-137.
59. Bartone PT, Adler AB, Vaitkus MA. Dimensions of psychological stress in peacekeeping operations. *Mil Med.* 1998;163(9):587-593.
60. Vogt DS, Pless AP, King LA, King DW. Deployment stressors, gender, and mental health outcomes among Gulf War I veterans. *J Trauma Stress.* 2005;18(2):115-127.
61. Lambert CE Jr, Lambert VA. Psychological hardness: state of the science. *Holist Nurs Pract.* 1999;13(3):11-19.
62. Dolan CA, Adler AB. Military hardness as a buffer of psychological health on return from deployment. *Mil Med.* 2006;171(2):93-98.
63. Shalev AY, Freedman S, Peri T, et al. Prospective study of posttraumatic stress disorder and depression following trauma. *Am J Psychiatry.* 1998;155(5):630-637.
64. Zatzick D, Jurkovich GJ, Rivara FP, et al. A national US study of posttraumatic stress disorder, depression, and work and functional outcomes after hospitalization for traumatic injury. *Ann Surg.* 2008;248(3):429-437.
65. Campbell DG, Felker BL, Liu CF, et al. Prevalence of depression-PTSD comorbidity: implications for clinical practice guidelines and primary care-based interventions. *J Gen Intern Med.* 2007;22(6):711-718.
66. Gerrity MS, Corson K, Dobscha SK. Screening for posttraumatic stress disorder in VA primary care patients with depression symptoms. *J Gen Intern Med.* 2007;22(9):1321-1324.
67. Grieger TA, Cozza SJ, Ursano RJ, et al. Posttraumatic stress disorder and depression in battle-injured soldiers. *Am J Psychiatry.* 2006;163(10):1777-1783, quiz 1860.
68. Oquendo M, Brent DA, Birmaher B, et al. Post-traumatic stress disorder comorbid with major depression: factors mediating the association with suicidal behavior. *Am J Psychiatry.* 2005;162(3):560-566.
69. Maes M, Mylle J, Delmeire L, Altamura C. Psychiatric morbidity and comorbidity following accidental man-made traumatic events: incidence and risk factors. *Eur Arch Psychiatry Clin Neurosci.* 2000;250(3):156-162.
70. North CS, Nixon SJ, Shariat S, et al. Psychiatric disorders among survivors of the Oklahoma City bombing. *JAMA.* 1999;282(8):755-762.
71. Koenen KC, Fu QJ, Ertel K, et al. Common genetic liability to major depression and posttraumatic stress disorder in men. *J Affect Disord.* 2008;105(1-3):109-115.
72. Scherrer JF, Xian H, Lyons MJ, et al. Posttraumatic stress disorder; combat exposure; and nicotine dependence, alcohol dependence, and major depression in male twins. *Compr Psychiatry.* 2008;49(3):297-304.
73. Riddle JR, Smith TC, Smith B, et al. Millennium Cohort: the 2001-2003 baseline prevalence of mental disorders in the U.S. military. *J Clin Epidemiol.* 2007;60(2):192-201.
74. Grieger TA, Fullerton CS, Ursano RJ, Reeves JJ. Acute stress disorder, alcohol use, and perception of safety among hospital staff after the sniper attacks. *Psychiatr Serv.* 2003;54(10):1383-1387.
75. Johnson EO, Breslau N. Is the association of smoking and depression a recent phenomenon? *Nicotine Tob Res.* 2006;8(2):257-262.
76. Rychnovsky JD. Postpartum fatigue in the active-duty military woman. *J Obstet Gynecol Neonatal Nurs.* 2007;36(1):38-46.
77. West M, Rose SM, Spreng S, Verhoef M, Bergman J. Anxious attachment and severity of depressive symptomatology in women. *Women Health.* 1999;29(1):47-56.
78. Levine S, Lyons DM, Schatzberg AF. Psychobiological consequences of social relationships. *Ann N Y Acad Sci.* 1997;807:210-218.
79. Hagerty BM, Williams RA. The effects of sense of belonging, social support, conflict, and loneliness on depression. *Nurs Res.* 1999;48(4):215-219.
80. Choenarom C, Williams RA, Hagerty BM. The role of sense of belonging and social support on stress and depression in individuals with depression. *Arch Psychiatr Nurs.* 2005;19(1):18-29.
81. Wells TS, LeardMann CA, Smith TC, et al. Self-reported adverse health events following smallpox vaccination in a large prospective study of US military service members. *Hum Vaccin.* 2008;4(2):127-133.

REPORT DOCUMENTATION PAGE

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB Control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. Report Date (DD MM YY) 28-01-08	2. Report Type New	3. DATES COVERED (from - to) Mar 2007 – Jan 2008
4. TITLE AND SUBTITLE A Prospective Study of Depression Following Combat Deployment in Support of the Wars in Iraq and Afghanistan		5a. Contract Number: 5b. Grant Number: 5c. Program Element: 5d. Project Number: 5e. Task Number: 5f. Work Unit Number: 60002
6. AUTHORS Timothy S. Wells, DVM, MPH, PhD; Cynthia A. LeardMann, MPH; Sarah O. Fortuna, MD; Besa Smith, MPH, PhD; Tyler C. Smith, MS, PhD; Margaret AK Ryan, MD, MPH; Edward J. Boyko, MD, MPH; Dan Blazer, MD, PhD; for the Millennium Cohort Study Team		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Commanding Officer Naval Health Research Center 140 Sylvester Road San Diego, CA 92106-3521		
8. SPONSORING/MONITORING AGENCY NAMES(S) AND ADDRESS(ES) Commanding Officer Naval Medical Research Center 503 Robert Grant Ave Silver Spring, MD 20910-7500		8. PERFORMING ORGANIZATION REPORT NUMBER Report 08-05
Commander Navy Medicine Support Command P.O. Box 140 Jacksonville, FL 32213-0140		10. Sponsor/Monitor's Acronyms(s) NMRC/NMSC
		11. Sponsor/Monitor's Report Number(s)

12 DISTRIBUTION/AVAILABILITY STATEMENT
Approved for public release; distribution is unlimited.

13. SUPPLEMENTARY NOTES

14. ABSTRACT (maximum 200 words)

Background: Previous studies have reported an association between deployment and depression; however, these studies have been limited by small sample size or lack of longitudinal design.

Results: Deployed men and women with combat exposures had the highest onset of depression at follow-up (5.6% and 15.7%, respectively), followed by those not deployed (3.9% and 7.7%, respectively), and those deployed without combat exposures (2.3% and 5.1%, respectively). After adjusting for demographic, behavioral, and military characteristics, combat-deployed men and women were at increased odds for new-onset depression compared with nondeployed men and women. Conversely, deployed men and women without combat exposures were at decreased odds for new-onset depression compared with nondeployed men and women.

Conclusions: To our knowledge, this is the first large, longitudinal study to examine the relationship between deployment and depression. Deployment with combat exposures is a significant risk factor for new-onset depression among US service members.

14. SUBJECT TERMS
depression, questionnaires, military psychiatry, combat disorders

16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UNCL	18. NUMBER OF PAGES 10	18a. NAME OF RESPONSIBLE PERSON Commanding Officer
a. REPORT UNCL	b. ABSTRACT UNCL	b. THIS PAGE UNCL			18b. TELEPHONE NUMBER (INCLUDING AREA CODE) COMM/DSN: (619) 553-8429