

Numerical Modeling of Acoustic Propagation In a Variable Shallow Water Waveguide

Daniel Rouseff
Applied Physics Laboratory
University of Washington
1013 NE 40th St.
Seattle, WA 98105

phone: (206) 685-3078 fax: (206) 543-6785 email: rouseff@apl.washington.edu

Grant Number: N00014-04-1-0148

<http://www.apl.washington.edu>

LONG-TERM GOALS

Random variability in shallow water will induce variability in a propagating acoustic field. The long-term goal of this research is to quantify how random variability in the ocean environment translates into random variability in the acoustic field and the associated signal processing algorithms propagation in the mid-frequency (1-10 kHz) band. In the present funding cycle, the emphasis is on the effects of the internal tide and linear internal waves.

OBJECTIVES

The first objective for the current funding cycle is to quantify how the slowing rising thermocline caused by the internal tide affects the mean acoustic intensity. Having successfully modeled the mean intensity, the primary objective is to then understand how the fluctuations in intensity about the mean are caused by linear internal waves.

APPROACH

Our approach is a mixture of data analysis and modeling. We use acoustic and environmental data collected during the Shallow Water 2006 Experiment (SW06).¹ These experimental observations guide the development of models. This work is done in close collaboration with other SW06 participants especially Dajun Tang, Jie Yang, and Frank Henyey.

Figure 1 shows the positioning of the SW06 assets on 18 August 2006. The acoustic source was deployed off the stern of the *R/V Knorr* at depth 40 m. Of present interest are 20 ms duration linear frequency-modulated (LFM) chirp signals in the 1.5 to 10 kHz band that were transmitted every 19 s. The signals were recorded on MORAY moored receiving system that had two short vertical sub arrays with the bottom four elements, one at depth 25 m and the other at 50 m. Two acoustic data sets were collected, each

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2009		2. REPORT TYPE Annual		3. DATES COVERED 00-00-2009 to 00-00-2009	
4. TITLE AND SUBTITLE Numerical Modeling Of Acoustic Propagation In A Variable Shallow Water Waveguide			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, Applied Physics Laboratory, 1013 NE 40th St., Seattle, WA, 98105			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Code 1 only					
14. ABSTRACT Random variability in shallow water will induce variability in a propagating acoustic field. The long-term goal of this research is to quantify how random variability in the ocean environment translates into random variability in the acoustic field and the associated signal processing algorithms propagation in the mid-frequency (1-10 kHz) band. In the present funding cycle, the emphasis is on the effects of the internal tide and linear internal waves.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 5	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

approximately 7 hours in duration, one with the *R/V Knorr* at fixed position 550 m away from MORAY, and the second with the *R/V Knorr* slowly transiting while towing the source. During the tow, the range to MORAY increased from 100 m to 8 km. Concurrent oceanographic data were collected on five moorings (labeled 14-17 and 54 in the figure). In addition, investigators onboard the *R/V Oceanus* collected oceanographic data 250 m east of the *R/V Knorr* as a non-linear internal wave first approached and then passed the two ships. The wave subsequently passed MORAY and the oceanographic moorings. X-band radar measurements estimated the bearing and speed of the wave. Note that the bearing of the internal wave nearly coincided with the bearing from the *R/V Knorr* to MORAY.




Figure 1. Experiment geometry. Acoustic source deployed off the stern of the *R/V Knorr*. Two acoustic data sets: fixed range 550 m and slow tow. Oceanographic data collected on *R/V Oceanus* and moorings labeled 14-17 and 54.

WORK COMPLETED

An analysis of the SW06 550 m path data taken while a non-linear internal wave passed was completed. In the current fiscal year, a previously submitted paper documenting this work was published.² The paper shows how non-linear internal waves can cause new acoustic paths to be generated. This year, a journal paper was written and accepted for publication that documents how the slowing rising thermocline caused by the internal tide affects the mean acoustic intensity for both the 550 m and towed-source data sets.³ A previously submitted journal paper looking at intensity fluctuations was accepted for publication in this fiscal year.⁴ New results showing intensity fluctuations in the SW06 towed-source data were presented in an invited talk at the Acoustical Society of America meeting in Portland.⁵

RESULTS

The internal tide causes both the generation of nonlinear internal waves and the subsequent slow rising of the high-gradient part of the thermocline. The latter has been studied in the current fiscal year both for the fixed range 550 m and towed-source SW06 data. For the fixed-range data, a 5 dB change in intensity was observed in the data and correctly modeled. For the towed-source data, among other effects, a 2 dB dip in transmission loss between ranges 5 and 7 km was observed in the data and correctly modeled using oceanographic mooring data as input. Water-column effects of this level may be significant as they could cause erroneous results if one is trying to predict the mean level of ensonification in an active sonar scenario. For the cases considered, however, the mean water column effects were predictable when data from nearby oceanographic moorings were incorporated into the numerical simulations.

While a deterministic acoustic model using nearby oceanographic mooring data as input is adequate for predicting the mean intensity, it fails to predict fluctuations in intensity. Figure 2 shows a model-data comparison for intensity in the 3 to 3.3 kHz band and range near 3 km. Both the model and data images have 20 dB dynamic range. The range-independent model result in the left panel shows a regular pattern of intensity striations, bands of high intensity. The striations are consistent with a value for the waveguide invariant⁶ β approximately equal to 2. The right panel shows SW06 towed-source results. The 300 m horizontal aperture (range) was synthesized by towing the acoustic source and took 20 minutes to generate. The experimental results are much more granular than the simulation; while striations are still evident, they are less distinct. The simulation, which ignores the temporal and spatial variability introduced by internal waves, over predicts the stability of the pattern. Ongoing research is attempting to quantify the “beta content” of the experimental results.




Figure 2. Intensity striation patterns in at mid-frequencies for SW06. Left: range-independent PE simulation. Right: Experimental results generated by synthesizing horizontal array. Plots have similar sampling in frequency (~ 1 Hz) and range (~ 6 m) and same 20 dB dynamic range.

IMPACT/APPLICATIONS

Internal waves are a ubiquitous feature of shallow water oceanography. Even when event-like non-linear internal waves are absent, random background internal waves will affect acoustic propagation in the mid-frequency regime relevant to Navy sonar systems. By studying data sets such as collected during the SW06 experiment, these effects can be quantified as a first step towards developing a predictive capability and possible mitigation strategies.

RELATED PROJECTS

Numerous ONR-supported investigators were involved in the SW06 Experiment and are analyzing the collected data.

REFERENCES

¹D. Tang, J. N. Moum, J. F. Lynch, P. Abbot, R. Chapman, P. H. Dahl, T. F. Duda, G. Gawarkiewicz, S. Glenn, J. A. Goff, H. Graber, J. Kemp, A. Maffei, J. D. Nash, and A. Newhall, “Shallow Water ’06, A Joint Acoustic Propagation/Nonlinear Internal Wave Physics Experiment,” *Oceanogr.* 20, 156-167, 2007.

²D. Rouseff, D. Tang, K. L. Williams, Z. Wang, and J. N. Moum, “Mid-frequency sound propagation through internal waves at short range with synoptic oceanographic observations,” *J. Acoust. Soc. Am.* 124, EL73-EL77 (2008).

³J. Yang, D. Rouseff, D. Tang, and F. S. Henyey, “Effect of the internal tide on acoustic transmission loss at mid-frequencies,” *IEEE J. Oceanic Eng.* [in press].

⁴A. Turgut, M. Orr, and D. Rouseff, “Broadband source localization using horizontal-beam acoustic intensity striations,” *J. Acoust. Soc. Am.* [in press].

⁵D. Rouseff and V. G. Petnikov, “Effect of shallow water variability on the waveguide invariant distribution,” Special session on waveguide invariant, 157th Mtg. of Acoustical Society of America, Portland, OR, May 18-22, 2009.

⁶L. M. Brekhovskikh and Y. P. Lysanov, *Fundamentals of Ocean Acoustics*, 2nd ed. (Springer-Verlag, New York, 1991).

PUBLICATIONS

D. Rouseff, “Physics-based modeling of coherent communications performance in environments with strong multi-pathing,” Pacific Rim Underwater Acoustics Conference (PRUAC), Xi’an, China, September 23-25, 2009 [invited, non-refereed].

J. Yang, D. Rouseff, D. Tang, and F. S. Henyey, "Effect of the internal tide on acoustic transmission loss at mid-frequencies," *IEEE J. Oceanic Eng.* [in press, refereed].

D. Rouseff, M. Badiey, and A. Song, "Effect of reflected and refracted signals on coherent underwater acoustic communication: Results from KauaiEx 2003." *J. Acoust. Soc. Am.* [in press, refereed].

A. Turgut, M. Orr, and D. Rouseff, "Broadband source localization using horizontal-beam acoustic intensity striations," *J. Acoust. Soc. Am.* [in press, refereed].

D. Rouseff and W. L. J. Fox, "Energy consumption model for a broadband shallow-water acoustic communications network," *IEEE J. Oceanic Eng.* 33, 335-340 (2008) [published, refereed].

D. Rouseff, D. Tang, K. L. Williams, Z. Wang, and J. N. Moum, "Mid-frequency sound propagation through internal waves at short range with synoptic oceanographic observations," *J. Acoust. Soc. Am.* 124, EL73-EL77 (2008) [published, refereed].

D. Tang, F. S. Henyey, Z. Wang, K. L. Williams, D. Rouseff, P. H. Dahl, J. Quijano, and J. W. Choi, "Mid-frequency acoustic propagation in shallow water on the New Jersey shelf: Mean intensity," *J. Acoust. Soc. Am.* 124, EL85-EL90 (2008) [published, refereed].

D. Tang, F. S. Henyey, Z. Wang, K. L. Williams, D. Rouseff, P. H. Dahl, J. Quijano, and J. W. Choi, "Mid-frequency acoustic propagation in shallow water on the New Jersey shelf. II. Intensity fluctuations," *J. Acoust. Soc. Am.* 124, EL91-EL96 (2008) [published, refereed].

HONORS/AWARDS/PRIZES

Best Paper Award Winner, SENSORCOMM 2009, Underwater Sensors Track, Athens, Greece, June 18-23, 2009. Sponsored by the International Academy, Research and Industry Association (IARIA). A. L. Matveev, P. I. Korotin, V. I. Turchin, D. Rouseff and R. C. Spindel, "Experimental study of acoustic forward scattering on a marine shelf."