

Optimal Combining Data for Improving Ocean Modeling

L.I.Piterbarg
University of Southern California
Center of Applied Mathematical Sciences
Kaprielian Hall, R108, 3620 Vermont Avenue
Los Angeles, CA 90089-2532
Phone (213) 740 2459, fax (213) 740 2424, e-mail piter@usc.edu
Award # N00014-08-1-0643
[http : //www.onr.navy.mil/sci_tech/ocean/onrpgahm.htm](http://www.onr.navy.mil/sci_tech/ocean/onrpgahm.htm)

LONG-TERM GOALS

The long range scientific objectives of the proposed research comprise: (1) developing rigorous approaches to optimal combining satellite and drifter data with an output of a regional circulation model for accurate estimating the upper ocean velocity field and mixing characteristics such as relative dispersion and finite size Lyapunov exponent, (2) constructing and comprehensive testing computationally efficient estimation algorithms based on alternative parameterizations of uncertainty, (3) processing real data in the Adriatic and Ligurian Sea (MREA coastal experiments) via new techniques

OBJECTIVES

The objectives for the first year of research were:

- Developing and verifying data fusion algorithms for optimal estimating surface velocities from tracer observations, Lagrangian data, and a circulation model output, based on the *fuzzy logic* approach [1,2].
- Carrying out a comprehensive error analysis via theoretical and Monte Carlo methods.
- Constructing and testing compatibility measures between data and model.
- Testing the developed fusion algorithms by 'twin' experiments via NCOM.
- Theoretical investigation of the absolute and relative dispersion (AD and RD) in the presence of shear flow, which would help in constructing realistic algorithms for estimating these characteristics from real data.
- Developing data fusion algorithms for estimating RD by combining model output, drifter observations and images.

APPROACH

I develop theoretical approaches to the data fusion problem in context of the possibility theory (fuzzy logic) and in the framework of the classical theory of random processes and fields covered by stochastic partial differential equations. I design computational algorithms derived from the theoretical findings. A significant part of the algorithm validation is their testing via stochastic simulations. Such an approach provides us with an accurate error analysis. Together with my collaborators from Rosenstiel School of Marine and Atmospheric Research (RSMAS), Consiglio Nazionale delle Ricerche (ISMAR, LaSpezia, Italy), Naval Research Laboratory (Stennis Space Center, Mississippi), ENEA (Rome, Italy), Koc University (Istanbul, Turkey) we implement the

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2008		2. REPORT TYPE Annual		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE Optimal Ccombining Data For Improving Ocean Modeling				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Southern California,Center of Applied Mathematical Sciences,Kaprielian Hall, R108, 3620 Vermont Avenue,Los Angeles,CA,90089-2532				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT The long range scientific objectives of the proposed research comprise: (1) developing rigorous approaches to optimal combining satellite and drifter data with an output of a regional circulation model for accurate estimating the upper ocean velocity field and mixing characteristics such as relativedispersionand finitesizeLyapunovexponent,(2) constructing and comprehensivetesting computationally efficient estimation algorithms based on alternative parameterizations of uncertainty,(3) processing realdataintheAdriaticandLigurianSea(MREA coastal experiments) via new techniques					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 6	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

algorithms in concrete ocean models such as QG, MICOM , and NCOM as well as carry out statistical analysis of real data sets by means of new methods.

WORK COMPLETED

1. *Data fusion for estimating surface velocities.*

Two efficient fuzzy logic based algorithms of combining tracer observations with a model output were developed and compared. Both methods appeal to the intersection $D = D_m \cap D_o$ of the credibility regions for the estimated velocity vector coming from the model and observations respectively. The first algorithm called GC takes the geometrical center of D as a combined estimate of the unknown velocity, while the second one (CM) addresses the center of mass of D . A theoretical error analysis was provided for the GC algorithm. The procedures were first tested on an idealized model of gyre superposed by a regular eddy structure, and then, in the framework of 'twin' experiment with NCOM at a specific area of the Ligurian Sea. A compatibility measure between tracer data and a model output was suggested and analyzed.

2. *Mixing in the presence of shear*

A theoretical investigation of the absolute and relative dispersion in stochastic flows with constant drift gradients was conducted. In particular, different regimes for the relative dispersion were identified depending on the type of the mean velocity stagnation points (hyperbolic or elliptic) and on the Hurst exponent characterizing the dynamics type (local or non-local).

3. *Fusion data for estimating RD.*

Theoretical relations between relative dispersion and statistical characteristics of a continuously distributed tracer were established to lay ground for data fusion procedures aiming at estimating RD from images. Similar relations were established between the finite size Lyapunov exponent (FSLE) and Eulerian characteristics of the underlying velocity field to estimate the former from tracer snapshots. A fusion algorithm has been developed for estimating RD by combining a model output, drifter data, and images which is based on ideas of fuzzy logic.

RESULTS

1. Principal results in developing and testing the GC and CM data fusion algorithms are as follows.
 - GC is computationally simpler, but less stable and less accurate than CM. The difference in stability and accuracy is especially manifested near the boundary of the circulation region of interest.
 - The estimation error of GC is in very good agreement with the theory which predicts the improvement with respect to the model output at least 30%.
 - Both algorithms demonstrated a high computational efficiency exceeding typical assimilation algorithms in several orders.
 - Satisfactory estimation results can be reached for the interval shorter than 6 hours between consecutive snapshots. For longer intervals the error goes out of control.
 - The data/model compatibility measure μ introduced as a complementary normalized distance between the credibility regions D_m and D_o turned out to be greatly helpful in interpreting estimation results. In particular, small values of μ in a particular place alarm on the model credibility in this place.

Fig.1 illustrates the CM performance in a 'twin' experiment with NCOM in a particular area of the Ligurian sea. Two outputs for May of 2007 separated by time interval of 7 days, were considered as a model and 'real' velocity field. The main feature of the 'real' circulation (first panel) comparatively

- We have shown that the anisotropy of $\mathbf{K}$ is determined by both factors, gradients of the mean flow and statistics of the velocity fluctuations. The crucial impact on the mixing ellipse orientation is produced by the normal component of the covariance. The growth rate Λ depends on the drift and fluctuation characteristics as well and can be partly investigated analytically.

In Fig.2 we give examples of the RD curves for both local and non-local dynamics in a vicinity of both hyperbolic ($\Omega < \gamma$, first panel) and elliptic (second panel) stagnation points as well as the dependence of Λ on the shear parameters Ω, γ (third panel) and on β, γ (fourth panel), where β is the characteristic wavenumber of the velocity fluctuations, i.e. $\beta = 1/d$ with d standing for the space correlation scale of the velocity.




Figure 2. Relative dispersion in shear flows from left to right: 1) RD vs time and different values of Hurst exponent $h = 0.1$ (black), $h = 1$ (red), $h = 2$ (blue) $\gamma = 0.1, \Omega = 0$, 2) Same for $\gamma = 0.1, \Omega = 2$). 3) RD vs Ω and γ . 4) RD vs β and γ for $\Omega = 2$.

3. The following result is in the base of the derived fusion algorithm illustrated in Fig.3 for estimating the relative diffusivity (RD) by combining a model output, tracer and drifter observations.




Figure 3: Schematic illustration of the fusion method for the RD estimate. Confidence regions for (D_x, D_y) defined from model experiments (yellow), drifter estimate (green) and tracer estimate (orange) have the common area (dark green) and the combined estimate is its center of mass

- Under wide conditions we proved that for small enough initial separation x_0, y_0

$$\frac{D_x}{d_x^2} + \frac{D_y}{d_y^2} \geq \frac{x_0^2 \langle \theta_x^2 \rangle + y_0^2 \langle \theta_y^2 \rangle}{\sigma_\theta^2} \quad (1)$$

In (1) D_x and D_y are the RD zonal and meridional components respectively, σ_θ^2 is the variance of the tracer $\theta(t, x, y)$, θ_x, θ_y its gradients, and d_x, d_y are its correlation scales in direction x and y respectively at $t = 0$, the dependence of $D_x, D_y, \langle \theta_x^2 \rangle, \langle \theta_y^2 \rangle$ on t is meant.

Relation (1) determines the orange confidence region. The center of mass ($\hat{D}_x, \hat{D}_y$) of the intersection (dark green) of the three confidence regions is taken as a combined estimate of (D_x, D_y) . Explicit expressions for estimates ($\hat{D}_x, \hat{D}_y$) were obtained as well.

IMPACT/APPLICATIONS

1. The suggested data fusion procedures for estimating surface velocities and relative dispersion from images, drifters, and a model output could be a supplement to existing assimilation algorithms explicitly involving OGCM.
2. The developed theory of the dispersion in stochastic flows leads to better understanding turbulent mixing in the presence of shear flow

TRANSITIONS

The developed velocity fusion algorithm was used in RSMAS to test it in NCOM circulation model. It is planned to apply the method to real data in the Mediterranean by the same RSMAS group in collaboration with ISAC-CNR remote sensing group (Rome, Italy).

RELATED PROJECTS

1. "Predictability of Particle Trajectories in the Ocean", ONR, PI T.Ozgokmen, RSMAS, N00014-05-1-0095
2. "Lagrangian turbulence and transport in semi-enclosed basins and coastal regions", ONR, PI A Griffa, RSMAS, N00014-05-1-0094

REFERENCES

1. D.Dubious and H. Prade, (1986), Possibility theory, Plenum Press, New York and London,
2. D. Dubois, H.Prode, and R.R. Yager, Fuzzy Information Engineering, 1997, John Wiley & Sons, Inc.
3. Veneziani M., A Griffa, A.M. Reynolds and A.J. Mariano, 2004: Oceanic turbulence and stochastic models from subsurface Lagrangian data for the North-West Atlantic Ocean, *J. Phys. Oceanogr.*, v.34, (8), 1884-1906.
4. Batchelor, G.K., 1952: Diffusion in a Field of Homogeneous Turbulence, II. The Relative Motion of Particles," *Proc. Camb. Philos. Soc.*, 48, 345-362.
5. Bennett, A.F., 1984: Relative Dispersion: Local and Nonlocal Dynamics, *J. Atmos. Sci.*, **41**, 1881-1886.

6. L.I. Piterbarg, (2008), Particle dispersion in stochastic flows with constant drift gradient, *International Journal of Pure and Applied Mathematics*, accepted
7. L.I. Piterbarg, (2005), Relative dispersion in 2D stochastic flows, *J. of Turbulence*, v.6, n.4, 1-19.

PUBLICATIONS

- L.I. Piterbarg, (2008), A simple method for computing velocities from tracer observations and a model output, *Applied Mathematical Modeling*, submitted
- L.I. Piterbarg, (2008), Particles in a stochastic flows with a linear drift, *2008 SIAM Annual Meeting, July 7-11, 2008*, 38
- L.I. Piterbarg, (2008), Particle dispersion in stochastic flows with constant drift gradient, *International Journal of Pure and Applied Mathematics*, accepted
- L.I. Piterbarg, (2008), Optimal estimation of Eulerian velocity field given Lagrangian observations, *Applied Mathematical Modeling*, 32, 2133 - 2148
- L.I. Piterbarg and M. Gaglar (2007), Research Trends in Stochastic Flows Modeling the Upper Ocean Turbulence, in *Leading-Edge Applied Mathematical Modeling Research*, ed. M.P. Alvarez, Nova Science Publishers, Inc., 2007, 67-110
- G. Falkovich, S. Musacchio, L. Piterbarg, M. Vucelja, (2007), Inertial particles driven by a telegraph noise, *Physical Review E*, v.76, n.2, 1-8
- L.I. Piterbarg, T. Ozgokmen, A. Griffa, and A. Mariano (2007), Predictability of Lagrangian trajectories, in *Lagrangian Analysis and Prediction of Coastal and Ocean Dynamics*, ed. A. Griffa, D.Kirwan, A. Mariano, T. Ozgokmen, and T. Rossby, Cambridge University Press, 2007, 136-171.
- A. Molcard, T. Ozgokmen, A. Griffa, L.I. Piterbarg, and M. Chin (2007), Lagrangian data assimilation in ocean general circulation models, in *Lagrangian Analysis and Prediction of Coastal and Ocean Dynamics*, ed. A. Griffa, D.Kirwan, A. Mariano, T. Ozgokmen, and T. Rossby, Cambridge University Press, 2007, 171-203.