

Understanding Lateral Mixing in the Ocean: 100 m to 1 km

Murray D. Levine

Oregon State University, 104 COAS Admin Bldg, Corvallis, OR 97331

phone: (541) 737-3047 fax: (541) 737-2064 email: Levine@coas.oregonstate.edu

Award Number: N00014-08-1-0552

LONG-TERM GOALS

The long-range goals of this research are to understand the submesoscale processes causing horizontal dispersion of momentum and scalar quantities over the continental shelf. Phenomena that may lead to horizontal dispersion and variability include: meandering coastal fronts and jets, stirring by the wind and tide, and nonlinear internal wave packets.

OBJECTIVES

The specific objectives of the proposed research include:

- participation in a planning study group to develop a coordinated multi-year program,
- refurbishing our dye tracking equipment and instrumentation, and
- field testing of equipment and tracking procedures during surface dye releases

APPROACH

The Approach for each of the three specific objectives is given below.

Planning for future coordinated multi-year program. We will contribute to the planning effort lead by ONR for the DRI: *Scalable Lateral Mixing and Coherent Turbulence*. The planning will involve discussions and meetings with others in the group selected by ONR.

Refurbish equipment and instrumentation. Some of the equipment has not been used for more than 5 years and requires a thorough check out. Since there have been changes in technical personnel since our last dye experiment, this effort will give the opportunity to familiarize new personnel with the equipment.

Field test. The first field test will be a day trip aboard the R/V Elakha. This will serve as an initial shakedown of the key elements of the dye tracking equipment, including the dye pumping system, the Minibat towing operation, and the real-time display software. The second field test will be a 2-day cruise aboard the R/V Wecoma off the Oregon coast, using both the Minibat and the ship's CTD for dye tracking.

WORK COMPLETED

An initial round of discussions for planning the DRI have taken place, including a two-day meeting in Boston in late May 2008.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2008		2. REPORT TYPE Annual		3. DATES COVERED 00-00-2008 to 00-00-2008	
4. TITLE AND SUBTITLE Understanding Lateral Mixing In The Ocean: 100 m To 1 km				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon State University,104 COAS Admin Bldg,Corvallis,OR,97331				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT The long-range goals of this research are to understand the submesoscale processes causing horizontal dispersion of momentum and scalar quantities over the continental shelf. Phenomena that may lead to horizontal dispersion and variability include: meandering coastal fronts and jets, stirring by the wind and tide, and nonlinear internal wave packets.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

In preparation for the field tests, significant overhaul of equipment and instrumentation was accomplished, including: revamping the dye pumping system, upgrading the system used to mix the fluorescein powder, testing sensitivity of new fluorometers, reconfiguring the CTD package used on the Minibat, checking Minibat wing operation, replacing mechanical hardware, upgrading the controlling computer system.

The day test on the 54' R/V Elakha occurred on 28 July about 10 miles off Newport, Oregon. Dye was release and was subsequently tracked with the Minibat CTD system. The two-day field test on the R/V Wecoma took place on 10-11 August. Two surface dye releases were done about 20 miles off Newport (see Figure 1). Tracking of the dye patch was done with both Minibat and the ship's CTD (tow-yo's). A fluorometer was also hooked up to the ship's flowthrough system to continuously sample water at 3 m depth.



Figure 1. View of surface release of fluorescein dye from R/V Wecoma off Newport, Oregon.

RESULTS

Much was learned regarding the conduct of a surface dye tracking experiment. We collected a significant amount of data (see example in Figure 2), and the analysis is just beginning. The first step, which is underway, is to determine the time lags between sensor responses so corrections can be made.

Our goal in further analysis is to determine if we are able to track a patch of dye adequately enough to learn something about the horizontal diffusion process. This analysis will also help us determine if

modifications should be made in our methods or equipment before conducting a future larger experimental effort.




Figure 2. Vertical profile of dye fluorescence with salinity, temperature and density from the R/V Wecoma off the Oregon coast. Note the dye has been mixed throughout the mixed layer where it was injected and is beginning to mix into the pycnocline below.

IMPACT/APPLICATIONS

During the two-day Wecoma cruise we took the opportunity to take 6 oceanography graduate students and one undergraduate REU student, who may not have had the opportunity to experience sea-going operations. A local high school teacher, who is developing the study of oceanography as part of the curriculum, also came along.

RELATED PROJECTS

The observation and modeling of the coastal ocean off Oregon continues to be of great interest at OSU. Ultimately, improving understanding of the lateral mixing process will help in our ability to model coastal circulation.

While there are many projects relating to the coastal ocean at OSU, I am currently involved in making continuous observations from a mooring at a site 10 miles off Newport (NH-10). This effort is funded by NOAA as part of the ocean observatory system, specifically through the Northwest Association of

Networked Ocean Observing Systems (www.nanoos.org and www.orcoos.org) and by NSF, through the Center for Coastal Margin Observation & Prediction (CMOP) (www.stccmop.org). Near-real time data are being collected and distributed on the web.

I am also involved in CMOP (as Co-Director) with a major goal of trying to improve understanding of the river to ocean system. Submesoscale processes are important in trying to understand the mixing of the river plume with the coastal ocean.