

AD _____

AWARD NUMBER: **W81XWH-09-1-0543**

TITLE: **Role of Cathepsin C During Breast Cancer Metastasis**

PRINCIPAL INVESTIGATOR: **Dr. Brian Ruffell**

CONTRACTING ORGANIZATION:
University of California, San Francisco
San Francisco, CA 94143-0962

REPORT DATE: **September 2010**

TYPE OF REPORT: **Annual** Summary

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY) 01-09-2010		2. REPORT TYPE Annual Summary		3. DATES COVERED (From - To) 1 SEP 2009 - 31 AUG 2010	
4. TITLE AND SUBTITLE Role of Cathepsin C During Breast Cancer Metastasis				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-09-1-0543	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Brian Ruffell, Ph.D. Email: brian.ruffell@ucsf.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of California, San Francisco San Francisco, CA 94143-0962				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The presence of some innate immune cell types in developing neoplasms provides a significant pro-tumor advantage. Myeloid-lineage immune cells, such as tumor-associated macrophages (TAMs) and immature myeloid-derived cells/monocytes, promote tumor development by exerting pro-tumor activities including activating angiogenic programs, suppressing anti-tumor immunity, and enhancing migratory and metastatic properties of malignant cells. Cathepsin C (CTSC) is a lysosomal cysteine-class hydrolase expressed in most mammalian tissues. In myeloid cells and cytotoxic lymphocytes, CTSC regulates catalytic activation of several important leukocyte-derived serine proteases including granzymes, chymases and elastases. Preliminary data from the Coussens' laboratory has revealed that in a mouse model of de novo mammary adenocarcinoma development, neither latency of progression to primary tumors, tumor burden, nor tumor histopathology is regulated by CSTC; however, in <i>cathepsin C (Ctsc)</i> -deficient mice, there is a significant reduction in the number of circulating malignant cells in peripheral blood, with a corresponding significant reduction in pulmonary metastasis development, thus indicating that CC plays a predominant role in metastatic dissemination of primary mammary tumors. Since CTSC expression in mammary adenocarcinomas is largely localized in TAMs, these provocative data indicate that macrophages play an important role in regulating intravasation and/or survival of malignant cells in peripheral blood, and that this function is at least partly dependent on CTSC.					
15. SUBJECT TERMS breast cancer, mouse models, inflammation, proteases, cathepsin protease, immune cells, leukocytes					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 25	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER

TABLE OF CONTENTS

	<u>Page</u>
INTRODUCTION.....	4
BODY	4
KEY RESEARCH ACCOMPLISHMENTS	13
REPORTABLE OUTCOMES	14
CONCLUSION	14
BIBLIOGRAPHY	15
APPENDICES	15

I. INTRODUCTION:

It is now clear that solid tumor development is profoundly regulated by infiltrating immune cells of both the innate and adaptive lineages (1, 2). Diverse populations of leukocytes infiltrate tumor microenvironments and are altered by the neoplastic cells and surrounding stroma in such a way that many of their bioactivities are co-opted to aid growth and/or metastasis of nascent tumors. Out of this realization emerges novel targets for therapy that may impact patient survival, i.e., immune targets regulating primary tumor development, and those regulating metastatic dissemination and growth. Metastasis is a highly complex process and involves multiple stages. First, malignant cells must intravasate into blood vessels and enter the peripheral circulation; second, they must survive in the blood, third, they must extravasate and leave the blood and enter a distal organ; and fourth, they must be able to survive and proliferate in ectopic sites. Preliminary analyses in the Coussens laboratory using a mouse model of breast cancer has revealed that absence of a single intracellular leukocyte lysosomal protease, i.e. cathepsin C (CTSC), reduces the ability of malignant mammary epithelial cells to metastasize to the lungs. As metastasis is a leading cause of mortality in cancer patients, these findings indicate that CTSC is a potential therapeutic target for breast cancer. To examine this possibility, the mechanism by which CTSC regulates pulmonary metastasis will be determined *in vivo*. In addition, we will reveal the therapeutic efficacy of targeting CTSC to inhibit lung metastasis *in vivo*. And finally, we will evaluate a large cohort of human breast cancer tissues for CTSC expression, and determine if this molecule can be used as a biomarker to predict clinical outcome and therefore guide therapy.


II. RESEARCH ACCOMPLISHMENTS BODY:


AIM 1: DETERMINE THE FUNCTIONAL ROLE OF CATHEPSIN C IN REGULATING LEUKOCYTE INFILTRATION AND BIOACTIVITY IN MAMMARY ADENOCARCINOMA DEVELOPMENT AND METASTASIS


Establish the profile of cells expressing CTSC during tumor development and metastasis:

Profile CTSC expression in mammary and lung tissue of MMTV-PyMT mice by flow cytometry. (months 1-6)

Despite using bone marrow derived mast cells from *Ctsc*^{+/-} and *Ctsc*^{-/-} mice, we were unable to detect a clear signal by flow cytometry following intracellular staining for CTSC. We therefore optimized staining for CC and other lineage markers for both immune and non-immune populations in PyMT tumors using PFA-fixed frozen tissue. As reported in our initial application, F4/80⁺ macrophages within the tumors were the dominant population expressing CTSC (**Figure 1**), with lower expression of CTSC in other tumor-infiltrating immune cells such as T cells and neutrophils (data not shown). Surprisingly however, the use of CTSC-deficient animals as a staining control revealed low, but consistent, expression of CTSC in other stromal cells such as fibroblasts and pericytes, in addition to the cancer cells themselves. Given the reduced metastatic phenotype in PyMT/*Ctsc*^{-/-} animals, we also stained for CTSC in the lung (**Figure 2**). While a similar expression pattern was observed in immune populations and in metastatic cancer cells, CTSC was not detected in the other stromal populations within the lung.







Evaluate CTSC activity using a selective probe, FY02, in lysates of purified leukocytes isolated from mammary glands, peripheral blood and lung from MMTV-PyMT mice. (months 3-6)

Detecting CTSC activity in specific leukocyte populations proved technically challenging, as approximately 1×10^6 CD45⁺ cells were required to either detect a signal with the FY02 probe (3), or western blot for CTSC. To demonstrate that CTSC protein expression determined by confocal microscopy corresponded to CTSC activity in leukocytes, CD45⁺ and CD45⁻ cells were sorted from PyMT tumors or metastatic lungs and cell lysates were analyzed using the FY02 probe. Both CTSC protein and activity were found only within the CD45⁺ population (**Figure 3B**), indicating that leukocytes express the majority of active CTSC within both tumors and lungs. Specific purification of F4/80⁺ macrophages also revealed CTSC activity within this specific leukocyte population (**Figure 3C**). As anticipated from increased gene expression of *Ctsc* in PyMT tumors compared to normal mammary glands (~7-fold, data not shown), higher CTSC activity was also observed (**Figure 3A**).

Quantitatively assess activity of CTSC enzymatic substrates in

purified leukocytes using selective protease activity assays (months 6-12)

As discussed below, we have yet to identify the specific leukocyte population responsible for the reduced metastatic phenotype in PyMT/*Ctsc*^{-/-} animals, and even have some preliminary data suggesting that it is expression of CTSC by the tumors cells themselves that is important for the phenotype. Pending identification of the cell type of interest, analysis of CTSC substrate activity is currently on hold.

Assess functional significance of CTSC in regulating macrophage phenotype:


Analyze polarization state of tumor-associated macrophages by intracellular staining for cytokine expression. (months 1-6)

Verify macrophage polarization by iNOS and arginase expression levels by real time PCR. (months 6-9)

Compare the ability of PyMT/*Ctsc*^{+/-} and PyMT/*Ctsc*^{-/-} tumor-associated macrophages to influence invasive properties of malignant MECs grown as organoids using the 3D organotypic model. (months 6-12)

While we originally proposed analyzing macrophage polarization and then examining whether altered polarization resulted in a functional difference in the ability of macrophages to promote invasion, we instead bypassed the early steps to directly determine whether CTSC expression by macrophages was important for this process. As shown in **Figure 4A**, expression of the M2 macrophage marker CD206 and the M1 marker MHCII were unchanged between PyMT/*Ctsc*^{+/-} and PyMT/*Ctsc*^{-/-} macrophages.

Furthermore, while as reported the IL-4-dependent polarization of the macrophages promoted the invasive properties of the cancer cells in the 3D organotypic model (4), invasion was unaffected by macrophage CTSC expression (**Figure 4B**). Based upon these results, and the preliminary data suggesting intrinsic CTSC expression by cancer cells is critical for lung colonization (see Aim 2), further analysis of macrophage polarization is currently on hold.




Evaluate leukocyte infiltration regulated by CTSC in PyMT mice:

Analyze leukocyte infiltration in mammary and lung tumors in PyMT/*Ctsc*^{+/-} and PyMT/*Ctsc*^{-/-} mice using multicolor flow cytometry. (months 1-6)

Measure cytokine expression levels in mammary and lung tumors in PyMT/*Ctsc*^{+/-} and PyMT/*Ctsc*^{-/-} mice via ELISA. (months 3-9)

Identify leukocyte populations with altered cytokine expression profiles in mammary and lung tumors in PyMT/*Ctsc*^{+/-} versus PyMT/*Ctsc*^{-/-} mice using intracellular cytokine staining and multicolor flow cytometry. (months 9-12)

12-color polychromatic flow cytometry was used to analyze tissue invasion by leukocytes and activation/polarization through extracellular markers. As shown in **Figure 5D**, leukocyte populations in both mammary glands and mammary tumors were unchanged between *Ctsc*^{+/-} and *Ctsc*^{-/-} mice, with total leukocyte infiltration was also unchanged (**Figure 5C**).



Leukocyte populations within the spleen, lymph node and peripheral blood; and expression of T cell activation markers were also unchanged (data not shown). Based upon our discovery that stromal populations and malignant epithelial cells also expression CTSC (Figure 1), we adapted our flow cytometric protocol to examine cellular composition of tumors. As show in **Figure 5A**, the relative frequency of fibroblasts (PDGFRα⁺), endothelial cells (CD31^{hi}), pericytes (SMA⁺ or Desmin⁺) and lymphatic endothelial cells (CD31^{lo}) was unchanged by CTSC expression. Based upon these results, and the preliminary data suggesting intrinsic CTSC expression by cancer cells is critical for lung colonization (see Aim 2), further analysis of cytokine expression by leukocytes is currently on hold.

AIM 2: DETERMINE AT WHICH STAGE OF CANCER DEVELOPMENT CATHEPSIN C FUNCTIONALLY REGULATES PULMONARY METASTASIS OF MAMMARY CARCINOGENESIS

Determine if cathepsin C regulates intravasation from primary mammary tumors into the circulation:

Measure the number of circulating neoplastic cells in the blood of PyMT/*Ctsc*^{+/-} and PyMT/*Ctsc*^{-/-} mice at day 85 and 95 using either flow cytometry or PCR. (months 12-18)




Figure 6: Cathepsin C expression by tumor cells affects lung metastasis but not intravasation from the primary tumor. **A)** Representative lung sections depicting metastatic tumor burden at day 110 visualized by H&E staining. The scale bar represents 1 mm. **B)** Quantitation of average number of metastatic foci/5.0 mm² lung section for PyMT/*Ctsc*^{+/-} (blue) and PyMT/*Ctsc*^{-/-} (red) mice at end stage. **C)** Real-time PCR quantitation of Keratin 18 and PyMT gene expression from circulating cells. Normalization was done with expression of control gene TBP (TATA box binding protein). **D)** Quantitation of circulating cytokeratin⁺CD45⁻ cells by flow cytometry. **E)** Lung colonization of *Ctsc*^{+/-} and *Ctsc*^{-/-} negative littermates following injection of carcinoma cells isolated from either PyMT/*Ctsc*^{+/-} or PyMT/*Ctsc*^{-/-} mice.

Preliminary data described in the initial application showed a reduction in the number of circulating carcinoma cells in the blood of PyMT/*Ctsc*^{-/-} mice (n=3). However, increasing the number of animals analyzed has revealed no significant difference in circulating carcinoma cells when measured by flow cytometry (**Figure 6D**), or gene expression of the PyMT transgene or keratin 18 (**Figure 6C**). These results suggest that the ability to extravasate into the blood and survive in circulation are unaffected by CTSC. This supports the data from Figure 4, demonstrating no defect in the ability of CTSC-deficient TAMs to induce invasion.

Examine directional growth of neoplastic cells in the 3D organotypic model during co-culture with the leukocyte populations identified in Aim 1. (months 12-18)

These studies have not yet been initiated.

Examine the ability of leukocyte populations to induce chemotaxis of neoplastic cells using a modified Boyden chamber (months 18-21)

These studies have not yet been initiated.

Adoptive transfer of splenic T cells into PyMT/Ctsc^{-/-} mice and evaluation of lung metastasis and the number of circulating MECs. (months 18-24)

These studies have not yet been initiated.

Determine if cathepsin C regulates MEC survival in blood, lung seeding, or outgrowth:
Generate 4 neoplastic cell lines from PyMT/Ctsc^{+/-} and PyMT/Ctsc^{-/-} mice that are also transgenic for GFP (2 from early non-invasive mammary tumors and 2 from late, invasive tumors). (months 1-6)

Neoplastic cells have been purified from two late stage PyMT/Ctsc^{+/-} and PyMT/Ctsc^{-/-} mice and one set of cells from each genotype has been used in a preliminary experiment (see below). Breeding issues have delayed generating PyMT/Ctsc^{+/-} and PyMT/Ctsc^{-/-} mice that also express GFP under control of the β -actin promoter, but these mice have now been generated and mice are aging out.

Intravenously inject neoplastic cell lines in Ctsc^{+/-} and Ctsc^{-/-} mice and measure MEC survival in the blood, lung infiltration, and outgrowth/tumor progression. (months 12-18)

One experiment has so far been completed wherein neoplastic cells from PyMT/Ctsc^{+/-} and PyMT/Ctsc^{-/-} mice were intravenously injected into Ctsc^{+/-} and Ctsc^{-/-} mice and the formation of metastatic lesions analyzed after 14 days. Surprisingly, CTSC-proficient neoplastic cells in 3/5 mice were able to form a substantial number of foci, irrespective of the host genotype (**Figure 6E**). CTSC-deficient neoplastic cells meanwhile, formed few lesions in all mice. Repetition of this experiment is currently underway to determine if this is a valid finding, but if true, it suggests that intrinsic expression of CTSC by tumor cells is important for the later stages of metastasis. Normal levels of circulating carcinoma cells in PyMT animals (Figure 6C and D) suggests this effect would not be due to survival in the blood, but either intravasation into the lung or survival/growth within the lung. GFP expressing neoplastic cells will be used in the future to differentiate between these possibilities.

If extravasation into the lung is affected by cathepsin C expression, then examine rolling, adhesion, and transmigration of neoplastic cell lines on immortalized mouse endothelial cells. (months 18-24)

These studies have not yet been initiated.

If tumor growth or development is altered, inject neoplastic cells subcutaneously in a xenograft model along with purified tumor-associated leukocytes and measure early growth. (months 18-24)

These studies have not yet been initiated.

Determine if metastasis of mammary tumors is regulated by leukocyte-derived CTSC:

Generate bone marrow chimeric mice by using irradiated PyMT/Ctsc^{+/-} and PyMT/Ctsc^{-/-} mice and the bone marrow from GFP⁺ and GFP⁺/Ctsc^{-/-} mice. (months 12-24)

Analyze tumor incidence, growth, progression and metastasis in the four combinations of chimeric mice. (months 18-36)

If tissue cathepsin C expression appears to be important, generate chimeric mice that do not express PyMT and measure tumor growth in a xenograft model and lung metastasis following intravenous injection. (months 18-36)

These studies have not yet been initiated.

AIM 3: DEVELOP DIAGNOSTIC AND THERAPEUTIC APPROACHES BASED ON CATHEPSIN C BIOACTIVITY TO IMPROVE BREAST CANCER PATIENT SURVIVAL

Develop a probe to measure cathepsin C activity in live mice:

Generate FY01-GdDTPA and FY01-FITC. (months 12-18)

Adjust animal protocol to include use of FY01-GdDTPA and JCP410. (months 12-18)

Evaluate entrance of labeled FY01 into purified TAMs (months 18-21)

Evaluate entrance of labeled FY01 into TAMs in vivo (months 18-21)

Analyze entrance of FY01-GdDTPA into tumors using MRI. (months 21-24)

These studies have not yet been initiated

Evaluate efficacy of a selective cathepsin C inhibitor in reducing pulmonary metastasis:

Examine the effect of the selective cathepsin C inhibitor JCP410 in the 3D organotypic co-culture model. (months 24-30)

Evaluate the ability of JCP410 to inhibit cathepsin C activity in mammary and lung tumors by analyzing post-injection activity by MRI or staining unfixed tissue sections. (months 24-36)

Evaluate the efficacy of JCP410 in inhibiting pulmonary metastasis through bi-weekly i.p injection in mice of age 60 and 80 days. (months 24-36)

These studies have not yet been initiated

Evaluate cathepsin C activity during human breast cancer progression:

Submit proposal for use of human tissue (months 12-24)

Analyze expression of cathepsin C in fixed sections of human mammary tumors by double staining for cathepsin C and leukocyte markers followed by confocal microscopy. (months 24-36)

If cathepsin C expression is observed, measure activity in unfixed tissue sections using FY02 (months 24-36)

Correlate cathepsin C expression and activity with information available about the clinical stage of the cancer and/or the outcome. (months 30-36)

Analyze breast cancer tissue microarray data with 10 year follow up data for a correlation between cathpesin C expression and clinical stage and/or outcome. (months 30-36)

Evaluate role of human leukocytes to promote invasive growth in the 3D organotypic co-culture model using human breast cancer cell lines (months 24-30)

Evaluate ability of JCP410 to inhibit the promotion of invasive growth by leukocytes (months 30-36)

These studies have not yet been initiated

III. KEY RESEARCH ACCOMPLISHMENTS:

Aim 1: Determine the functional role of cathepsin C in regulating leukocyte infiltration and bioactivity in mammary adenocarcinoma development and metastasis

Months 1-12

- Predominant expression of CTSC by TAMs in mammary tumors and lungs.
- Low expression levels of CTSC by stromal and neoplastic cells in tumors.
- CTSC activity and protein levels increased in tumors versus mammary glands.
- CTSC activity localized to leukocytes and TAMs in mammary tumors.
- CTSC expression by TAMs does not affect their ability to induce invasion by MECs.
- CTSC expression does not affect the composition of leukocytes within the tumor.
- CTSC expression does not affect the composition of the stromal compartment.

Aim 2: Determine at which stage of cancer development cathepsin C functionally regulates pulmonary metastasis of mammary carcinogenesis

Months 1-12

- CTSC expression does not influence the number of circulating neoplastic cells
- Preliminary data suggesting CTSC expression by neoplastic cells is important in mediating the later stages of lung metastasis.

Aim 3: Develop diagnostic and therapeutic approaches based on cathepsin C bioactivity to improve breast cancer patient survival

Months 1-12

- These studies have not yet been initiated

IV. REPORTABLE OUTCOMES:

Manuscripts:

Ruffell, B., D. G. Denardo, N. I. Affara, and L. M. Coussens. 2010. Lymphocytes in cancer development Polarization towards pro-tumor immunity. *Cytokine Growth Factor Rev* 21:3-10.

Abstracts/Posters:

- 'Role of Cathepsin C During Breast Cancer Metastasis', Breast Oncology Program Retreat. San Francisco, CA. January 29-30, 2009.
- 'Role of Cathepsin C During Breast Cancer Metastasis', Diller Building Scientific Symposium. San Francisco, CA. August 27-28, 2009.

Presentations:

- 'Role of Cathepsin C During Breast Cancer Metastasis', Stanford Univ. School of Medicine. August, 2009.
- 'Role of Cathepsin C in Cancer Progression', Pacific Coast Protease Meeting, Borrego Springs, CA. April 2010.

V. CONCLUSION:

It is now clear that infiltrating immune cells profoundly regulate solid tumor development. Diverse populations of cells infiltrate tumor microenvironments and are altered by cancer cells and surrounding stroma in such a way that many of their bioactivities are co-opted to aid growth and/or metastasis of tumors. Out of this realization emerges novel targets for therapy that may impact patient survival, i.e., immune targets regulating primary tumor development, and those regulating metastatic dissemination and growth. It is thus hypothesized that manipulating the immune response and neutralizing its effect on neoplastic cells represents an efficacious alternative approach to current disease management. Preliminary analyses using a mouse model of breast cancer supports this hypothesis, by demonstrating that absence of a single immune cell enzyme reduces the ability of malignant breast cancer cells to metastasize to the lungs. Results from this study will therefore prove significant for elucidation of an important regulatory cascade affecting pulmonary metastasis. Furthermore, as metastasis is a leading cause of mortality in breast cancer, this research may lead to the identification of a novel therapeutic target for reducing breast cancer mortality. This is an innovative approach to formulating a novel anti-cancer therapy, as it targets a molecule expressed by normal cells. This contrasts with the traditional development of anti-cancer therapies, which are most often based upon molecules expressed by cancer cells that are important for their survival or growth. This innovative approach has two potential advantages over the traditional method: it avoids selecting for treatment-resistant neoplastic cells by instead targeting the pro-tumor activities of the microenvironment; and by targeting a specific pro-tumor pathway, there is a strong possibility of developing a treatment with low toxicity.

VI. BIBLIOGRAPHY:

1. Coussens, L. M., and Z. Werb. 2002. Inflammation and cancer. *Nature* 420:860-867.
2. Ruffell, B., D. G. Denardo, N. I. Affara, and L. M. Coussens. 2010. Lymphocytes in cancer development: Polarization towards pro-tumor immunity. *Cytokine Growth Factor Rev* 21:3-10.
3. Yuan, F., S. H. Verhelst, G. Blum, L. M. Coussens, and M. Bogoy. 2006. A selective activity-based probe for the papain family cysteine protease dipeptidyl peptidase I/cathepsin C. *J Am Chem Soc* 128:5616-5617.
4. DeNardo, D. G., J. B. Baretto, P. Andreu, L. Vasquez, N. Kolhatkar, D. Tawfik, and L. M. Coussens. 2009. CD4⁺ T cells regulate pulmonary metastasis of mammary carcinomas by enhancing protumor properties of macrophages. *Cancer Cell* 16:91-102.

VII. APPENDICES:

- A. Complete academic Curriculum vitae for Dr. Brian Ruffell
- B. Collected publications from months 1-12.

BIOGRAPHICAL SKETCH

NAME Brian Howard Ruffell	POSITION TITLE Postdoctoral Scholar University of California, San Francisco
-------------------------------------	---

EDUCATION/TRAINING <i>(Begin with baccalaureate or other initial professional education, such as nursing, and include postdoctoral)</i>			
INSTITUTION AND LOCATION	DEGREE <i>(if applicable)</i>	YEAR(s)	FIELD OF STUDY
University of British Columbia, Vancouver, BC	BSc	2001	Cell Biology/Genetics
University of British Columbia, Vancouver, BC	PhD	2008	Immunology
University of California, San Francisco, CA	n/a	Current	Cancer Immunology

POSITIONS AND HONORS:

- 1997 - 2001 Undergraduate Student: Department of Science. University of British Columbia, Vancouver, BC, Canada
- 2002 - 2008 Graduate Student: Department of Microbiology and Immunology. University of British Columbia, Vancouver, BC, Canada
- 2008 – pres. Postdoctoral Scholar: Department of Pathology, University of California, San Francisco, San Francisco, CA.

FELLOWSHIPS AND AWARDS:

- 2009 DOD Breast Cancer Research Program Postdoctoral Award
- 2006 Heart and Stroke Foundation of Canada Doctoral Research Award
- 2006 Robert Emmanuel and Mary Day Endowment Award
- 2005 John Richard Turner Fellowship in Microbiology
- 2005 Armauer-Hansen Memorial Scholarship
- 1998 Outstanding Student Initiative Scholarship
- 1997 British Columbia Government Scholarship
- 1997 Canadian Armed Forces Memorial Scholarship
- 1997 Outstanding Student Initiative Scholarship

PEER REVIEWED PUBLICATIONS:

- Ruffell B**, DeNardo DG, Affara, NI, Coussens LM. (2010) Lymphocytes in cancer development: polarization towards pro-tumor immunity. Cytokine Growth Factor Rev. 21(1): 3-10.
- Johnson P, **Ruffell B**. (2009) CD44 and its role in inflammation and inflammatory diseases. Inflamm. Allergy Drug Targets. 8(3): 208-220.
- Ruffell B**, Johnson P. (2009) The regulation and function of hyaluronan binding by CD44 in the immune system. Invited review for "Glycoforum" www.glycoforum.gr.jp
- Ruffell B**, Johnson P. (2008) Hyaluronan induces cell death in activated T cells through CD44. J. Immunol. 181: 7044-7054.
- Ruffell B**, Johnson P. (2005) Chondroitin sulfate addition to CD44H negatively regulates hyaluronan binding. Biochem. Biophys. Res. Commun. 334(2): 306-312.
- Khan AI, Kerfoot SM, Heit B, Liu L, Andonegui G, **Ruffell B**, Johnson P, Kubes P. (2004) Role of CD44 and hyaluronan in neutrophil recruitment. J. Immunol. 173(12): 7594-7601.

MANUSCRIPTS IN PREPARATION:

7. **Ruffell B**, Brown KL, Tjew SL, Johnson P. Chondroitin sulfate addition to CD44 regulates hyaluronan binding in macrophages.

PUBLISHED ABSTRACTS:

1. **Ruffell B**, Johnson P. (2004) Chondroitin sulfate chains on CD44 negatively regulate its ability to bind hyaluronan. Glycobiology 14(11): 1108.
2. **Ruffell B**, Johnson P. (2006) Hyaluronan induces apoptosis through CD44 in activated lymphoma cells. EJC Supplements 4(12): 147.

POSTER PRESENTATIONS:

1. US/Japan Glycobiology 2004. Honolulu, HI. November 17-20, 2004
2. 18th EORTC-NCI-AACR: Symposium on Molecular Targets and Cancer Therapeutics. Prague, Czech Republic. November 7-10, 2006
3. Breast Oncology Program Retreat. San Francisco, CA. January 29-30, 2009.
Winner of “Best-in-Show Poster Presentation” Award
4. Diller Building Scientific Symposium. San Francisco, CA. August 27-28, 2009.

CONFERENCES:

1. US/Japan Glycobiology 2004. Honolulu, HI. November 17-20, 2004
2. 18th EORTC-NCI-AACR: Symposium on Molecular Targets and Cancer Therapeutics. Prague, Czech Republic. November 7-10, 2006
3. Breast Oncology Program Retreat. San Francisco, CA. January 29-30, 2009.
4. 6th International Symposium on the Intraductal Approach to Breast Cancer. Santa Monica, CA. February 19-21, 2009.
5. Diller Building Scientific Symposium. San Francisco, CA. August 27-28, 2009.
6. Molecular and Cellular Biology of Immune Escape in Cancer. Keystone, CO. February 7-12, 2010

ORAL PRESENTATIONS:

1. Life Sciences Institute, Vancouver, BC (Jan, 2008)
2. BC Cancer Research Centre, Vancouver, BC (May, 2008)
3. Deeley Research Centre, Victoria, BC (May, 2008)
4. Cancer Research Institute, San Francisco, CA (June 2008)
5. Susan Love Research Foundation, Los Angeles, CA (Jan, 2009)
6. Stanford University School of Medicine, Stanford, CA (Aug, 2009)
7. Pacific Coast Protease Meeting, Borrego Springs, CA (April, 2010)

ADDITIONAL TRAINING:

1. BD LSR^{II} Operator Training Course (August, 2009)

TEACHING:

2003 - 2006 Teaching Assistant, Department of Microbiology and Immunology, University of British Columbia, Vancouver, BC, Canada

MENTORING:

<u>Dates</u>	<u>Name</u>	<u>Program</u>	<u>Role</u>	<u>Current Position</u>
2003	Patrick Yang	Summer Student	Supervisor	PhD/MD Program
2006-2008	Grace Poon	Co-op / Graduate Student	Mentor	PhD Program
2010	Lucia Cottone	PhD Exchange Student	Supervisor	PhD Program
2010	Heather Chan	Undergrad Summer Student	Supervisor	Undergraduate



Contents lists available at ScienceDirect

Cytokine & Growth Factor Reviews

journal homepage: www.elsevier.com/locate/cytogfr



Lymphocytes in cancer development: Polarization towards pro-tumor immunity

Brian Ruffell^a, David G. DeNardo^a, Nesrine I. Affara^a, Lisa M. Coussens^{a,b,*}

^a Department of Pathology, University of California, San Francisco, CA 94143, United States

^b Helen Diller Family Comprehensive Cancer Center, University of California, San Francisco, CA 94143, United States

ARTICLE INFO

Article history:
Available online xxx

Keywords:
Lymphocytes
Inflammation
Cancer
Tumors

ABSTRACT

The classic view that the role of immune cells in cancer is primarily one of tumor rejection has been supplanted by a more complex view of leukocytes having both pro- and anti-tumor properties. This shift is due to the now well recognized capabilities of several myeloid cell types that foster pro-tumor programming of premalignant tissue, as well as the discovery that subsets of leukocytes also suppress development and effector functions of lymphocytes important for mediating anti-tumor immunity. In this review, we focus on the underappreciated role that T lymphocytes play in promoting tumor development. This includes, in addition to the role of T regulatory cells, a role for natural killer T cells and CD4⁺ T helper cells in suppressing anti-tumor immunity and promoting cancer growth and metastasis.
© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Leukocyte infiltration into developing tumors is now considered one of the hallmarks of cancer development [1]. It is thought that the initial immune response to an early neoplasm mirrors the response to acute tissue injury, with sequential infiltration by various myeloid populations leading to eventual infiltration by lymphocytes [2]. However, as the kinetics of tumor development and the neoplastic cells themselves alter the local immune microenvironment, making inferences between an immune response to injury/infection and tumor development is difficult. Regardless, if clearance of the would-be cancer cells is not achieved and the initial acute inflammatory response fails to resolve, there inevitably results a state of chronic inflammation within the local tissue. It is now well established that chronic inflammation fosters early cancer development through a number of mechanisms mediated primarily by myeloid-lineage cells, including tumor-associated macrophages, immature myeloid cells that can possess suppressive activity, and Tie2-expressing monocytes [3,4]. The immune microenvironment of a neoplastic tissue encompasses not only the composition of infiltrating leukocytes, but also the bioeffector function of these cells within the tissue. Thus, both the presence of a cell within a tumor and expression of tissue-specific cytokines, chemokines and other immune mediators profoundly influence whether an anti-tumor or pro-tumor immune response is elicited [4,5].

Although responding to tissue damage in the form of inflammatory cues, tumor-infiltrating myeloid cells rapidly respond to soluble and insoluble signals emanating from the neoplastic microenvironment. Responses take the form of dramatically altered gene expression programs that alter bioeffector functions of the immune cells. These often result in increased expression of factors/mediators that enhance growth and survival of neoplastic cells, as well as activating and sustaining angiogenic responses, furthering tissue remodeling, and squelching anti-tumor immune programs [4]. Chronic inflammation in tissue resulting from infection or autoimmune disease can also alter the risk of cancer development by providing an environment permissive for initiated preneoplastic cell survival and subsequent proliferation, as well as through production of DNA damaging compounds such as reactive oxygen and nitrogen species that increase mutation frequency [6]. While all of these aspects of solid tumor development are susceptible to regulation by infiltrating immune cells, in the context of this review, we will focus on aspects of carcinogenesis regulated by infiltrating lymphocytes, as mechanisms regulated by myeloid cells have been reviewed elsewhere [5–9].

1.1. T lymphocytes

T cells develop in the thymus from a common lymphoid progenitor and are defined by expression of a T cell receptor (TCR) that is responsible for recognizing antigens presented by the major histocompatibility complex (MHC) family of genes (also called human leukocyte antigen or HLA). T cells are classically divided into either CD8⁺ cytotoxic lymphocytes (CTL) or CD4⁺ T helper (T_H) cells that recognize peptides presented by MHCI or MHCII, respectively (Fig. 1). T_H cells are further divided into interferon

* Corresponding author at: Department of Pathology, 513 Parnassus Avenue, HSX-450C, University of California, San Francisco, CA 94143, United States.
E-mail address: lisa.coussens@ucsf.edu (L.M. Coussens).




Fig. 1. T cell lineages and subsets. Successful rearrangement and expression of a TCR determines lineage commitment between $\gamma\delta$ and $\alpha\beta$ T cells. Recognition by $\alpha\beta$ T cells of MHC I, MHC II, or CD1d drives CD8⁺, CD4⁺ or NKT cell development, respectively. Strong recognition of peptide:MHCII complexes by CD4⁺ cells drives natural T_{Reg} cell development in the thymus, otherwise CD4⁺ T cells differentiate into T_H1, T_H2, T_H17 or inducible T_{Reg} cells following activation in the periphery, with polarization directed by IL-4, IL-6, IL-12 and TGF- β . Type I NKT cells are defined by expression of specific α -chain regions (V α 14-J α 18 in mice, V α 24-J α 18 in humans), but the reason for functional differences between type I and type II NKT cells is unclear.

(IFN)- γ and tumor necrosis factor (TNF)- α expressing T_H1 cells and interleukin (IL)-4, IL-5 and IL-13 expressing T_H2 cells. This simplified view of the T cell compartment has been expanded upon by the identification of a range of additional subtypes, including T follicular helper cells (T_{FH}), IL-17 expressing T_H cells (T_H17), and regulatory T cells (T_{Reg}) [10]. Paralleling these subtypes in the CD4⁺ T cell compartment, type 1, type 2, and type 17 CD8⁺ T cells (T_C1, T_C2, T_C17), as well as regulatory CD8⁺ cells, have all been described [11–13]. There also exist two ‘innate-like’ T cell subsets that can be activated either by cytokines or TCR stimulation. Natural killer T (NKT) cells recognize glycolipids presented by the non-classical MHC molecule CD1d [14], while $\gamma\delta$ T cells are not MHC restricted and recognize a diverse range of molecules, including soluble non-protein antigens [15]. All of these T lymphocyte subsets have been examined for their role in tumor development and anti-tumor immunity, each with unique roles in directing the immune response.

1.2. Cytotoxic T lymphocytes

Mice harboring specific immune-based genetic deficiencies are more susceptible to formation of carcinogen-induced sarcomas, and depending on the specific defect, are also more prone to develop certain spontaneous tumors and lymphomas [16,17]. The ability of immune-deficient mice to reject and/or inhibit the

growth of many, but not all cell lines is also impaired. Numerous studies have shown that, due to their ability to produce IFN γ and directly kill target cells, both CD8⁺ CTLs and natural killer (NK) cells are the critical mediators of the anti-tumor response [16]. $\gamma\delta$ T cells, which share characteristics with both CTLs and NK cells, are also involved in the anti-tumor response in epithelial tissues such as the skin [18], where they can be the dominant T cell population [15]. The relative importance of CTLs, NK cells, and $\gamma\delta$ T cells is highly dependent upon the cancer model being used. Even in the skin, genetic deficiency of $\alpha\beta$ T cells increases sarcoma formation following administration of methylcholanthrene (MCA), but not 7,12-dimethylbenz[a]anthracene (DMBA) or 12-O-tetradecanoylphorbol-13-acetate (TPA) [18], while the absence of CD8⁺ T cells does not influence the development of neoplasms in a mouse model of de novo squamous carcinogenesis, e.g., K14-HPV16 mice [19].

Epidemiological studies of cancer incidence in acquired immune deficiency syndrome (AIDS) and organ transplant patients reveal that the relative risk (RR) for cancer development varies considerably depending upon organ site and cancer etiology [17,20] where viral-associated cancers, in particular Human Herpes Virus 8-associated Kaposi’s sarcoma, Epstein-Barr virus-associated Non-Hodgkin’s lymphoma and HPV-associated squamous carcinoma, are elevated in immune suppressed individuals due largely to lack of protection against viral infections or viral reactivation in the absence of T cells [21]. That said, some cancer types occur with increased frequency in selected groups of immune compromised patients for reasons unrelated to infection for example, chronic exposure to carcinogens (tobacco) for thoracic malignancies, whereas head and neck, esophageal, gastrointestinal and pancreatic cancers are increased in liver transplant patients associated with prior history of alcohol (and tobacco) use [22,23]. On the other hand, the RR for the most common non-viral-associated solid tumors of epithelial origin, including breast and prostate, are decreased in immune suppressed patients, with some of these having a RR less than 1.0 [17,20].

Examination of T cell infiltrates in tumors reveals cells that can display activation markers and are able to recognize tumor antigens [16], indicating that some tumors are indeed immunogenic and can induce an anti-tumor immune response. Tumor antigens encompass both neo and overexpressed antigens, e.g., c-myc, HER-2/neu and p53 [24], as well as host/stromal cell-derived antigens unique to individual tumors. Clinical studies have even reported accumulation of autoantibodies against extracellular matrix (ECM) components, including anti-collagen type I, III and V, as well as anti-fibronectin antibodies that accumulate in lung cancer and nasopharyngeal patients [25]. How then do neoplastic cells, expressing mutant proteins in an *inflammatory* microenvironment that seemingly engenders a robust T cell response, avoid killing by cytotoxic cells? Importantly, lymphocytes do not act in isolation, and their effector functions are largely dependent upon the release of cytokines and binding of inhibitory and activating receptors to ligands expressed by other leukocytes, stromal cells, and even neoplastic cells. For example, NK cells are potent regulators of CD8⁺ T cell responses through their release of IFN γ , which provides a maturation signal for tissue resident DCs and assists in CD8⁺ T cell effector function; meanwhile, cytokines released by mature DCs and activated T cells are important for promoting NK cell effector function [26]. These interweaving regulatory pathways are necessary to initiate, direct, maintain and eventually shutdown an appropriate immune response. Such pathways are also necessary to prevent an inappropriate immune response: despite central tolerance through negative selection of self-reactive lymphocytes, peripheral tolerance mediated by cytokines, inhibitory receptors and immune regulatory cell types is necessary to prevent autoimmune disorders. As cancer cells are

largely recognized as 'self', it is not surprising they utilize similar mechanisms that effectively dampen anti-tumor immunity; thus, T lymphocytes are intimately involved in regulating both pro- and anti-tumor immunity and chronic inflammation within the tumor microenvironment.

1.3. Regulatory T cells

Since the rediscovery of suppressor T cells as CD4⁺CD25⁺ regulatory T cells, and their further characterization as cells expressing glucocorticoid-induced tumor necrosis factor receptor (GITR), cytotoxic T lymphocyte antigen (CTLA)-4, and uniquely, the transcription factor forkhead box P3 (FoxP3), this T cell subset has become an intense focus of cancer research. T_{Reg} cells can develop in the thymus or can be converted in the periphery by exposure to transforming growth factor (TGF)- β [27]. These 'natural' and 'inducible' T_{Reg} cells, respectively, utilize the same mechanisms to mediate immune suppression and may perform overlapping functions [28]. A number of regulatory T cells that do not conform to the CD4⁺CD25⁺FoxP3⁺ phenotype have also been described, but these remain poorly characterized [29] and evidence of a role for these cells in cancer has so far been limited to the isolation of IL-10 producing CD8⁺ T cells from human ovarian tumors [30]. Immune suppression of CD4⁺ and CD8⁺ T cell responses by T_{Reg} cells is mediated both in the secondary lymphoid organs where T cell activation occurs, and in the tissues [31]. Interestingly, in a pancreatic islet allograft model, T_{Reg} cells were shown to enter the inflamed tissue and then migrate to the lymph nodes, where these sequential series of steps was found to be associated with increasing graft survival time [32].

While a host of molecules have been described as important for mediating T_{Reg} cell suppression, these molecules have been broadly classified as acting in one of four ways [33]: (i) cytokine inhibition, such as with TGF- β , IL-35 and IL-10; (ii) direct cytotoxicity of effector T cells through perforin and granzyme; (iii) metabolic disruption, such as IL-2 deprivation and cyclic adenosine monophosphate (cAMP) transfer; and (iv) inhibition of DC function, such as through binding of CTLA-4 to CD80/86 and the induction of indoleamine 2,3-dioxygenase (IDO). These mechanisms appear to have overlapping but non-redundant roles, with the degree of importance being tissue and model dependent [34]. This may be significant in cancer, as CD4⁺CD25^{high} tumor-infiltrating T lymphocytes from patients with head and neck squamous cell carcinoma (HNSCC) were found to mediate suppression through IL-10 and TGF- β , while the same population from peripheral blood did not express these cytokines and were less able to suppress proliferation [35]. A role for IL-10 and TGF- β has also been revealed in mice following injection of a fibrosarcoma cell line [36], while a role for perforin- and granzyme B-dependent killing of NK and CD8⁺ T cells was found in mice after injection of a lymphoma cell line [37]. Other than these studies, the mechanisms by which T_{Reg} cells mediate suppression in cancer have been largely ignored. Instead, research has been focused on correlating T_{Reg} cell infiltration with prognosis and attempts to deplete T_{Reg} cells with anti-CD25 antibody based therapies [29].

Suppression by T cells in solid tumors was first suggested by Fujimoto et al. [38], cumulating in work by North and colleagues, who showed that the ability to reject a second subcutaneous injection of a fibrosarcoma cell line was inversely correlated to increased suppressor activity of CD3⁺CD4⁺ cells over time [39,40]. These findings have since been expanded to show that depletion of CD25⁺ cells, which are largely CD4⁺FoxP3⁺, reduces tumor growth of some tumor cell lines [29], as well as MCA-induced fibrosarcomas [41,42]. Infiltration of T_{Reg} cells into the tumor is observed in all of these models, while an increased percentage of T_{Reg} cells in the periphery is observed only in some instances. In one study

using a fibrosarcoma cell line, depletion of CD4⁺ cells resulted in CD8⁺ T cell-dependent tumor regression in 50% of the mice [36], that was increased to 100% following either CD4⁺ or CD25⁺ cell depletion when another fibrosarcoma-derived cell line expressing a strong antigen was used [36]. Together with other studies, experimental systems such as these indicate that initiation of a CD8⁺ T cell response is possible during tumor development, depending on the immunogenicity of the tumor antigens involved, but that local and/or systemic immune suppression by T_{Reg} cells can limit their effectiveness. Unfortunately, these results have been limited in scope to transplantable tumor models representing few tumor types, and thus further investigation is required to determine whether T_{Reg}-dependent immune suppression is applicable to a wide range of spontaneous tumors, as might be expected.

Increased number of tumor-infiltrating FoxP3⁺ cells is associated with poor prognosis in several cancers, including hepatocellular carcinoma [43], ovarian carcinoma [41], pancreatic ductal carcinoma [44], cervical cancer [45], non-small cell lung carcinoma [46], HNSCC [35], and breast cancer [47,48]; with varying degrees of prognostic value regarding patient outcome. As with some mouse models, increased frequency of T_{Reg} cells in peripheral blood of some cancer patients has been reported [49]. Peripheral blood T_{Reg} cells from patients with ovarian cancer displayed equal suppressive capacity compared to tumor-derived T_{Reg} cells [41]. This contrasts with the study of patients with HNSCC [35], highlighting the potential for differences in the immune response based on cancer type and/or etiology.


Attempts to translate T_{Reg} research into the clinic have focused around CD25⁺ cell depletion using denileukin diftitox. Known commercially as Ontak, this compound composed of IL-2 fused to a portion of diphtheria toxin has been approved for treating CD25⁺ cutaneous T cell leukemia and lymphoma [29]. Ontak administration has been demonstrated to reduce the numbers of peripheral T_{Reg} cells and improve T cell activation in a small number of patients with either lung, ovarian, breast, or renal cancer; either alone or in combination with DC-based vaccination [50,51]. Other small studies have confirmed these findings, but as before, objective clinical responses were rare. Furthermore, a larger study with NSCLC patients observed no objective clinical responses, and almost half of the patients suffered side effects usually associated with IL-2 immunotherapy [52]. Dosage optimization thus remains an issue, as does route of administration, as intratumoral injection of anti-CD25 antibodies demonstrated efficacy in mouse models [36]. Determination of which cancer types are most suitable for T_{Reg} depletion is also required, as the ability of CD25⁺ cell depletion to improve anti-tumor immunity is known to depend on the tumor type [53]. Finally, combinatorial therapies, such as CD25⁺ cell depletion with blockage of immunosuppressive molecules such as CTLA-4 and programmed death (PD)-1 [54], have great potential to overcome immune suppression within the tumor, although the induction of autoimmune diseases will likely remain an issue for patients.

1.4. T helper cells

Lineage commitment between CD4⁺ and CD8⁺ T lymphocytes occurs during development within the thymus. Further differentiation of the CD4⁺ lineage, with the exception of natural T_{Reg} cell development, requires T cell activation through MHCII and co-stimulatory molecules, as well as cytokine dependent signaling that is responsible for directing the cell towards a particular lineage (Fig. 1). Classical differentiation of CD4⁺ T cells into T_H1 and T_H2 cells, mediated by IL-12 and IL-4 respectively, was recently updated to include a new T_H17 lineage [10,55]. T_H17 cells are induced by a combination of IL-6 and TGF- β and mediate their

However, as with inflammation in general, IL-17-dependent inflammation may have both positive and negative effects on tumor growth, depending on the tumor model. MC38 colon cancer cell growth is enhanced in IL-17-deficient mice [71], while adoptive transfer of *in vitro* polarized Th17 cells specific for a B16 melanoma antigen can induce tumor regression [72]. This effect appears to depend upon IFN γ , as blocking antibodies prevent Th17 cell transfer from causing tumor regression [72], while reduced frequency of IFN γ producing cells were observed in IL-17-deficient tumors [71]. Adding to the confusion is that both IL-21 and IL-22 activate Stat3, and IL-21 can promote Th17 differentiation [56]. Meanwhile, CD8⁺ T cells primed *in vitro* in the presence of

By virtue of reduced IFN γ production, T_H2 polarization is likely to be detrimental to the anti-tumor response. T_H2 polarization is dependent upon, and leads to, production of IL-4. This differs from T_H1 and T_H17 polarization, which are not induced by their respective cytokines, although these are involved in lineage stabilization [10,55]. In addition to reducing T_H1 polarization, IL-4 may have direct immunosuppressive effects on CD8⁺ T cells, as *in vitro* activation of naïve CD8⁺ T cells in the presence of IL-4 reduces effectiveness of adoptive transfer of tumor-specific transgenic T cells [94,95]. It should be noted however, that these cells, termed T_C2 (as opposed to T_C1 CD8⁺ T cells activated in the presence of IL-12), were still able to improve survival from B16 melanoma cells in the lung when transferred in greater quantities [95]. Subsequent work by the same group showed that IL-4 and IL-5 expression by the adoptively transferred T_C2 cells was important in mediating this effect [96]. Recombinant expression of IL-4 by several tumor lines also greatly improves clearance [97], and one



Please cite this article in press as: Ruffell B, et al. Lymphocytes in cancer development: Polarization towards pro-tumor immunity. Cytokine Growth Factor Rev (2009), doi:[10.1016/j.cytogfr.2009.11.002](https://doi.org/10.1016/j.cytogfr.2009.11.002)

study found that ovalbumin (OVA)-specific T_H2 polarized cells helped clear B16 melanoma lung colonies expressing OVA through recruitment of eosinophils [98]. Is IL-4 production, and by extension T_H2 polarization, therefore protective? Probably not, since the increased anti-tumor response does not depend upon endogenous IL-4, but instead upon an effective T_H1 and CTL response by infiltrating leukocytes [96,99]. By promoting inflammation and leukocyte recruitment, IL-4 production does appear to improve anti-tumor immunity. However, this protective effect is time-dependent, with later stage tumors more resistant to the adoptive transfer of IL-4 producing $CD8^+$ cells [95]. Adoptive transfer of IL-4 producing T_H2 cells, or administration of IL-4 intravenously, also increases lung colonization of B16 melanoma cells injected intravenously [100]. Thus, while IL-4 has the potential to improve anti-tumor immunity, its use may be limited therapeutically to inducing acute inflammation prior to the development of a chronically inflamed tumor microenvironment, with a T_H2 type response itself being detrimental for anti-tumor immunity.

Also produced by T_H2 cells, IL-13 affects the immune response in many of the same ways as IL-4 through activation of Stat6 [101]. As with IL-4, IL-13 expression by tumors can improve the anti-tumor response [102], while endogenous IL-13 can inhibit anti-tumor immunity [103]. T cells do not express the type II IL-4 receptor necessary for binding to IL-13 however, and IL-13 instead appears to inhibit the CTL response indirectly by increasing TGF- β production by myeloid cells in the tumor [104]. Notably however, the source of this IL-13 did not appear to be T_H2 cells in this model, but instead NKT cells [103]. Both IL-4 and IL-13 are considered to be the inducers of the M2 phenotype in monocytes/macrophages, reducing inflammatory cytokine expression and increasing expression of immune suppressive cytokines, resulting in indirect immunosuppression [7,105]. We have recently reported that IL-4 also affects late-stage mammary cancer development by mediating the activity of tumor-associated macrophages (TAMs) [77]. In the MMTV-PyMT mouse model of mammary carcinogenesis, lung metastasis was dramatically reduced by genetic deficiency of either CD4 or IL-4R α (which would prevent IL-4 and IL-13 signaling). *In vitro*, IL-4 expression by T_H2 cells was found to increase expression of epidermal growth factor (EGF) by TAMs, enhancing the ability of neoplastic cells to extravasate into the circulation. IL-4 and IL-13 may also affect tumor growth independent of their effects on the immune system, as many cell types express the type II IL-4 receptor that binds both IL-4 and IL-13 [106]. This has not been addressed by many studies, but IL-4 has been shown to reduce angiogenesis by inhibiting endothelial cell migration [107,108]. IL-4 and IL-13 also inhibit proliferation of several epithelial cancers, although IL-13 can also promote proliferation and/or inhibit apoptosis of some hematological malignancies [101] as well as breast carcinomas in some models [109].

1.5. Natural killer T cells

NKT cells play a key regulatory role in directing a T_H1 or T_H2 polarized immune response through the rapid production of IFN γ , TNF α , IL-4, and IL-13 following stimulation. This is observed in mice deficient for NKT cells, as infections in these mice, particularly bacterial or parasitic, are often more severe [110]. As with conventional CTL and T_H cells, NKT cells develop in the thymus and express the α and β chains of the TCR. Instead of recognizing a peptide presented by MHC class I or II molecules however, the TCR expressed by NKT cells recognizes glycolipid antigens presented by CD1d, a non-classical member of the MHC family [14]. Type I, or invariant, NKT cells (iNKT) express a specific alpha chain variable (V) and joining (J) region (V α 14-J α 18 in mice, V α 24-J α 18 in

humans) in combination with a limited number of β chains, and were identified for their ability to recognize α -galactosylceramide (α -GalCer). Type II NKT cells, while also recognizing CD1d, express a variety of $\alpha\beta$ TCR chains and are activated by glycolipids that remain poorly defined.

Interest in targeting NKT cells for anti-cancer therapy began with the discovery that treatment with α -GalCer increased the survival time of mice injected with B16 melanoma cells [111]. The anti-tumor effects of α -GalCer [112] and IL-12 treatment [113] were subsequently found to depend upon the presence of iNKT cells. J α 18-deficient mice, that lack iNKT cells, were also found to be more susceptible to MCA-induced fibrosarcomas [84]. Although capable of directly lysing tumor cells in a perforin-dependent manner, a series of studies by Godfrey and colleagues utilizing MCA-induced fibrosarcomas demonstrated that both NKT-dependent immune surveillance and protection provided by IL-12/ α -GalCer therapy was dependent upon IFN γ production by NKT cells, leading to CTL-dependent and NK-dependent anti-tumor responses, respectively [84,114,115]. It has yet to be determined if these same mechanisms are at play in solid tumors of epithelial origin or in models of de novo cancer development.

In addition to possible direct effects of IFN γ production by NKT cells on CTL and NK cells, α -GalCer treatment has been shown to act as an adjuvant through NKT-dependent DC activation and IL-12 production [116,117]. Pulsing DCs *in vitro* with α -GalCer prior to adoptive transfer also more effectively prevents liver metastasis of B16 melanoma cells than injection of α -GalCer [118], possibly by improving long-term IFN γ production and limiting T_H2 cytokine expression [119,120]. Unfortunately, despite these successes in murine tumor transplantation models, injection of α -GalCer, α -GalCer pulsed DCs, or transfer of α -GalCer activated NKT cells all proved ineffectual in early clinical trials in patients with a range of cancer types, even though NKT activation was evident [121].

Using transplantable sarcoma models in which immune surveillance was evident, work by Berzofsky and colleagues found increased resistance of CD1d-deficient mice to tumor development [122]. This was due to the absence of IL-13 producing NKT cells in these mice [103], which were responsible for promoting TGF β production by splenic CD11b $^+$ Gr-1 $^+$ immature myeloid cells in a model using NIH/3T3-derived cell lines [104]. By comparing J α 18-deficient (lacking type I iNKT cells) to CD1d-deficient mice (lacking both type I and type II NKT cells), they concluded that type II NKT cells were responsible for inhibiting immune surveillance in several models where CD25-depletion had no effect [123]. Importantly, in agreement with other studies, α -GalCer treatment increased protection in these models [124]. Meanwhile, stimulation of at least a portion of type II NKT cells with a sulfatide compound reduced protection, and could even counteract the protection offered by α -GalCer treatment [124]. The most parsimonious explanation for these observations is that IFN γ production by type I NKT cells improves the CTL and NK cell responses, while IL-13 production by type II NKT cells inhibits the immune response (Fig. 2). As T, NK, and NKT cells are unresponsive to IL-13, the results also suggest that the effects of NKT cells depend on an intermediate cell, such as DCs [125]. It should be noted however that studies using other tumor injection models found that improved protection of CD1d-deficient mice was not related to IL-13 [126,127]. Protection through adoptive transfer of NKT cells is also dependent on whether the NKT cells are CD4 $^+$ or CD4 $^-$, and upon the tissue used to isolate NKT cells, which relates at least partially to production of IL-4 [128]. In humans, peripheral blood CD4 $^-$ NKT cells expressed only IFN γ and TNF α , whereas CD4 $^+$ NKT cells produced both T_H1 and T_H2 type cytokines following stimulation [129,130]. A possible role for cytotoxic effector functions by NKT cells can also not be ruled out. Song and colleagues found that human NKT cells can directly kill monocytes

pulsed with lysate from human neuroblastoma cell lines [131]. Using immunocompromised mice, the authors found that human NKT cells reduced the number of monocytes in the tumors and inhibited growth of the xenograft, indicating that NKT cells also inhibit pro-tumor immunity by killing tumor-associated macrophages. Tissue localization and tumor type may therefore greatly affect cytokine expression by NKT cells and determine whether they promote or inhibit the anti-tumor response.

2. Conclusions

Although first described over 20 years ago [132], the complexity by which the immune system directs a T_H1 or T_H2 response is only now being appreciated. The concept of polarized populations of immune cells has now been expanded to include $CD8^+$ CTLs, macrophages, and NKT cells. Whether these populations can be defined by genetic programs remains to be determined, but the effects of this polarization on the anti-tumor response demonstrate the importance for further research. Inhibiting immune suppression by blocking the activity of regulatory immune cells, or blocking self-suppression of the CTL response by inhibitory molecules such as PD-1 and CTLA-4, holds great potential for improving anti-tumor responses. These approaches will likely be enhanced by therapies that also dampen the effect of pro-tumor immune molecules released by other leukocytes that enhance angiogenesis, tissue remodeling and cell survival pathways, and in combination, may increase clinical efficacy of adoptive transfer therapies to engender durable anti-tumor immunity. Early clinical results have shown success in some of these areas, although the ability to inhibit peripheral tolerance and improve the anti-tumor response appears directly related to the severity of autoimmunity that is also induced. Being able to release the full potential of the immune response, while also being able to appropriate direct and control that response, is key to the future of immunotherapy.

Acknowledgements

The authors would like to acknowledge all of the scientists who made contributions to the areas of research reviewed here that were not cited due to space constraints. The authors acknowledge support from Postdoctoral Awards from the Department of Defense Breast Cancer Research Program (BR), the American Cancer Society (DGN), and the American Association for Cancer Research (NIA), as well as grants from the National Institutes of Health, the Dr. Susan Love Research Foundation, and a Department of Defense Era of Hope Scholar Award (LMC).

References

- Colotta F, Allavena P, Sica A, Garlanda C, Mantovani A. Cancer-related inflammation, the seventh hallmark of cancer: links to genetic instability. *Carcinogenesis* 2009;30:1073–81.
- Clark CE, Hingorani SR, Mick R, Combs C, Tuveson DA, Vonderheide RH. Dynamics of the immune reaction to pancreatic cancer from inception to invasion. *Cancer Res* 2007;67:9518–27.
- De Palma M, Coussens LM. Immune cells and inflammatory mediators as regulators of tumor angiogenesis. In: Figg WD, Folkman J, editors. *Angiogenesis: an integrative approach from science to medicine*. New York: Springer; 2008. p. 225–38.
- Mantovani A, Allavena P, Sica A, Balkwill F. Cancer-related inflammation. *Nature* 2008;454:436–44.
- Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002;420:860–7.
- Balkwill F, Charles KA, Mantovani A. Smoldering and polarized inflammation in the initiation and promotion of malignant disease. *Cancer Cell* 2005;7:211–7.
- Pollard JW. Trophic macrophages in development and disease. *Nat Rev Immunol* 2009;9:259–70.
- Joyce JA, Pollard JW. Microenvironmental regulation of metastasis. *Nat Rev Cancer* 2009;9:239–52.
- Lewis CE, Pollard JW. Distinct role of macrophages in different tumor microenvironments. *Cancer Res* 2006;66:605–12.
- Wilson CB, Rowell E, Sekimata M. Epigenetic control of T-helper-cell differentiation. *Nat Rev Immunol* 2009;9:91–105.
- Croft M, Carter L, Swain SL, Dutton RW. Generation of polarized antigen-specific CD8 effector populations: reciprocal action of interleukin (IL)-4 and IL-12 in promoting type 2 versus type 1 cytokine profiles. *J Exp Med* 1994;180: 1715–28.
- Hamada H, Garcia-Hernandez Mde L, Reome JB, Misra SK, Strutt TM, McKinsty KK, et al. Tc17, a unique subset of CD8 T cells that can protect against lethal influenza challenge. *J Immunol* 2009;182:3469–81.
- Lu L, Cantor H. Generation and regulation of CD8(+) regulatory T cells. *Cell Mol Immunol* 2008;5:401–6.
- Godfrey DI, Berzins SP. Control points in NKT-cell development. *Nat Rev Immunol* 2007;7:505–18.
- Carding SR, Egan PJ, Gammadelta. T cells: functional plasticity and heterogeneity. *Nat Rev Immunol* 2002;2:336–45.
- Dunn GP, Old LJ, Schreiber RD. The three Es of cancer immunoediting. *Annu Rev Immunol* 2004;22:329–60.
- de Visser KE, Eichten A, Coussens LM. Paradoxical roles of the immune system during cancer development. *Nat Rev Cancer* 2006;6:24–37.
- Girardi M, Oppenheim DE, Steele CR, Lewis JM, Glusac E, Filler R, et al. Regulation of cutaneous malignancy by gammadelta T cells. *Science* 2001;294:605–9.
- Daniel D, Meyer-Morse N, Bergsland EK, Dehne K, Coussens LM, Hanahan D. Immune enhancement of skin carcinogenesis by CD4+ T cells. *J Exp Med* 2003;197:1017–28.
- Patel P, Hanson DL, Sullivan PS, Novak RM, Moorman AC, Tong TC, et al. Incidence of types of cancer among HIV-infected persons compared with the general population in the United States, 1992–2003. *Ann Intern Med* 2008;148:728–36.
- Boshoff C, Weiss R. AIDS-related malignancies. *Nat Rev Cancer* 2002;2:373–82.
- Herrero JL, Lorenzo M, Quiroga J, Sangro B, Pardo F, Rotellar F, et al. De Novo neoplasia after liver transplantation: an analysis of risk factors and influence on survival. *Liver Transpl* 2005;11:89–97.
- Fung JJ, Jain A, Kwak EJ, Kusne S, Dvorchik I, Eghtesad B. De novo malignancies after liver transplantation: a major cause of late death. *Liver Transpl* 2001;7:5109–18.
- Abu-Shakra M, Buskila D, Ehrenfeld M, Conrad K, Shoenfeld Y. Cancer and autoimmunity: autoimmune and rheumatic features in patients with malignancies. *Ann Rheum Dis* 2001;60:433–41.
- Bei R, Masuelli L, Palumbo C, Modesti M, Modesti A. A common repertoire of autoantibodies is shared by cancer and autoimmune disease patients: Inflammation in their induction and impact on tumor growth. *Cancer Lett* 2009;281:8–23.
- Vivier E, Tomasello E, Baratin M, Walzer T, Ugolini S. Functions of natural killer cells. *Nat Immunol* 2008;9:503–10.
- Josefowicz SZ, Rudensky A. Control of regulatory T cell lineage commitment and maintenance. *Immunity* 2009;30:616–25.
- Curotto de Lafaille MA, Lafaille JJ. Natural and adaptive foxp3+ regulatory T cells: more of the same or a division of labor? *Immunity* 2009;30:626–35.
- Zou W. Regulatory T cells, tumour immunity and immunotherapy. *Nat Rev Immunol* 2006;6:295–307.
- Wei S, Kryczek I, Zou L, Daniel B, Cheng P, Mottram P, et al. Plasmacytoid dendritic cells induce CD8+ regulatory T cells in human ovarian carcinoma. *Cancer Res* 2005;65:5020–6.
- Rudensky AY, Campbell DJ. In vivo sites and cellular mechanisms of T reg cell-mediated suppression. *J Exp Med* 2006;203:489–92.
- Zhang N, Schroppel B, Lal G, Jakubczik C, Mao X, Chen D, et al. Regulatory T cells sequentially migrate from inflamed tissues to draining lymph nodes to suppress the alloimmune response. *Immunity* 2009;30:458–69.
- Vignali DA, Collison LW, Workman CJ. How regulatory T cells work. *Nat Rev Immunol* 2008;8:523–32.
- Tang Q, Bluestone JA. The Foxp3+ regulatory T cell: a jack of all trades, master of regulation. *Nat Immunol* 2008;9:239–44.
- Strauss L, Bergmann C, Szczepanski M, Gooding W, Johnson JT, Whiteside TL. A unique subset of CD4+CD25highFoxp3+ T cells secreting interleukin-10 and transforming growth factor-beta1 mediates suppression in the tumor micro-environment. *Clin Cancer Res* 2007;13:4345–54.
- Yu P, Lee Y, Liu W, Krausz T, Chong A, Schreiber H, Fu YX. Intratumor depletion of CD4+ cells unmasks tumor immunogenicity leading to the rejection of late-stage tumors. *J Exp Med* 2005;201:779–91.
- Cao X, Cai SF, Fehniger TA, Song J, Collins LI, Pniwnica-Worms DR. Granzyme B and perforin are important for regulatory T cell-mediated suppression of tumor clearance. *Immunity* 2007;27:635–46.
- Fujimoto S, Greene M, Sehon AH. Immunosuppressor T cells in tumor bearing host. *Immunol Commun* 1975;4:201–17.
- Berend MJ, North RJ. T-cell-mediated suppression of anti-tumor immunity. An explanation for progressive growth of an immunogenic tumor. *J Exp Med* 1980;151:69–80.
- Awwad M, North RJ. Immunologically mediated regression of a murine lymphoma after treatment with anti-L3T4 antibody. A consequence of removing L3T4+ suppressor T cells from a host generating predominantly Lyt-2+ T cell-mediated immunity. *J Exp Med* 1988;168:2193–206.
- Curiel TJ, Coukos G, Zou L, Alvarez X, Cheng P, Mottram P, et al. Specific recruitment of regulatory T cells in ovarian carcinoma fosters immune privilege and predicts reduced survival. *Nat Med* 2004;10:942–9.

- [42] Betts G, Twhog J, Van den Broek M, Sierro S, Godkin A, Gallimore A. The impact of regulatory T cells on carcinogen-induced sarcogenesis. *Br J Cancer* 2007;96:1849–54.
- [43] Gao Q, Qiu SJ, Fan J, Zhou J, Wang XY, Xiao YS, et al. Intratumoral balance of regulatory and cytotoxic T cells is associated with prognosis of hepatocellular carcinoma after resection. *J Clin Oncol* 2007;25:2586–93.
- [44] Hiraoka N, Onozato K, Kosuge T, Hirohashi S. Prevalence of FOXP3+ regulatory T cells increases during the progression of pancreatic ductal adenocarcinoma and its premalignant lesions. *Clin Cancer Res* 2006;12:5423–34.
- [45] Jordanova ES, Gorter A, Ayachi O, Prins F, Durrant LG, Kenter GG, et al. Human leukocyte antigen class I. MHC class I chain-related molecule A, and CD8+ regulatory T-cell ratio: which variable determines survival of cervical cancer patients? *Clin Cancer Res* 2008;14:2028–35.
- [46] Petersen RP, Campa MJ, Sperlazza J, Conlon D, Joshi MB, Harpole Jr DH, Patz Jr EF. Tumor infiltrating Foxp3+ regulatory T-cells are associated with recurrence in pathologic stage I NSCLC patients. *Cancer* 2006;107:2866–72.
- [47] Bates GJ, Fox SB, Han C, Leek RD, Garcia JF, Harris AL, Banham AH. Quantification of regulatory T cells enables the identification of high-risk breast cancer patients and those at risk of late relapse. *J Clin Oncol* 2006;24:5373–80.
- [48] Merlo A, Casalini P, Carcangiu ML, Malventano C, Triulzi T, Menard S, et al. FOXP3 expression and overall survival in breast cancer. *J Clin Oncol* 2009;27:1746–52.
- [49] Betts GJ, Clarke SL, Richards HE, Godkin AJ, Gallimore AM. Regulating the immune response to tumours. *Adv Drug Deliv Rev* 2006;58:948–61.
- [50] Barnett B, Kryczek I, Cheng P, Zou W, Curiel TJ. Regulatory T cells in ovarian cancer: biology and therapeutic potential. *Am J Reprod Immunol* 2005;54:369–77.
- [51] Dannull J, Su Z, Rizzieri D, Yang BK, Coleman D, Yancey D, et al. Enhancement of vaccine-mediated antitumor immunity in cancer patients after depletion of regulatory T cells. *J Clin Invest* 2005;115:3623–33.
- [52] Gerena-Lewis M, Crawford J, Bonomi P, Maddox AM, Hainsworth J, McCune DE, et al. A Phase II Trial of Denileukin Diftitox in Patients With Previously Treated Advanced Non-Small Cell Lung Cancer. *Am J Clin Oncol* 2009.
- [53] Onizuka S, Tawara I, Shimizu J, Sakaguchi S, Fujita T, Nakayama E. Tumor rejection by in vivo administration of anti-CD25 (interleukin-2 receptor alpha) monoclonal antibody. *Cancer Res* 1999;59:3128–33.
- [54] Fong L, Small EJ. Anti-cytotoxic T-lymphocyte antigen-4 antibody: the first in an emerging class of immunomodulatory antibodies for cancer treatment. *J Clin Oncol* 2008;26:5275–83.
- [55] Weaver CT, Harrington LE, Mangan PR, Gavrieli M, Murphy KM. Th17: an effector CD4 T cell lineage with regulatory T cell ties. *Immunity* 2006;24:677–88.
- [56] Dong C. TH17 cells in development: an updated view of their molecular identity and genetic programming. *Nat Rev Immunol* 2008;8:337–48.
- [57] Ouyang W, Kolls JK, Zheng Y. The biological functions of T helper 17 cell effector cytokines in inflammation. *Immunity* 2008;28:454–67.
- [58] Bettelli E, Korn T, Oukka M, Kuchroo VK. Induction and effector functions of T(H)17 cells. *Nature* 2008;453:1051–7.
- [59] Miyahara Y, Odunsi K, Chen W, Peng G, Matsuzaki J, Wang RF. Generation and regulation of human CD4+ IL-17-producing T cells in ovarian cancer. *Proc Natl Acad Sci USA* 2008;105:15505–10.
- [60] Sfanos KS, Bruno TC, Maris CH, Xu L, Thoburn CJ, DeMarzo AM, et al. Phenotypic analysis of prostate-infiltrating lymphocytes reveals TH17 and Treg skewing. *Clin Cancer Res* 2008;14:3254–61.
- [61] Zhang B, Rong G, Wei H, Zhang M, Bi J, Ma L, et al. The prevalence of Th17 cells in patients with gastric cancer. *Biochem Biophys Res Commun* 2008;374:533–7.
- [62] Zhang JP, Yan J, Xu J, Pang XH, Chen MS, Li L, et al. Increased intratumoral IL-17-producing cells correlate with poor survival in hepatocellular carcinoma patients. *J Hepatol* 2009;50:980–9.
- [63] Numasaki M, Watanabe M, Suzuki T, Takahashi H, Nakamura A, McAllister F, et al. IL-17 enhances the net angiogenic activity and in vivo growth of human non-small cell lung cancer in SCID mice through promoting CXCR-2-dependent angiogenesis. *J Immunol* 2005;175:6177–89.
- [64] Numasaki M, Fukushi J, Ono M, Narula SK, Zavodny PJ, Kudo T, et al. Interleukin-17 promotes angiogenesis and tumor growth. *Blood* 2003;101:2620–7.
- [65] Nam JS, Terabe M, Kang MJ, Chae H, Voong N, Yang YA, et al. Transforming growth factor beta subverts the immune system into directly promoting tumor growth through interleukin-17. *Cancer Res* 2008;68:3915–23.
- [66] Xiao M, Wang C, Zhang J, Li Z, Zhao X, Qin Z. IFN-gamma promotes papilloma development by up-regulating Th17-associated inflammation. *Cancer Res* 2009;69:2010–7.
- [67] Wang L, Yi T, Kortylewski M, Pardoll DM, Zeng D, Yu H. IL-17 can promote tumor growth through an IL-6-Stat3 signaling pathway. *J Exp Med* 2009;206:1457–64.
- [68] Yu H, Kortylewski M, Pardoll D. Crosstalk between cancer and immune cells: role of STAT3 in the tumour microenvironment. *Nat Rev Immunol* 2007;7:41–51.
- [69] Kortylewski M, Yu H. Role of Stat3 in suppressing anti-tumor immunity. *Curr Opin Immunol* 2008;20:228–33.
- [70] Chen Z, Laurence A, Kanno Y, Pachter-Zavisin M, Zhu BM, Tato C, et al. Selective regulatory function of Socs3 in the formation of IL-17-secreting T cells. *Proc Natl Acad Sci USA* 2006;103:8137–42.
- [71] Kryczek I, Wei S, Szeliga W, Vatan L, Zou W. Endogenous IL-17 contributes to reduced tumor growth and metastasis. *Blood* 2009;114:357–9.
- [72] Muranski P, Boni A, Antony PA, Cassard L, Irvine KR, Kaiser A, et al. Tumor-specific Th17-polarized cells eradicate large established melanoma. *Blood* 2008;112:362–73.
- [73] Spolski R, Leonard WJ. Interleukin-21: basic biology and implications for cancer and autoimmunity. *Annu Rev Immunol* 2008;26:57–79.
- [74] Koebel CM, Vermi W, Swann JB, Zerafa N, Rodig SJ, Old LJ, et al. Adaptive immunity maintains occult cancer in an equilibrium state. *Nature* 2007;450:903–7.
- [75] Girardi M, Oppenheim D, Glusac EJ, Filler R, Balmain A, Tigelaar RE, Hayday AC. Characterizing the protective component of the alpha-beta T cell response to transplantable squamous cell carcinoma. *J Invest Dermatol* 2004;122:699–706.
- [76] Daniel D, Chiu C, Giraud E, Inoue M, Mizzen LA, Chu NR, Hanahan D. CD4+ T cell-mediated antigen-specific immunotherapy in a mouse model of cervical cancer. *Cancer Res* 2005;65:2018–25.
- [77] DeNardo DG, Baretto JB, Andreu P, Vasquez L, Tawfik D, Kolhatkar N, Coussens LM. CD4+ T cells regulate pulmonary metastasis of mammary carcinomas by enhancing protumor properties of macrophages. *Cancer Cell* 2009;16:91–102.
- [78] Mantovani A, Allavena P, Sica A. Tumour-associated macrophages as a prototypic type II polarised phagocyte population: role in tumour progression. *Eur J Cancer* 2004;40:1660–7.
- [79] Johansson M, Denardo DG, Coussens LM. Polarized immune responses differentially regulate cancer development. *Immunol Rev* 2008;222:145–54.
- [80] Dunn GP, Bruce AT, Ikeda H, Old LJ, Schreiber RD. Cancer immunoediting: from immunosurveillance to tumor escape. *Nat Immunol* 2002;3:991–8.
- [81] Street SE, Cretney E, Smyth MJ. Perforin and interferon-gamma activities independently control tumor initiation, growth, and metastasis. *Blood* 2001;97:192–7.
- [82] Kaplan DH, Shankaran V, Dighe AS, Stockert E, Aguet M, Old LJ, Schreiber RD. Demonstration of an interferon gamma-dependent tumor surveillance system in immunocompetent mice. *Proc Natl Acad Sci USA* 1998;95:7556–61.
- [83] Street SE, Trapani JA, MacGregor D, Smyth MJ. Suppression of lymphoma and epithelial malignancies effected by interferon gamma. *J Exp Med* 2002;196:129–34.
- [84] Smyth MJ, Thia KY, Street SE, Cretney E, Trapani JA, Taniguchi M, et al. Differential tumor surveillance by natural killer (NK) and NKT cells. *J Exp Med* 2000;191:661–8.
- [85] Langowski JL, Zhang X, Wu L, Mattson JD, Chen T, Smith K, et al. IL-23 promotes tumour incidence and growth. *Nature* 2006;442:461–5.
- [86] Noguchi Y, Jungbluth A, Richards EC, Old LJ. Effect of interleukin 12 on tumor induction by 3-methylcholanthrene. *Proc Natl Acad Sci USA* 1996;93:11798–801.
- [87] Nastala CL, Edington HD, McKinney TC, Tahara H, Nalesnik MA, Brunda MJ, et al. Recombinant IL-12 administration induces tumor regression in association with IFN-gamma production. *J Immunol* 1994;153:1697–706.
- [88] Dunn GP, Koebel CM, Schreiber RD. Interferons, immunity and cancer immunoediting. *Nat Rev Immunol* 2006;6:836–48.
- [89] Gao Y, Yang W, Pan M, Scully E, Girardi M, Augenlicht LH, et al. Gamma delta T cells provide an early source of interferon gamma in tumor immunity. *J Exp Med* 2003;198:433–42.
- [90] Kacha AK, Fallarino F, Markiewicz MA, Gajewski TF. Cutting edge: spontaneous rejection of poorly immunogenic P1.HTR tumors by Stat6-deficient mice. *J Immunol* 2000;165:6024–8.
- [91] Ostrand-Rosenberg S, Grusby MJ, Clements VK. Cutting edge: STAT6-deficient mice have enhanced tumor immunity to primary and metastatic mammary carcinoma. *J Immunol* 2000;165:6015–9.
- [92] Fallarino F, Gajewski TF. Cutting edge: differentiation of antitumor CTL in vivo requires host expression of Stat1. *J Immunol* 1999;163:4109–13.
- [93] Bui JD, Carayannopoulos LN, Lanier LL, Yokoyama WM, Schreiber RD. IFN-dependent down-regulation of the NKG2D ligand H60 on tumors. *J Immunol* 2006;176:905–13.
- [94] Kemp RA, Ronchese F. Tumor-specific Tc1, but not Tc2, cells deliver protective antitumor immunity. *J Immunol* 2001;167:6497–502.
- [95] Dobrzanski MJ, Reome JB, Dutton RW. Therapeutic effects of tumor-reactive type 1 and type 2 CD8+ T cell subpopulations in established pulmonary metastases. *J Immunol* 1999;162:6671–80.
- [96] Dobrzanski MJ, Reome JB, Hollenbaugh JA, Dutton RW. Tc1 and Tc2 effector cell therapy elicit long-term tumor immunity by contrasting mechanisms that result in complementary endogenous type 1 antitumor responses. *J Immunol* 2004;172:1380–90.
- [97] Olver S, Apte S, Baz A, Kienzie N. The duplicious effects of interleukin 4 on tumour immunity: how can the same cytokine improve or impair control of tumour growth? *Tissue Antigens* 2007;69:293–8.
- [98] Mattes J, Hulett M, Xie W, Hogan S, Rothenberg ME, Foster P, Parish C. Immunotherapy of cytotoxic T cell-resistant tumors by T helper 2 cells: an eotaxin and STAT6-dependent process. *J Exp Med* 2003;197:387–93.
- [99] Eguchi J, Kuwashima N, Hatano M, Nishimura F, Dusak JE, Storkus WJ, Okada H. IL-4-transfected tumor cell vaccines activate tumor-infiltrating dendritic cells and promote type-1 immunity. *J Immunol* 2005;174:7194–201.
- [100] Kobayashi M, Kobayashi H, Pollard RB, Suzuki F. A pathogenic role of Th2 cells and their cytokine products on the pulmonary metastasis of murine B16 melanoma. *J Immunol* 1998;160:5869–73.
- [101] Wynn TA. IL-13 effector functions. *Annu Rev Immunol* 2003;21:425–56.

- [102] Lebel-Binay S, Laguerre B, Quintin-Colonna F, Conjeaud H, Magazin M, Miloux B, et al. Experimental gene therapy of cancer using tumor cells engineered to secrete interleukin-13. *Eur J Immunol* 1995;25:2340–8.
- [103] Terabe M, Matsui S, Noben-Trauth N, Chen H, Watson C, Donaldson DD, et al. NKT cell-mediated repression of tumor immunosurveillance by IL-13 and the IL-4R-STAT6 pathway. *Nat Immunol* 2000;1:515–20.
- [104] Terabe M, Matsui S, Park JM, Mamura M, Noben-Trauth N, Donaldson DD, et al. Transforming growth factor-beta production and myeloid cells are an effector mechanism through which CD1d-restricted T cells block cytotoxic T lymphocyte-mediated tumor immunosurveillance: abrogation prevents tumor recurrence. *J Exp Med* 2003;198:1741–52.
- [105] Pollard JW. Tumour-educated macrophages promote tumour progression and metastasis. *Nat Rev Cancer* 2004;4:71–8.
- [106] Terabe M, Park JM, Berzofsky JA. Role of IL-13 in regulation of anti-tumor immunity and tumor growth. *Cancer Immunol Immunother* 2004;53:79–85.
- [107] Volpert OV, Fong T, Koch AE, Peterson JD, Waltenbaugh C, Pepper RI, Bouck NP. Inhibition of angiogenesis by interleukin 4. *J Exp Med* 1998;188:1039–46.
- [108] Schuler T, Kornig S, Blankenstein T. Tumor rejection by modulation of tumor stromal fibroblasts. *J Exp Med* 2003;198:1487–93.
- [109] Aspod C, Pedroza-Gonzalez A, Gallegos M, Tindle S, Burton EC, Su D, et al. Breast cancer instructs dendritic cells to prime interleukin 13-secreting CD4+ T cells that facilitate tumor development. *J Exp Med* 2007;204:1037–47.
- [110] Tupin E, Kinjo Y, Kronenberg M. The unique role of natural killer T cells in the response to microorganisms. *Nat Rev Microbiol* 2007;5:405–17.
- [111] Kobayashi E, Motoki K, Uchida T, Fukushima H, Koezuka Y. KRN7000, a novel immunomodulator, and its antitumor activities. *Oncol Res* 1995;7: 529–34.
- [112] Kawano T, Cui J, Koezuka Y, Toura I, Kaneko Y, Motoki K, et al. CD1d-restricted and TCR-mediated activation of valpha14 NKT cells by glycosylceramides. *Science* 1997;278:1626–9.
- [113] Cui J, Shin T, Kawano T, Sato H, Kondo E, Toura I, et al. Requirement for Valpha14 NKT cells in IL-12-mediated rejection of tumors. *Science* 1997;278:1623–6.
- [114] Crowe NY, Smyth MJ, Godfrey DI. A critical role for natural killer T cells in immunosurveillance of methylcholanthrene-induced sarcomas. *J Exp Med* 2002;196:119–27.
- [115] Smyth MJ, Crowe NY, Pellicci DG, Kyparissoudis K, Kelly JM, Takeda K, et al. Sequential production of interferon-gamma by NK1.1(+) T cells and natural killer cells is essential for the antitumorigenic effect of alpha-galactosylceramide. *Blood* 2002;99:1259–66.
- [116] Kitamura H, Iwakabe K, Yahata T, Nishimura S, Ohta A, Ohmi Y, et al. The natural killer T (NKT) cell ligand alpha-galactosylceramide demonstrates its immunopotentiating effect by inducing interleukin (IL)-12 production by dendritic cells and IL-12 receptor expression on NKT cells. *J Exp Med* 1999;189:1121–8.
- [117] Fujii S, Shimizu K, Smith C, Bonifaz L, Steinman RM. Activation of natural killer T cells by alpha-galactosylceramide rapidly induces the full maturation of dendritic cells in vivo and thereby acts as an adjuvant for combined CD4 and CD8 T cell immunity to a coadministered protein. *J Exp Med* 2003;198:267–79.
- [118] Toura I, Kawano T, Akutsu Y, Nakayama T, Ochiai T, Taniguchi M. Cutting edge: inhibition of experimental tumor metastasis by dendritic cells pulsed with alpha-galactosylceramide. *J Immunol* 1999;163:2387–91.
- [119] Burdin N, Brossay L, Kronenberg M. Immunization with alpha-galactosylceramide polarizes CD1-reactive NK T cells towards Th2 cytokine synthesis. *Eur J Immunol* 1999;29:2014–25.
- [120] Fujii S, Shimizu K, Kronenberg M, Steinman RM. Prolonged IFN-gamma-producing NKT response induced with alpha-galactosylceramide-loaded DCs. *Nat Immunol* 2002;3:867–74.
- [121] Motohashi S, Nakayama T. Clinical applications of natural killer T cell-based immunotherapy for cancer. *Cancer Sci* 2008;99:638–45.
- [122] Berzofsky JA, Terabe M. NKT cells in tumor immunity: opposing subsets define a new immunoregulatory axis. *J Immunol* 2008;180:3627–35.
- [123] Terabe M, Swann J, Ambrosino E, Sinha P, Takaku S, Hayakawa Y, et al. A nonclassical non-Valpha14Jalpha18 CD1d-restricted (type II) NKT cell is sufficient for down-regulation of tumor immunosurveillance. *J Exp Med* 2005;202:1627–33.
- [124] Ambrosino E, Terabe M, Halder RC, Peng J, Takaku S, Miyake S, et al. Cross-regulation between type I and type II NKT cells in regulating tumor immunity: a new immunoregulatory axis. *J Immunol* 2007;179:5126–36.
- [125] Halder RC, Aguilera C, Maricic I, Kumar V. Type II NKT cell-mediated anergy induction in type I NKT cells prevents inflammatory liver disease. *J Clin Invest* 2007;117:2302–12.
- [126] Ostrand-Rosenberg S, Clements VK, Terabe M, Park JM, Berzofsky JA, Disanayake SK. Resistance to metastatic disease in STAT6-deficient mice requires hemopoietic and nonhemopoietic cells and is IFN-gamma dependent. *J Immunol* 2002;169:5796–804.
- [127] Terabe M, Khanna C, Bose S, Melchionda F, Mendoza A, Mackall CL, et al. CD1d-restricted natural killer T cells can down-regulate tumor immunosurveillance independent of interleukin-4 receptor-signal transducer and activator of transcription 6 or transforming growth factor-beta. *Cancer Res* 2006;66:3869–75.
- [128] Crowe NY, Coquet JM, Berzins SP, Kyparissoudis K, Keating R, Pellicci DG, et al. Differential antitumor immunity mediated by NKT cell subsets in vivo. *J Exp Med* 2005;202:1279–88.
- [129] Lee PT, Benlagha K, Teyton L, Bendelac A. Distinct functional lineages of human V(alpha)24 natural killer T cells. *J Exp Med* 2002;195:637–41.
- [130] Gumperz JE, Miyake S, Yamamura T, Brenner MB. Functionally distinct subsets of CD1d-restricted natural killer T cells revealed by CD1d tetramer staining. *J Exp Med* 2002;195:625–36.
- [131] Song L, Asgharzadeh S, Salo J, Engell K, Wu HW, Spoto R, et al. Valpha24-invariant NKT cells mediate antitumor activity via killing of tumor-associated macrophages. *J Clin Invest* 2009;119:1524–36.
- [132] Mosmann TR, Cherwinski H, Bond MW, Giedlin MA, Coffman RL. Two types of murine helper T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J Immunol* 1986;136:2348–57.

Brian Ruffell obtained his Ph.D. from the University of British Columbia in 2008 following his investigations into the role of the cell adhesion molecule CD44 in the immune response. He joined the laboratory of Dr. Lisa Coussens at the University of California, San Francisco shortly thereafter as a postdoctoral fellow to pursue his research interests in the role of inflammation in cancer progression and metastasis, with a particular interest in immune suppression within the tumor microenvironment.

David Denardo earned his Ph.D. from Baylor College of Medicine in 2005 where he evaluated mechanisms of non-classical estrogen-induced breast cancer growth. His postdoctoral studies at USC have focused on the role of adaptive immunity in regulating metastatic breast cancer. His recent studies have elucidated a novel role for T_H2 CD4+ T lymphocytes in potentiating pulmonary metastasis through cross-talk with tumor-associated macrophages, as well as utilizing this realization to develop an immune based signature that predicts breast cancer survival.

Nesrine I. Affara obtained her Ph.D. in Molecular, Cellular, and Developmental Biology from Ohio State University. She joined the laboratory of Dr. Lisa Coussens at the University of California, San Francisco in 2006 where she is currently a postdoctoral fellow. Her current research is focused on identifying molecular mechanisms by which recruitment of innate immune cells into premalignant tissues is regulated by humoral immunity in a transgenic mouse model of epithelial carcinogenesis.

Lisa M. Coussens. Dr. Coussens is a Professor in the Pathology Department and Helen Diller Family Comprehensive Cancer Center at the University of California, San Francisco. The Coussens lab focuses on the role of immune cells and leukocyte proteases as critical regulators of solid tumor development, including skin, lung (non-small cell and mesothelioma) and breast cancer development and is interested in understanding the molecular mechanisms that regulate leukocyte recruitment into neoplastic tissue and the subsequent regulation those leukocytes exert on evolving cancer cells. Dr. Coussens received her Ph.D. in Biological Chemistry from UCLA in 1993, and completed her postdoctoral fellowship in Cancer Biology at the University of California, San Francisco in Douglas Hanahan's laboratory.