

Analysis of Scale Separation Procedures

Peter Muller

University of Hawaii

Department of Oceanography

1000 Pope Road, MSB 429

Honolulu, HI 96822

phone: (808)956-8081 fax: (808)956-9164 email: pmuller@soest.hawaii.edu

Bo Qiu

Co-Principal Investigator

University of Hawaii

Department of Oceanography

1000 Pope Road, MSB 431

Honolulu, HI 96822

phone: (808)956-4098 fax: (808)956-9225 email: bo@soest.hawaii.edu

Grant Number: N00014-96-1-0412

LONG-TERM GOAL

Improving the parameterization of subgrid-scale motions in oceanic general circulation models.

SCIENTIFIC OBJECTIVES

Establish the fraction of subgrid-scale motions that is caused by propagating remotely forced waves, especially wind forced Rossby waves. Standard parameterization schemes (eddy diffusion, eddy induces transport velocities) assume that the subgrid scale fluxes are determined by local mean flow quantities. For propagating remotely forced waves this assumption becomes inadequate and non-local and perhaps stochastic parameterization schemes need to be developed.

APPROACH

Theoretical analysis and numerical calculation of the oceanic Rossby wave response to wind forcing and comparison with observations. The theoretical analysis is based on the linear quasi-geostrophic potential vorticity equation with idealized geometry and forcing. The numerical calculations use a 1-1/2 layer reduced gravity model. Observations are drawn primarily from BEMPEX (Barotropic Electro-magnetic and Pressure Experiment) and TOPEX/POSEIDON.

WORK COMPLETED

Two tasks were completed.

1. We analyzed theoretically the information content of coherence maps. These maps are an analytical tool first introduced by Brink (1989). They display the coherence between the oceanic current (or pressure) at a point (the mooring location) and the windstress curl field at other locations as a function of separation for a specific frequency band. We calculated such

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 1998		2. REPORT TYPE		3. DATES COVERED 00-00-1998 to 00-00-1998	
4. TITLE AND SUBTITLE Analysis of Scale Separation Procedures				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Hawaii, Department of Oceanography, 1000 Pope Rd, Honolulu, HI, 96822				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES See also ADM002252.					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 3	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

coherence maps for a simple linear quasi-geostrophic model forced by statistically stationary and homogeneous wind fields.

2. We calculated the baroclinic forced and free Rossby wave response in a 1-½ layer reduced gravity model of the North Pacific and rationalized the results in terms of simple analytic models.

RESULTS

Coherence maps can be used to estimate the Green's function of the linear quasi-geostrophic potential vorticity equation by multiple regression analysis. The presence of noise or non-linearities can be inferred from the multiple coherence, which is a number. The results can be generalized to other systems where response and forcing are related by a Green's function. Comparison of calculated coherence maps with observed maps confirms the notion, derived earlier, that part of the barotropic sub-inertial variability in the ocean is directly forced by the atmospheric windstress.

The calculations with the 1-½ layer reduced gravity model show that the interplay of wind-forced Rossby waves and free Rossby waves originating at the coastal and topographic boundaries leads to apparent phase speeds between one and two times the free Rossby wave speed. The details depend on dissipation and latitude. Overall, the findings are consistent with TOPEX/POSEIDON observations (Chelton and Schlax, 1996).

IMPACT/APPLICATION

Remotely forced propagating Rossby waves constitute part of the oceanic subgrid-scale variability. The parameterization of these radiative subgrid-scale motions must be expected to require non-local and perhaps stochastic parameterization schemes.

TRANSITIONS

None

RELATED PROJECTS

None

REFERENCES

Brink, K.H., 1989: Evidence for wind-driven current fluctuations in the western North Atlantic. *J. Geophys. Res.*, **94**, 1029-2044.

Chelton, D.B., and M.G. Schlax, 1996: Global observations of oceanic Rossby waves. *Science*, **272**, 234-238.

PUBLICATIONS

Müller, P. and A. Lippert, 1998: Reply to comments on ‘Direct atmospheric forcing of geostrophic eddies. Part II: Coherence maps’ by K.H. Brink and R.M. Samelson. *J. Phys. Oceanogr.*, **28**, 1010-1013.

Qiu, B., W. Miao and P. Müller, 1997: Propagation and decay of forced and free baroclinic Rossby waves in off-equatorial oceans. *J. Phys. Oceanogr.*, **27**, 2405-2417.

Müller, P., 1997: Coherence maps for wind forced quasi-geostrophic flows. *J. Phys. Oceanogr.*, **27**, 1927-1936.

Müller, P., 1997: Stochastic forcing of oceanic motions. In: *Stochastic Models in Geosystems*. IMA Volume 85, Springer-Verlag, 219-237.

PATENTS

None