

Long-Term Health Effects of Embedded Depleted Uranium

John F. Kalinich, Ph.D.

Program Advisor
Internal Contamination and Metal Toxicity Program
Armed Forces Radiobiology Research Institute
Uniformed Services University

kalinich@afrrri.usuhs.mil

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE NOV 2010		2. REPORT TYPE		3. DATES COVERED 00-00-2010 to 00-00-2010	
4. TITLE AND SUBTITLE Long-Term Health Effects of Embedded Depleted Uranium				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Uniformed Services University, Armed Forces Radiobiology Research Institute, 8901 Wisconsin Avenue, Bethesda, MD, 20889-5603				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES Presented at the Depleted Uranium Symposium, held November 4, 2010, at the Armed Forces Radiobiology Research Institute					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 22	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

Disclaimer

The views expressed in this presentation are those of the author and do not reflect the official policy or position of the Armed Forces Radiobiology Research Institute, the Uniformed Services University, the Department of Defense, or the United States Government.

The Embedded Pellet Rat Model




X-ray of pellets implanted in gastrocnemius muscle


Experimental Approach

- Two-year longevity study to determine whether intramuscularly implanted DU or tungsten alloy pellets are carcinogenic.
- Six treatment groups of Fisher 344 rats
 - **Two groups with 4 or 20 DU pellets**
 - Two groups with 4 or 20 tungsten alloy pellets (91% W, 6% Ni, 3% Co)
 - One nickel group (positive control)
 - One tantalum group (negative control)
- One set of pellet-implanted rats for duration of study. Second set includes rats euthanized at selected times after pellet implantation to provide tissues for histopathology, assessment for metal content and immunotoxicity testing.
- USAMRMC Award DAMD17-01-1-0821


Body Weight Gain After Pellet Implantation




Survival After Pellet Implantation




DU pellet implants: new and 12 weeks




Spleen Uranium Levels




Fibrotic Capsule from DU Implantation Site (13 weeks)



DU Implantation Site – 13 weeks




Popliteal Lymph Node Uranium Levels




Tumor Distribution Based on Gross Necropsy Examination

	6 Month (n = 20)	12 Month (n = 20)	18 Month (n = 10)	24 Month (n = 16)
Non-surgical	None	1-abdominal	8-testicle	7-testicle 3-abdominal
Tantalum	2-abdominal	None	8-testicle 2-abdominal	9-testicle 5-abdominal 1-muscle (leg)
DU Low Dose	None	2-abdominal	8-testicle 1-abdominal	10-testicle 1-abdominal 1-adrenal 1-kidney
DU High Dose	None	2-abdominal 1-lung	6-testicle 2-abdominal 1-kidney	9-testicle 2-lung 8-kidney 1-muscle (leg)


Renal Tubule Carcinoma – High-Dose DU (104 weeks)




Renal Tubule Carcinoma – High-Dose DU (104 weeks)




Urine Uranium Levels




Kidney Uranium Levels



Correlation of Kidney and Urinary Uranium Levels



Uranium Levels in Rats with and without Renal Tumors



Summary

- DU-implanted rats did not exhibit tumors at the pellet implantation sites.
- High-dose DU rats, in the 24 month group, had an increased incidence of renal neoplasias.
- Urine uranium levels in DU-implanted rats increased over time in a dose-dependent manner.
- Uranium levels in the kidney also increased over time, reaching 3 $\mu\text{g/g}$ tissue by 18 months in the high-dose DU group.

Current Work

- Continue histopathology assessment of renal carcinomas

Future Directions

- Identify early serum or urinary biomarkers of DU-induced neoplastic renal changes
- Investigate molecular mechanisms associated with DU-induced renal effects
- Tier-testing approach for assessing potential health effects of embedded metal fragments

The “Team”

Christy Emond

Vernieda Vergara

Histopathology

Steven Mog, DVM (U.S. Army, retired) – FDA

Eric Lombardini, MAJ, VMD

QUESTIONS

???

