

AD _____
(Leave blank)

Award Number:
W81-XWH-08-1-0738

TITLE:
Characterization of the pathological and biochemical markers that correlate to the clinical features of autism

PRINCIPAL INVESTIGATOR:
Jerzy Wegiel, Ph.D., D.V.M.

CONTRACTING ORGANIZATION:
Research Foundation for Mental Hygiene, Inc.,
Staten Island, New York 10314

REPORT DATE:
October 2010

TYPE OF REPORT:
Annual

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

X Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE			<i>Form Approved</i> OMB No. 0704-0188		
1. REPORT DATE 21-OCT-2010		2. REPORT TYPE Annual		3. DATES COVERED (From - To) 22 Sept 2009 –21 Sept 2010	
4. TITLE AND SUBTITLE Characterization of the pathological and biochemical markers that correlate to the clinical features of autism.			5a. CONTRACT NUMBER 5b. GRANT NUMBER W81XWH-08-1-0738		
			5c. PROGRAM ELEMENT NUMBER		
			5d. PROJECT NUMBER		
6. AUTHOR(S) Jerzy Wegiel, Ph.D., D.V.M., Email: j_wegiel@msn.com			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Research Foundation for Mental Hygiene, Inc. 1050 Forest Hill Road Staten Island, New York 10314 USA			8. PERFORMING		
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and And Medical Command Fort Detrick, MD. 21702-5012			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S NUMBER(S)		
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Multi-regional qualitative and quantitative study of autistic subjects brains revealed a broad spectrum of developmental alterations suggesting that there is no one brain structure, neuronal population, neurotransmitter or neuropeptide defect which might be considered as the single cause of the autistic phenotype. Delay of neuronal growth, appears as a common developmental alteration detected in many brain regions. However, the study of major regulatory systems of the brain: nucleus basalis of Meynert (cholinergic system), the substantia nigra (dopaminergic system), hypothalamic NPV and NSO (oxytocin and vasopressin) suggest that the neuronal populations playing a key role in production of neurotransmitters or neuropeptides are not affected (substantia nigra) or show rather limited distortion of developmental trajectories (NBMC, NSO, NPV). However, brain structures that are regulated by these centers reveal much broader and more significant developmental delays of neuronal growth. Global pattern of developmental alterations reflects the complexity of desynchronized development of neurons, neuronal networks, and brain structures most likely contributing to the complex clinical manifestations of autism. Significant interindividual differences in pattern of qualitative developmental abnormalities, topography, and severity of delayed neuronal growth appear to be the major factors determining broad spectrum of interindividual differences observed in clinical studies.					
15. SUBJECT TERMS Autism, Developmental Delay of Neuronal Growth, Desynchronization of Brain Development					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION UU	18. NUMBER 82	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER (include area code)

**Table of Contents
Subproject 2
Annual Report**

	<u>Page</u>
Introduction.....	1
Body.....	2
Key Research Accomplishments.....	16
Reportable Outcomes.....	17
Conclusion.....	18
References.....	19
Appendices.....	22

**Annual Report #2
October 22, 2010**

**Program Project Title: Characterization of the Pathological and Biochemical
Markers that Correlate to the Clinical Features of Autism**

Program Project PI: Jerzy Wegiel, Ph.D.; Co-PI: W. Ted Brown, M.D., Ph.D.

The overall aim of this multidisciplinary program project is to establish correlations between morphological and biochemical markers of autism and the clinical symptoms of the disorder.

SUBPROJECT 2

**Contribution of significant delay of neuronal development and metabolic shift of neurons
to clinical phenotype of autism**

Subproject 2 P.I.: Jerzy Wegiel, Ph.D.

INTRODUCTION

This Program Project is focused on detection of:

- (a) mechanisms leading to morphological changes and clinical autism phenotype,
- (b) morphological and biochemical markers of autism,
- (c) correlations between pathology and clinical manifestations of autism, and
- (d) those pathological domains that might be a target for treatment.

In the first year of this Program Project we concentrated efforts on establishing of inclusion and exclusion criteria, to reduce the risk of distortion of results of quantitative morphometric studies.

- a. Inclusion criteria were based on Autism Diagnostic Interview- Revised (ADI-R).
- b. Exclusion criteria were based on expanded neuropathological protocol.

Clinical diagnosis was provided by the Autism Tissue Program. Neuropathological criteria were established in cooperation with Project 1. Cases not meeting ADI-R criteria and cases with signs of comorbidity, perimortem and postmortem changes affecting brain structure were eliminated from morphometric studies. The products of this neuropathological qualitative studies were characteristics of the type, topography, severity and prevalence of developmental brain abnormalities in autistic and control cohorts selected for morphometric studies.

Morphometric studies (Project 2) were concentrated on:

- a. Characterization of developmental delay of neuronal growth in brain structures linked to functional deficits and modifications observed in autism and
- b. Detection of markers of desynchronization of development of brain structures, neuronal populations and neuronal networks.

In the second year, consistent with the Statement of Work the study focused on:

- a. brain structures and neuronal populations interacting with structures examined in the first year, and
- b. structures regulating development and function of other brain structures or neuronal populations.

BODY

The study in the second year was designed to determine what types of interactions, neurotransmitters or neuropeptides contribute to morphological, biochemical and functional changes leading to clinical phenotype of autism.

- a. Claustrum was selected as a brain structure integrating function of several brain modalities with major contribution to cognition.
- b. Substantia nigra was selected as a source of dopamine controlling motor, reward and other systems.
- c. Nucleus basalis of Maynert was selected as the source of brain acetylcholine.
- d. Hypothalamus was selected as the major source of oxytocin and vasopressin regulating social memory and interactions, communication, ritualistic behaviors, aggression and anxiety, sleep and other features of autism.

Material and methods.

Nine brain hemispheres of autistic subjects from 4 to 36 years of age and nine control subjects 4 to 32 years of age were examined. In the majority of the cases, the right hemisphere was examined (16 cases, 89%). Four age-matched pairs of autistic and control subjects represented the youngest group (4/4, 5/4, 7/7, and 8/8 years of age). Five age-matched pairs represented late childhood and adulthood (13/14, 17/20, 21/23, 23/28, and 36/32).

Brain hemispheres were fixed in 10% formalin, dehydrated in ascending concentrations of ethyl alcohol for 19 days (dehydration results in the reduction of brain hemisphere weight on average by 47%), and embedded in celloidin. Free-floating celloidin 200- μ m-thick sections were stained with cresyl violet.

Clinical inclusion criteria. Selection of the autistic cases was based on clinical evaluation including Autism Diagnostic Interview-Revised (ADI-R) scores. Subjects with atypical autism were excluded from this morphometric study.

Neuropathological exclusion criteria. An expanded neuropathological evaluation was based on the examination of 120 coronal hemispheric CV-stained serial sections. Brains affected with premortem pathology (ischemia, anoxia-related neuronal degeneration/loss, gliosis, and neovascularization) and severe postmortem degradation (autolysis) affecting brain region volume; cytoarchitecture; and the number, shape, and size of neurons were excluded from this study.

Morphometric equipment. Image analyzer with computer-controlled automatic stage installed on Axiophot II microscope (Zeiss) and Stereo-Investigator and Nucleator software (Microbrightfield, VA, USA).

Morphometric methods. The Cavalieri method was used to estimate the volume of the examined brain structures, the disector method was used to determine the number of neurons,

and the nucleator to determine the volume of neurons and neuronal nuclei. The coefficient of error was less than 0.01 (Schaffer).

Statistical analysis. The volume of brain structures, number of neurons, neuronal density, mean neuron size and mean nuclear size were compared between autistic and control subjects in repeated-measures ANOVA.

PART I: DELAYED DEVELOPMENT OF CLAUSTRUM IN AUTISM

Evidence for failure of integration of interactions between brain structural and functional modalities

In 2005, Crick and Koch proposed the claustrum as the ideal candidate for the brain "consciousness" center, because it acts like a conductor in an orchestra to integrate information from all of the various modalities.

Clastrum anatomy. The claustrum is a thin ribbon of gray matter located between the insular cortex and the striatum, closely associated with the amygdala and built of 16 million cells of both cortical and subcortical origin. 94% of claustrum neurons are large projection spiny neurons with numerous coarse pigment granules (Braak and Braak 1982).

Clastrum connectivity. The claustrum has an extensive reciprocal connections to and from almost all brain regions (Fernandez-Miranda et al 2008). The information from the various brain regions is segmented topographically in the claustrum and in an overlapping manner.

- The anterior claustrum projects to and receives signals from the frontal cortex.
- The middle claustrum is associated with the parietal cortex.
- The posterior/inferior claustrum is associated with the temporal and occipital cortices.- The dorsal claustrum forms a visual loop with BA17 (Fernandez-Miranda et al 2008, Morys et al 1993, Pearson et al 1982).
- The claustrum also projects to the amygdala (Amaral and Insausti 1992), the hippocampus (Amaral and Cowan 1980), and the striatum (Arikuni and Kubota 1985, Crick and Koch 2005).
- Social brain circuitry (orbito-frontal cortex, superior temporal gyrus and amygdala) (Brothers et al 1990) projects to the claustrum and receives claustral connections (Fernandez-Miranda et al 2008).

Clastrum function in normal and pathological conditions. Higher order cognitive functions such as:

- Fear recognition (Stein et al. 2007)
- Experiential dread (Berns et al 2006)
- Cognitive impairment (Dubroff et al 2008)
- Memory storage (Morys et al 1996)
- Associative learning (Chachi and Powell 2004)
- Repetitive behaviors and addiction (Morys et al 1996; Naqvi et al 2007)
- Multimodal processing of olfactory, auditory, visual, and tactile information.
- Emotional and behavioral responses (Bennet and Baired 2006)
- Suppression of natural urges (Lerner et al 2008)
- Seizures (Zhang et al 2001)

- Contribution to psychoses (Sperner et al 1996)

Hypothesis

The severely compromised ability to adapt that is observed in individuals with autism, suggests that numerous high order functions are not processed properly and that developmental defects in the claustrum may contribute to the cognitive impairment, repetitive behaviors, deficits of social interactions, altered processing of sensory signals and other components of the autistic phenotype.

Aim

The aim of this morphometric study of the claustrum of autistic and control subjects was to determine whether and how the development of claustrum, known to be responsible for integration of the functions of the social and somatosensory circuits, is affected in autism.

Results

Clastrum volume The claustrum volume was 22% less in autistic (388 mm^3) than in control (494 mm^3) subjects from 4 to 8 years of age. In autistic subjects 13 to 36 years of age, the volume of claustrum increased by 18% (to 472 mm^3) and was insignificantly higher than in age-matched controls (452 mm^3). This observation is consistent with MRI studies of 16 high-functioning (IQ 85) autistic and 14 control males from 7 to 12 years of age which revealed a lower volume of the claustrum of autistic (555 mm^3) than control subjects (701 mm^3) (Davies 2008). The lower volumes detected in postmortem study than in clinical MRI study are an effect of tissue shrinkage during dehydration.

Number of neurons. The mean numerical density of neurons was almost identical in the autistic ($29,156/\text{mm}^3$; $n = 9$) and control subjects ($29,260/\text{mm}^3$; $n = 9$). The mean total number of neurons in the nine autistic subjects (12.5 millions) was insignificantly less (by 8%) than in the control subjects (13.6 millions).

Mean neuron volume. The mean volume of the neuronal body in 4 to 8-year old autistic children was less by 32% ($1,465 \mu\text{m}^3$, $n = 4$) than in control group ($2,154 \mu\text{m}^3$; $n = 4$; $p < 0.05$). However, the mean volume of neuronal body in 13 to 36-year-old autistic subjects was less only by 19% ($1,640 \mu\text{m}^3$; $n = 5$) than in control group ($2,010 \mu\text{m}^3$; $n = 5$; $p < 0.02$).

Mean volume of neuron nucleus. The mean volume of neuron nucleus in 4 to 8-year old autistic children ($252 \mu\text{m}^3$; $n = 4$), was 41% less than in the control group ($427 \mu\text{m}^3$; $n = 4$; $p < 0.009$). The mean volume of the neuronal nucleus in 13 to 36-year-old autistic subjects was insignificantly less ($269 \mu\text{m}^3$; $n = 5$) than in the control group ($358 \mu\text{m}^3$; $n = 5$).

Desynchronization of development of claustral circuitry. The study of 14 brain subdivisions in 13 autistic and 14 control subjects revealed:


- (a) brain structure specific delay in rate of neurons growth and
- (b) desynchronization of neurons growth in the brain of autistic subjects.

Desynchronization is reflected in striking differences of neuron volume deficit in early childhood in eight examined structures and nonsignificant volume deficit in four other structures including the:

- nucleus accumbens (-37%; $p < 0.004$),
- Purkinje cells (-32%; $p < 0.018$),
- neurons in thalamus (-31%; $p < 0.001$),
- globus pallidus (-31%; $p < 0.008$),
- entorhinal cortex (-28%; $p < 0.011$),
- amygdala (-27%; $p < 0.006$),
- caudate nucleus (-22%; $p < 0.025$),
- putamen (-22%; $p < 0.042$),
- dentate nucleus (not significant reduction of cell body size),
- cornu Ammonis (not significant reduction of cell body size),
- lateral geniculate nucleus (not significant reduction of cell body size), and
- inferior olive (not significant reduction of cell body size).

The function of the examined structures suggests that delayed and desynchronized development of affected networks contributes to all three diagnostic domains of autism, including disrupted social and communication development, restricted repetitive and stereotyped patterns of behavior, and intellectual deficits.

The striking differences in the developmental volume deficits of neurons in the claustrum and in brain structures projecting to the claustrum that were detected in 4- to 8-year-old autistic children are signs of desynchronization of claustral circuitry development. The observed pattern of developmental delays suggests that a number of physiologic processes that collectively ensure the maintenance of undisturbed contact with the environment are altered in the early childhood of autistic subjects.



Underconnectivity. Smaller volume of claustrum and reduced size of neurons could be indicative of underconnectivity in brains of autistic subjects. Underconnectivity of claustrum may result in defects/deficits of integration of cortical and subcortical modalities and may contribute to complex autistic phenotype.

Conclusions

1. Stereological study revealed significant delay of neuronal growth in claustrum that is integrating a number of physiological processes that collectively maintain contact between people and environment.
2. Examination of older children and adults with autism shows that in claustrum of teenagers, acceleration of neuronal growth results in partial correction of neuronal size.
3. Detected desynchronization of neuronal growth in claustrum and interacting brain modalities including social brain, sensorimotor system and memory system appears to be significant contributor to the clinical symptoms of autism.

PART II. DOPAMINERGIC SYSTEM IN AUTISM

Evidence for intact substantia nigra but desynchronized development of structures within dopaminergic system

Anatomical substrate of repetitive and stereotyped behaviors. Repetitive and stereotyped behaviors, defined as recurring, nonfunctional activities or interests that occur regularly and interfere with daily functioning, are diagnostic signs of autism (Gabriels et al., 2005). Several studies have implicated the role of the basal ganglia and fronto-striatal circuitry in the pathophysiology of autism, especially in repetitive and stereotyped behaviors. The substantia nigra, together with the caudate nucleus, putamen, n. accumbens and globus pallidus, belong to the basal ganglia. These structures form the cortico-subcortical circuits that play a key role (a) in programming and execution of movements, and (b) in the reward system.

Partial anatomical and functional overlap of motor and reward systems. The reward system includes (a) the prefrontal cortex, (b) the limbic system components with the amygdala, hippocampus, and anterior cingulate, (c) the substantia nigra and limbic striatum with the nucleus accumbens, ventral caudate nucleus, and putamen (Lopes de Silva et al., 1990; Kotter and Stephan, 1997).

Substantia nigra and brain reward pathway. Dopamine has been associated with pleasure, and it has been called the "anti-stress molecule" and "pleasure molecule" (Bloom et al., 2000). Released dopamine stimulates a number of receptors (D1-D5), which results in increased feelings of well-being and stress reduction. A reduced number of dopamine receptor sites in n. accumbens and the amygdala may result in repetitive behavior in an attempt to stimulate dopamine release.

Reward deficiency syndrome is associated with abnormalities of the basal ganglia, including the substantia nigra, caudate, putamen, and nucleus accumbens, as well as the prefrontal cortex, hypothalamus, and central nucleus of the amygdala. Our studies indicate that almost all of these structures are affected by developmental neuronal delay in people with autism. The substantia nigra appears to play a key role in both the

motor and reward systems, but the morphology of the substantia nigra dopaminergic neurons is not known.

Nucleus accumbens dopaminergic innervation and role in reward system. The nucleus accumbens is at the center of network processing of reward-related information. The shell of the nucleus accumbens (Heimer et al., 1997) is a part of the extended amygdala involved in integrating emotions, whereas, the core belongs to the striato-pallidal system integrating motor responses. The nucleus accumbens projections to the motor areas, such as the ventral pallidum, turn reward information into motivated action (Mogenson et al., 1980). Dopaminergic projections from the substantia nigra to the nucleus accumbens have a critical role in stimulus-reward learning. Repetitive behaviors and stereotypies may reflect altered motor responses modulated by the altered reward system.

Substantia nigra structure. The substantia nigra pars compacta forms a shallow layer of approximately 1.4 million pigment-positive, dopamine-producing neurons organized into ventral and dorsal tiers, and the lateral part. Neurons from both the ventral and dorsal tiers project to the striatum and cerebral cortex. The dorsal tier also projects to the amygdala. Numerous reciprocal connections feedback within basal ganglia networks.

Hypotheses

1. Development of the dopaminergic system contributing to repetitive and stereotyped behaviors is altered in autism.
2. The clinical outcome of altered brain development is a complex product of changes in different dopaminergic system components.
3. The absence of clinical signs of dopamine deficit or excess suggests that the substantia nigra is not altered in autistic subjects, and that modifications of the development of striatal components are the main contributor to repetitive and stereotyped behaviors.

Aim

The aim of this morphometric study was to determine whether the developmental trajectories of neurons within the dorsal, ventral, and lateral parts of the substantia nigra, the source of dopaminergic innervation of the human striatum, are similar to the patterns of development of four striatal subdivisions, including the caudate, putamen, nucleus accumbens, and globus pallidus.

Results

No difference was noted between the volume of pigmented neurons and their nuclei in the substantia nigra of control and autistic subjects.

Neuron volume in control cohort. The mean volume of the neurons in the ventral and dorsal tiers and the lateral part of the substantia nigra in 4 to 64-year-old control subjects was similar (9,904 μm^3 , 9,624 μm^3 , and 10,041 μm^3 , respectively). The lack of significant difference


suggests the homogeneity of pigmented neurons within different portions of the pars compacta, with a mean volume of $9,856 \mu\text{m}^3$ in the control group. There was no statistically significant difference in neuron volume between 4 to 8-year-old and 14 to 64-year-old control subjects

Neuron volume in autistic cohort. The study of autistic subjects from 4 to 56-years of age did not reveal any difference in the volume of the pigmented neurons in the ventral and dorsal tiers, and in the lateral part of the substantia nigra ($9,949 \mu\text{m}^3$, $9,903 \mu\text{m}^3$, and $10,281 \mu\text{m}^3$, respectively). The mean volume of neurons in the substantia nigra of autistic subjects ($10,044 \mu\text{m}^3$) was comparable with those observed in the control cohort ($9,856 \mu\text{m}^3$). In both the autistic and control cohorts, there was no difference in the volume of pigmented neurons between subjects younger than 8 years of age and in the teenagers/adults group.

Nucleus volume in neurons of autistic and control subjects. The nucleus volume in neurons in the ventral and dorsal tiers and in the lateral part of the substantia nigra is comparable across anatomical subdivisions, age subgroups, and autistic and control cohorts.

Desynchronized development of nigro-striatal circuitry in autistic subjects: a potential contribution to repetitive and stereotyped behaviors. In the brains of autistic children 4 to 8 years of age, we found a significant delay of neuron growth, with the volume of cell bodies smaller than in control subjects by: 22% in the caudate nucleus ($p < 0.025^*$), 21% in the putamen ($p < 0.042^*$), 31% in the globus pallidus ($p < 0.008^{**}$) and 37% in the nucleus accumbens ($p < 0.004^{**}$). The volume of the neuronal nucleus was also significantly reduced. Much less, or lack of, significant difference between the volume of neuronal soma and nucleus in autistic and control subjects older than 8 years of age suggests that a correction of neuronal growth occurs in late childhood and adulthood.

In the brains of autistic children 4 to 8 years of age, we found a significant delay of neuron growth, with the volume of cell bodies smaller than in control subjects by: 22% in the caudate nucleus ($p < 0.025^*$), 21% in the putamen ($p < 0.042^*$), 31% in the globus pallidus ($p < 0.008^{**}$) and 37% in the nucleus accumbens ($p < 0.004^{**}$). The volume of the neuronal nucleus was also significantly reduced. Much less, or lack of, significant difference between the volume of neuronal soma and nucleus in autistic and control subjects older than 8 years of age suggests that a correction of neuronal growth occurs in late childhood and adulthood.



The unaffected size of the neurons of the substantia nigra providing dopamine to the motor and reward system, but the significant structure-specific delay of neuronal growth in striatal subdivisions indicates that development of the nigrostriatal system is desynchronized and that differences are most striking in 4- to 8-year-old autistic subjects. The observed pattern of developmental desynchronization of the nigrostriatal system might be a key contributor to the repetitive and stereotyped behaviors of autistic subjects.

Conclusions

1. The observed pattern indicates that in autistic subjects the growth of dopaminergic neurons within the ventral, dorsal and lateral part of the substantia nigra is not altered.
2. This pattern may suggest that dopaminergic projections to other components of the nigrostriatal system are not deficient.
3. However, normal development of neurons producing dopamine and significantly delayed development of neurons using dopamine may result in incoherence of dopaminergic system with clinical consequences.
4. No difference in the volume of pigmented neurons in the substantia nigra of autistic and control subjects, but the significant delay of the growth of neurons in the striatal subdivisions of autistic children, indicates that development of different components of the motor and award systems is desynchronized, most likely by neuron type-specific regulatory mechanisms.
5. This study suggests that the repetitive and stereotyped behaviors seen in autism are driven by a delayed development of neurons within the striatum but not in the substantia nigra.

PART III. OXYTOCINERGIC AND VASOPRESSINERGIC SYSTEM IN AUTISM.

Moderate developmental change in the hypothalamus of autistic subjects

Biochemical studies of hypothalamic neuropeptides in people with autism, and human and animal studies of the role of hypothalamic neuroendocrine system in brain development and behavior, suggest that dysfunction of neuroendocrine system could be a significant contributor to all three types of behavioral impairments and abnormalities observed in autism.

Clinical symptoms of autism. Deficit of social attachment is the first symptom noticed in infants diagnosed later with autism. Impairments in reciprocal social interactions are persistent feature of autistic children and adults. Social interaction involves multi-sensory processing including a broad spectrum of verbal and nonverbal communication. Communication impairment is the second life-long lasting feature of autistic individuals. Restricted repetitive and stereotyped patterns of behavior, interests, and activities are the third diagnostic feature of autism.

Behavior regulation by hypothalamic neuropeptides – oxytocin and vasopressin.

Hypothalamic nuclei, nucleus paraventricularis and supraopticus are the main source of brain oxytocin and vasopressin.

Social memory deficits. Oxytocin is indispensable for the development of social memory and facilitates learning of social interactions. Oxytocin is involved in the formation of associations specifically related to the mother. In oxytocin KO mice, loss of social memory could be rescued by oxytocin treatment (Fergusson et al 2000). Vasopressin is involved in the formation of social memory, while oxytocin is necessary for the retention of these newly formed memories (Popik et al 1992a, Popik and Van Ree 1992).

Anxiety, aggression. Vasopressin enhances aggressiveness, anxiety, and consolidates fear memory (Griebel et al 2002, Bielsky et al 2004, Landgraf and Neuman 2004, Kovacs and deWied 1994). Oxytocin decreases anxiety and stress, and facilitates social encounters, maternal care and the extinction of conditioned avoidance behavior (Kovacs and deWied 1994, Windle et al 1997, Champagne et al 2001, Bale et al 2001).

Repetitive behaviors. Oxytocin and vasopressin are known to play a role in repetitive behaviors. Subjects with autism spectrum disorders show a significant reduction in repetitive behaviors following oxytocin infusion (Hollander et al 2003).

Sleep disorder. In about 60% of subjects with autism abnormal sleep patterns, such as never sleeping throughout the night, trouble going to sleep, or getting up for the day at 2:00 am are observed. Vasopressin is a day/night cycle regulator (Swaab 1997).

Oxytocin synthesis and processing. Oxytocin is synthesized as a prohormone neurophysin I (Rao et al 1992). Neurophysin I is sequentially processed to peptides including: carboxy-terminally extended peptides: OT-Gly, OT-Gly-Lys and OT-Gly-Lys-Arg. Oxytocin extended forms are further cleaved enzymatically to yield the nine-amino-acid active peptide, oxytocin (Gainer et al 1995).

Defect of oxytocin prohormone processing in autistic children. The study of male autistic subjects (6-12 year of age) and age-matched control subjects revealed: decrease in plasma oxytocin, deficit in oxytocine prohormone processing with an increase in OT-X (inactive oxytocin), increase in the ratio of C-terminally extended (inactive) forms of oxytocin. These

results indicate that while in the control children nearly all available OT-X is metabolized to active oxytocine, in autistic children the immature (extended) oxytocin forms serve as the primary circulating molecule in the absence or in compensation for biologically active oxytocin (Green et al 2001).

Genetic defects of oxytocin convertases in autistic subjects. Deficient conversion of OT-X to OT in autism could be the result of alterations in the level of pro-hormone convertases (PC2 and/or PC5) associated with genetic defects. Faulty OT-X/OT processing could result from defects in the area of 20p11-13 affecting PC2 and the oxytocin gene detected in autistic subjects (Cook 1998, Szatmari et al 1998).

Justification to explore morphology of developing hypothalamus in autism. Strong evidence of hypothalamic involvement in autism such as:

- Detection of abnormal levels of hypothalamic neuropeptides in sera,
- Abnormal processing of oxytocine precursor,
- Improvement of behavior after treatment with hypothalamic neuropeptides,
- Links between genes encoding hypothalamic prohormones, enzymes involved in prohormone processing, and neuropeptides receptors
- Experimental studies demonstrating that hypothalamic abnormalities contribute to impaired social interactions, deficits in communication, and stereotypic behaviors.

Hypothesis: Biochemical and clinical data suggest developmental defects of neurons producing oxytocin and vasopressin. We assumed that:

- (a) Hypothalamic neurons belong to brain neuronal populations with a significant delay of growth and maturation, and that
- (b) This delay results in impairment of conversion of inactive forms of oxytocin into biologically active oxytocin.

Morphological study of hypothalamus may help to identify mechanisms leading to clinical expression of autism and help to prevent and treat hypothalamus related pathology.

Aim: The aim of this morphometric study is to determine whether development of neurons in the nucleus paraventricularis (NPV) and supraopticus (NSO), known as the main brain source of oxytocin and vasopressin, is modified in comparison to control subjects.

Preliminary results.

Results are based on examination of the hypothalamus of 5 autistic subjects from 4 to 36 years of age and 8 control cases. The morphometrist was blind to examined subject diagnosis. The study is in progress. Results are missing for 7 autistic and 4 control subjects).

The volume of neuron and neuronal nucleus. The mean volume of neurons in the nucleus paraventricularis was 20% less in autistic subjects ($2,193 \pm 82 \mu\text{m}^3$) than in control group ($2,714 \pm 297 \mu\text{m}^3$). The volume of neuronal nuclei was insignificantly less (by 13%) in autistic subjects ($319 \pm 26 \mu\text{m}^3$) than in controls ($367 \pm 28 \mu\text{m}^3$).

The mean volume of neurons in the nucleus supraopticus was 22% less in autism ($3,700 \pm 467 \mu\text{m}^3$) than in control ($4,721 \pm 349 \mu\text{m}^3$). The mean volume of neuronal nuclei was also less (by 26%) in autistic ($476 \pm 81 \mu\text{m}^3$) than in control subjects ($643 \pm 62 \mu\text{m}^3$).




Fig. The mean volume of neurons in the nucleus paraventricularis was 20% less in autistic subjects ($2,193 \pm 82 \mu\text{m}^3$) than in control group ($2,714 \pm 297 \mu\text{m}^3$).




Fig. The mean volume of neurons in the nucleus supraopticus was 22% less in autism ($3,700 \pm 467 \mu\text{m}^3$) than in control ($4,721 \pm 349 \mu\text{m}^3$).

Number of neurons. The numerical density of neurons (number of neurons/mm³) was insignificantly less (-12%) in the NSO of autistic ($28,248 \pm 935 \text{mm}^3$) than in control subjects ($32,157 \pm 1877 \text{mm}^3$). Similar trend was observed in the NPV with insignificantly lower numerical density (by 12%) in autism ($58,252 \pm 5,763 \text{mm}^3$) than in control ($65,785 \pm 3,455 \text{mm}^3$).

The total number of neurons in the NPV of autistic subjects ($322,233 \pm 30,211$) was insignificantly (-14%) less than in controls ($375,794 \pm 39,080$). The 6% difference between total number of neurons in the NSO in autistic ($110,242 \pm 9,760$) and control subjects ($116,863 \pm 11,585$) was also not significant.

Conclusions.

1. Results of morphometric study of hypothalamic nuclei suggests that in both examined regions (NPV and NSO) neurons are smaller in autistic subjects than in control subjects.
2. The trend to reduced numerical density and total number of neurons in VPV and NSO may represent other signs of developmental defects within these two hypothalamic nuclei.
3. Reduced size of the cell body and cell nucleus may reflect moderate delay of neuronal growth and maturation of hypothalamic neurons in autistic subjects from 4 to 36 year of age.
4. Detected developmental alteration in these two sources of brain oxytocin and vasopressin may contribute to (a) developmental delay of processing of neurophysin into mature (active) form of oxytocin revealed in clinical studies, (b) reduced level of oxytocin and the imbalance of the ratio between oxytocin and vasopressin leading to behavioral changes including social and communication deficits, stereotypic behavior, anxiety and sleep disorder.

Interindividual differences in the level of developmental delays in the NSO and NPV within autistic group may reflect interindividual differences in clinical phenotype including ADI-R scores.

PART IV: CHOLINERGIC SYSTEM IN AUTISM

A week trend to delayed neuronal development

Anatomical subdivisions and connectivity of the nucleus basalis of Meynert complex (NBMC). Four subdivisions could be distinguished on behalf of their topography and projection patterns including:

Ch1 group: Medial septal nucleus. 10% of its neurons are cholinergic. Neurons are primarily small, hyperchromatic. Projections to the: cornu Ammonis, dentate gyrus, subicular complex, entorhinal cortex.

Ch2 group: Vertical limb nucleus of diagonal band of Broca. About 70% of neurons are cholinergic. Neurons are also hyperchromatic but larger with clearly visible ellipsoid nucleus. Projections to the hippocampus, cingulate cortex, thalamus, parahippocampal gyrus.

Ch3 group: Horizontal limb nucleus of diagonal band of Broca. Only 1 % of these neurons are cholinergic. Projections to the: hippocampus, amygdala, caudate nucleus thalamus, brainstem.

Ch4 group: Nucleus basalis of Meynert (more than 90% is cholinergic). Hyperchromatic, large and very large neurons with large and excentric nucleus. They project to the: temporal, frontal and parietal lobe, parahippocampal gyrus and the entorhinal cortex, amygdala, thalamus, and caudate nucleus.

NBMC function. NBMC function as cholinergic relay for transmitting predominantly limbic and paralimbic information to the neocortex and is involved in integration of emotional and motor responses, learning and memory, emotional and motivational states

Results

In all examined subregions of the NBMC, the mean volume of neuronal soma, as well as the neuronal nucleus, was less in autistic subjects than in controls (Fig). In magnocellular group (CH4) with the highest percentage of cholinergic neurons the volume of cells was significantly less in autistic subjects ($9,706; SE \pm 296; p < 0.01$) than in controls ($10,681; SE \pm 426$). The mean volume of neuronal nuclei was also significantly less in CH4 of autistic subjects ($677 \pm 31; p < 0.01$) than in controls (791 ± 38). The volume of noncholinergic neurons in CH3 (only 1% of neurons in CH4 is cholinergic) is 9.4% less in autistic subjects ($7,404 \pm 310$) than in control group ($8,175 \pm 321$) but the difference is not significant (Table).




Fig. Mean volume of neuronal soma and neuronal nucleus in autistic subjects, NBMC is 9% less in autistic subjects than in control subjects. However, statistically significant differences are observed only in CH4 groups (See Table)

Table The volume of neuronal body and nucleus in four groups of neurons of the NBMC (CH1 – CH4).

Group of neurons within NBMC	Neuron body volume		Neuron nucleus volume	
	Control	Autism	Control	Autism
CH1	6,421 ± 460	5,926 ± 294 -7.7%	423 ± 35	403 ± 29 -4.6%
CH2	6,885 ± 327	6,278 ± 241 -8.8%	479 ± 29	454 ± 32 -3.8%
CH3	8,175 ± 321	7,404 ± 310 -9.4%	567 ± 35	508 ± 27 -10.4%
CH4	10,681 ± 426	9,706 ± 296 (p<0.04) -9.1%	791 ± 38	677 ± 31 (p<0.01) -14.4%

Conclusions

1. The NBMC, which is the main controller of brain cholinergic neurotransmission, shows only a weak trend of delayed growth of neurons.
2. The neuron volume deficit is statistically significant only in the CH4 group.
3. Reduced volume of neurons both, in CH4 with 90% of cholinergic neurons and in CH3 with only 1% of cholinergic neurons, suggests that factor regulating growth of neurons in the NBMC does not recognize the type of neurotransmission of neuronal populations in CH3 and CH4.
4. In general, one may assume that developmental delays of the cholinergic system may not have contribution to the clinical phenotype of autism or this contribution is rather limited.

KEY RESEARCH ACCOMPLISHMENTS

1. This project established criteria for clinical inclusion and neuropathological exclusion criteria. It reduced the risk of distortion of morphometric studies results by the exclusion of cases with comorbidity modifying autism-related pathology, and premortem, perimortem and postmortem changes, that may lead to false results and conclusions.
2. Mapping of qualitative and quantitative changes results in a map of the brain regions affected by developmental alterations associated with autism and contributing to the clinical phenotype.
3. Comparative study of neurons of the dopaminergic and cholinergic system, neurons producing oxytocin and vasopressin, with neurons in basal ganglia and cortex reflects relatively less or non-affected regulatory systems. In contrast, the study of the neuronal populations in basal ganglia or cortex demonstrate a broad spectrum of developmental delays.
4. The major research accomplishment is:
 - (a) Detection of pattern of focal developmental alterations affecting individual brain regions, structures, and neuronal populations,
 - (b) Establishing that global pattern of desynchronization of neuronal networks is the major contributor to the clinical phenotype of autism, and
 - (c) Revealing that regulatory brain centers are less affected or not affected by developmental delay.
5. Limited contribution of major regulatory centers to autism clinical phenotype is consistent with current concepts of “fractionation” of clinical features and “localizing models” linking regional brain defects with functional deficits/alterations.

REPORTABLE OUTCOMES

1. The neurobiological and neuropathological background of this Program Project is summarized in our book chapter (see Appendices):

Wegiel J, Wisniewski T, Chauhan A, Chauhan V, Kuchna I, Nowicki K, Imaki H, Wegiel J, Ma SY, Wierzba Bobrowicz T, Cohen IL, London E, Brown WT. Type, topography and sequelae of neuropathological changes shaping clinical phenotype of autism. In: Autism: Oxidative Stress, Inflammation, and Immune Abnormalities. Ed.: Abha Chauhan, Ved Chauhan and W. Ted Brown. Taylor & Francis/CRC Press, Boca Raton, FL. 2010, pp 1-34

2. Results of neuropathological evaluation of brains of 13 autistic and 14 control subjects are summarized in a paper published in *Acta Neuropathologica* (see Appendices):

Wegiel J, Kuchna I, Nowicki K, Imaki H, Wegiel J, Marchi E, Ma SY, Chauhan A, Chauhan V, Wierzba Bobrowicz T, de Leon M, Saint Louis LA, Cohen IL, London E, Brown WT, Wisniewski T. The Neuropathology of autism: defects of neurogenesis and neuronal migration, and dysplastic changes. *Acta Neuropathol* 2010, 119, 755-770

3. Oral presentation. 9th Annual International Meeting for Autism Research (IMFAR), Philadelphia, PA, May 20-22, 2010

Wegiel J, Kuchna I, Nowicki K, Imaki H, Wegiel J, Marchi E, Ma SY, Chauhan A, Chauhan V, Cohen IL, London E, Brown WT, Wisniewski T. Defects of neurogenesis, neuronal migration, and dysplastic changes in the brains of autistic subjects.

4. Three posters. 9th Annual International Meeting for Autism Research (IMFAR), Philadelphia, PA, May 20-22, 2010

Ma S-Y, Kuchna I, Nowicki K, Wegiel J, Imaki H, Cohen IL, London E, Flory M, Brown WT, Wisniewski T, Wegiel J. Neuronal growth delay within the claustrum of autistic subjects.

Nowicki K, Wisniewski T, Kuchna I, Wegiel J, Imaki H, Ma S-Y, Cohen IL, London E, Brown WT, Wegiel J. Repetitive and stereotyped behaviors in autism are driven by abnormal development of the striatum but not of the substantia nigra.

Kuchna I, Nowicki K, Imaki H, Wegiel J, Ma SY, Marchi E, Cohen IL, London E, Brown WT, Wisniewski T, Wegiel J. Dystrophy with calcification within brains of autistic and control subjects.

CONCLUSIONS

2. **There is no single cause for the autistic phenotype.** Multi-regional qualitative and quantitative study of autistic subjects brains demonstrates a broad spectrum of developmental alterations and indicates that there is no one brain structure, neuronal population, neurotransmitter or neuropeptide defect which might be considered as the single cause of the autistic phenotype.
3. **Delay of neuronal growth.** Delay of neuronal growth, immaturity of some neurons and neuronal networks appears as a common developmental defect detected in the brain of autistic subjects by unbiased morphometric methods.
4. **Neurotransmitters and neuropeptides regulatory systems in autism.** The study of major regulatory systems of the brain: nucleus basalis of Meynert (cholinergic system), the substantia nigra (dopaminergic system), hypothalamic NPV and NSO (oxytocin and vasopressin) suggest that the neuronal populations playing a key role in production of neurotransmitters or neuropeptides are not affected (substantia nigra) or show rather limited distortion of developmental trajectories (NBMC, NSO, NPV). However, brain structures that are regulated by these centers reveal much broader and more significant developmental delays of neuronal growth.
3. **Desynchronized development of brain structures.** Brain structure and neuronal population-specific delay of neuronal growth with a broad range of differences in developmental delay, different trajectory of partial or complete correction of neuron size during late childhood and adulthood, reflect the complexity of desynchronized development of neurons, neuronal networks, and brain structures most likely contributing to the complex clinical manifestations of autism.
4. **Interindividual differences.** Significant interindividual differences in type, topography and severity of qualitative developmental abnormalities (altered neurogenesis, migration and dysplasia), and significant interindividual differences in topography and severity of delayed neuronal growth appear to be the major factor determining typical for autism extremely broad spectrum of interindividual differences.
5. **Hypothesis unifying qualitative and quantitative developmental alterations.** The study suggests that neuropathological qualitative alterations and morphometric qualitative alterations are not completely independent. Further studies may show that mechanisms triggering abnormal neurogenesis, migration and maturation delay neuronal growth.

REFERENCES

- Amaral D, Insausti R. Retrograde transport of D-[3H]-aspartate injected into the monkey amygdaloid complex. *Experimental Brain Research*, 1992, 88, 2, 375-388
- Amaral DG, Cowan WM, Subcortical afferents to the hippocampal formation in the monkey. *Journal of Comparative Neurology*, 1980, 189, 4, 573-591
- Arikuni T, Kubota K, Claustral and amygaloid afferents to the head of the caudate nucleus in macaque monkeys. *Neuroscience Research*, 1985, 2, 4, 239-254
- Bale TL, Davis AM, Auger AP, Dorsa DM, McCarthy MM, CNS region-specific oxytocine receptor expression: importance in regulation of anxiety and sex behavior. *J Neurosci* 2001, 21, 2546-2552
- Bennett CM, Baird AA. Anatomical changes in the emerging adult brain: a voxel-based morphometry study. *Hum Brain Mapp* 2006, 27, 9, 766-77
- Berns GS, Chappelow J, Cekic M, Zink CF, Pagnoni G, Martin-Skurski ME. Neurobiological substrates of dread. *Science*, 2006, 312, 5774, 754-758
- Bielsky IF, Hu SB, Szegda KL, Westphal H, Young LJ. Profound impairment in social recognition and reduction in anxiety-like behavior in vasopressin V1a receptor knockout mice. *Neuropsychopharmacology*, 2004, 29, 3, 483-93.
- Bloom P, German T. Two reasons to abandon the false belief task as a test of mind theory. *Cognition*, 2000, 7, B25-B31
- Brothers L, Ring B, Kling A, Response to neuronsss in the macaque amygdala to complex social stimuli. *Behavioral Brain Research*, 1990, 41, 3, 199-213
- Chachich ME, Powell DA. The role of claustrum in Pavlovian heart rate conditioning in the rabbit (*Oryctolagus cuniculus*): anatomical, electrophysiological, and lesion studies. *Behav Neuroscience*. 2004, 118, 3, 514-25
- Champagne F, Diorio J, Sharma S, Meaney MJ. Naturally occurring variations in maternal behavior in the rat are associated with differences in estrogen-inducible central oxytocine receptors. *PNAS* 2001,98, 12736-12741
- Cook EH Jr, Courchesne RY, Cox NJ, Lord C, Gonen D, Guter SJ, Lincoln A, Nix K, Haas R, Leventhal BL, Courchesne E. Linkage-disequilibrium mapping of autistic disorder, with 15q11-13 markers. *Am J Hum Genet* 1998, 62, 1077-1083
- Crick FC, Koch C. What function is the claustrum? *Philosophical Transactions of the Royal Society of London: Series B-Biological Sciences*, 2005,360,1458, 1271-1279

- Dubroff JG, Ficicioglu C, Segal S, Wintering N, Alavi A, Newberg A. FDG-PET finding of patients with galactosemia. *Journal of Inherited Metabolic Diseases*, 2008, 31, 533-339.
- Ferguson JN, Young LJ, Hearn EF, Matzuk MM, Insel TR, Winslow JT. Social amnesia in mice lacking the oxytocin gene. *Nat Genet* 2000, 25, 284–288
- Fernandez-Miranda JC, Rhoton AL, Kakizawa Y, Choi C, Alvarez-Linera J. The Claustrum and its projection system in the human brain: a microsurgical and tractographic anatomical study. *Journal of Neurosurgery*, 2008, 108, 4, 764-774
- Gabriels, R. L., M. L. Cuccaro, D. E. Hill, B. J. Ivers, and E. Goldson. Repetitive behaviors in autism: relationships with associated clinical features. *Res Dev Disabil*. 2005, 26, 169-181.
- Griebel G, Simiand J, Le Gal CS, Wagnon J, Pascal M, Scatton B, Maffrand JP, Soubrie P. Anxiolytic- and antidepressant-like effects of the non-peptide vasopressin V1b receptor antagonist, SSR149415, suggest an innovative approach for the treatment of stress-related disorders. *PNAS*, 2002, 99, 9, 6370-6375
- Heimer L, Alheid, G, Groenewegen H, Haber S, Harlan R, Zahm D. The Accumbens: beyond the core-shell dichotomy. *Journal of Neuropsychiatric & Clinical Neuroscience*, 1997, 9, 354-381
- Hollander, E., S. Novotny, M. Hanratty, R. Yaffe, C. M. DeCaria, B. R. Aronowitz, and S. Mosovich. Oxytocin infusion reduces repetitive behaviors in adults with autistic and Asperger's disorders. *Neuropsychopharmacology* 2003, 28, 193-198.
- Kovács GL, De Wied D. Peptidergic modulation of learning and memory processes. *Pharmacol Rev*. 1994, 46, 269-91
- Kötter R, Stephan KE, Useful or helpful? The “limbic system” concept. 1997, *Rev Neuroscience*, 1997, 8, 2, 139-145
- Landgraf R, Neumann ID. Vasopressin and oxytocin release within the brain: a dynamic concept of multiple and variable modes of neuropeptide communication. *Front Neuroendocrinol*. 2004, 25, 150-76.
- Lerner A, Bagic A, Hanakawa T, Boudreau E, Pagan F, Mari Z, Bara-Jimenez W, Aksu M, Sato S, Murphy D, Hallet M. Involvement of insula and cingulate cortices in control and suppression of natural urges. *Cereb Cortex*. 2009, 19, 1, 218-223
- Lopes de Silva. Anatomical organization and physiology of the limbic cortex. *Physiol. Rev*. 1990, 70, 453-511
- Lopes da Silva FH, Witter M, Boeijinga PH, Lohman. AH. Anatomic organization and physiology of the limbic cortex. *Physiol Rev*, 1990, 70, 2, 453-511
- Mogenson GJ, Jones DL, and Yim CY. From motivation to action: functional interface between the limbic system and the motor system. *Prog Neurobiol*, 1980, 14, 69-97

Morys J, Bobinski M, Wegiel J, Wisniewski HM, Narkiewicz. Alzheimer's disease severely affects areas of the claustrum connected with the entorhinal cortex. *J Hirnforsch*, 1996, 37, 2, 173-180

Morys J, Narkiewicz O, Wisniewski HM. Neuronal loss in the human claustrum following ulegyria. *Brain Research*, 1993, 616, 11, 176-180

Naqvi N, Durant D, Admission H, Benchers A. Damage to the insula disrupts addition to cigarette smoking. *Science*, 2007, 315, 5811, 531-534

Pearson R, Bridal O, Gutter KC, Powell TP. The organization of the connections between the cortex and claustrum in the monkey. *Brain Research*, 1982, 234, 2, 435-441

Popik P, Vetulani J, van Ree JM. Low doses of oxytocin facilitate social recognition in rats. *Psychopharmacology* 1992, 106, 1, 71-74

Popik P, Vos P, van Ree JM. Neurohypophyseal hormone receptors in the septum are implicated in social recognition in the rat. *Behavioral Pharmacology* 1992, 3, 4, 351-358

Schwab JJ. Psychiatric aspects of infectious diseases. *Current Psychiatric* 1982, 225-239

Spencer J, Sander B, Lau S, Krude H, Scheffner D. Severe transitory encephalopathy with reversible lesions of the claustrum. *Pediatric Radiology*, 1996, 26,11,769-771

Stein MB, Simmons AN, Feinstein JS, Paulus MP. Increased amygdala and insula activation during emotion processing in anxiety-prone subjects. *American Journal of Psychiatry* 2007, 164, 2, 318-327

Szatmari P, Jones MB, Zwaigenbaum L, MacLean ME. Genetics of autism: overview and new directions. *J Autism Dev Disord* 1998, 28, 351-368.

Windle RC, Windle M. An investigation of adolescents' substance use behaviors, depressed affect, and suicidal behaviors. *J Child Psychol Psychiatry*. 1997, 38, 8, 921-9.

Zhang X, Hannesson D, Saucier D, Wallace A, Howland J, Corcoran ME. Susceptibility to kindling and neuronal connections of the anterior Claustrum. *Journal of Neuroscience* 2001, 21, 10, 3674-3687

APPENDICES

1. Wegiel J, Wisniewski T, Chauhan A, Chauhan V, Kuchna I, Nowicki K, Imaki H, Wegiel J, Ma SY, Wierzbica Bobrowicz T, Cohen IL, London E, Brown WT. Type, topography and sequelae of neuropathological changes shaping clinical phenotype of autism. In: Autism: Oxidative Stress, Inflammation, and Immune Abnormalities. Ed.: Abha Chauhan, Ved Chauhan and W. Ted Brown. Taylor & Francis/CRC Press, Boca Raton, FL. 2010, pp 1-34
2. Wegiel J, Kuchna I, Nowicki K, Imaki H, Wegiel J, Marchi E, Ma SY, Chauhan A, Chauhan V, Wierzbica Bobrowicz T, de Leon M, Saint Louis LA, Cohen IL, London E, Brown WT, Wisniewski T. The Neuropathology of autism: defects of neurogenesis and neuronal migration, and dysplastic changes. *Acta Neuropathol* 2010, 119, 755-770
3. Abstracts- Three posters. 9th Annual International Meeting for Autism Research (IMFAR), Philadelphia, PA, May 20-22, 2010

Ma S-Y, Kuchna I, Nowicki K, Wegiel J, Imaki H, Cohen IL, London E, Flory M, Brown WT, Wisniewski T, Wegiel J. Neuronal growth delay within the claustrum of autistic subjects.

Nowicki K, Wisniewski T, Kuchna I, Wegiel J, Imaki H, Ma S-Y, Cohen IL, London E, Brown WT, Wegiel J. Repetitive and stereotyped behaviors in autism are driven by abnormal development of the striatum but not of the substantia nigra.

Kuchna I, Nowicki K, Imaki H, Wegiel J, Ma SY, Marchi E, Cohen IL, London E, Brown WT, Wisniewski T, Wegiel J. Dystrophy with calcification within brains of autistic and control subjects.

Oral Presentation

Wegiel J, Kuchna I, Nowicki K, Imaki H, Wegiel J, Marchi E, Ma SY, Chauhan A, Chauhan V, Cohen IL, London E, Brown WT, Wisniewski T. Defects of neurogenesis, neuronal migration, and dysplastic changes in the brains of autistic subjects.

4. Posters 9th Annual International Meeting for Autism Research (IMFAR), Philadelphia, PA, May 20-22, 2010

Ma S-Y, Kuchna I, Nowicki K, Wegiel J, Imaki H, Cohen IL, London E, Flory M, Brown WT, Wisniewski T, Wegiel J. Neuronal growth delay within the claustrum of autistic subjects.

Nowicki K, Wisniewski T, Kuchna I, Wegiel J, Imaki H, Ma S-Y, Cohen IL, London E, Brown WT, Wegiel J. Repetitive and stereotyped behaviors in autism are driven by abnormal development of the striatum but not of the substantia nigra.

Kuchna I, Nowicki K, Imaki H, Wegiel J, Ma SY, Marchi E, Cohen IL, London E, Brown WT, Wisniewski T, Wegiel J. Dystrophy with calcification within brains of autistic and control subjects.

1 Type, Topography, and Sequelae of Neuropathological Changes Shaping Clinical Phenotype of Autism

*Jerzy Wegiel, Thomas Wisniewski,
Abha Chauhan, Ved Chauhan, Izabela Kuchna,
Krzysztof Nowicki, Humi Imaki, Jarek Wegiel,
Shuang Yong Ma, Teresa Wierzba Bobrowicz,
Ira L. Cohen, Eric London, and W. Ted Brown*

CONTENTS

1.1	Introduction	2
1.2	Clinical, Etiological, and Neuropathological Diversity in Autism	3
1.2.1	Clinic	3
1.2.2	Etiology	4
1.2.3	Neuropathology	4
1.3	Deregulation of Brain Growth in Early Childhood	5
1.3.1	Developmental Heterochronicity	7
1.3.2	Functional Consequences of Abnormal Brain Development	8
1.4	Cortical and Subcortical Neuropathology	8
1.4.1	Cortical Dysgenesis, Lamination Defects, Migration Disturbances	8
1.4.2	Brain Structure–Specific Delay of Neuronal Growth	9
1.4.3	Minicolumnar Abnormalities in Autism	9
1.5	Neuronal Oxidative Stress and Metabolic Changes	10
1.5.1	Oxidative Stress in Autism	10
1.5.2	Lipofuscin in Autism	11

1.5.3	β -Amyloid Precursor Protein and Intraneuronal Amyloid β in Autism	11
1.6	Clinicopathological Correlations	12
1.6.1	Speech, Language, and Verbal and Nonverbal Communication.....	12
1.6.2	Face Perception.....	13
1.6.3	Social Attachment—The Role of the Hypothalamus in Behavioral Deficits	14
1.6.4	Sensorimotor Deficits, and Repetitive and Stereotyped Behaviors	15
1.6.5	Cognitive Deficits.....	16
1.6.6	Epilepsy-Associated Pathology.....	17
1.7	Mechanisms Affecting Brain Development.....	18
1.7.1	BDNF and Neurotrophins in Autism.....	18
1.7.2	Brain Stem and the Role of Serotonin in Brain Development and Clinical Features of Autism	19
1.8	Closing Remarks.....	20
	Acknowledgments.....	20
	References.....	20

1.1 INTRODUCTION

The aim of this chapter is to identify the type, topography, and sequelae of neuropathological changes that contribute to the clinical phenotype of autism. Results of recent magnetic resonance imaging (MRI) and postmortem neuropathological and stereological studies of autism brain suggest a dynamic model of sequential subdivision of age- and brain-specific structural and functional changes. Acceleration of brain growth in the first year of life and deceleration in the second and third years appear to play a pivotal role in the onset of clinical signs of autism (Courchesne et al., 2001, 2003; Courchesne and Pierce, 2005b; Dawson et al., 2007; Dementieva et al., 2005; Gillberg and de Souza, 2002; Redcay and Courchesne, 2005). The range of deviation from the normal trajectory of brain growth may be a factor determining the severity of the disease (Courchesne et al., 2003). Developmental heterochronicity (differential rates of growth of various brain regions compared to controls), resulting in selective overgrowth of some brain regions, appears to be a key factor determining topography and brain region-specific type of cytoarchitectonic changes (Carper and Courchesne, 2005; Carper et al., 2002; Courchesne et al., 2001; Hazlett et al., 2005; Sparks et al., 2002). Topographic developmental heterochronicity may result in impairment of both local and global connectivity, leading to local overconnectivity and impairment of long-distance connectivity (Baron-Cohen, 2004; Casanova et al., 2006; Courchesne and Pierce, 2005a). Stereological studies have revealed neuronal developmental heterochronicity in early childhood, resulting in selective developmental delay of the growth of neurons in some subcortical structures and the cerebellum during the most critical stage of development of social behaviors and communication skills (Wegiel et al., 2008). Distortions of brain and neuronal development are reflected in abnormal

cortical minicolumn organization (Casanova et al., 2002, 2006), local dysgenesis, and ectopias (Bauman and Kemper, 1985; Bauman et al., 1997; Kemper and Bauman, 1993, 1998). These complex developmental abnormalities appear to lay the foundation for secondary and tertiary metabolic, structural, and functional changes, including seizures and risk of sudden unexpected death; signs of oxidative stress, early and enhanced accumulation of products of cell organelle degradation with lipofuscin deposition; modified processing of β -amyloid precursor protein with accumulation of truncated amyloid beta; and other as of yet unidentified changes. Secondary pathologic changes appear to be indicators of the susceptibility of abnormally developing neurons to further modifications during cell maturation and aging. The pattern of morphological changes emerging from these multidisciplinary studies appears to represent a major trend. However, modifications of the course of disease and sub-patterns of developmental changes result in a broad spectrum of morphological and clinical interindividual differences.

1.2 CLINICAL, ETIOLOGICAL, AND NEUROPATHOLOGICAL DIVERSITY IN AUTISM

Autism is the prototype of a pervasive developmental disorder (PDD) and is characterized by (a) qualitative impairments in reciprocal social interactions, (b) qualitative impairments in verbal and nonverbal communication, (c) restricted repetitive and stereotyped patterns of behavior, interests, and activities, and (d) onset prior to the age of 3 years. PDD also includes childhood disintegrative disorder, Asperger's disorder, Rett syndrome, and pervasive developmental disorder—not otherwise specified (PDD-NOS). The common features of all these disorders are qualitative deficits in social behavior and communication (American Psychiatric Association, 2000).

1.2.1 CLINIC

AQ2

In most cases (90%–95%), it is not presently possible to detect a known or specific etiology. These cases are referred to as idiopathic or nonsyndromic autism (Boddaert et al., 2009; Gillberg and Coleman, 1996). In 6% (Fombonne, 2003), 5% (Tuchman et al., 1991), or 10% (Rutter et al., 1994) of cases, autism was diagnosed in association with other disorders. About 30% of children with idiopathic autism have complex autism, defined by the presence of dysmorphic features, microcephaly and/or a structural brain malformation (Miles et al., 2005). About 70% of children with autism have essential autism, defined by the absence of physical abnormalities. For most children, the onset of autism is gradual. However, a multisite study revealed significant regression at ages of 18 to 33 months (regressive autism) in about 13.8% (Colorado) to 31.6% (Utah) of autistic subjects (Department of Health and Human Services, 2007). Moreover, the manifestations of autism vary greatly, depending on developmental level and chronological age of the affected individual. The majority of patients exhibit serious social and communicative impairments throughout life but some improve enough to be able to live relatively independently as adults. In 44.6% of children, autism is associated with cognitive impairment (defined as having intelligence quotient scores of <70; Department of Health and Human Services, 2007). Expressive language function in

individuals with autism may vary from mutism to verbal fluency (Rapin, 1996; Stone et al., 1997; Wetherby et al., 1998). Sensorimotor deficits also show significant interindividual differences, with more frequent and severe impairments of gross and fine motor function (motor stereotypes, hypotonia, limbic apraxia) in subjects with lower IQ (Rogers et al., 1996). Hand mannerisms and body rocking are reported in 37% to 95% of individuals with autism (Lord and Rutter, 1995; Rapin, 1996; Rogers et al., 1996), whereas preoccupation with sensory features of objects, abnormal responsiveness to environmental stimuli, or paradoxical responses to sensory stimuli are seen in 42% to 88% of people with autism (Kientz and Dunn, 1997). Epilepsy is a comorbid complication, occurring in up to 33% of individuals with autism (Tuchman and Rapin, 2002).

1.2.2 ETIOLOGY

The clinical diversity of autism reflects the etiologic heterogeneity of this disorder. Genetic factors; pre-, peri-, and postnatal pathological factors; and concurrent diseases may contribute to autism (Muhle et al., 2004; Newschaffer et al., 2002; Rutter et al., 1994). About 5% to 10% of cases are associated with several distinct genetic conditions including fragile X syndrome, tuberous sclerosis, phenylketonuria, Rett syndrome, and chromosomal anomalies such as Down syndrome (DS) (Folstein and Rosen-Scheidley, 2001; Fombonne, 2003; Smalley et al., 1988; Yonan et al., 2003). Autism spectrum disorders (ASDs) in people with DS have been described in several reports (Ghaziuddin et al., 1992; Howlin et al., 1995; Pracher and Clarke, 1996; Wakabayashi, 1979), and the prevalence of autism in boys with DS was estimated as at least 7% (Kent et al., 1999). The prevalence of autism in the fragile X syndrome is estimated as 15%–28% (Hagerman, 2002). Cytogenetic abnormalities (partial duplications, deletions, inversions) in the 15q11-q13 region account for 1% to 4% of autism cases (Cook, 1998; Gillberg, 1998). Several potential candidate genes have been identified in both autosomes and X chromosomes, including the tuberous sclerosis gene on chromosomes 9 and 16; serotonin transporter on chromosome 17; gamma-aminobutyric acid receptor-beta 3 on chromosome 15; neuroligins on the X chromosome (see Vorstman et al., 2006); and possibly *PTEN* on chromosome 10 (Butler et al., 2005). Modifications in the tryptophan hydroxylase gene may play a modest role in autism susceptibility (Coon et al., 2005).

1.2.3 NEUROPATHOLOGY

While knowledge of the clinical and genetic factors in autism is based on examination of thousands of patients, postmortem neuropathological studies are based on reports of a very small number of brains. A review by Palmen et al. (2004) revealed that between 1980 and 2003, only 58 brains of individuals with autism have been examined, and results of only a few neuropathological and stereological studies were published. Usually, neuropathological reports and morphometric reports were based on evaluation of one or several brains. Due to the broad age spectrum and the etiological and clinical diversity in autism, the pattern of neuropathological changes reported is incomplete and often inconsistent. As a result, the morphological markers and

neuropathological diagnostic criteria of autism have not yet been established (Lord et al., 2000; Pickett and London, 2005). In the past, the contribution of postmortem studies to the detection and characterization of neuropathological changes and mechanisms leading to structural and functional manifestations of autism was limited because of (a) the deficit of autism brains, resulting in a lack of statistical power, (b) the lack of efficient mechanisms for sharing the limited tissue resources, (c) the lack of complex studies of the dynamic of changes during the life span, (d) the infrequent application of unbiased morphometric methods to detect quantitative differences, and (e) the averaging of results from subjects with different clinical and morphological manifestations of autism. Heterogeneity within the autism spectrum is the major obstacle to autism research at all levels (Newschaffer et al., 2002), including neuropathological studies and attempts at detection of clinicopathological correlations. Recent evidence of genetic fractionation of social impairment, communication difficulties, and rigid and repetitive behaviors indicates that heterogeneity in ASD could be an unavoidable consequence of the contribution of nonoverlapping genes. If different features of autism are caused by different genes associated with different brain regions and related to different core cognitive impairments (Happé et al., 2006), it seems likely that many brain networks are involved in the pathology of autism. The diversity of neuropathological findings and the commonly reported inconsistencies in regional findings correspond to developmental impairments in many interacting brain networks and to expansion from “local” abnormalities to “nonlocal” effects of the emerging cognitive system. In spite of these limitations, “localizing” models are still the main approach to the identification of pathological changes as a component of the structural and functional abnormalities of the networks (Müller, 2007).

The possibility that autism is associated with neuropathological changes was explored in the first studies reported between 1980 and 1989 (Bauman and Kemper, 1985; Courchesne et al., 1987, 1988; Damasio et al., 1980; Gaffney et al., 1987; Hashimoto et al., 1989, 1993; Murakami et al., 1989; Ritvo et al., 1986). Expansion of these studies through examination of larger cohorts and application of stereology, functional and structural MRI, and biochemistry resulted in the identification of several major forms of pathology contributing to the clinical phenotype including abnormal acceleration of brain growth in early childhood (Redcay and Courchesne, 2005), delay of neuronal growth (Wegiel et al., 2008), changes in brain cytoarchitecture (Bailey et al., 1998; Bauman and Kemper, 1985; Casanova et al., 2002, 2006), metabolic modifications with abnormal amyloid precursor protein (APP) processing (Bailey et al., 2008; Brown et al., 2008; Sokol et al., 2006), enhanced oxidative stress (reviewed in Chauhan and Chauhan, 2006), and turnover of cell organelle with pigment accumulation and glial activation (Lopez-Hurtado and Prieto, 2008).

1.3 DEREGULATION OF BRAIN GROWTH IN EARLY CHILDHOOD

The major measures of age-related changes are head circumference, MRI-based volumetry of the brain and brain structures, and postmortem brain weight and volume of brain subdivisions. Between 1990 and 2000, several groups reported increased head circumference (Bailey et al., 1995; Bolton et al., 1995; Davidovitch et al., 1996;

Fidler et al., 2000; Fombonne et al., 1999; Lainhart et al., 1997; Miles et al., 2000; Steg and Rapoport, 1975; Stevenson et al., 1997), whereas MRI-based studies revealed increased brain volume (Piven et al., 1995, 1996). According to Fombonne et al. (1999), the prevalence of macrocephaly in autism is about 20%. In a report by Bailey et al. (1998), four of six subjects with autism 4 to 24 years of age had macrocephaly. Increased brain weight was reported in postmortem studies by Bailey et al. (1993) and Kemper and Bauman (1998). Increase in the volume was regional and not generalized, with the greatest enlargement in the occipital and parietal lobes (Filipek, 1996; Filipek et al., 1999; Piven et al., 1995, 1996). However, in several studies, an increase in brain size was not detected (Garber and Ritvo, 1992; Haznedar et al., 2000). Inconsistency in detection of abnormal head and brain size can be associated with interindividual differences, the age of examined individuals, and the methods applied. Courchesne et al. (2003) integrated their work and that of other researchers into the concept of four phases of modified brain growth, described below.

At birth, the head circumference of neonates later diagnosed with autism is normal or slightly less than that observed in normally developing children (Courchesne et al., 2003; Dawson et al., 2007; Dementieva et al., 2005; Dissanayake et al., 2006; Gillberg and de Souza, 2002; Hazlett et al., 2005; Lainhart et al., 1997; Mason-Brothers et al., 1990; Stevenson et al., 1997). Slight undergrowth is independent of body growth and may be a reflection of prenatal neural developmental defects corresponding to pathology detected in postmortem studies of the brains of autistic adults (Bailey et al., 1998; Casanova et al., 2002; Courchesne et al., 2003; Kemper and Bauman, 1998). In only 5% of neonates diagnosed later as autistic was the head circumference more than that in normally developing infants (Courchesne et al., 2003; Dementieva et al., 2005).

In the second phase, by 1 or 2 years of age, a rapid and large increase in head circumference distinguished children diagnosed later with autism from normally developing children (Courchesne et al., 2003; Dawson et al., 2007; Dementieva et al., 2005; Dissanayake et al., 2006; Hazlett et al., 2005). Ninety percent of 2- and 3-year-old children with autism had brain volumes larger than those of control children (Courchesne et al., 2001). According to Dawson et al. (2007), a period of exceptionally rapid head growth is limited to the first year of life, and head growth decelerates after 12 months of age. Acceleration of growth in head circumference appears to begin at about 4 months (Courchesne and Pierce, 2005a; Gillberg and de Souza, 2002; Redcay and Courchesne, 2005). Using meta-analysis based on evaluation of head circumference converted to brain volume, brain volume measured from MRI, and brain weight from postmortem studies, Redcay and Courchesne (2005) revealed that brain size increases from 13% smaller than in control subjects at birth to 10% larger than in control infants at 1 year, but only 2% greater by adolescence. The greater growth rate of head circumference in the first year, and its return to normal rates thereafter, is not accounted for by an overall growth in stature. Studies of behavioral development in infants later diagnosed with autism suggest that the period of acceleration of head growth precedes and overlaps with the onset of behavioral changes, and that the period of deceleration coincides with a period of behavioral decline or worsening of symptoms in the second year of life (Dawson et al., 2007). Coincidence of acceleration of brain growth rate with onset and

worsening of clinical symptoms may indicate that structural developmental changes critical for a lifelong phenotype occur in early infancy. Acceleration of brain growth in the first year and deceleration in the second year of life suggest that failure of the mechanism controlling brain growth in the first year of life plays an essential role in the onset of clinical features of autism. Identification of these mechanisms may lead to conceptualization of early preventive treatments.

In the third phase, of 2 to 4 years, the overall rate of brain growth slows but is still 10% more than in normally developing children (Carper et al., 2002; Courchesne et al., 2001; Hazlett et al., 2005; Sparks et al., 2002). In 4- to 5-year-old autistic children, MRI-based estimated brain volume is 1350 mL, whereas in normally developing children, a comparable volume (1360 mL) is reached about 8 years later. In post-mortem studies, the brain weight of 3- to 5-year-old autistic males was 15% higher (1451 g) than in control males of this age (1259 g) (Redcay and Courchesne, 2005).

In the fourth phase, the volume of the brain decreases, and this trend extends from middle/late childhood through adulthood. Head (Aylward et al., 2002) or brain enlargement (Bailey et al., 1998; Hardan et al., 2001; Lainhart et al., 1997; Piven et al., 1995, 1996) has also been observed in studies of older populations of autistic individuals. However, by adolescence and adulthood, the average size of the brain is only 1% to 3% greater in autistic than in control cohorts (Redcay and Courchesne, 2005).

Moreover, the pattern of brain growth reflects the severity of clinical manifestation of autism (Courchesne et al., 2003). Among infants who have the more severe form of autism, 71% showed increases during their first year of more than 1.5 S.D., with 59% showing increases between 2.0 and 4.3 S.D. In children with a less severe form of autism, PDD-NOS, acceleration of brain growth is observed later, and the increase is less pronounced. Later onset and slower rate of progression of autism appear to be associated with a better outcome.

1.3.1 DEVELOPMENTAL HETEROCHRONICITY

Developmental heterochronicity studies indicate that autism is a disorder involving a transient period of pathological acceleration of brain growth. Developmental heterochronicity, with different rates of growth for different brain regions/structures, appears to be the second major factor contributing to the clinical phenotype. MRI studies showed that overgrowth of the frontal and temporal lobes and amygdala, brain regions that are involved in cognitive, social, and emotional functions as well as language development, is synchronized with brain overgrowth in 2- to 4-year-old autistic children in contrast to a different rate of growth of the occipital cortex (Carper and Courchesne, 2005; Carper et al., 2002; Courchesne et al., 2001; Hazlett et al., 2005; Sparks et al., 2002). The reduced size of the body and posterior subregions of the corpus callosum noted in subjects with autism may indicate disproportions in brain subregions development (Piven et al., 1997b). The cellular and molecular basis for transient acceleration of brain growth and enhanced growth of some brain regions is not known, but Courchesne et al. (2003) proposed that the observed pattern is associated with an excessive number of neurons, enhanced rate of growth of size of neurons, and increased number of minicolumns as well as excessive and premature expansion of the dendritic tree.

1.3.2 FUNCTIONAL CONSEQUENCES OF ABNORMAL BRAIN DEVELOPMENT

Using a computational model analogue of autism, Cohen (2007) has argued that an interaction between stochastic and above-average or “excessive” numbers of neural connection factors has implications for understanding the disorder. In particular, a relative excess of connections could lead to enhanced recognition of complex patterns in the environment. In Cohen’s chapter, it was noted that if large and complex brains are in part familial (Courchesne et al., 2003; Fidler et al., 2000), and brain size is heritable (Pfefferbaum et al., 2000) and positively correlated with IQ (Pennington et al., 2000), then behavioral outcomes both within and across generations of family members could result in (a) individuals who may be unusually gifted in their ability to handle complex nonlinear problems such as mathematics or computer science, (b) individuals with autism, or (c) individuals with a combination of autism or autistic-like behavior and giftedness (many typical Asperger’s cases). These trends are detected among relatives of subjects with autism (Folstein and Rutter, 1988).

The effect of an abnormal trajectory of brain development observed in autism are well-validated characteristics of the learning style of children with autism, including (a) greater attention to idiosyncratic than socially relevant stimuli, (b) stimulus overselectivity or a lack of drive for central coherence, (c) problems with acquiring fuzzy concepts, (d) development of savant skills, (e) problems with generalization of previously acquired skills, (f) rigidity and resistance to change, (g) social and communication deficits, and (h) difficulty in learning complex higher-order concepts (Cohen, 2007).

1.4 CORTICAL AND SUBCORTICAL NEUROPATHOLOGY

1.4.1 CORTICAL DYSGENESIS, LAMINATION DEFECTS, MIGRATION DISTURBANCES

The fundamental characteristics of the neuropathological changes described by Kemper and Bauman (1993, 1998), Bauman and Kemper (1985, 1996), and Bauman et al. (1997) suggest three major neuropathologies in the brain of people with autism: (a) curtailed development of neurons in the structures that are substrates for memory and emotions—the entorhinal cortex, hippocampus, subiculum, anterior cingulate gyrus and mamillary body; (b) a congenital decrease in the number of Purkinje cells in the cerebellum; and (c) age-related differences in cell size and number of neurons in the cerebellar nuclei and in the inferior olivary nucleus. Microdysgenesis is represented by increased neuronal density in the cortical layer, clustering of cortical neurons, disorganization of cortical layers, neuron cytomegaly, ectopic neurons, and nodular heterotopias. A detailed study of serial sections from the brain of a 29-year-old man with autism revealed reduced neuronal size and increased cell-packing density (Bauman and Kemper, 1985), both features of an immature brain (Friede, 1975). Cell-packing density was increased by 66% in the hypothalamus and mamillary body, and by 54% in the medial septal nucleus, with smaller nerve cells. The reduced size of neurons and the selective increase in cell-packing density were seen in central (40%), medial (28%), and cortical nuclei (35%). Atrophy of the neocerebellar cortex, with marked loss of Purkinje cells and, to a lesser extent, of granule cells, was present in gracile, tonsil, and inferior semilunar lobules. Changes were not detected in

the anterior lobe or the vermis. Reduced numbers of cells were noticed in fastigial, globose, and emboliform nuclei, and cells were small and pale. The dentate nucleus was distorted. Retrograde neuronal loss in the inferior olive related to neuronal loss in cerebellar cortex was not found, but olivary neurons were small and pale. Brain cytoarchitecture abnormalities were not associated with gliosis. In a 21-year-old female with autism, Rodier et al. (1996) found that the brain was smaller than a control brain, and the length of facial nerve nucleus was less than 500 μm as compared to 2610 μm in the control subject.

1.4.2 BRAIN STRUCTURE–SPECIFIC DELAY OF NEURONAL GROWTH

The reduced size of neurons and their nuclei in the cortex of autistic subjects reported by Casanova et al. (2006) could be an indicator of reduced or impaired functional connectivity between distant cortical regions (Casanova et al., 2006; Just et al., 2004; Koshino et al., 2005). Our ongoing studies of series of brains from age-matched autistic and control subjects (Wegiel et al., 2008) indicate that reduced size of neurons is a brain structure–specific marker. In 4- to 7-year-old autistic children, Purkinje cells were smaller by 38%. Neurons in the dentate nucleus were reduced by 26%; in the amygdala, by 24%; in the nucleus accumbens, by 41%; in caudate, by 20%; and in the putamen, by 27%. Neurons in the nucleus of the facial nerve and the nucleus olivaris did not show a significant difference from controls. The second significant feature of the pattern of neuronal size abnormalities is the partial or complete correction of the size of neurons (for example, in the nucleus accumbens) observed in late childhood or adulthood. This study indicates that the delay of growth of neurons is the most consistent pathology detected in the brains of examined people with autism. Pathology is brain structure–specific. Changes may range from no delay to severe developmental delay. The youngest examined children (4 to 7 years old) show the most severe deficit in the volume of the neuronal body and nucleus. Partial correction of cell volume is observed in late childhood and adulthood, which indicates that brain structure and function undergo modifications during the life span. The study of basal ganglia and cerebellum supports the hypothesis that clinical manifestations of autism are the result of regional neuronal maldevelopment.

One may assume that mechanisms regulating growth of the neuron in early childhood are the target of factors that are the cause of autism. The result of deregulation of these mechanisms could be (a) significantly delayed growth of neuronal body, nucleus, dendritic tree, spines, and reduced number of synapses and (b) functional deficits corresponding to these structural developmental delays. These abnormalities of very early childhood might be the major contributor to clinical deficits that are the basis for the clinical diagnosis of autism at the age of 3 years.

1.4.3 MINICOLUMNAR ABNORMALITIES IN AUTISM

The next significant contribution to detection of neocortical developmental pathology is the result of studies of minicolumns by Casanova's group (Buxhoeveden and Casanova, 2002; Casanova et al., 2002, 2006). Malformations of cortical development are observed in heterogeneous disorders caused by abnormalities of cell

proliferation, apoptosis, cell migration, cortical organization, and axon pathfinding (Hevner, 2007). Clinically malformations of cortical development are significant causes of mental retardation, seizures, cerebral palsy, and neuropsychiatric disorders (Barkovich et al., 2005; Guerrini and Marini, 2006; Sarnat and Flores-Sarnat, 2004). Minicolumns are considered a basic architectonic and functional unit of the human neocortex (Buxhoeveden and Casanova, 2002; Casanova et al., 2002). Increased neuron density by 23%, reduced size of neurons in minicolumns, and a concomitant increase in the total number of minicolumns appears to illustrate the bias of local rather than global information processing (Casanova et al., 2002, 2006), resulting in a “hyper-specific brain” (McClelland, 2000). Synchronization of interactions requiring the involvement of distant brain regions is impaired in autism as a result of developmental connectivity deficits (underconnectivity) of smaller neurons (Just et al., 2004; Koshino et al., 2005; Zilbovicius et al., 1995). Structural imaging studies also suggest the overrepresentation of short association fibers in autism, with a regional increase in the volume of white matter (Herbert et al., 2004) favoring the local information processing observed in autistic subjects (Happé, 1999).

1.5 NEURONAL OXIDATIVE STRESS AND METABOLIC CHANGES

An increasing body of evidence suggests that the abnormal rate of development of neurons and neuronal networks in early infancy is followed by metabolic changes, with signs of oxidative stress, enhanced autophagocytosis, and lipofuscin accumulation, leading to early selective neuronal structural and functional changes.

1.5.1 OXIDATIVE STRESS IN AUTISM

Oxidative stress is known to be associated with premature aging of cells and can lead to inflammation, damaged cell membranes, autoimmunity, and cell death. The brain is highly vulnerable to oxidative stress due to its limited antioxidant capacity, higher energy requirement, and high amounts of unsaturated lipids and iron (Juurlink and Peterson, 1998). The brain makes up about 2% of body mass but consumes 20% of metabolic oxygen. The vast majority of energy is used by the neurons (Shulman et al., 2004). Glutathione (GSH) is the most important antioxidant for detoxification and elimination of environmental toxins. Due to the lack of glutathione-producing capacity by neurons, the brain has a limited capacity to detoxify reactive oxygen species (ROS). Therefore, neurons are the first cells to be affected by the increase in ROS and shortage of antioxidants and, as a result, they are most susceptible to oxidative stress. Antioxidants are required for neuronal survival during the early critical period (Perry et al., 2004). Children are more vulnerable than adults to oxidative stress because of their naturally low glutathione levels from conception through infancy (Erden-Inal et al., 2002; Ono et al., 2001). The risk created by this natural deficit in detoxification capacity in infants is increased by the fact that some environmental factors that induce oxidative stress are found at higher concentrations in developing infants than in their mothers, and accumulate in the placenta.

Accumulating evidence from our and other groups suggests increased oxidative stress in autism (reviewed in Chauhan and Chauhan, 2006). Lipid peroxidation is a chain reaction between polyunsaturated fatty acids and ROS, producing lipid peroxides

and hydrocarbon polymers that are both highly toxic to the cell. We have reported that levels of malondialdehyde (MDA), a marker of lipid peroxidation, are increased in the plasma from children with autism (Chauhan et al., 2004). Other studies on erythrocytes (Zoroglu et al., 2004) and urine samples (Ming et al., 2005) have also indicated increased levels of lipid peroxidation markers in autism, thus confirming an increased oxidative stress in autism. Recent studies with the postmortem brain samples from autism and control subjects have provided further evidence on increased oxidative stress in autism. Increased levels of lipid-derived oxidative protein modifications, i.e., carboxyethylpyrrole and iso[4]levuglandin E₂-protein adducts, and heme-oxygenase-1 (an inducible antioxidant enzyme) have been reported in the autistic brain, primarily in the white matter (Evans et al., 2008). Sajdel-Sulkowska et al. (2008) have reported elevated levels of 3-nitrotyrosine (a specific marker for oxidative damage to proteins) in the cerebella of subjects with autism. In addition, we have observed increased lipid peroxidation in cerebellum and temporal cortex of brain in autism (Chauhan et al., 2009). MDA levels were significantly increased by 124% in the cerebellum, and by 256% in the temporal cortex in autism as compared to control subjects.

1.5.2 LIPOFUSCIN IN AUTISM

Lopez-Hurtado and Prieto (2008) revealed a significant increase in the number of lipofuscin-containing cells in the brain of 7- to 14-year-old autistic subjects (by 69% in area 22, 149% in area 39, and 45% in area 44). The increase in the number of lipofuscin-containing cells was paralleled by neuronal loss and glial proliferation. Lipofuscin accumulation is a component of aging (Brunk and Terman 2002a,b; Brunk et al., 1992; Szweda et al., 2003), the neurodegeneration observed in Alzheimer's (Stojanovic et al., 1994) and Parkinson's diseases (Tórsdóttir et al., 1999), developmental syndromes such as Rett syndrome (Jellinger et al., 1988), and autism (Lopez-Hurtado and Prieto, 2008), and such psychiatric disorders as bipolar affective disorder (Yanik et al., 2004) and schizophrenia (Akyol et al., 2002; Herken et al., 2001).

Lipofuscin is an intralysosomal deposit of products of autophagocytosis and degradation of cytoplasmic components, including mitochondria, which cannot be degraded further or exocytosed. Oxidative stress is considered the factor contributing to lipid and protein damage and degradation, resulting in lipofuscin production and accumulation (Brunk et al., 1992; Sohal and Brunk, 1989). The presence of oxidatively modified proteins and lipids in lipofuscin supports the causative link between enhanced oxidative stress, autophagocytosis, and deposition of products of degradation in the lysosomal pathway and lipofuscin (Brunk and Terman, 2002a,b; Szweda et al., 2003; Terman and Brunk, 2004) and suggests that in autism, abnormal development is associated with early signs of oxidative stress and enhanced degradation and, possibly, turnover of cytoplasmic components.

1.5.3 β -AMYLOID PRECURSOR PROTEIN AND INTRANEURONAL AMYLOID β IN AUTISM

Sokol et al. (2006) detected signs of overexpression of APP in about 40% of autistic subjects. The levels of secreted APP in plasma in children with severe autistic behavior and aggression were two or more times the levels in children without autism, and up to

fourfold more than in children with mild autism. The trend observed in autistic children, with higher levels of secreted β -APP and nonamyloidogenic secreted β -APP, and lower levels of A β 1–40 compared to controls, suggests an increased α -secretase pathway in autism (anabolic nonamyloidogenic APP processing). Enzyme-linked immunosorbent assay (ELISA) study of blood plasma from 25 autistic children 2–4 years of age and 25 age-matched control children revealed significantly increased level of secreted amyloid precursor protein alpha (sAPP- α) in 60% of autistic children (Bailey et al., 2008). Western blotting analysis confirmed higher levels of sAPP- α in autistic children.

AQ3 Amino-terminally truncated intraneuronal amyloid (A β) is present in the neurons of control subjects, and the amount of intraneuronal A β increases with age (Wegiel et al., 2007). This author's study of 10 brains of autistic people revealed enhanced intraneuronal accumulation of amino-terminally truncated A β in 50% of autistic subjects, including in one 5-year-old child and four adults 20, 23, 52, and 62 years of age. A similar pattern was also found in four examined brains of people with autism and isodicentric chromosome 15 (idic15) (Brown et al., 2008). In idic15, excessive accumulation of intraneuronal A β might be related to an extra copy of one of the amyloid precursor protein-binding protein (APBA-2) genes localized on chromosome 15. In many brain regions, A β is accumulated in large cytoplasmic granules corresponding to deposits of lipofuscin. Numerous large lipofuscin deposits with very strong A β immunoreactivity in the neurons of several children and adults with autism appear to reflect severe metabolic stress affecting all of the neurons in the amygdala, all large neurons in the caudate/putamen, a majority of Purkinje cells, and the neurons in the dentate nucleus and nucleus olivaris, but only about 30%–40% of cortical pyramidal neurons. Accumulation of truncated A β appears to be a by-product of enhanced degradation of transmembrane APP. The aggregated intracellular A β induces the production of ROS and lipid peroxidation products and ultimately results in leakage of the lysosomal membrane (Glabbe, 2001). This process appears to affect many neuronal populations, not only in young and old adults, but also in children diagnosed with autism. A metabolic shift with A β accumulation in neurons in these brain areas that are involved in the expression of emotions, stereotypic behaviors, and social deficits, such as the amygdala, hippocampus, some striatal subdivisions, and cerebellum, may contribute to cellular dysfunction and the clinical expression of autism.

1.6 CLINICOPATHOLOGICAL CORRELATIONS

Studies of clinicopathological correlations cover several domains of functional deficits in people with autism, including (a) speech, language, and verbal and nonverbal communication, (b) social deficits and face perception, (c) sensorimotor deficits, and (d) cognitive deficits.

1.6.1 SPEECH, LANGUAGE, AND VERBAL AND NONVERBAL COMMUNICATION

Expressive language function of individuals with autism ranges from complete mutism to verbal fluency. Verbal abilities are often accompanied by errors in word meaning (semantics) or language and communicative deficits in social context (social pragmatics) (Rapin, 1996; Stone et al., 1997; Wetherby et al., 1998). Studies of the language-related

neocortex, including Wernicke's area (BA 22, speech recognition), Broca's area (BA 44, speech production) and the gyrus angularis (BA 39, reading) of 7- to 44-year-old autistic and 8- to 56-year-old control individuals, revealed reduced numerical density of neurons by 38% in area 22, and by 24% in area 39 in autistic subjects, as well as an increased numerical density of lipofuscin-containing neurons by 50% in BA 22, and 44% in BA 44. These neuronal changes were paralleled by an increase of numerical density of glial cells in all three examined regions. Lopez-Hurtado and Prieto (2008) hypothesized that structural alteration in one or more of these cortical areas may contribute to the communication impairment observed in autism.

1.6.2 FACE PERCEPTION

All subjects with ASDs have disturbance of social behavior, including abnormalities in social reciprocity and difficulties in use of eye contact, facial expression, and social motivation. Social functioning includes eye contact, processing of faces, identification of individuals, and monitoring of face expression (Baron-Cohen et al., 1994). Patients with autism reveal deficits in face-processing (Grelotti et al., 2001), perception (Schultz, 2005), and recognition (Joseph and Tanaka, 2003).

The face-processing network includes the visual cortex (BA17), which projects via the inferior occipital gyrus to the fusiform gyrus. Fibers from the fusiform gyrus project to the amygdala, and inferior frontal gyrus and orbital cortex (Fairhall and Ishai, 2007; van Kooten et al., 2008). Functional magnetic resonance imaging (fMRI) identified the fusiform gyrus and other cortical regions as supporting face-processing in control subjects, and hypoactivity of the fusiform gyrus in autistic patients (Bolte et al., 2006; Kanwisher et al., 1999; Pierce et al., 2004). Hypoactivation of the fusiform gyrus is believed to be associated with the failure to make direct eye contact in autism (Dalton et al., 2005). Results of imaging-based fusiform volume estimation are inconsistent. Increased (Waiter et al., 2004) and unchanged (Pierce et al., 2001) volume in both hemispheres and increased fusiform gyrus in the left hemisphere (Herbert et al., 2002) were reported. Morphometric studies of the brain of 7 autistic and 10 control subjects revealed a reduced number of neurons in layers III, V, and VI, and reduced volume of neuronal soma in layers V and VI in the fusiform gyrus. No alterations in Brodman area 17 in these autistic individuals suggest that the input from the visual cortex to the fusiform gyrus is intact. These results indicate the underdevelopment of connections in the fusiform gyrus that may contribute to abnormal face perception in autism (van Kooten et al., 2008).

Bailey et al. (1998) noted abnormalities in cytoarchitectonic organization and neuronal density in the superior frontal cortex and superior temporal gyrus in autism. Neurons in the superior temporal sulcus are sensitive to the angle of gaze (Perrett et al., 1985). Neurons that are attuned to particular facial expressions were found in the inferior and superior temporal lobes (Hasselmo et al., 1989). Cortical areas responsive to faces, facial expressions, and angle of gaze send direct projections to the amygdala (Stefanacci and Amaral, 2000). Pathological changes in the amygdala may play a central role in the dysfunction seen in autism, including disturbed components of social cognition such as attention to and interpretation of facial expressions. fMRI studies show that judging from the expression of another

person's eyes what the other person might be thinking or feeling is associated with activation in the superior temporal gyrus, frontal cortex, and amygdala, whereas in subjects with autism, activation appears in the temporal and frontal cortex but not in the amygdala (Baron-Cohen et al., 1999).

1.6.3 SOCIAL ATTACHMENT—THE ROLE OF THE HYPOTHALAMUS IN BEHAVIORAL DEFICITS

Experimental studies revealed that the hypothalamic nucleus paraventricularis (NPV) and the nucleus supraopticus (NSO), producing oxytocin (OT) and vasopressin (VAS), regulate emotional responses, social attachment, cognitive functions, sleep, and appetite (Barden, 2004; Ehler et al., 2001; Manaye et al., 2005). OT and VAS are relayed from the human brain into the bloodstream via the posterior pituitary. The presence of receptors for both peptides throughout the forebrain, limbic system, thalamus, brain stem, and spinal cord (Raggenbass, 2001) indicates that hypothalamic neuropeptides modulate the function of many brain regions. Developmental changes in the distribution and expression of receptors suggest that the hypothalamic peptides play a significant role both in brain development and function (Shapiro and Insel, 1989). OT is required for the development of social memory. In OT knockout mice, the loss of social memory could be rescued by OT treatment (Ferguson et al., 2000). VAS is necessary for the formation of social memory and OT for retention of newly formed social memories (Popik and Van Ree, 1992; Popik et al., 1992). OT facilitates the learning of social interactions and the formation of associations that are specifically related to the mother (Nelson and Panksepp, 1996).

The initial product of oxytocin mRNA is a polypeptide containing both nanopeptide OT and neurophysin I, separated by tripeptide glycine-lysine-arginine. The result of enzymatic cleavage are intermediate forms containing 10, 11, or 12 amino acids, collectively referred to as carboxy-extended forms of OT (OT-X), and oxytocin (Gabreels et al., 1998; Gainer et al., 1995; Mitchell et al., 1998; Rao et al., 1992). In 5.8- to 11.5-year-old autistic individuals, reduced plasma OT level, deficits in OT prohormone processing (increase in OT-X), and an increase in the ratio of C-terminal extended forms to OT were found. In control children, nearly all OT-X is metabolized to OT, whereas in autistic children, the immature OT forms serve as the primary circulating molecule in the absence of or in compensation for OT (Green et al., 2001). However, experimental studies show that OT-X is not an effective agonist at OT-sensitive sites (Mitchell et al., 1998). Deficient conversion of OT-X to OT in autism could be the result of alterations in the level of prohormone convertases associated with genetic defects (Cook, 1998; Szatmari et al., 1998). The identification of four single nucleotide polymorphisms located within the OT receptor gene of 195 Chinese autistic subjects indicates that abnormal modulation of the OT receptor results in autism (Wu et al., 2005). OT and VAS are known to play a role in repetitive behaviors. Patients with ASDs show a significant reduction in repetitive behaviors following OT infusion (Hollander et al., 2003). In about 60% of subjects with autism, abnormal sleep patterns are observed. VAS is involved in the control of circadian rhythmicity (Swaab, 1997). VAS enhances aggressiveness, anxiety, stress levels, and the consolidation of fear memory (Bielsky et al., 2004; Griebel et al., 2002; Landgraf and Neumann, 2004). OT decreases anxiety

and stress; facilitates social encounters, maternal care, and the extinction of conditioned avoidance behavior (Bale et al., 2001; Champagne et al., 2001; Windle et al., 1997); reduces activation of the amygdala and modulates fear processing (Kirsch et al., 2005). The presence of abnormal levels of hypothalamic neuropeptides in patients with autism provides strong evidence that an abnormality in OT, VAS and other hypothalamic neuropeptides may have a significant contribution to the behavioral features of autism. However, the morphology and biochemistry of the hypothalamus of autistic subjects remains unknown. The only study of the hypothalamic mammillary body of a 26-year-old autistic man revealed that the cell-packing density was increased by 66% (Bauman and Kemper, 1985).

1.6.4 SENSORIMOTOR DEFICITS, AND REPETITIVE AND STEREOTYPED BEHAVIORS

In individuals with autism, impairments of gross and fine motor function recognized as hypotonia, limbic apraxia, and motor stereotypes are common findings and are more severe in subjects with lower IQ (Rogers et al., 1996). Hand mannerisms, body rocking, or unusual posturing are reported in 37% to 95% of individuals (Lord, 1995; Rapin, 1996; Rogers et al., 1996). In 42% to 88% of subjects with autism, aberrant sensory processing results in a preoccupation with sensory features of objects, over- or underresponsiveness to environmental stimuli or paradoxical responses to sensory stimuli (Kientz and Dunn, 1997). Sensorimotor deficits may be associated with pathological changes in both the nigrostriatal system (basal ganglia) and the cerebellum (Bailey et al., 1998; Kemper and Bauman, 1998; Ritvo et al., 1986; Saitoh and Courchesne, 1998; Sears et al., 1999). Cerebellar abnormality with a deficit/loss of Purkinje cells (Bailey et al., 1998; Kemper and Bauman, 1993, 1998; Ritvo et al., 1986) has been a common finding. Individuals with autism have been classified as affected by cerebellar hyper- or hypoplasia (Saitoh and Courchesne, 1998). A reduced number of Purkinje cells without significant glial activation and a reduced size of Purkinje cells were noticed in the majority of cerebellar reports (Bailey et al., 1998; Fehlow et al., 1993; Kemper and Bauman, 1993; Lee et al., 2002; Ritvo et al., 1986) in 21 of 29 examined cases (Palmen et al., 2004).

Results of evaluation of the size of the cerebellum using MRI are inconsistent. In several MRI studies, smaller cerebellar hemispheres (Gaffney et al., 1987; Murakami et al., 1989) and vermis (Ciesielski et al., 1997; Courchesne et al., 1988; Hashimoto et al., 1995) were reported. In a study by Piven et al. (1997a), the total cerebellar volume was found to be greater in subjects with autism than in the control group, and the increase was proportional to the increased total brain volume. In the cerebellum, boys with autism had less gray matter, a smaller ratio of gray to white matter, and smaller lobules VI and VII than did controls. Despite the inconsistency of reports characterizing topographic autism-associated vermian hypoplasia (Hashimoto et al., 1993; Kaufmann et al., 2003; Levitt et al., 1999; Piven et al., 1997a; Schaefer et al., 1996), several reports show associations between the size of the vermis and deficits in attention-orienting (Harris et al., 1999; Townsend et al., 1999), stereotypic behavior, and reduced exploration in autism (Pierce and Courchesne, 2001).

The reduced size of the pons, midbrain, and medulla in autism reported by Hashimoto et al. (1992, 1993, 1995) was not confirmed in other studies (Hsu et al., 1991; Piven et al., 1992).

Changes in neurons in the deep cerebellar nuclei were noticed by some authors (Kemper and Bauman, 1998) but not by others (Bailey et al., 1998). Structural MRI shows variable patterns of changes. Volumetry of the cerebellum may show no change, hypoplasia, or hyperplasia. Courchesne et al. (1988) reported selective hypertrophy of lobules VI and VII, but these results were not confirmed in other subjects. In part, the pattern may correspond to the functional status of subjects. In highly functioning subjects with autism, hypoplasia of the cerebellum has not been detected (Holtum et al., 1992).

A decrease in the number of GABAergic Purkinje cells is considered the most consistent neuropathological finding in autism, as it was detected in at least 50% of examined cases (Arin et al., 1991; Bailey et al., 1998). Recent studies indicate that preserved Purkinje cells reveal a 40% decrease in the expression of glutamic acid decarboxylase 67 (GAD67) mRNA in autistic subjects relative to control patients (Yip et al., 2007). Moreover, in autism, the basket cells likely provide increased GABAergic feed-forward inhibition to Purkinje cells. The result may include disruption in the timing of Purkinje cell firing and altered inhibition of the cerebellar nuclei, which could directly affect cerebello-cortical output, leading to changes in motor behavior and cognition (Yip et al., 2008).

Repetitive and stereotyped behaviors defined as recurring, nonfunctional activities, or interests that occur regularly and interfere with daily functioning are a defining signs of autism. These behaviors include lower-order repetitive motor behavior, intense circumscribed patterns of interests, and higher-order rituals and compulsions (Gabriels et al., 2005). Several studies implicated the role of basal ganglia and frontostriatal circuitry in the pathophysiology of autism, especially in repetitive and stereotyped behaviors. Increased volume of the basal ganglia was reported in several MRI studies (Herbert et al., 2003; Hollander et al., 2005; Langen et al., 2007; Sears et al., 1999). Sears et al. (1999) and Hollander et al. (2005) observed a positive correlation between caudate volumes and repetitive behavior scores. A significant increase in caudate nucleus volume, disproportional to brain volume, was detected in MRI studies in two independent samples of medication-naïve subjects with autism (21 high-functioning children and adolescents, and 21 typically developing subjects; 21 high-functioning adolescents and young adults, and 21 healthy subjects) (Langen et al., 2007). Our studies showing a significantly smaller size of neurons in the caudate, putamen, and nucleus accumbens, especially in the brains of children 4–7 years of age suggest a developmental delay in the growth of neurons in the basal ganglia of autistic subjects, which may contribute to basal ganglia dysfunction (Wegiel et al., 2008). MRI and postmortem morphometric studies support the hypothesis that developmental abnormalities in frontostriatal circuitry contribute to repetitive and stereotyped behaviors, which are one of three defining symptoms of autism.

1.6.5 COGNITIVE DEFICITS

Many individuals with autism demonstrate a particular pattern on intellectual tests that is characteristic of autism. Performance IQ is usually higher than verbal IQ, and block design is the highest subtest, whereas comprehension is usually the lowest (Siegel et al., 1996). Individuals with autism have poorer adaptive function than would be predicted by IQ alone (Volkmar et al., 1993).

Cognitive deficits might be related in part to the memory system and limbic region abnormalities. Reduced volume of both the hippocampal formation and amygdala were noticed in subjects examined by Aylward et al. (1999), but not in populations examined by other researchers (Piven et al., 1998). Neurons in the hippocampus have reduced complexity of dendritic arbors. They are smaller and more densely packed in various portions of the hippocampal formation, entorhinal cortex, medial nuclei of the amygdala, medial septal nucleus, mammillary nuclei, and anterior cingulate gyrus (Bauman and Kemper, 1985; Kemper and Bauman, 1993). Haznedar et al. (1997) observed reduced volume of the anterior cingulate gyrus and decreased positron emission tomography (PET) activity in subjects with autism. Because the cerebellum is involved in a variety of cognitive and affective processes, abnormalities of both the limbic system and the cerebellum may be linked to the core social and communicative deficits in autism.

The caudate nucleus is an integral component of the frontostriatal network involved in cognitive functions (Chow and Cummings, 1999; Voelbel et al., 2006), including learning (Poldrack et al., 1999), short- and long-term memory (Fuh and Wang, 1995), and planning and problem-solving (Mendez et al., 1989; Schmidtke et al., 2002). The increased volume of the caudate observed in autistic children may be indicative of impaired neuronal pruning, contributing to a decrease in executive function (Voelbel et al., 2006).

1.6.6 EPILEPSY-ASSOCIATED PATHOLOGY

The 1% prevalence of epilepsy in the general population increases to 8% in DS, 10% in AD (Menendez, 2005; Risse et al., 1990; Velez and Selwa, 2003), and 33% in autism (Tuchman and Rapin, 2002). The interpretation of developmental changes in autism has been challenged by the need to differentiate among lesions that are not associated with epilepsy, that cause epilepsy, and that are produced by epilepsy (Sutula and Pitkanen, 2001). Recent studies support the hypothesis that epilepsy induces brain alterations that contribute to changes in circuitry, which potentiates the seizure-genic focus (Armstrong, 2005).

Studies of nonautistic subjects indicate that epilepsy-associated pathology includes patchy or laminar neuronal loss and gliosis in the cerebral cortex in one or both hemispheres. In temporal epilepsy, abnormalities were reported in 75% of the specimens examined, and hippocampal sclerosis was found in 50% (Bruton, 1988). Loss of hippocampal neurons correlates with the frequency of tonic-clonic seizures and the total duration of epilepsy (Dam, 1980; Tasch et al., 1999). Loss is accentuated in the CA4 sector and is observed in the granule cell layer in the dentate gyrus. Dispersion of dentate gyrus granular neurons might be a result of seizure-related, disturbed migration of neurons (Bengzon et al., 1997), or epilepsy-enhanced neurogenesis (Ericksson et al., 1998). Ammon horn sclerosis is a progressive lesion that can be induced and propagated by seizures (Armstrong et al., 2005).

In nearly all cases with hippocampal pathology, changes are also observed in other brain regions. In about 25%, the amygdala, thalamus and mammillary body are affected with neuronal loss. More severe neuronal loss and gliosis in the hippocampus is paralleled by severe neuronal loss and gliosis in the lateral nucleus in the amygdala (Bruton, 1988; Hudson et al., 1993; Thom et al., 1999). Ectopias

AQ4

with more than 8 neurons per 2 sq. mm of white matter occurred in 43% of epileptic patients but in none of the controls (Hardiman et al., 1988). In 45% of severely affected epileptics, significant neuronal loss and astrocytosis spreading out into the overlying molecular layer is observed in the cerebellar cortex. The severity of the cerebellar damage may range from gross atrophy of most or many folia to the restricted neuronal loss in some folia, especially at their basal portion (Gessaga and Urich, 1985).

Central apnea, asphyxia, and pulmonary edema occurring during a seizure (Nashef et al., 1996) as well as life-threatening cardiac arrhythmias during seizures (Earnest et al., 1992; Jallon, 1997; Nashef et al., 1996; Reeves et al., 1996; Saussu et al., 1998) have been suggested as possible causes of sudden unexpected death in epilepsy (Thom et al., 1999).

Enhanced electric activity of neurons and/or increased cell synaptic transmission with enhanced vesicle exocytosis, both in normal and in disease-affected brains are a common cause of modifications of APP processing and A β levels. Epilepsy is associated with an elevation of APP expression (Sheng et al., 1994) and occurs in 10 of 11 examined subjects with diffuse nonfibrillar A β plaque formation (mean age 47.9 ± 8.8 years of age) (Mackenzie and Miller, 1994; Mackenzie et al., 1996).

1.7 MECHANISMS AFFECTING BRAIN DEVELOPMENT

1.7.1 BDNF AND NEUROTROPHINS IN AUTISM

The neurotrophins, a related family of growth factors, including nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF), and neurotrophins (NT) NT-3 and NT-4/5, have a major role in the survival, growth, and differentiation of neurons (Conner et al., 1997). During typical brain development, only neurons making the appropriate connections survive and form synapses, whereas neurons that fail to obtain adequate neurotrophins die (Oppenheim, 1991). BDNF is broadly distributed throughout the human central nervous system (CNS) and provides neurotrophic support for many neuronal populations in the cortex, amygdala, hippocampus, and striatum (Murer et al., 2001; Schmidt-Kastner et al., 1996; Tapia-Arancibia et al., 2004). The hypothalamus is the brain structure that contains the highest BDNF protein levels (Katoh-Semba et al., 1997; Nawa et al., 1995; Yan et al., 1997) and BDNF mRNA (Castren et al., 1995; Kawamoto et al., 1996; Yan et al., 1997). In the cerebellum, immunoreactivity was observed in Purkinje cells and the olivary complex of the nuclei (Kawamoto et al., 1996; Murer et al., 2001).

In the basal forebrain of autistic individuals, the level of BDNF was three times higher than in controls (Perry et al., 2001). Miyazaki et al. (2004) observed a higher level of BDNF in the blood samples of young children with autism than in adult control subjects. The mean BDNF levels in sera of children diagnosed with autism and childhood disintegrative disorder were about four times higher than in control children (Connolly et al., 2006). Children with autism and childhood disintegrative disorder have both elevated BDNF levels and levels of autoantibodies against BDNF. Serum BDNF has been shown to be increased after seizures (Binder et al., 2001; Chavko et al., 2002).

1.7.2 BRAIN STEM AND THE ROLE OF SEROTONIN IN BRAIN DEVELOPMENT AND CLINICAL FEATURES OF AUTISM

Because 5-hydroxytryptamine (5-HT; serotonin) serves as both a neurotransmitter and an important developmental signal in the brain, dysregulation of the 5-HT system during development may be responsible for many of the abnormalities seen in autism (Whitaker-Azmitia, 2005). In fact, all known chemical inducers of autism including cocaine, thalidomide, valproate, and alcohol modulate 5-HT levels in the brain (Harris et al., 1995; Kramer et al., 1994; Narita et al., 2002; Rathbun and Druse, 1985; Stromland et al., 1994; Williams et al., 2001). A high proportion of children with autism exhibit elevated blood 5-HT levels (hyperserotonemia) and specific alterations in 5-HT biosynthesis. The severity of hyperserotonemia is correlated with the severity of autistic behaviors (Chandana et al., 2005; Chugani et al., 1999; Kuperman et al., 1987). A causal role for serotonergic abnormalities in the etiology of autism is also suggested by studies indicating autism-specific genetic polymorphisms in 5-HT metabolizing enzyme, transporter, or receptor genes (Cohen et al., 2003; Sutcliffe et al., 2005). Also, gender-specific differences in serotonergic regulation during development (Chandana et al., 2005; Chugani et al., 1999), combined with a 52% higher rate of 5-HT biosynthesis in the male than female brain (Nishizawa et al., 1997), and the increased susceptibility of males to early insults imposed by elevated levels of 5-HT (Johns et al., 2002), may contribute to the fourfold higher propensity of males to develop autism compared to females.

As a result of the regulatory role of serotonin affecting the size of neurons, the size of dendritic tree and the number of synapses in innervated cortical and subcortical structures and cerebellum, developmental abnormalities in the serotonergic system may contribute to structural and functional changes in target brain regions and structures. Virtually all regions of the brain receive serotonergic afferents from raphe system neurons. The rostral raphe nuclei form ascending pathways of axons mainly to the forebrain. The caudal raphe system innervates the lower brain stem and the spinal cord (Aitken and Törk, 1988; Lidov and Molliver, 1982). The functions of serotonin are mediated by 14 subtypes of 5-HT receptors in the nervous system (Hoyer et al., 1994; Martin and Humphrey, 1994; Saudou and Hen, 1994a,b). The serotonin_{2A} (5-HT_{2A}) receptor is known to be one of the major subtypes and is associated with psychological and mental events (Roth, 1994). The 5-HT_{2A} receptor plays a role in facilitating the formation and maintenance of synapses (Niitsu et al., 1995). Staining for 5-HT_{2A} shows the entire somata and dendritic tree of Purkinje cells in a rat cerebellum (Maeshima et al., 1998). In vitro studies have shown that 5-HT inhibits the growth and arborization of Purkinje cell dendrites through 5-HT_{2A} receptors and stimulates them through the 5-HT_{1A} receptor (Kondoh et al., 2004). 5-HT promotes the formation of synapses in developing and mature brain and spinal cord (Chen et al., 1997; Niitsu et al., 1995; Okado et al., 1993), and this process is mediated by the 5-HT_{2A} receptor in the spinal cord (Niitsu et al., 1995). Biochemical studies support the hypothesis that developmental defects of the raphe nuclei may make a major contribution to the structural and functional defects of cortical and subcortical structures. However, raphe nuclei have not yet been examined in autistic subjects.

1.8 CLOSING REMARKS

The detected brain structure-specific patterns of structural aberrations in a majority of examined anatomic subdivisions in autism brain may contribute to deficits in expression of emotions, processing of social stimuli, learning of social behaviors, verbal and nonverbal communication, and stereotypic behaviors. Pathological acceleration of brain growth and immaturity of neurons and neuronal networks in early childhood indicate that (a) a significant portion of structural/functional defects starts in early infancy and (b) causative factors dysregulate the mechanisms controlling brain/neuron development. The deceleration of brain growth in the second year of life and the increase of neuronal size in late childhood/adulthood suggests delayed activation of correcting mechanisms. However, the delayed correction of brain and neuronal size does not result in functional recovery. Analysis of the detected pattern of abnormal brain development in autism indicates that early diagnosis and early treatment may prevent or reduce developmental delay, reduce/eliminate secondary structural and functional changes, and improve clinical status throughout the life span.

ACKNOWLEDGMENTS

This study was supported in part by funds from the New York State Office of Mental Retardation and Developmental Disabilities and grants from Autism Speaks and the Department of Defense Autism Spectrum Disorders Research Program (AS073234).

REFERENCES

- Aitken, A. R. and I. Tork (1988). Early development of serotonin-containing neurons and pathways as seen in wholemount preparations of the fetal rat brain. *J. Comp. Neurol.* 274:32–47.
- Akyol, O., H. Herken, E. Uz, E. Fadillioglu, S. Unal, S. Sogut, H. Ozyurt, and H. A. Savas (2002). The indices of endogenous oxidative and antioxidative processes in plasma from schizophrenic patients: The possible role of oxidant/antioxidant imbalance. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 26:995–1005.
- American Psychiatric Association (2000). *Diagnostic and Statistical Manual of Mental Disorders DSM-IV-TR*. American Psychiatric Association, Washington, DC.
- AQ5 Arin, D. M., M. L. Bauman, and T. L. Kemper (1991). The distribution of Purkinje cell loss in the cerebellum in autism. 307.
- Armstrong, D. D. (2005). Epilepsy-induced microarchitectural changes in the brain. *Pediatr. Dev. Pathol.* 8:607–614.
- Aylward, E. H., N. J. Minshew, K. Field, B. F. Sparks, and N. Singh (2002). Effects of age on brain volume and head circumference in autism. *Neurology* 59:175–183.
- Aylward, E. H., N. J. Minshew, G. Goldstein, N. A. Honeycutt, A. M. Augustine, K. O. Yates, P. E. Barta, and G. D. Pearlson (1999). MRI volumes of amygdala and hippocampus in non-mentally retarded autistic adolescents and adults. *Neurology* 53:2145–2150.
- Bailey, A., C. A. Le, I. Gottesman, P. Bolton, E. Simonoff, E. Yuzda, and M. Rutter (1995). Autism as a strongly genetic disorder: Evidence from a British twin study. *Psychol. Med.* 25:63–77.
- Bailey, A., P. Luthert, P. Bolton, C. A. Le, M. Rutter, and B. Harding (1993). Autism and megalencephaly. *Lancet* 341:1225–1226.

- Bailey, A., P. Luthert, A. Dean, B. Harding, I. Janota, M. Montgomery, M. Rutter, and P. Lantos (1998). A clinicopathological study of autism. *Brain* 121 (Pt 5):889–905.
- Bailey, A. R., B. N. Giunta, D. Obregon, W. V. Nikolic, J. Tian, C. D. Sanberg, D. T. Sutton, and J. Tan (2008). Peripheral biomarkers in autism: Secreted amyloid precursor protein- α as a probable key player in early diagnosis. *Int. J. Clin. Exp. Med.* 1:338–344.
- Bale, T. L., A. M. Davis, A. P. Auger, D. M. Dorsa, and M. M. McCarthy (2001). CNS region-specific oxytocin receptor expression: Importance in regulation of anxiety and sex behavior. *J. Neurosci.* 21:2546–2552.
- Barden, N. (2004). Implication of the hypothalamic-pituitary-adrenal axis in the physiopathology of depression. *J. Psychiatry Neurosci.* 29:185–193.
- Barkovich, A. J., R. I. Kuzniecky, G. D. Jackson, R. Guerrini, and W. B. Dobyns (2005). A developmental and genetic classification for malformations of cortical development. *Neurology* 65:1873–1887.
- Baron-Cohen, S. (2004). The cognitive neuroscience of autism. *J. Neurol. Neurosurg. Psychiatry* 75:945–948.
- Baron-Cohen, S., H. Ring, J. Moriarty, B. Schmitz, D. Costa, and P. Ell (1994). Recognition of mental state terms. Clinical findings in children with autism and a functional neuroimaging study of normal adults. *Br. J. Psychiatry* 165:640–649.
- Baron-Cohen, S., H. A. Ring, S. Wheelwright, E. T. Bullmore, M. J. Brammer, A. Simmons, and S. C. Williams (1999). Social intelligence in the normal and autistic brain: An fMRI study. *Eur. J. Neurosci.* 11:1891–1898.
- Bauman, M. L., P. A. Filipek, and T. L. Kemper (1997). Early infantile autism. *Int. Rev. Neurobiol.* 41:367–386.
- Bauman, M. L. and T. L. Kemper (1985). Histoanatomic observations of the brain in early infantile autism. *Neurology* 35:866–867.
- Bauman, M. L., and T. L. Kemper (1996). Observation on the Purkinje cells in the cerebellar vermis in autism. *J. Neuropath. Exp. Neurol.* 55:613.
- Bengzon, J., Z. Kokaia, E. Elmer, A. Nanobashvili, M. Kokaia, and O. Lindvall (1997). Apoptosis and proliferation of dentate gyrus neurons after single and intermittent limbic seizures. *Proc. Natl. Acad. Sci. USA* 94:10432–10437.
- Bielsky, I. F., S. B. Hu, K. L. Szegda, H. Westphal, and L. J. Young (2004). Profound impairment in social recognition and reduction in anxiety-like behavior in vasopressin V1a receptor knockout mice. *Neuropsychopharmacology* 29:483–493.
- Binder, D. K., S. D. Croll, C. M. Gall, and H. E. Scharfman (2001). BDNF and epilepsy: Too much of a good thing? *Trends Neurosci.* 24:47–53.
- Boddaert N., M. Zilbovicius, A. Philipe, L. Robel, M. Bourgeois, C. Barthelemy, D. Seidenwurm, I. Meresse, L. Laurier, I. Desguerre, N. Bahi-Buisson, F. Brunelle, A. Munnich, Y. Samson, M-C. Mouren, and N. Chabane (2009). MRI findings in 77 children with non-syndromic autism. *PLOS one* (www.plosone.com) 4:4415e.
- Bolte, S., D. Hubl, S. Feineis-Matthews, D. Prvulovic, T. Dierks, and F. Poustka (2006). Facial affect recognition training in autism: Can we animate the fusiform gyrus? *Behav. Neurosci.* 120:211–216.
- Bolton, P., J. Powell, M. Rutter, V. Buckle, J. R. Yates, Y. Ishikawa-Brush, and A. P. Monaco (1995). Autism, mental retardation, multiple exostoses and short stature in a female with 46,X,t(X;8)(p22.13;q22.1). *Psychiatr. Genet.* 5:51–55.
- Brown, W. T., T. Wisniewski, I. L. Cohen, E. London, M. Flory, H. Imaki, I. Kuchna, J. Wegiel, S. Y. Ma, K. Nowicki, J. Wang, and J. Wegiel (2008). Neuropathologic changes in chromosome 15 duplication and autism. In 7th Annual International Meeting for Autism Research (IMFAR).
- Brunk, U. T., C. B. Jones, and R. S. Sohal (1992). A novel hypothesis of lipofuscinogenesis and cellular aging based on interactions between oxidative stress and autophagocytosis. *Mutat. Res.* 275:395–403.

- Brunk, U. T. and A. Terman (2002a). Lipofuscin: Mechanisms of age-related accumulation and influence on cell function. *Free Radic. Biol. Med.* 33:611–619.
- Brunk, U. T. and A. Terman (2002b). The mitochondrial–lysosomal axis theory of aging: Accumulation of damaged mitochondria as a result of imperfect autophagocytosis. *Eur. J Biochem.* 269:1996–2002.
- Bruton, C. J. (1988). *The Neuropathology of Temporal Lobe Epilepsy*. In Maudsley Monographs. Oxford University Press, New York.
- AQ6 Buitelaar, J. K. and S. H. Willemsen-Swinkels (2000). Autism: Current theories regarding its pathogenesis and implications for rational pharmacotherapy. *Paediatr. Drugs* 2:67–81.
- Butler, M. G., M. J. Dasouki, X. P. Zhou, Z. Talebizadeh, M. Brown, T. N. Takahashi, J. H. Miles, C. H. Wang, R. Stratton, R. Pilarski, and C. Eng (2005). Subset of individuals with autism spectrum disorders and extreme macrocephaly associated with germline PTEN tumour suppressor gene mutations. *J. Med. Genet.* 42:318–321.
- Buxhoeveden, D. P. and M. F. Casanova (2002). The minicolumn and evolution of the brain. *Brain Behav. Evol.* 60:125–151.
- Carper, R. A. and E. Courchesne (2005). Localized enlargement of the frontal cortex in early autism. *Biol. Psychiatry* 57:126–133.
- Carper, R. A., P. Moses, Z. D. Tigue, and E. Courchesne (2002). Cerebral lobes in autism: Early hyperplasia and abnormal age effects. *Neuroimage* 16:1038–1051.
- Casanova, M. F., D. P. Buxhoeveden, A. E. Switala, and E. Roy (2002). Minicolumnar pathology in autism. *Neurology* 58:428–432.
- Casanova, M. F., K. van, I. A. E. Switala, H. van Engeland, H. Heinsen, H. W. Steinbusch, P. R. Hof, J. Trippe, J. Stone, and C. Schmitz (2006). Minicolumnar abnormalities in autism. *Acta Neuropathol.* 112:287–303.
- Castren, E., H. Thoenen, and D. Lindholm (1995). Brain-derived neurotrophic factor messenger RNA is expressed in the septum, hypothalamus and in adrenergic brain stem nuclei of adult rat brain and is increased by osmotic stimulation in the paraventricular nucleus. *Neuroscience* 64:71–80.
- Champagne, F., J. Diorio, S. Sharma, and M. J. Meaney (2001). Naturally occurring variations in maternal behavior in the rat are associated with differences in estrogen-inducible central oxytocin receptors. *Proc. Natl. Acad. Sci. USA* 98:12736–12741.
- Chandana, S. R., M. E. Behen, C. Juhasz, O. Muzik, R. D. Rothermel, T. J. Mangner, P. K. Chakraborty, H. T. Chugani, and D. C. Chugani (2005). Significance of abnormalities in developmental trajectory and asymmetry of cortical serotonin synthesis in autism. *Int. J. Dev. Neurosci.* 23:171–182.
- Chauhan, A., V. Chauhan, W. T. Brown, and I. Cohen (2004). Oxidative stress in autism: Increased lipid peroxidation and reduced serum levels of ceruloplasmin and transferrin—the antioxidant proteins. *Life Sci.* 75:2539–2549.
- Chauhan, A. and V. Chauhan (2006). Oxidative stress in autism. *Pathophysiology* 13:171–181.
- Chauhan, A., B. Muthaiyah, M. M. Essa, T. W. Brown, J. Wegiel, and V. Chauhan (2009). Increased lipid peroxidation in cerebellum and temporal cortex in autism. International Meeting for Autism Research, Chicago, IL, May 8, 2009.
- Chavko, M., N. S. Nadi, and D. O. Keyser (2002). Activation of BDNF mRNA and protein after seizures in hyperbaric oxygen: Implications for sensitization to seizures in re-exposures. *Neurochem. Res.* 27:1649–1653.
- Chen, L., K. Hamaguchi, S. Hamada, and N. Okado (1997). Regional differences of serotonin-mediated synaptic plasticity in the chicken spinal cord with development and aging. *J. Neural Transplant. Plast.* 6:41–48.
- Chow, T. W. and J. L. Cummings (1999). Frontal-subcortical circuits. In B. L. Miller and J. L. Cummings, (eds.), *The Human Frontal Lobes: Functions and Disorders*. Guilford Press, New York, pp. 3–26.

- Chugani, D. C., O. Muzik, M. Behen, R. Rothermel, J. J. Janisse, J. Lee, and H. T. Chugani (1999). Developmental changes in brain serotonin synthesis capacity in autistic and nonautistic children. *Ann. Neurol.* 45:287–295.
- Ciesielski, K. T., R. J. Harris, B. L. Hart, and H. F. Pabst (1997). Cerebellar hypoplasia and frontal lobe cognitive deficits in disorders of early childhood. *Neuropsychologia* 35:643–655.
- Cohen, I. L. (2007). A neural network model of autism: Implications for theory and treatment. In D. Mareschal, S. Sirois, G. Westerman, and M. H. Johnson (eds.), *Neuroconstructivism*. Oxford University Press, Oxford.
- Cohen, I. L., X. Liu, C. Schutz, B. N. White, E. C. Jenkins, W. T. Brown, and J. J. Holden (2003). Association of autism severity with a monoamine oxidase A functional polymorphism. *Clin. Genet.* 64:190–197.
- Conner, J. M., J. C. Lauterborn, Q. Yan, C. M. Gall, and S. Varon (1997). Distribution of brain-derived neurotrophic factor (BDNF) protein and mRNA in the normal adult rat CNS: Evidence for anterograde axonal transport. *J. Neurosci.* 17:2295–2313.
- Connolly, A. M., M. Chez, E. M. Streif, R. M. Keeling, P. T. Golumbek, J. M. Kwon, J. J. Riviello, R. G. Robinson, R. J. Neuman, and R. M. Deuel (2006). Brain-derived neurotrophic factor and autoantibodies to neural antigens in sera of children with autistic spectrum disorders, Landau-Kleffner syndrome, and epilepsy. *Biol. Psychiatry* 59:354–363.
- Cook, E. H. Jr. (1998). Genetics of autism. *Mental Retardation Dev. Disabil. Res. Rev.* 4:113–120.
- Coon, H., D. Dunn, J. Lainhart, J. Miller, C. Hamil, A. Battaglia, R. Tancredi, M. F. Leppert, R. Weiss, and W. McMahon (2005). Possible association between autism and variants in the brain-expressed tryptophan hydroxylase gene (TPH2). *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 135B:42–46.
- AQ7 Courchesne, E. (2001). Unusual brain growth patterns in early life in patients with autistic disorder: An MRI study. *Neurology* 57:245–254.
- Courchesne, E., R. Carper, and N. Akshoomoff (2003). Evidence of brain overgrowth in the first year of life in autism. *JAMA* 290:337–344.
- Courchesne, E., J. R. Hesselink, T. L. Jernigan, and R. Yeung-Courchesne (1987). Abnormal neuroanatomy in a nonretarded person with autism. Unusual findings with magnetic resonance imaging. *Arch. Neurol.* 44:335–341.
- Courchesne, E., C. M. Karns, H. R. Davis, R. Ziccardi, R. A. Carper, Z. D. Tigue, H. J. Chisum, P. Moses, K. Pierce, C. Lord, A. J. Lincoln, S. Pizzo, L. Schreibman, R. H. Haas, N. A. Akshoomoff, and R. Y. Courchesne, E., R. Yeung-Courchesne, G. A. Press, J. R. Hesselink, and T. L. Jernigan (1988). Hypoplasia of cerebellar vermal lobules VI and VII in autism. *N. Engl. J. Med.* 318:1349–1354.
- Courchesne, E. and K. Pierce (2005a). Why the frontal cortex in autism might be talking only to itself: Local over-connectivity but long-distance disconnection. *Curr. Opin. Neurobiol.* 15:225–230.
- Courchesne, E. and K. Pierce (2005b). Brain overgrowth in autism during a critical time in development: Implications for frontal pyramidal neuron and interneuron development and connectivity. *Int. J. Dev. Neurosci.* 23:153–170.
- Dalton, K. M., B. M. Naciewicz, T. Johnstone, H. S. Schaefer, M. A. Gernsbacher, H. H. Goldsmith, A. L. Alexander, and R. J. Davidson (2005). Gaze fixation and the neural circuitry of face processing in autism. *Nat. Neurosci.* 8:519–526.
- Dam, A. M. (1980). Epilepsy and neuron loss in the hippocampus. *Epilepsia* 21:617–629.
- Damasio, H., R. G. Maurer, A. R. Damasio, and H. C. Chui (1980). Computerized tomographic scan findings in patients with autistic behavior. *Arch. Neurol.* 37:504–510.
- Davidovitch, M., B. Patterson, and P. Gartside (1996). Head circumference measurements in children with autism. *J. Child Neurol.* 11:389–393.

- Dawson, G., J. Munson, S. J. Webb, T. Nalty, R. Abbott, and K. Toth (2007). Rate of head growth decelerates and symptoms worsen in the second year of life in autism. *Biol. Psychiatry* 61:458–464.
- Dementieva, Y. A., D. D. Vance, S. L. Donnelly, L. A. Elston, C. M. Wolpert, S. A. Ravan, G. R. DeLong, R. K. Abramson, H. H. Wright, and M. L. Cuccaro (2005). Accelerated head growth in early development of individuals with autism. *Pediatr. Neurol.* 32:102–108.
- Department of Health and Human Services, Centers for Disease Control and Prevention (2007). *Morbidity and Mortality Weekly Report* 56:1–28.
- Dissanayake, C., Q. M. Bui, R. Huggins, and D. Z. Loesch (2006). Growth in stature and head circumference in high-functioning autism and Asperger disorder during the first 3 years of life. *Dev. Psychopathol.* 18:381–393.
- Earnest, M. P., G. E. Thomas, R. A. Eden, and K. F. Hossack (1992). The sudden unexplained death syndrome in epilepsy: Demographic, clinical, and postmortem features. *Epilepsia* 33:310–316.
- Ehlert, U., J. Gaab, and M. Heinrichs (2001). Psychoneuroendocrinological contributions to the etiology of depression, posttraumatic stress disorder, and stress-related bodily disorders: The role of the hypothalamus-pituitary-adrenal axis. *Biol. Psychol.* 57:141–152.
- Erden-Inal, M., E. Sunal, and G. Kanbak (2002). Age-related changes in the glutathione redox system. *Cell Biochem. Funct.* 20:61–66.
- Ericksson, P. S., E. Perfilieva, T. Bjork-Eriksson, A. M. Alborn, C. Nordborg, D. A. Peterson, and F. H. Gage (1998). Neurogenesis in the adult human hippocampus. *Nat. Med.* 4:1313–1317.
- Evans, T., S. L. Siedlak, L. Lu, X. Fu, Z. Wang, W. R. McGinnis, E. Fakhoury, R. J. Castellani, S. L. Hazen, W. L. Walsh, A. T. Levis, R. G. Salomon, M. A. Smith, G. Perry, and X. Zhu (2008). The autistic phenotype exhibits a remarkably localized modification of brain protein by products of free radical-induced lipid oxidation. *Am J Biochem. Biotechnol., Special Issue on Autism Spectrum Disorders* 4:61–72.
- Fairhall, S. L. and A. Ishai (2007). Effective connectivity within the distributed cortical network for face perception. *Cereb. Cortex* 17:2400–2406.
- Fehlow, P., K. Bernstein, A. Tennstedt, and F. Walther (1993). Early infantile autism and excessive aerophagy with symptomatic megacolon and ileus in a case of Ehlers-Danlos syndrome. *Pediatr. Grenzgeb.* 31:259–267.
- Ferguson, J. N., L. J. Young, E. F. Hearn, M. M. Matzuk, T. R. Insel, and J. T. Winslow (2000). Social amnesia in mice lacking the oxytocin gene. *Nat. Genet.* 25:284–288.
- Fidler, D. J., J. N. Bailey, and S. L. Smalley (2000). Macrocephaly in autism and other pervasive developmental disorders. *Dev. Med. Child Neurol.* 42:737–740.
- Filipek, P. A. (1996). Brief report: Neuroimaging in autism: The state of the science 1995. *J. Autism Dev. Disord.* 26:211–215.
- Filipek, P. A., P. J. Accardo, G. T. Baranek, E. H. Cook Jr., G. Dawson, B. Gordon, J. S. Gravel, C. P. Johnson, R. J. Kallen, S. E. Levy, N. J. Minshew, S. Ozonoff, B. M. Prizant, I. Rapin, S. J. Rogers, W. L. Stone, S. Teplin, R. F. Tuchman, and F. R. Volkmar (1999). The screening and diagnosis of autistic spectrum disorders. *J. Autism Dev. Disord.* 29:439–484.
- Folstein, S. E. and B. Rosen-Sheidley (2001). Genetics of autism: Complex aetiology for a heterogeneous disorder. *Nat. Rev. Genet.* 2:943–955.
- Folstein, S. E. and M. L. Rutter (1988). Autism: Familial aggregation and genetic implications. *J. Autism Dev. Disord.* 18:3–30.
- Fombonne E. (2003). Epidemiological surveys of autism and other pervasive developmental disorders. *J. Autism Dev. Disord.* 33:365–382.
- Fombonne, E., B. Roge, J. Claverie, S. Courty, and J. Fremolle (1999). Microcephaly and macrocephaly in autism. *J. Autism Dev. Disord.* 29:113–119.
- Friede, R. L. (1975). *Developmental Neuropathology*. Springer Verlag, Berlin.

- Fuh, J. L. and S. J. Wang (1995). Caudate hemorrhage: Clinical features, neuropsychological assessments and radiological findings. *Clin. Neurol. Neurosurg.* 97:296–299.
- Gabreels, B. A., D. F. Swaab, D. P. de Kleijn, N. G. Seidah, J. W. Van de Loo, d. Van, V. G. J. Martens, and F. W. Van Leeuwen (1998). Attenuation of the polypeptide 7B2, pro-hormone convertase PC2, and vasopressin in the hypothalamus of some Prader-Willi patients: Indications for a processing defect. *J. Clin. Endocrinol. Metab.* 83:591–599.
- Gabriels, R. L., M. L. Cuccaro, D. E. Hill, B. J. Ivers, and E. Goldson (2005). Repetitive behaviors in autism: Relationships with associated clinical features. *Res. Dev. Disabil.* 26:169–181.
- Gaffney, G. R., L. Y. Tsai, S. Kuperman, and S. Minchin (1987). Cerebellar structure in autism. *Am. J. Dis. Child* 141:1330–1332.
- Gainer, H., M. O. Lively, and M. Morris (1995). Immunological and related techniques for studying neurohypophyseal peptide-processing pathways. *Methods Neurosci.* 23:195–207.
- Garber, H. J. and E. R. Ritvo (1992). Magnetic resonance imaging of the posterior fossa in autistic adults. *Am. J. Psychiatry* 149:245–247.
- Gessaga, E. C. and H. Urich (1985). The cerebellum of epileptics. *Clin. Neuropathol.* 4:238–245.
- Ghaziuddin, M., L. Y. Tsai, and N. Ghaziuddin (1992). Autism in Down's syndrome: Presentation and diagnosis. *J. Intellect. Disabil. Res.* 36 (Pt 5):449–456.
- Gillberg, C. (1998). Chromosomal disorders and autism. *J. Autism Dev. Disord.* 28:415–425.
- Gillberg C. and M. Coleman (1996). Autism and medical disorders: A review of the literature. *Dev. Med. Child. Neurol.* 38:191–202.
- Gillberg, C. and L. de Souza (2002). Head circumference in autism, Asperger syndrome, and ADHD: A comparative study. *Dev. Med. Child Neurol.* 44:296–300.
- Glabe, C. (2001). Intracellular mechanisms of amyloid accumulation and pathogenesis in Alzheimer's disease. *J. Mol. Neurosci.* 17:137–145.
- Green, L. A., D. Fein, C. Modahl, C. Feinstein, L. Waterhouse, and M. Morris (2001). Oxytocin and autistic disorder: Alteration in peptide forms. *Biol. Psychiatry* 50:609–613.
- Grelotti, D., I. Gauthier, and R. T. Shultz (2001). Social interest and the development of cortical face specialization: What autism teaches us about face processing. *Dev. Psychobiol.* 40:213–225.
- Griebel, G., J. Simiand, G. C. Serradeil-Le, J. Wagnon, M. Pascal, B. Scatton, J. P. Maffrand, and P. Soubrie (2002). Anxiolytic- and antidepressant-like effects of the non-peptide vasopressin V1b receptor antagonist, SSR149415, suggest an innovative approach for the treatment of stress-related disorders. *Proc. Natl. Acad. Sci. USA* 99:6370–6375.
- Guerrini, R. and C. Marini (2006). Genetic malformations of cortical development. *Exp. Brain Res.* 173:322–333.
- Hagerman, R. J. (2002). The physical and behavioral phenotype. In R. J. Hagerman and P. J. Hagerman (eds.), *Fragile X Syndrome: Diagnosis, Treatment, and Research*. Johns Hopkins University Press, Baltimore, MD, pp. 206–248.
- Happe, F. (1999). Autism: Cognitive deficit or cognitive style? *Trends Cogn. Sci.* 3:216–222.
- Happe, F., A. Ronald, and R. Plomin (2006). Time to give up on a single explanation for autism. *Nat. Neurosci.* 9:1218–1220.
- Hardan, A. Y., N. J. Minshew, M. Mallikarjunn, and M. S. Keshavan (2001). Brain volume in autism. *J. Child Neurol.* 16:421–424.
- Hardiman, O., T. Burke, J. Phillips, S. Murphy, B. O'Moore, H. Staunton, and M. A. Farrell (1988). Microdysgenesis in resected temporal neocortex: Incidence and clinical significance in focal epilepsy. *Neurology* 38:1041–1047.
- Harris, N. S., E. Courchesne, J. Townsend, R. A. Carper, and C. Lord (1999). Neuroanatomic contributions to slowed orienting of attention in children with autism. *Brain Res. Cogn. Brain Res.* 8:61–71.

- Harris, S. R., L. L. MacKay, and J. A. Osborn (1995). Autistic behaviors in offspring of mothers abusing alcohol and other drugs: A series of case reports. *Alcohol Clin. Exp. Res.* 19:660–665.
- Hashimoto, T., M. Tayama, M. Miyazaki, K. Murakawa, and Y. Kuroda (1993). Brainstem and cerebellar vermis involvement in autistic children. *J. Child Neurol.* 8:149–153.
- Hashimoto, T., M. Tayama, M. Miyazaki, N. Sakurama, T. Yoshimoto, K. Murakawa, and Y. Kuroda (1992). Reduced brainstem size in children with autism. *Brain Dev.* 14:94–97.
- Hashimoto, T., M. Tayama, K. Mori, K. Fujino, M. Miyazaki, and Y. Kuroda (1989). Magnetic resonance imaging in autism: Preliminary report. *Neuropediatrics* 20:142–146.
- Hashimoto, T., M. Tayama, K. Murakawa, T. Yoshimoto, M. Miyazaki, M. Harada, and Y. Kuroda (1995). Development of the brainstem and cerebellum in autistic patients. *J. Autism Dev. Disord.* 25:1–18.
- Hasselmo, M. E., E. T. Rolls, and G. C. Baylis (1989). The role of expression and identity in the face-selective responses of neurons in the temporal visual cortex of the monkey. *Behav. Brain Res.* 32:203–218.
- Hazlett, H. C., M. Poe, G. Gerig, R. G. Smith, J. Provenzale, A. Ross, J. Gilmore, and J. Piven (2005). Magnetic resonance imaging and head circumference study of brain size in autism: Birth through age 2 years. *Arch. Gen. Psychiatry* 62:1366–1376.
- Haznedar, M. M., M. S. Buchsbaum, M. Metzger, A. Solimando, J. Spiegel-Cohen, and E. Hollander (1997). Anterior cingulate gyrus volume and glucose metabolism in autistic disorder. *Am. J. Psychiatry* 154:1047–1050.
- Haznedar, M. M., M. S. Buchsbaum, T. C. Wei, P. R. Hof, C. Cartwright, C. A. Bienstock, and E. Hollander (2000). Limbic circuitry in patients with autism spectrum disorders studied with positron emission tomography and magnetic resonance imaging. *Am. J. Psychiatry* 157:1994–2001.
- Herbert, M. R., G. J. Harris, K. T. Adrien, D. A. Ziegler, N. Makris, D. N. Kennedy, N. T. Lange, C. F. Chabris, A. Bakardjiev, J. Hodgson, M. Takeoka, H. Tager-Flusberg, and V. S. Caviness Jr. (2002). Abnormal asymmetry in language association cortex in autism. *Ann. Neurol.* 52:588–596.
- Herbert, M. R., D. A. Ziegler, C. K. Deutsch, L. M. O'Brien, N. Lange, A. Bakardjiev, J. Hodgson, K. T. Adrien, S. Steele, N. Makris, D. Kennedy, G. J. Harris, and V. S. Caviness Jr. (2003). Dissociations of cerebral cortex, subcortical and cerebral white matter volumes in autistic boys. *Brain* 126:1182–1192.
- Herbert, M. R., D. A. Ziegler, N. Makris, P. A. Filipek, T. L. Kemper, J. J. Normandin, H. A. Sanders, D. N. Kennedy, and V. S. Caviness Jr. (2004). Localization of white matter volume increase in autism and developmental language disorder. *Ann. Neurol.* 55:530–540.
- Herken, H., E. Uz, H. Ozyurt, S. Sogut, O. Virit, and O. Akyol (2001). Evidence that the activities of erythrocyte free radical scavenging enzymes and the products of lipid peroxidation are increased in different forms of schizophrenia. *Mol. Psychiatry* 6:66–73.
- Hevner, R. F. (2007). Layer-specific markers as probes for neuron type identity in human neocortex and malformations of cortical development. *J. Neuropathol. Exp. Neurol.* 66:101–109.
- Hollander, E., E. Anagnostou, W. Chaplin, K. Esposito, M. M. Haznedar, E. Licalzi, S. Wasserman, L. Soorya, and M. Buchsbaum (2005). Striatal volume on magnetic resonance imaging and repetitive behaviors in autism. *Biol. Psychiatry* 58:226–232.
- Hollander, E., S. Novotny, M. Hanratty, R. Yaffe, C. M. DeCaria, B. R. Aronowitz, and S. Mosovich (2003). Oxytocin infusion reduces repetitive behaviors in adults with autistic and Asperger's disorders. *Neuropsychopharmacology* 28:193–198.
- Holtum, J. R., N. J. Minshew, R. S. Sanders, and N. E. Phillips (1992). Magnetic resonance imaging of the posterior fossa in autism. *Biol. Psychiatry* 32:1091–1101.

- Howlin, P., L. Wing, and J. Gould. 1995. The recognition of autism in children with Down syndrome—Implications for intervention and some speculations about pathology. *Dev. Med. Child Neurol.* 37: 406–414.
- Hoyer, D., D. E. Clarke, J. R. Fozard, P. R. Hartig, G. R. Martin, E. J. Mylecharane, P. R. Saxena, and P. P. Humphrey (1994). International Union of Pharmacology classification of receptors for 5-hydroxytryptamine (Serotonin). *Pharmacol. Rev.* 46:157–203.
- Hsu, M., R. Yeung-Courchesne, E. Courchesne, and G. A. Press (1991). Absence of magnetic resonance imaging evidence of pontine abnormality in infantile autism. *Arch. Neurol.* 48:1160–1163.
- Hudson, L. P., D. G. Munoz, L. Miller, R. S. McLachlan, J. P. Girvin, and W. T. Blume (1993). Amygdaloid sclerosis in temporal lobe epilepsy. *Ann. Neurol.* 33:622–631.
- Jallon, P. (1997). Epilepsy and the heart. *Rev. Neurol.* 153:173–184.
- Jellinger, K., D. Armstrong, H. Y. Zoghbi, and A. K. Percy (1988). Neuropathology of Rett syndrome. *Acta Neuropathol.* 76:142–158.
- Johns, J. M., D. A. Lubin, J. A. Lieberman, and J. M. Lauder (2002). Developmental effects of prenatal cocaine exposure on 5-HT1A receptors in male and female rat offspring. *Dev. Neurosci.* 24:522–530.
- Joseph, R. M. and J. Tanaka (2003). Holistic and part-based face recognition in children with autism. *J. Child Psychol. Psychiatry* 44:529–542.
- Just, M. A., V. L. Cherkassky, T. A. Keller, and N. J. Minshew (2004). Cortical activation and synchronization during sentence comprehension in high-functioning autism: Evidence of underconnectivity. *Brain* 127:1811–1821.
- Juurink, B. H. and P. G. Paterson (1998). Review of oxidative stress in brain and spinal cord injury: Suggestions for pharmacological and nutritional management strategies. *J. Spinal Cord. Med.* 21:309–334.
- Kanwisher, N., D. Stanley, and A. Harris (1999). The fusiform face area is selective for faces not animals. *Neuroreport* 10:183–187.
- Katoh-Semba, R., I. K. Takeuchi, R. Semba, and K. Kato (1997). Distribution of brain-derived neurotrophic factor in rats and its changes with development in the brain. *J. Neurochem.* 69:34–42.
- Kaufmann, W. E., K. L. Cooper, S. H. Mostofsky, G. T. Capone, W. R. Kates, C. J. Newschaffer, I. Bukelis, M. H. Stump, A. E. Jann, and D. C. Lanham (2003). Specificity of cerebellar vermian abnormalities in autism: A quantitative magnetic resonance imaging study. *J. Child Neurol.* 18:463–470.
- Kawamoto, Y., S. Nakamura, S. Nakano, N. Oka, I. Akiguchi, and J. Kimura (1996). Immunohistochemical localization of brain-derived neurotrophic factor in adult rat brain. *Neuroscience* 74:1209–1226.
- AQ8 Kemper, T. L. and M. L. Bauman (1992). Neuropathology of infantile autism. In H. Naruse and E. M. Ornitz (eds.), *Neurobiology of Infantile Autism*. Elsevier Science, Amsterdam, pp. 43–57.
- Kemper, T. L. and M. L. Bauman (1993). The contribution of neuropathologic studies to the understanding of autism. *Neurol. Clin.* 11:175–187.
- Kent, L., J. Evans, M. Paul, and M. Sharp (1999). Comorbidity of autistic spectrum disorders in children with Down syndrome. *Dev. Med. Child Neurol.* 41:153–158.
- Kientz, M. A. and W. Dunn (1997). A comparison of the performance of children with and without autism on the sensory profile. *Am. J. Occup. Ther.* 51:530–537.
- Kirsch, P., C. Esslinger, Q. Chen, D. Mier, S. Lis, S. Siddhanti, H. Gruppe, V. S. Mattay, B. Gallhofer, and A. Meyer-Lindenberg (2005). Oxytocin modulates neural circuitry for social cognition and fear in humans. *J. Neurosci.* 25:11489–11493.
- Kondoh, M., T. Shiga, and N. Okado (2004). Regulation of dendrite formation of Purkinje cells by serotonin through serotonin 1A and serotonin 2A receptors in culture. *Neurosci. Res.* 48:101–109.

- Koshino, H., P. A. Carpenter, N. J. Minshew, V. L. Cherkassky, T. A. Keller, and M. A. Just (2005). Functional connectivity in an fMRI working memory task in high-functioning autism. *Neuroimage* 24:810–821.
- Kramer, K., E. C. Azmitia, and P. M. Whitaker-Azmitia (1994). In vitro release of [³H]5-hydroxytryptamine from fetal and maternal brain by drugs of abuse. *Brain Res. Dev. Brain Res.* 78:142–146.
- Kuperman, S., J. Beechly, T. Burns, and L. Tsai (1987). Association of serotonin concentration to behavior and IQ in autistic children. *J. Autism Dev. Disord.* 17:133–140.
- Lainhart, J. E., J. Piven, M. Wzorek, R. Landa, S. L. Santangelo, H. Coon, and S. E. Folstein (1997). Macrocephaly in children and adults with autism. *J. Am. Acad. Child Adolesc. Psychiatry* 36:282–290.
- Landgraf, R. and I. D. Neumann (2004). Vasopressin and oxytocin release within the brain: A dynamic concept of multiple and variable modes of neuropeptide communication. *Front. Neuroendocrinol.* 25:150–176.
- Langen, M., S. Durston, W. G. Staal, S. J. Palmen, and H. van Engeland (2007). Caudate nucleus is enlarged in high-functioning medication-naïve subjects with autism. *Biol. Psychiatry* 62:262–266.
- Lee, M., C. Martin-Ruiz, A. Graham, J. Court, E. Jaros, R. Perry, P. Iversen, M. Bauman, and E. Perry (2002). Nicotinic receptor abnormalities in the cerebellar cortex in autism. *Brain* 125:1483–1495.
- Levitt, J. G., R. Blanton, L. Capetillo-Cunliffe, D. Guthrie, A. Toga, and J. T. McCracken (1999). Cerebellar vermis lobules VIII-X in autism. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 23:625–633.
- Lidov, H. G. and M. E. Molliver (1982). Immunohistochemical study of the development of serotonergic neurons in the rat CNS. *Brain Res Bull.* 9:559–604.
- Lopez-Hurtado, E. and J. J. Prieto (2008). A microscopic study of language-related cortex in autism. *Am. J. Biochem. Biotechnol. Special Issue on Autism Spectrum Disorders*, 4:130–145.
- Lord, C., S. Risi, L. Lambrecht, E. H. Cook Jr., B. L. Leventhal, P. C. DiLavore, A. Pickles, and M. Rutter (2000). The autism diagnostic observation schedule-generic: A standard measure of social and communication deficits associated with the spectrum of autism. *J. Autism Dev. Disord.* 30:205–223.
- Lord, C. and M. Rutter (1995). Autism and pervasive developmental disorders. In M. Rutter, E. Taylor, and L. Hersov (eds.), *Child and Adolescent Psychiatry, Modern Approaches*. Blackwell Science, Oxford, pp. 569–593.
- Mackenzie, I. R., R. S. McLachlan, C. S. Kubu, and L. A. Miller (1996). Prospective neuropsychological assessment of nondemented patients with biopsy proven senile plaques. *Neurology* 46:425–429.
- Mackenzie, I. R. and L. A. Miller (1994). Senile plaques in temporal lobe epilepsy. *Acta Neuropathol.* 87:504–510.
- Maeshima, T., F. Shutoh, S. Hamada, K. Senzaki, K. Hamaguchi-Hamada, R. Ito, and N. Okado (1998). Serotonin_{2A} receptor-like immunoreactivity in rat cerebellar Purkinje cells. *Neurosci. Lett.* 252:72–74.
- Manaye, K. F., D. L. Lei, Y. Tizabi, M. I. Vila-Garcia, P. R. Mouton, and P. H. Kelly (2005). Selective neuron loss in the paraventricular nucleus of hypothalamus in patients suffering from major depression and bipolar disorder. *J. Neuropathol. Exp. Neurol.* 64:224–229.
- Martin, G. R. and P. P. Humphrey (1994). Receptors for 5-hydroxytryptamine: Current perspectives on classification and nomenclature. *Neuropharmacology* 33:261–273.
- Mason-Brothers, A., E. R. Ritvo, C. Pingree, P. B. Petersen, W. R. Jenson, W. M. McMahon, B. J. Freeman, L. B. Jorde, M. J. Spencer, and A. Mo (1990). The UCLA-University of Utah epidemiologic survey of autism: Prenatal, perinatal, and postnatal factors. *Pediatrics* 86:514–519.

- McClelland, J. L. (2000). The basis of hyperspecificity in autism: A preliminary suggestion based on properties of neural nets. *J. Autism Dev. Disord.* 30:497–502.
- Mendez, M. F., N. L. Adams, and K. S. Lewandowski (1989). Neurobehavioral changes associated with caudate lesions. *Neurology* 39:349–354.
- Menendez, M. (2005). Down syndrome, Alzheimer's disease and seizures. *Brain Dev* 27:246–252.
- Miles, J. H., L. L. Hadden, T. N. Takahashi, and R. E. Hillman (2000). Head circumference is an independent clinical finding associated with autism. *Am. J. Med. Genet.* 95:339–350.
- Miles, J. H., T. N. Takahashi, S. Bagby, P. K. Sahota, D. F. Vaslow, C. H. Wang, R. E. Hillman, and J. E. Farmer (2005). Essential versus complex autism: Definition of fundamental prognostic subtypes. *Am. J. Med. Genet. A* 135:171–180.
- Ming, X., T. P. Stein, M. Brimacombe, W. G. Johnson, G. H. Lambert, and G. C. Wagner (2005). Increased excretion of a lipid peroxidation biomarker in autism. *Prostaglandins Leukot. Essent. Fatty Acids* 73:379–384.
- Mitchell, B. F., X. Fang, and S. Wong (1998). Role of carboxy-extended forms of oxytocin in the rat uterus in the process of parturition. *Biol. Reprod.* 59:1321–1327.
- Miyazaki, K., N. Narita, R. Sakuta, T. Miyahara, H. Naruse, N. Okado, and M. Narita (2004). Serum neurotrophin concentrations in autism and mental retardation: A pilot study. *Brain Dev.* 26:292–295.
- Muhle, R., S. V. Trentacoste, and I. Rapin (2004). The genetics of autism. *Pediatrics* 113:e472–e486.
- Muller, R. A. (2007). The study of autism as a distributed disorder. *Ment. Retard. Dev. Disabil. Res. Rev.* 13:85–95.
- Murakami, J. W., E. Courchesne, G. A. Press, R. Yeung-Courchesne, and J. R. Hesselink (1989). Reduced cerebellar hemisphere size and its relationship to vermal hypoplasia in autism. *Arch. Neurol.* 46:689–694.
- Murer, M. G., Q. Yan, and R. Raisman-Vozari (2001). Brain-derived neurotrophic factor in the control human brain, and in Alzheimer's disease and Parkinson's disease. *Prog. Neurobiol.* 63:71–124.
- Narita, N., M. Kato, M. Tazoe, K. Miyazaki, M. Narita, and N. Okado (2002). Increased monoamine concentration in the brain and blood of fetal thalidomide- and valproic acid-exposed rat: Putative animal models for autism. *Pediatr. Res.* 52:576–579.
- Nashef, L., F. Walker, P. Allen, J. W. Sander, S. D. Shorvon, and D. R. Fish (1996). Apnoea and bradycardia during epileptic seizures: Relation to sudden death in epilepsy. *J. Neurol. Neurosurg. Psychiatry* 60:297–300.
- Nawa, H., J. Carnahan, and C. Gall (1995). BDNF protein measured by a novel enzyme immunoassay in normal brain and after seizure: Partial disagreement with mRNA levels. *Eur. J. Neurosci.* 7:1527–1535.
- Nelson, E. and J. Panksepp (1996). Oxytocin mediates acquisition of maternally associated odor preferences in preweanling rat pups. *Behav. Neurosci.* 110:583–592.
- Newschaffer, C. J., D. Fallin, and N. L. Lee (2002). Heritable and nonheritable risk factors for autism spectrum disorders. *Epidemiol. Rev.* 24:137–153.
- Niitsu, Y., S. Hamada, K. Hamaguchi, M. Mikuni, and N. Okado (1995). Regulation of synapse density by 5-HT_{2A} receptor agonist and antagonist in the spinal cord of chicken embryo. *Neurosci. Lett.* 195:159–162.
- Nishizawa, S., C. Benkelfat, S. N. Young, M. Leyton, S. Mzengeza, M. C. de, P. Blier, and M. Diksic (1997). Differences between males and females in rates of serotonin synthesis in human brain. *Proc. Natl. Acad. Sci. USA.* 94:5308–5313.
- Okado, N., L. Cheng, Y. Tanatsugu, S. Hamada, and K. Hamaguchi (1993). Synaptic loss following removal of serotonergic fibers in newly hatched and adult chickens. *J. Neurobiol.* 24:687–698.

- Ono, H., A. Sakamoto, and N. Sakura (2001). Plasma total glutathione concentrations in healthy pediatric and adult subjects. *Clin. Chim. Acta* 312:227–229.
- Oppenheim, R. W. (1991). Cell death during development of the nervous system. *Annu. Rev. Neurosci.* 14:453–501.
- Palmen, S. J., H. van Engeland, P. R. Hof, and C. Schmitz (2004). Neuropathological findings in autism. *Brain* 127:2572–2583.
- Pennington, B. F., P. A. Filipek, D. Lefly, N. Chhabildas, D. N. Kennedy, J. H. Simon, C. M. Filley, A. Galaburda, and J. C. DeFries (2000). A twin MRI study of size variations in human brain. *J. Cogn. Neurosci* 12:223–232.
- Perrett, D. I., P. A. Smith, D. D. Potter, A. J. Mistlin, A. S. Head, A. D. Milner, and M. A. Jeeves (1985). Visual cells in the temporal cortex sensitive to face view and gaze direction. *Proc. R. Soc. Lond B Biol. Sci.* 223:293–317.
- Perry, E. K., M. L. Lee, C. M. Martin-Ruiz, Court JA, S. G. Volsen, J. Meritt, E. Folly, P. E. Iversen, M. L. Bauman, R. H. Perry, and G. L. Wenk (2001). Cholinergic activity in autism: Abnormalities in the cerebral cortex and basal forebrain. *Am. J. Psychiatry* 158:1058–1066.
- Perry, S. W., J. P. Norman, A. Litzburg, and H. A. Gelbard (2004). Antioxidants are required during the early critical period, but not later, for neuronal survival. *J. Neurosci. Res.* 78:485–492.
- Pfefferbaum, A., E. V. Sullivan, G. E. Swan, and D. Carmelli (2000). Brain structure in men remains highly heritable in the seventh and eighth decades of life. *Neurobiol. Aging* 21:63–74.
- Pickett, J. and E. London (2005). The neuropathology of autism: A review. *J. Neuropathol. Exp. Neurol.* 64:925–935.
- Pierce, K. and E. Courchesne (2001). Evidence for a cerebellar role in reduced exploration and stereotyped behavior in autism. *Biol. Psychiatry* 49:655–664.
- Pierce, K., R. A. Muller, J. Ambrose, G. Allen, and E. Courchesne (2001). Face processing occurs outside the fusiform “face area” in autism: Evidence from functional MRI. *Brain* 124:2059–2073.
- Pierce, K., F. Haist, F. Sedaghat, and E. Courchesne (2004). The brain response to personally familiar faces in autism: Findings of fusiform activity and beyond. *Brain* 127:2703–2716.
- Piven, J., E. Nehme, J. Simon, P. Barta, G. Pearlson, and S. E. Folstein (1992). Magnetic resonance imaging in autism: Measurement of the cerebellum, pons, and fourth ventricle. *Biol. Psychiatry* 31:491–504.
- Piven, J., S. Arndt, J. Bailey, S. Havercamp, N. C. Andreasen, and P. Palmer (1995). An MRI study of brain size in autism. *Am. J. Psychiatry* 152:1145–1149.
- Piven, J., S. Arndt, J. Bailey, and N. Andreasen (1996). Regional brain enlargement in autism: A magnetic resonance imaging study. *J. Am. Acad. Child Adolesc. Psychiatry* 35:530–536.
- Piven, J., K. Saliba, J. Bailey, and S. Arndt (1997a). An MRI study of autism: The cerebellum revisited. *Neurology* 49:546–551.
- Piven, J., J. Bailey, B. J. Ranson, and S. Arndt (1997b). An MRI study of the corpus callosum in autism. *Am. J. Psychiatry* 154:1051–1056.
- Piven, J., J. Bailey, B. J. Ranson, and S. Arndt (1998). No difference in hippocampus volume detected on magnetic resonance imaging in autistic individuals. *J. Autism Dev. Disord.* 28:105–110.
- Poldrack, R. A., V. Prabhakaran, C. A. Seger, and J. D. Gabrieli (1999). Striatal activation during acquisition of a cognitive skill. *Neuropsychology*. 13:564–574.
- Popik, P. and J. M. Van Ree (1992). Long-term facilitation of social recognition in rats by vasopressin related peptides: A structure–activity study. *Life Sci.* 50:567–572.
- Popik, P., J. Vetulani, and J. M. Van Ree (1992). Low doses of oxytocin facilitate social recognition in rats. *Psychopharmacology* 106:71–74.

- AQ9 Pracher, V. P. and D. E. J. Clarke (1996). Case report: Challenging behaviour in a young adult with Down's syndrome and autism. *Brit. J. Learning Disab.* 24:167–169.
- Ragenbass, M. 2001. Vasopressin- and oxytocin-induced activity in the central nervous system: Electrophysiological studies using in-vitro systems. *Prog. Neurobiol.* 64:307–326.
- Rao, V. V., C. Löffler, J. Battey, and I. Hansmann (1992). The human gene for oxytocin-neurophysin I (OXT) is physically mapped to chromosome 20p13 by in situ hybridization. *Cytogenet. Cell Genet.* 61:271–273.
- Rapin, I. (1996). *Preschool Children with Inadequate Communication: Developmental Language Disorder, Autism, Low IQ*. MacKeith Press, London.
- Rathbun, W. and M. J. Druse (1985). Dopamine, serotonin, and acid metabolites in brain regions from the developing offspring of ethanol-treated rats. *J. Neurochem.* 44:57–62.
- Redcay, E. and E. Courchesne (2005). When is the brain enlarged in autism? A meta-analysis of all brain size reports. *Biol. Psychiatry* 58:1–9.
- Reeves, A. L., K. E. Nolle, D. W. Klass, F. W. Sharbrough, and E. L. So (1996). The ictal bradycardia syndrome. *Epilepsia* 37:983–987.
- Risse, S. C., T. H. Lampe, T. D. Bird, D. Nochlin, S. M. Sumi, T. Keenan, L. Cubberley, E. Peskind, and M. A. Raskind (1990). Myoclonus, seizures, and paratonia in Alzheimer disease. *Alzheimer Dis. Assoc. Disord.* 4:217–225.
- Ritvo, E. R., B. J. Freeman, A. B. Scheibel, T. Duong, H. Robinson, D. Guthrie, and A. Ritvo (1986). Lower Purkinje cell counts in the cerebella of four autistic subjects: Initial findings of the UCLA-NSAC Autopsy Research Report. *Am. J. Psychiatry* 143:862–866.
- Rodier, P. M., J. L. Ingram, B. Tisdale, S. Nelson, and J. Romano (1996). Embryological origin for autism: Developmental anomalies of the cranial nerve motor nuclei. *J. Comp. Neurol.* 370:247–261.
- Rogers, S. J., L. Bennetto, R. McEvoy, and B. F. Pennington (1996). Imitation and pantomime in high-functioning adolescents with autism spectrum disorders. *Child Dev.* 67:2060–2073.
- Roth, B. L. (1994). Multiple serotonin receptors: Clinical and experimental aspects. *Ann. Clin. Psychiatry* 6:67–78.
- Rutter, M., A. Bailey, P. Bolton, and C. A. Le (1994). Autism and known medical conditions: Myth and substance. *J. Child Psychol. Psychiatry* 35:311–322.
- Saitoh, O. and E. Courchesne (1998). Magnetic resonance imaging study of the brain in autism. *Psychiatry Clin. Neurosci.* 52 Suppl:S219–S222.
- Sajdel-Sulkowska, E. M., B. Lipinski, H. Windom, T. Audhya, and W. McGinnis (2008). Oxidative stress in autism: Cerebellar 3-nitrotyrosine levels. *Am. J. Biochem. Biotechnol. Special Issue on Autism Spectrum Disorders*, 4:73–84.
- Sarnat, H. B. and L. Flores-Sarnat (2004). Integrative classification of morphology and molecular genetics in central nervous system malformations. *Am. J. Med. Genet. A* 126A:386–392.
- Saudou, F. and R. Hen (1994a). 5-Hydroxytryptamine receptor subtypes in vertebrates and invertebrates. *Neurochem. Int.* 25:503–532.
- Saudou, F. and R. Hen (1994b). 5-Hydroxytryptamine receptor subtypes: Molecular and functional diversity. *Adv. Pharmacol.* 30:327–380.
- Saussu, F., K. van Rijkevorsel, and T. de Barse (1998). [Bradycardia: An unrecognized complication of some epileptic crises]. *Rev Neurol.* 154:250–252.
- Schaefer, G. B., J. N. Thompson, J. B. Bodensteiner, J. M. McConnell, W. J. Kimberling, C. T. Gay, W. D. Dutton, D. C. Hutchings, and S. B. Gray (1996). Hypoplasia of the cerebellar vermis in neurogenetic syndromes. *Ann. Neurol.* 39:382–385.
- Schmidt-Kastner, R., C. Wetmore, and L. Olson (1996). Comparative study of brain-derived neurotrophic factor messenger RNA and protein at the cellular level suggests multiple roles in hippocampus, striatum and cortex. *Neuroscience* 74:161–183.
- Schmidtke, K., H. Manner, R. Kaufmann, and H. Schmolck (2002). Cognitive procedural learning in patients with fronto-striatal lesions. *Learn. Mem.* 9:419–429.

- Schultz, R. T. (2005). Developmental deficits in social perception in autism: The role of the amygdala and fusiform face area. *Int. J. Dev. Neurosci.* 23:125–141.
- Sears, L. L., C. Vest, S. Mohamed, J. Bailey, B. J. Ranson, and J. Piven (1999). An MRI study of the basal ganglia in autism. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 23:613–624.
- Shapiro, L. E. and T. R. Insel (1989). Ontogeny of oxytocin receptors in rat forebrain: A quantitative study. *Synapse* 4:259–266.
- Sheng, J. G., F. A. Boop, R. E. Mrak, and W. S. Griffin (1994). Increased neuronal beta-amyloid precursor protein expression in human temporal lobe epilepsy: Association with interleukin-1 alpha immunoreactivity. *J. Neurochem.* 63:1872–1879.
- Shulman, R. G., D. L. Rothman, K. L. Behar, and F. Hyder (2004). Energetic basis of brain activity: Implications for neuroimaging. *Trends Neurosci.* 27:489–495.
- Siegel, D. J., N. J. Minshew, and G. Goldstein (1996). Wechsler IQ profiles in diagnosis of high-functioning autism. *J. Autism Dev. Disord.* 26:389–406.
- Smalley, S. L., R. F. Asarnow, and M. A. Spence (1988). Autism and genetics. A decade of research. *Arch. Gen. Psychiatry* 45:953–961.
- Sohal, R. S. and U. T. Brunk (1989). Lipofuscin as an indicator of oxidative stress and aging. *Adv. Exp. Med. Biol.* 266:17–26.
- Sokol, D. K., D. Chen, M. R. Farlow, D. W. Dunn, B. Maloney, J. A. Zimmer, and D. K. Lahiri (2006). High levels of Alzheimer beta-amyloid precursor protein (APP) in children with severely autistic behavior and aggression. *J. Child Neurol.* 21:444–449.
- Sparks, B. F., S. D. Friedman, D. W. Shaw, E. H. Aylward, D. Echelard, A. A. Artru, K. R. Maravilla, J. N. Giedd, J. Munson, G. Dawson, and S. R. Dager (2002). Brain structural abnormalities in young children with autism spectrum disorder. *Neurology* 59:184–192.
- Stefanacci, L. and D. G. Amaral (2000). Topographic organization of cortical inputs to the lateral nucleus of the macaque monkey amygdala: A retrograde tracing study. *J. Comp. Neurol.* 421:52–79.
- Steg, J. P. and J. L. Rapoport (1975). Minor physical anomalies in normal, neurotic, learning disabled, and severely disturbed children. *J. Autism Child Schizophr.* 5:299–307.
- Stevenson, R. E., R. J. Schroer, C. Skinner, D. Fender, and R. J. Simensen (1997). Autism and macrocephaly. *Lancet* 349:1744–1745.
- Stojanovic, A., A. E. Roher, and M. J. Ball (1994). Quantitative analysis of lipofuscin and neurofibrillary tangles in the hippocampal neurons of Alzheimer disease brains. *Dementia* 5:229–233.
- Stone, W. L., O. Y. Ousley, P. J. Yoder, K. L. Hogan, and S. L. Hepburn (1997). Nonverbal communication in two- and three-year-old children with autism. *J. Autism Dev. Disord.* 27:677–696.
- Stromland, K., V. Nordin, M. Miller, B. Akerstrom, and C. Gillberg (1994). Autism in thalidomide embryopathy: A population study. *Dev. Med. Child Neurol.* 36:351–356.
- Sutcliffe, J. S., R. J. Delahanty, H. C. Prasad, J. L. McCauley, Q. Han, L. Jiang, C. Li, S. E. Folstein, and R. D. Blakely (2005). Allelic heterogeneity at the serotonin transporter locus (SLC6A4) confers susceptibility to autism and rigid-compulsive behaviors. *Am. J. Hum. Genet.* 77:265–279.
- Sutula, T. P. and A. Pitkanen (2001). More evidence for seizure-induced neuron loss: Is hippocampal sclerosis both cause and effect of epilepsy? *Neurology* 57:169–170.
- Swaab, D. F. (1997). Prader-Willi syndrome and the hypothalamus. *Acta Paediatr. Suppl.* 423:50–54.
- AQ10 Szatmari, P., M. B. Jones, L. Tuff, G. Bartolucci, S. Fisman, and W. Mahoney (1993). Lack of cognitive impairment in first-degree relatives of children with pervasive developmental disorders. *J. Am. Acad. Child Adolesc. Psychiatry* 32:1264–1273.
- Szatmari, P., M. B. Jones, L. Zwaigenbaum, and J. E. MacLean (1998). Genetics of autism: Overview and new directions. *J. Autism Dev. Disord.* 28:351–368.

- Szweda, P. A., M. Camouse, K. C. Lundberg, T. D. Oberley, and L. I. Szweda (2003). Aging, lipofuscin formation, and free radical-mediated inhibition of cellular proteolytic systems. *Ageing Res. Rev.* 2:383–405.
- Tapia-Arancibia, L., F. Rage, L. Givalois, and S. Arancibia (2004). Physiology of BDNF: Focus on hypothalamic function. *Front. Neuroendocrinol.* 25:77–107.
- Tasch, E., F. Cendes, L. M. Li, F. Dubeau, F. Andermann, and D. L. Arnold (1999). Neuroimaging evidence of progressive neuronal loss and dysfunction in temporal lobe epilepsy. *Ann. Neurol.* 45:568–576.
- Terman, A. and U. T. Brunk (2004). Lipofuscin. *Int. J. Biochem. Cell Biol.* 36:1400–1404.
- Thom, M., B. Griffin, J. W. Sander, and F. Scaravilli (1999). Amygdala sclerosis in sudden and unexpected death in epilepsy. *Epilepsy Res* 37:53–62.
- Torsdottir, G., J. Kristinsson, S. Sveinbjornsdottir, J. Snaedal, and T. Johannesson (1999). Copper, ceruloplasmin, superoxide dismutase and iron parameters in Parkinson's disease. *Pharmacol. Toxicol.* 85:239–243.
- Townsend, J., E. Courchesne, J. Covington, M. Westerfield, N. S. Harris, P. Lyden, T. P. Lowry, and G. A. Press (1999). Spatial attention deficits in patients with acquired or developmental cerebellar abnormality. *J. Neurosci.* 19:5632–5643.
- Tuchman, R. and I. Rapin (2002). Epilepsy in autism. *Lancet Neurol.* 1:352–358.
- van Kooten, I., S. J. Palmen, P. von Cappeln, H. W. Steinbusch, H. Korr, H. Heinsen, P. R. Hof, H. van Engeland, and C. Schmitz (2008). Neurons in the fusiform gyrus are fewer and smaller in autism. *Brain* 131:987–999.
- Velez, L. and L. M. Selwa (2003). Seizure disorders in the elderly. *Am. Fam. Physician* 67:325–332.
- Voelbel, G. T., M. E. Bates, J. F. Buckman, G. Pandina, and R. L. Hendren (2006). Caudate nucleus volume and cognitive performance: Are they related in childhood psychopathology? *Biol. Psychiatry* 60:942–950.
- Volkmar, F. R., A. Carter, S. S. Sparrow, and D. V. Cicchetti (1993). Quantifying social development in autism. *J. Am. Acad. Child Adolesc. Psychiatry* 32:627–632.
- Vorstman, J. A. S., W. G. Staal, E. van Daalen, H. van Engeland, P. F. R. Hochstenbach, and L. Franke (2006). *Mol. Psych.* 11:18–28.
- Waiter, G. D., J. H. Williams, A. D. Murray, A. Gilchrist, D. I. Perrett, and A. Whiten (2004). A voxel-based investigation of brain structure in male adolescents with autistic spectrum disorder. *Neuroimage.* 22:619–625.
- Wakabayashi, S. (1979). A case of infantile autism associated with Down's syndrome. *J. Autism Dev. Disord.* 9:31–36.
- Wegiel, J., I. Kuchna, K. Nowicki, J. Frackowiak, B. Mazur-Kolecka, H. Imaki, J. Wegiel, P. D. Mehta, W. P. Silverman, B. Reisberg, M. Deleon, T. Wisniewski, T. Pirttilla, H. Frey, T. Lehtimäki, T. Kivimäki, F. E. Visser, W. Kamphorst, A. Potempska, D. Bolton, J. R. Currie, and D. L. Miller (2007). Intraneuronal Abeta immunoreactivity is not a predictor of brain amyloidosis-beta or neurofibrillary degeneration. *Acta Neuropathol.* 113:389–402.
- Wegiel, J., E. London, I. L. Cohen, M. Flory, T. Wisniewski, H. Imaki, I. Kuchna, J. Wegiel, S. Y. Ma, K. Nowicki, J. Wang, and W. T. Brown (2008). Detection of leading developmental defects in brains of autistic subjects. In 7th Annual International Meeting for Autism Research (IMFAR).
- Wetherby, A. M., B. M. Prizant, and T. Hutchinson (1998). Communicative, social-affective, and symbolic profiles of young children with autism and pervasive developmental disorder. *Am. J. Speech Language Pathol.* 7:79–91.
- Whitaker-Azmitia, P. M. (2005). Behavioral and cellular consequences of increasing serotonergic activity during brain development: A role in autism? *Int. J. Dev. Neurosci.* 23:75–83.
- Williams, J. H., A. Whiten, T. Suddendorf, and D. I. Perrett (2001). Imitation, mirror neurons and autism. *Neurosci. Biobehav. Rev.* 25:287–295.

AQ11

- Windle, R. J., N. Shanks, S. L. Lightman, and C. D. Ingram (1997). Central oxytocin administration reduces stress-induced corticosterone release and anxiety behavior in rats. *Endocrinology* 138:2829–2834.
- Wu, S. P., M. X. Jia, Y. Ruan, Y. Q. Guo, M. Shuang, X. H. Gong, Y. B. Zhang, X. L. Yang, and D. Zhang (2005). Positive association of the oxytocin receptor gene (OXTR) with autism in the Chinese Han population. *Biol. Psychiatry* 58:74–77.
- Yan, Q., R. D. Rosenfeld, C. R. Matheson, N. Hawkins, O. T. Lopez, L. Bennett, and A. A. Welcher (1997). Expression of brain-derived neurotrophic factor protein in the adult rat central nervous system. *Neuroscience* 78:431–448.
- Yanik, M., H. Vural, H. Tutkun, S. S. Zoroglu, H. A. Savas, H. Herken, A. Kocyigit, H. Keles, and O. Akyol (2004). The role of the arginine-nitric oxide pathway in the pathogenesis of bipolar affective disorder. *Eur. Arch. Psychiatry Clin. Neurosci.* 254:43–47.
- Yip, J., J. J. Soghomonian, and G. J. Blatt (2007). Decreased GAD67 mRNA levels in cerebellar Purkinje cells in autism: Pathophysiological implications. *Acta Neuropathol.* 113:559–568.
- Yip, J., J. J. Soghomonian, and G. J. Blatt (2008). Increased GAD67 mRNA expression in cerebellar interneurons in autism: Implications for Purkinje cell dysfunction. *J. Neurosci. Res.* 86:525–530.
- Yonan, A. L., M. Alarcon, R. Cheng, P. K. Magnusson, S. J. Spence, A. A. Palmer, A. Grunn, S. H. Juo, J. D. Terwilliger, J. Liu, R. M. Cantor, D. H. Geschwind, and T. C. Gilliam (2003). A genomewide screen of 345 families for autism-susceptibility loci. *Am. J. Hum. Genet.* 73:886–897.
- Zilbovicius, M., B. Garreau, Y. Samson, P. Remy, C. Barthelemy, A. Syrota, and G. Lelord (1995). Delayed maturation of the frontal cortex in childhood autism. *Am. J. Psychiatry* 152:248–252.
- Zoroglu, S. S., F. Armutcu, S. Ozen, A. Gurel, E. Sivasli, O. Yetkin, and I. Meram (2004). Increased oxidative stress and altered activities of erythrocyte free radical scavenging enzymes in autism. *Eur. Arch. Psychiatry Clin. Neurosci* 254:143–147.

AUTHOR QUERIES

- [AQ1] Running head is edited. Please check.
- [AQ2] Tuchman et al. (1991) is not listed. Please supply the reference.
- [AQ3] Please suggest which author. There are many authors for this chapter.
- [AQ4] Armstrong (2005) in the reference list. Please reconcile.
- [AQ5] Please provide missing details for Arin et al. (1991).
- [AQ6] Buitelaar and Willemsen-Swinkels (2000) is not cited in the text.
- [AQ7] Courchesne et al. (2001) in the text. Please reconcile.
- [AQ8] Kemper and Bauman (1992) is not cited in the text.
- [AQ9] Please verify the author names in Pracher et al. (1996).
- [AQ10] Szatmari et al. (1993) is not cited in the text.
- [AQ11] Year added from citation. Please confirm.

The neuropathology of autism: defects of neurogenesis and neuronal migration, and dysplastic changes

Jerzy Wegiel · Izabela Kuchna · Krzysztof Nowicki · Humi Imaki · Jarek Wegiel · Elaine Marchi · Shuang Yong Ma · Abha Chauhan · Ved Chauhan · Teresa Wierzbza Bobrowicz · Mony de Leon · Leslie A. Saint Louis · Ira L. Cohen · Eric London · W. Ted Brown · Thomas Wisniewski

Received: 12 November 2009 / Revised: 5 February 2010 / Accepted: 9 February 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract Autism is characterized by a broad spectrum of clinical manifestations including qualitative impairments in social interactions and communication, and repetitive and stereotyped patterns of behavior. Abnormal acceleration of brain growth in early childhood, signs of slower growth of neurons, and minicolumn developmental abnormalities suggest multiregional alterations. The aim of this study was to detect the patterns of focal qualitative developmental defects and to identify brain regions that are prone to developmental alterations in autism. Formalin-fixed brain hemispheres of 13 autistic (4–60 years of age) and 14 age-matched control subjects were embedded in celloidin and cut into 200- μ m-thick coronal sections, which were stained with cresyl violet and used for neuropathological evaluation. Thickening of the subependymal cell layer in two brains and subependymal nodular dysplasia in one brain is

indicative of active neurogenesis in two autistic children. Subcortical, periventricular, hippocampal and cerebellar heterotopias detected in the brains of four autistic subjects (31%) reflect abnormal neuronal migration. Multifocal cerebral dysplasia resulted in local distortion of the cytoarchitecture of the neocortex in four brains (31%), of the entorhinal cortex in two brains (15%), of the cornu Ammonis in four brains and of the dentate gyrus in two brains. Cerebellar flocculonodular dysplasia detected in six subjects (46%), focal dysplasia in the vermis in one case, and hypoplasia in one subject indicate local failure of cerebellar development in 62% of autistic subjects. Detection of flocculonodular dysplasia in only one control subject and of a broad spectrum of focal qualitative neuropathological developmental changes in 12 of 13 examined brains of autistic subjects (92%) reflects multiregional dysregulation

Jerzy Wegiel (✉) · I. Kuchna · K. Nowicki · H. Imaki · Jarek Wegiel · E. Marchi · S. Y. Ma · T. Wisniewski
Department of Developmental Neurobiology,
NYS Institute for Basic Research in Developmental Disabilities
(IBR), 1050 Forest Hill Road, Staten Island, NY 10314, USA
e-mail: jerzy.wegiel@omr.state.ny.us

M. de Leon · L. A. S. Louis · T. Wisniewski
Department of Neurology,
New York University School of Medicine, New York, NY, USA

M. de Leon · L. A. S. Louis · T. Wisniewski
Department of Pathology,
New York University School of Medicine, New York, NY, USA

M. de Leon · L. A. S. Louis · T. Wisniewski
Department of Psychiatry,
New York University School of Medicine, New York, NY, USA

A. Chauhan · V. Chauhan
Department of Neurochemistry, IBR, Staten Island, NY, USA

T. W. Bobrowicz
Department of Neuropathology,
Institute of Psychiatry and Neurology, Warsaw, Poland

I. L. Cohen · E. London
Department of Psychology, IBR, Staten Island, NY, USA

W. T. Brown
Department of Human Genetics, IBR, Staten Island, NY, USA

L. A. S. Louis
Corinthian Diagnostic Radiology, New York, NY, USA

M. de Leon
Nathan S. Kline Institute for Psychiatric Research,
Orangeburg, NY, USA

of neurogenesis, neuronal migration and maturation in autism, which may contribute to the heterogeneity of the clinical phenotype.

Keywords Autism · Developmental neuropathology · Subependymal nodular dysplasia · Heterotopia · Dysplasia

Introduction

Autism is characterized by a broad spectrum of clinical manifestations, including (a) qualitative impairments in reciprocal social interactions, (b) qualitative impairments in verbal and nonverbal communication, (c) restricted repetitive and stereotyped patterns of behavior, interests and activities and (d) onset prior to the age of 3 years [1]. In most cases, the etiology is unknown, and patients are diagnosed with idiopathic or non-syndromic autism [10, 43]. About 70% of individuals with idiopathic autism have essential autism, defined by the absence of physical abnormalities, but in 30%, complex autism with dysmorphic features such as microcephaly and/or a structural brain malformation is diagnosed [79]. In 5–10% of cases, autism is diagnosed in association with other disorders such as fragile X syndrome, Rett syndrome, Down syndrome, and tuberous sclerosis [94, 105]. Intellectual impairments, defined as intelligence quotient (IQ) scores less than 70, were reported in 44.6% of children diagnosed with autism [28]. Epilepsy is observed in up to 33% of individuals with autism [106].

The phenotypic heterogeneity is a major obstacle in all areas of autism research [83] and may be the result of a contribution of non-overlapping gene effects. The genetic fractionation of social impairment, communication difficulties and rigid and repetitive behaviors suggests that different features of autism are caused by different genes associated with different brain regions and are related to different cognitive impairments and functional abnormalities [48].

In spite of the broad spectrum of clinical manifestations and striking inter-individual differences, studies of thousands of children have resulted in establishing the clinical diagnostic criteria of pervasive developmental disorder [1]; however, corresponding neuropathological diagnostic criteria do not exist. One of the reasons for the disproportionate progress in clinical and neuropathological studies is the limited tissue resources available for post-mortem studies. Between 1980 and 2003, only 58 brains of individuals with autism were examined [85]. Due to the diversity of research aims, of protocols for tissue preservation and of methods of sampling and examination, and the small number of brains examined in an individual

project, the pattern of neuropathological changes emerging from these studies remains incomplete and inconsistent.

The hypothesis that autism is associated with neuropathological changes was explored in the first reports published between 1980 and 1993 [7, 21, 22, 27, 42, 50, 51, 82, 90]. Since then, implementation of broader diagnostic terms such as Autism Spectrum Disorder (ASD), examination of larger cohorts, applications of stereology, and functional and structural magnetic resonance imaging (MRI) have resulted in the detection of several major types of pathology, most likely contributing to the clinical phenotype. An emerging concept of autism-related brain pathology integrates evidence of (a) abnormal acceleration of brain growth in early childhood [89], (b) minicolumn pathology [13, 14], (c) curtailed neuronal development [7, 108] and brain structure-specific delays of neuronal growth [111] with indications of abnormalities in brain cytoarchitecture [4, 7], metabolic modifications with abnormal amyloid protein precursor (APP) processing [5, 101], enhanced oxidative stress [17] and enhanced turnover of cell organelles with pigment accumulation and glial activation [68].

In spite of the conceptual limitations, “localizing” models are still the main approach to the identification of pathological changes as a component of the networks’ structural and functional abnormalities [81]. We hypothesize that dysregulation of neurogenesis, neuronal migration and maturation is also reflected in qualitative, focal, developmental alterations of brain microarchitecture. The aim of this study is to detect the pattern of focal, qualitative, developmental defects in autism brain, including their type, topography and severity, and to identify the structures and brain regions that are prone to developmental alterations in autism.

Materials and methods

The autistic cohort study consisted of 13 subjects (4–62 years of age), including 9 males (69%) and 4 females (31%), while the control cohort consisted of 14 subjects (4–64 years of age), including 9 males and 5 females (Table 1).

Clinical and genetic characteristics of the autistic subjects

The source of our clinical data was the medical records of the autistic subjects, which consisted of psychological, behavioral, neurological and psychiatric evaluation reports. All of the records were obtained after the subjects’ deaths. The Autism Diagnostic Interview-Revised (ADI-R) was administered to each donor family as a standardized assessment tool in order to confirm the diagnosis on a postmortem basis. Inclusion of the subject in this study was based on a summary of scores of four domains:

Table 1 Material examined

#	Group	Brain bank #	Sex	Age (years)	Cause of death	PMI (h)	H	Brain weight (g)
1	A	IBR425-02	M	4	Drowning	30	R	1,280
2	A	UMB-1627	F	5	Traumatic multiple injuries	13.2	R	1,390
3	A	B-6403	M	7	Drowning	25	R	1,610
4	A	B-5666	M	8	Rhabdomyosarcoma	22.2	R	1,570
5	A	B-5342	F	11	Seizure-related drowning	12.9	L	1,460
6	A	B-5535	M	13	Seizure-related	8	L	1,470
7	A	B-6115	F	17	Cardiac arrest related to cardiomyopathy	25	L	1,580
8	A	UMB-1638	F	21	Seizure-related respiratory failure	50	R	1,108
9	A	B-6337	M	22	Seizure-related	25	R	1,375
10	A	IBR93-01	M	23	Status epilepticus-related respiratory failure	14	R	1,610
11	A	B-6212	M	36	Cardiac arrest	24	R	1,480
12	A	B-6276	M	56	Cardiac arrest	3.35	R	1,570
13	A	B-7090	M	60	Pancreatic cancer	26.5	R	1,210
1	C	B-6736	F	4	Acute bronchopneumonia	17	R	1,530
2	C	UMB-1499	F	4	Lymphocytic myocarditis	21	R	1,222
3	C	UMB-4898	M	7	Drowning	12	R	1,240
4	C	UMB-1708	F	8	Traumatic multiple injuries	20	R	1,222
5	C	BTB-3638	M	14	Electrocution	20	R	1,464
6	C	UMB-1843	F	15	Traumatic multiple injuries	9	R	1,250
7	C	UMB-1846	F	20	Traumatic multiple injuries	9	R	1,340
8	C	UMB-1646	M	23	Ruptured spleen	6	R	1,520
9	C	UMB-4543	M	29	Traumatic multiple injuries	13	R	1,514
10	C	UMB-1576	M	32	Traumatic compressional asphyxia	24	R	1,364
11	C	BTB-3899	M	48	Atherosclerotic heart disease	24	L	1,412
12	C	IBR252-02	M	51	Myocardial infarct	18	L	1,450
13	C	BTB-3983	M	52	Heart atherosclerosis	13	R	1,430
14	C	B-6874	M	64	Cardiac arrest	28	R	1,250

PMI postmortem interval, H hemisphere, R right, L left

(a) qualitative abnormalities in reciprocal social interaction; (b) qualitative abnormalities in verbal and nonverbal communication; (c) restricted, repetitive and stereotyped patterns of behavior; and (d) abnormality of development evident at or before 36 months [69]. All 13 autistic subjects met ADI-R criteria for autism. For some subjects, the intellectual evaluation was available and was based on the Wechsler Intelligence Scale for Children III and the Woodcock-Johnson Tests of Achievement-Revised (Table 2). Eight subjects were diagnosed with intellectual disability, usually in the range from mild to severe (61%). Six of 13 autistic subjects had seizures (46%). In five cases, the age of onset of seizures was from 14 months to 5 years of age. A 23-year-old autistic male had only one seizure, which was reported as the cause of his death. In one child, an abnormal EEG was detected, but without seizures.

Several forms of challenging behaviors and behavioral disorders were noted, including self-injurious behavior (six cases, 46%), aggression (four cases, 31%), hyperactivity

(three cases, 23%), obsessive compulsive disorder (two cases, 12%) and depression and mania (a single case of each).

For three of the 13 autistic subjects, the list of high-confidence copy number variations identified both by quantiSNAP and Partek HMM computational algorithm was posted on the ATP portal by Drs. Steve Scherer and Richard Wintle from The Center for Applied Genomics, Toronto. The copy number variations detected in the three autistic subjects do not differ from those commonly observed [75], except for the loss of 25,505 kb within Neuropeptide S Receptor 1 (NPSR1) gene at 7p15–p14 detected in a 22-year-old autistic male (B-6337). NPSR1 has not been linked to autism in the genomic reports [103, 112]; however, an association of NPSR1 copy number variation with allergies has been reported [11] that might be linked to the patient's history of allergies.

Originally, 38 brains, including 20 brains of autistic and 18 brains of control subjects, were assigned to this project. However, application of the clinical and neuropathological

Table 2 Behavioral and neurological signs, and the type and topography of developmental abnormalities

Brain bank #	Psychiatric disorders and neurological symptoms	Mental retardation (MR)	Seizures age of onset	Type and topography of developmental abnormalities
IBR425-02	Hyperactivity. Tantrums. Self-injurious behavior	–	–	No changes
UMB-1627	Aggression	–	–	Focal neuronal heterotopia in white matter of the anterior cingulate gyrus
B-6403	–	–	14 months	Subependymal nodular dysplasia in the wall of the occipital horn of the lateral ventricle. Two periventricular nodular heterotopias (2 and 4 mm in diameter) near the frontal horn of the lateral ventricle. Tuber-like expansion of the tail of caudate nucleus into the lumen of the ventricle. Flocculonodular dysplasia
B-5666	–	–	Abnormal EEG; no seizures	Cortical dysplasia in the middle and inferior temporal gyri with focal dyslamination, clustering of dystrophic neurons and severe local neuronal deficits. Several focal dysplastic changes within CA. Flocculonodular dysplasia affecting almost entire lobe
B-5342	Pervasive developmental disorder. Hyperlexia	Mild MR	4.5 months	Focal cortical dysplasia. Dysplasia of the granule layer of the dentate gyrus. Subcortical heterotopia in the inferior frontal gyrus. Heterotopia in vermis and in cerebellar white matter
B-5535	Hyperactivity. Self-injurious behavior including head-banging	Moderate to severe MR	2 years	Thickening of the subependymal cell layer. Focal dysplasia within CA1 pyramidal layer with neuronal deficit, abnormal neuron morphology and spatial orientation. Multifocal dysplasia of the dentate gyrus with distortion of the shape of granule and molecular cell layers. Focal dysplasia within vermis
B-6115	Sensory integration disorder	–	–	Flocculonodular dysplasia affecting the majority of lobe volume. Cortical angioma
UMB-1638	ADHD	Moderate MR	5 years	Focal dysplasia within CA1 with diffuse neuronal deficit but without glial activation
B-6337	Obsessive compulsive disorder. Mania. Tourette syndrome. Self-injurious behavior	MR	–	Minor focal flocculonodular dysplasia
IBR93-01	Hyperactivity. Aggressive and self-injurious behavior	Severe MR	23 years	Focal dysplasia within islands in the entorhinal cortex. Pineal gland cysts
B-6212	Obsessive compulsive disorder. Depression, aggression, and anxiety	Severe MR	–	Several areas of focal cortical dysplasia within frontal cortex and insula with local loss of vertical and horizontal organization. Merger of ventral portion of the claustrum with insula. Flocculonodular dysplasia
B-6276	Aggression and self-injurious behavior, anxiety and agitation	Moderate MR	–	Focal dysplasia within CA1 sector with focal neuronal deficit. Heterotopia within stratum oriens. Flocculonodular dysplasia affecting approximately 70% of the lobe
B-7090	Disturbed movement coordination (walking like drunk)	MR	3 years	Three focal dysplasias in the frontal cortex. Dysplasia of layers 1–3 in the entorhinal cortex with missing numerous islands of the stellate neurons. Severe hypoplasia of cerebellar lobes 1–4. Reduced convolutions within dentate nucleus

Developmental abnormalities in brains of autistic subjects

exclusion criteria reduced the size of the cohort to 27 brains. Based on the results of the ADI-R, two cases were excluded, including one case diagnosed with atypical autism, and one that did not meet ADI-R criteria. Based on postmortem evaluation, five more autistic cases were excluded: one due to severe postmortem autolytic changes, three due to severe global hypoxic encephalopathy related to the mechanism of death, and one due to multiple microinfarcts. Moreover, four brains of control subjects were disqualified due to severe postmortem autolysis. In all these brains, neuronal loss, changes of neuronal size and shape, and gliosis were so severe that they masked and distorted the qualitative and quantitative characteristics of the developmental alterations associated with autism.

Brain tissue preservation

Brains of 13 autistic and 14 age-matched control subjects were examined by postmortem MRI and neuropathologically. The postmortem interval (PMI) varied, ranging from 6 to 27.8 h in the control group (16 h on average; SD 6 h) and from 8 to 30 h in the autistic group (20 h on average; SD 12 h). The median PMI was 15 h.

The brain hemispheres were removed using standard techniques, exercising extra care to avoid damaging the brain tissue. The brain was weighed in the fresh state. The fresh brain was sagittally cut through the corpus callosum and brainstem. Half of the brain was fixed in 10% buffered formalin. Following at least 3 weeks of fixation, the brain hemisphere was scanned using MRI. The aim of the MRI application was to determine the type of developmental changes detectable by MRI and to microscopically characterize MRI findings. All brains within this project were scanned (L.A.S.L.) using a standardized protocol (established and implemented for this and for other postmortem MRI studies by L.A.S.L. and M.L.). MRI scans were acquired on a 1.5 T GE Signa Imager (General Electric, Milwaukee, USA). The research scan consisted of a 124-slice T1-weighted fast gradient echo acquired in a coronal orientation perpendicular to the long axis of the hippocampus with a 1.5-mm slice thickness, which encompassed the entire brain hemisphere without gaps or wrap artifacts (FOV = 25 cm; NEX = 1; matrix = 256×192 ; TR = 35 ms; FA = 60°). All file names were assigned sequential code numbers, and demographic information was removed from image headers [9]. MRI scans were first screened in a diagnosis-blind manner, and the brains with abnormalities were re-evaluated by both radiologists and neuropathologists to determine the topography, type, and size of lesions detected with both methods.

The brain hemisphere was fixed with 10% buffered formalin. Formalin was washed out from the tissue during an overnight tap water rinsing. Brains were dehydrated

using a series of increasing ethyl alcohol concentrations (50% ethanol 3 days; 70% ethanol 4 days; 80% ethanol 3 days; 95% ethanol 4 days). The brain hemisphere was embedded in 8% celloidin [53]. During hardening, celloidin blocks were exposed to chloroform vapors for approximately 2.5 weeks, and celloidin blocks were then stored in 70% ethanol. For sectioning, the block was attached to the block holder with 10–15 ml of 8% celloidin. To fasten adhesion of the block to the holder, the block with the holder attached was immersed in 70% ethanol overnight. Serial 200- μ m-thick sections were separated with filter paper and stored in 70% ethanol. For the four control and four brains of autistic subjects, alternative series of 200- and 50- μ m-thick sections were preserved. To ensure the same probability of detection of changes in each case, every 200- μ m-thick section, with a distance 1.2 mm, was used in this project. Sections were washed in water for 2–3 h, stained with cresyl violet (CV) and mounted with Acrytol.


One neuropathologist (I.K.) examined, in a blind-to-diagnosis fashion, on average 120 hemispheric CV-stained sections per case with a 1.2-mm distance between sections. Two-step screening included examination at low magnification ($28\times$) using Zeiss DL2 Documator and microscopic examination using objective lenses from $5\times$ to $100\times$. Two other neuropathologists (T.W. and J.W.) examined all histological slides for which pathology was detected during the primary screening. The defects of neurogenesis, neuronal migration, and dysplastic changes that they detected were summarized in this report.

Tissue acquisition for this program project is based on individual tissue transfer agreements between the program project's principal investigator and several tissue banks: (a) the NICHD Brain and Tissue Bank for Developmental Disorders at the University of Maryland, (b) the Harvard Brain Tissue Resource Center and (c) the Brain Bank for Developmental Disabilities and Aging of the NYS Institute for Basic Research in Developmental Disabilities. Each brain hemisphere number given by the institution that received the donation was used as the only identifier of clinical records and tissue samples. Brain Bank identification of tissue samples is listed in Tables 1 and 2 to keep non-overlapping records of results of examination of brains in different projects and research groups. The Institutional Review Board of the New York State Institute for Basic Research in Developmental Disabilities approved the methods applied in this study.

Results

Neuropathological evaluation of serial coronal hemispheric sections from the cerebral and cerebellar hemispheres of

Fig. 1 Nodules in the wall of the lateral ventricle detected in postmortem MRI (**a**) in the brain of a 7-year-old male diagnosed with autism (B-6403) revealed features of subependymal nodular dysplasia (SND; **b**) in examination of CV-stained sections. **c** Numerous large and small nodules (*arrow*) dispersed within subependymal cell layer. They contained a few pyramidal-like neurons (**d**) and numerous poorly differentiated cells (**e**). Tuber-like expansion of the caudate nucleus (*arrow*) into the ventricle lumen is shown in MRI (**f**) and in CV-stained section (**g**). **g** A thick subependymal cell layer above and below (*two arrows*) the caudate nucleus (CN), and the absence of matrix in the tuber-like area. Under ependymal (E) cap of the caudate nucleus (CN) tuber-like expansion, small poorly differentiated neurons are present (**h**)



13 autistic and 14 control subjects revealed more details characterizing the topography and severity of changes than did standard sampling of brains for routine neuropathological evaluation. A broad range of changes was found. Developmental abnormalities included subependymal nodular dysplasia, heterotopia and very common dysplastic changes within the neo- and archicortex, hippocampus and cerebellum in 12 of 13 examined brains of the autistic subjects (92%) (Table 2). The general result of these developmental defects was a multifocal disorganization of gray and white matter. The developmental pathology observed in control brains was limited to one cerebellar dysplasia.

Alterations of the subependymal cell layer and subependymal nodular dysplasia

In two autistic subjects, there was a several-fold local increase in the thickness of the subependymal cell layer. Numerous subependymal nodules were found within a pathologically thickened subependymal cell layer, in the wall of the occipital horn of the lateral ventricle of a 7-year-old male, which reflects a subependymal nodular dysplasia (Fig. 1a–e). Nodules occupied 13.3 mm of the caudal portion of the occipital horn of the lateral ventricle. The diameter of round/oval nodules varied in size from 285 to 3,310 μm . While the smallest nodules were dispersed

within the subependymal cell layer, the large nodules expanded partially into the white matter, and partially into the lumen of the ventricles and were detectable on MRI (Fig. 1a) and CV-stained histological sections (Fig. 1b, c). The effect was narrowing of the ventricle and the tuberous appearance of the ventricular wall. There were large tubers that contained dysplastic neurons with a partially modified morphology of pyramidal, multipolar or bipolar large neurons (Fig. 1d) and irregularly shaped medium and small neurons. Neurons in the small nodules were small and poorly differentiated (Fig. 1e). In the large nodules, several hypocellular areas were observed. The nodules were free of oval or polygonal giant cells or ballooned glial cells, as well as signs of calcification.

In the brain with the subependymal nodular dysplasia, an abnormal tuberous expansion of the caudate nucleus was detected on MRI (Fig. 1f) and in histological sections (Fig. 1g, h). Only the ependyma separated the tuber-like expansion of the caudate from the ventricle lumen. The very thick subependymal cell layer that was present close to the caudate was substituted by loosely arranged and poorly differentiated neurons in the affected area.

Heterotopia

Heterotopias were found in the brains of four autistic subjects and no control subjects. The topography of the lesions was different in each case. Subcortical heterotopias were detected in the white matter of the anterior cingulate gyrus of a 5-year-old (Fig. 2a, b) and in the inferior frontal gyrus in an 11-year-old subject. Periventricular heterotopias were detected near the wall of lateral ventricle in 7-year old subject (Fig. 2c, d). A single heterotopia was noted in the stratum oriens of the hippocampus. In the cerebellum of the 11-year-old subject, heterotopias were detected in the vermis and the cerebellar white matter close to the dentate nucleus (Fig. 2f–h). These defects of migration were observed in two brains as a single aggregate of gray matter, in one brain as two aggregates and in one brain as three lesions measuring from 1 to 3 mm in diameter. Subcortical and periventricular heterotopias comprised poorly differentiated oval or multipolar neurons without spatial orientation (Fig. 2a) or had a distorted laminar organization (Fig. 2e). Cerebellar heterotopias had a distorted morphology of the granule and molecular layers with a few dispersed Purkinje cells (Fig. 2g, h).

Dysplasia within neocortex and archicortex, hippocampus and cerebellum

The multifocal neocortical dysplasia detected in four brains of autistic subjects (31%) was associated with a local loss

of vertical and horizontal organization of the neocortex, formation of abnormal layers, loss of orientation of neurons (Fig. 3a, b) and thickening of the affected portion of the cortical ribbon. A focal dysplasia in the entorhinal cortex, observed mainly in the second layer with a local absence of islands and/or reduced number of neurons, was found in the 23-year-old and the 60-year-old autistic subjects (15%) (Fig. 3c, d). A lack of giant multinuclear neurons and large, ballooned glial cells typical of focal cortical dysplasia indicated that the observed developmental changes in neocortex and archicortex reflect a more subtle cortical malformation, classified usually as focal cortical microdysgenesis.

Two types of changes were observed in the dentate gyrus. An abnormal migration of granule neurons into the molecular layer resulted in the formation of an additional fragmentary granule cell layer (Fig. 3e). In other areas, granule cells formed irregular circles and loops (Fig. 3f).

In the CA1 sector of a 13-year-old male, several areas of dysplastic changes with a significant deficit of pyramidal neurons without gliosis were found (Fig. 3g). In affected areas, the size and shape of neurons varied over a wide range. Pyramidal neurons were very rare, whereas small irregular or oval-shaped, poorly differentiated neurons prevailed (Fig. 3h). In the dysplastic area in the CA1 sector of the 56-year-old autistic subject, an opposite trend was present, with thickening of the pyramidal layer and an increased packing of dysplastic neurons (not shown).

The most common developmental abnormality within the cerebellum was dysplasia, which was detected in seven autistic subjects (54%) and in the cerebellum of one control subject.

Flocculonodular dysplasia (Fig. 4a, b), usually affecting the entire nodule, was found in the cerebellum in six autistic subjects (46%). In the dysplastic areas, a thin granule layer formed the labyrinth, which was mixed with irregular islands of the molecular layer. Clusters of granule cells and a few Purkinje cells were dispersed within the distorted molecular layer. The only developmental abnormality detected in the control group was flocculonodular dysplasia in the cerebellum of the 51-year-old control subject (not shown). Local cortical dysplasia was also detected within the vermis of the 13-year-old autistic male. In the affected area, the cytoarchitecture of the molecular and granule layers and the Purkinje cells was completely disorganized (Fig. 4c, d).

In the cerebellum of the 60-year-old autistic male, severe hypoplasia affected lobes 1–4 (Fig. 4e). The thickness of the molecular and granular layer was decreased by almost 50% in comparison to that of the non-affected areas (Fig. 4e, f). The number of Purkinje cells was significantly reduced in the hypoplastic area. Hypoplastic changes within the portion of cerebellar cortex were observed,




Fig. 2 Large subcortical heterotopia within anterior cingulate gyrus in a 5-year-old autistic child (UMB-1627) (**a**) contained dysplastic neurons without spatial orientation (**b**). Periventricular heterotopia near the frontal horn of the lateral ventricle (**c** MRI, **d**, **e** CV-stained section) shows a structure resembling molecular, granule and

pyramidal layers in a 7-year-old autistic subject (B-6403). MRI (**f**), low (**g**) and large (**h**) magnification of heterotopia (*arrow*) with dysplastic granule (*G*) and molecular layer (*M*) detected within cerebellar white matter in an 11-year-old autistic subject (B-5342)

together with a significantly reduced convolution of the dentate nucleus (Fig. 4g).

Discussion

This neuropathological study revealed a broad spectrum of focal developmental abnormalities and pre- and perinatally acquired lesions in 92% of the brains of autistic subjects and striking inter-individual differences in the type and topography of changes. Evidence that different features of autism are caused by different genes associated with different brain regions [48] suggests a link between regional

developmental alterations in the brain and different components of the autistic phenotype.

Altered neurogenesis in autism

Increased brain mass in autistic children and some autistic adults [89], increase in the numerical density of neurons [13, 14], reduced size of neurons [7] and brain structure-specific delay of neuronal growth [111] indicate alterations in neuronal and brain growth in autistic individuals. The subventricular zone of the lateral ventricles [26] and the dentate gyrus [33] are active sites of neurogenesis in adult humans. Several of our findings support the hypothesis of





Fig. 3 Dysplastic changes within neocortex (**a, b**), entorhinal cortex (**c, d**), dentate gyrus (**e, f**) and the cornu Ammonis (**g, h**). Focal dysplasia in frontal cortex with loss of vertical and horizontal cytoarchitecture (*two arrows*) and abnormal (*arrowhead*) laminar organization (**a**). Dysplastic neurons within affected area (B-6212) (**b**). Microdysgenesis within the entorhinal cortex with deficit of stellate neurons in the islands (**c**) and normal morphology of islands in adjacent cortex (**d**) in 60-year-old autistic subject (B-7090).

Microdysgenesis of the dentate gyrus with dispersion of granule cells within the molecular layer (**e, arrow**) and distortion of the granule cell layer shape (**f, arrows**) in 13-year-old autistic male (B-5535). CA1 sector microdysgenesis with local deficit of pyramidal neurons (**g, arrow**) without markers of gliosis but with signs of poor differentiation of dysplastic abnormally arranged neurons (**h**) in 13-year-old autistic subject (B-5535)

altered neurogenesis in autistic subjects. The increased thickness of the subependymal cell layer, subependymal nodular dysplasia, abnormal growth of the dentate nucleus and dysplasia of the granule layer in the dentate gyrus, detected in this study, appear to be signs of abnormal neurogenesis in the brains of three autistic subjects.

Subependymal nodules were reported in approximately 80% of patients with tuberous sclerosis, a disorder that is highly associated with epilepsy, autism and mental retardation [73]. Tuberous sclerosis nodules were detected in one fetus [12], suggesting that fetal development of subependymal nodules can lead to the early onset of epilepsy

Fig. 4 Flocculonodular dysplasia in cerebellum of 56-year-old autistic subject (B-6276) (**a**) with thin irregular granule (*G*) and molecular (*M*) layer. **b** Dysplastic granule layer (*G*), ectopic granule cells (arrow) in the molecular layer, and loosely dispersed Purkinje cells (*P*) (B-6276). Cortical dysplasia within vermis of 13-year-old autistic male (**c**) with dysplastic granule neurons mixed with heterotopic (arrow) large cells (**d**) (B-5535). **e** Severe hypoplasia of cerebellar lobe 3 and unmodified lobe 6 (**f**), respectively, within the cerebellum of a 60-year-old autistic male (B-7090). In the affected region, the thickness of the hypoplastic molecular and granule cell layer was reduced by about 50%. Almost half of the dentate nucleus (*DN*) was less convoluted than the unaffected part (**g**)



that was diagnosed at the age of 14 months in a neuropathologically examined autistic male. The subependymal nodules detected in this autistic male's brain are partially similar to tubers seen in subjects diagnosed with tuberous sclerosis [24]. The cause of subependymal nodular dysplasia in the examined subject is unknown. In the reported subjects, bilateral periventricular nodules are linked to mutations of the filamin A (FLNA) gene located on chromosome Xp28. Filamin A is an actin-crosslinking protein that is essential for cell locomotion [16], and nodule formation might be related to a defect in cell migration. The presence of miniature nodules that were built of poorly differentiated small neurons within the subependymal cell layer and an increase in nodular size with signs of growth and differentiation of neurons suggests that neurogenesis, differentiation and maturation of neurons were in progress within the subependymal germinal matrix of the 7-year-old autistic child. This interpretation of subependymal nodule genesis is consistent with lineage studies demonstrating that cells in nodules express cellular markers that are typical for progenitors derived from the subventricular germinal zone [35, 67]. However, in contrast to the subependymal nodules seen in subjects with tuberous sclerosis,

in the examined autistic subject, the nodules seen were small (from 258 to 3,310 μm in diameter), and did not have the characteristic ovoid or polygonal giant cells, 80–150 μm in diameter, giant cells with multiple and peripherally displaced nuclei [25], or balloon cells, which are considered the sine qua non histopathological features of the cortical tubers and subependymal nodules observed in tuberous sclerosis [73].

The enlarged caudate nucleus detected in the brain of the 7-year-old autistic subject is consistent with MRI reports documenting an increased volume of basal ganglia, including the caudate, in autism [54, 55, 66, 99]. A disproportionate increase of the caudate nucleus volume [66] suggests that in brains of some autistic individuals, extended neurogenesis within the subependymal cell layer may contribute to abnormal growth of the caudate nucleus. A similar process has been observed in the brains of people with Huntington disease, showing enhanced neurogenesis in the subependymal layer and suggesting renewal of the neuronal population in a degenerating caudate nucleus [26]. The caudate nucleus is a part of the fronto-striatal network involved in several functional domains that are impaired in autism, including lower order repetitive motor

behavior; intense circumscribed patterns of interests and higher order rituals and compulsions [41], and defects in cognitive functions [19, 109], planning and problem-solving skills [78, 98], short- and long-term memory [40] and learning [88].

Defective migration in autism

Heterotopia is a sign of altered migration leading to an abnormal distribution of gray matter nodular masses with disorganized or rudimentary lamination within the periventricular area (periventricular heterotopia) or subcortical white matter (subcortical heterotopia) [2]. In the examined cohorts, heterotopias were detected in the brains of four autistic subjects and in the brain of one control subject. Heterotopias are associated with mutations in the filamin 1 gene (FLNA1) [39, 46] and the chromosome X-linked DCX gene that codes for doublecortin, a protein expressed during brain development in migrating neurons, and in the cortical plate [29, 44, 45], which is involved in the formation of the microtubules necessary for neuronal migration [15]. Periventricular nodular heterotopia has been reported to be associated with pharmaco-resistant seizures in 80–90% of patients [31]. In the examined cohort, two periventricular heterotopias were detected in the brain of a child with subependymal nodular dysplasia and seizures diagnosed at 14 months of age (B-6403). Early onset epilepsy, diagnosed at the age of 4.5 months, might be related to the multiple heterotopias found within the frontal inferior gyrus, vermis and cerebellar white matter, coexisting with a focal cortical dysplasia and dentate gyrus dysplasia (B-5342).

Cortical, hippocampal and cerebellar dysplasia in autism

The most common form of developmental changes detected in the examined brains was focal dysplasia, which was observed in 11 (85%) of the autistic subjects. The morphology of focal dysplasias appears to reflect signs of abnormal migration, neuronal immaturity and altered cell arrangement, resulting in focal distortion of cytoarchitecture. In spite of similarities, the dysplastic changes in the neocortex and archicortex, dentate gyrus and cornu Ammonis and cerebellum also reveal a brain structure-specific pattern of dysplastic changes in autism.

Dysplasias encompass a spectrum of changes ranging from a mild form of cortical disruption, without cellular abnormalities, to the most severe form with cortical dyslamination, with abnormal morphology of neurons and astrocytes [93, 96, 107]. Focal cortical dysplasias with giant neurons and balloon cells [107, 113] are histopathologically similar to tubers containing giant cells in tuberous

sclerosis complex [25, 73], suggesting a common pathogenic basis [113]. However, activation of the mammalian target of rapamycin (mTOR) pathway observed in the tuberous sclerosis complex is not present in focal cortical dysplasia [8, 80]. The giant neurons and ballooned cells, which are histopathological features of tuberous sclerosis and focal cortical dysplasia, were absent both in the subependymal nodules and in the focal cortical dysplasia observed in the examined autistic cohort. These findings suggest that in spite of similarities, the pathomechanisms of developmental alterations are different in the examined autistic subjects than those in tuberous sclerosis heterotopias or focal cortical dysplasia. The development of the giant neuron- and balloon cell-free dysplasias observed in the autistic subjects might be related to differences in cause and/or mechanism. The detection of changes similar to focal cortical dysplasia in association with prenatal ischemia [65] or in shaken infant syndrome [74] may support these speculations.

Ectopias and dysplastic changes were reported in the brains of autistic subjects, by several groups [4, 62–64, 91]. Bailey et al. [4] detected olivary dysplasia in the brain of three of the five autistic subjects, and ectopic neurons related to the olivary complex in two cases. Moreover, in the brains of four autistic subjects, cortical dysgenesis was found. In the brains of the autistic subjects, a strikingly consistent finding was cingulate cortex disordered lamination [62–64, 100]. A recent study of the cingulate cortex of nine autistic subjects revealed a developmental malformation with irregular lamination in three cases, and an increased number of neurons within the subcortical white matter in two [100]. Simms et al. [100] suggest that the excessive number of neurons in the subcortical white matter reflects the lack of proper resolution of the transient zone in the developing brain of autistic subjects. Studies by Fatemi et al. [37, 38] link the migration and lamination defects to a striking reduction of reelin (by 40%) and Bcl-2 (by 34–51%) in the brains of autistic subjects. Our studies along with others' suggest that in the majority of autistic subjects, heterotopias and dysplastic changes are the local sign of general developmental defects of migration with a broad spectrum of topographic, morphological, and functional outcomes.

In the examined brains of autistic subjects, signs of neuronal immaturity were a common finding. Failure of maturation of neuronal precursors caused by altered expression of cytoskeletal proteins and loss of neuronal polarity results in defects in migration to the destined layer and in incorrect vertical and horizontal orientation [93]. The immaturity of dysplastic neurons is reflected in the expression of a variety of proteins and mRNA that are not present in mature neurons an altered expression of developmentally regulated cytoskeletal elements [3, 23, 61, 76],

which are known to be crucial for dendrite arborization, spine formation, axon outgrowth and maintenance of cell size and shape. Reduced cell size, dendritic arborization and spine expression are characteristic of dysplastic neurons [6, 93]. Cortical dysplasias are the most epileptogenic lesions of the brain [107] and are observed in up to 25% of all epileptic surgeries [102]. More subtle cortical malformations or dysgenesis encountered in adults with epilepsy may lack the histological criteria for focal cortical dysplasia. They have been described as mild cortical dysplasia or microdysgenesis [77].

Microdysgenesis within the entorhinal cortex of the 23- and the 60-year-old autistic subjects in the examined cohort is unique because the selective deficit of neurons was limited almost exclusively to the stellate neurons in the second layer. It is possible that the observed dysgenesis is a result of defective migration of neurons to their intended destinations. The presence of a thicker molecular layer and the deeper location of islands in the entorhinal cortex of subjects with schizophrenia were previously interpreted as evidence that the stellate neurons do not reach their destinations during development, probably due to abnormal migration [36, 57]. Studies indicating the involvement of *reelin* and *Bcl2* genes in the pathogenesis of schizophrenia [37, 47, 60] and the reduced expression of *reelin* and *Bcl2* in people with autism suggest that these two genes play a role in abnormal brain development and contribute to the structural and functional anomalies seen in autism and schizophrenia [37].

The distortion of dentate gyrus development detected in two autistic subjects was reflected in granule cell migration into the molecular layer and formation of an additional granule cell layer. Distortion of the shape of the dentate granule cell layer with the formation of irregular circles and loops appears to be another piece of evidence suggesting abnormal neuronal migration and networking. Numerous factors up-regulate neurogenesis in the hippocampus [32], including seizures [70, 71], antidepressant drugs [59, 72] and lithium [18]. Several areas of dysplastic changes with significant deficits of pyramidal neurons were found in the CA1 sector in three autistic subjects, but thickening of the pyramidal layer and an increased packing of dysplastic neurons in the CA1 sector of the 56-year-old subject suggests a diversity of CA dysplasia patterns in autism. The lack of gliosis indicates that the observed pathology is a sign of microdysgenesis rather than an effect of hypoxic neuronal loss. A significant deficit of mature pyramidal neurons and the presence of small irregular or poorly differentiated oval neurons suggest the defect of neuronal maturation in autism.

We report a spectrum of focal developmental changes seen in the cerebellum of eight autistic subjects, including nodular (lobe X) [97] dysplasia in the cerebellum in five,

vermal dysplasia in one, severe focal hypoplasia in one, and heterotopias in one other subject. The presence of heterotopias only in one control subject is evidence of a strong tendency for focal developmental changes of cerebellar microarchitecture that were present in 61% of the autistic subjects. Flocculonodular dysplasia affecting almost the entire lobe indicates that mechanisms leading to focal dysplasia, which were present in five (38%) of the autistic subjects, show extremely strong topographic predilection. The observed focal dysplasia was associated with profound local disorganization of granule cells, Purkinje cells and molecular layers limited to a small cerebellar compartment receiving major projections from the vestibular complex involved in the oculomotor and postural system. Similar cerebellar dysplastic changes classified as heterotaxias (clusters of poorly organized mixed cells) were identified in 14% of normal infants but in 83% of infants with trisomy of different chromosomes [92]. The presence within the dysplastic nodule of both GABAergic Purkinje cells produced from the cerebellar ventricular zone, and the glutamatergic granule neurons produced from the rhombic lip, and the preservation of the cytoarchitecture in the adjacent cerebellar folia suggest that the final steps of migration and networking are disturbed mainly or exclusively in the nodule of the majority of autistic subjects. The characteristic feature distinguishing lobule X from the other lobules is the abundance of the transcription factor *Tbr2* positive unipolar brush cells (UBCs) [30, 34], which amplify inputs from vestibular ganglia and nuclei, by spreading and prolonging excitation within the internal granular layer [84]. Abnormal networking of Purkinje cells, granule neurons, and UBCs may contribute to altered cerebellar coordination of locomotion and motor learning and planning, as well as of higher cognitive processing [58]. Flocculonodular dysplasia appears to be another sign of the mosaic of local developmental defects, most likely predetermined by the spatial patterning of germinal zones in developing rhombic lip [110], and coexisting with more general developmental defects resulting in the accelerated growth of the brain in early childhood [89], minicolumn pathology [13, 14], reduced neuron volume [7, 108, 111], and desynchronized neuronal growth in many brain regions [111] observed in autism.

Identification of sub-groups with signs of hyperplasia, hypoplasia and normal-sized cerebellum [95] reflects the heterogeneity of the autistic population. Piven et al. [87] reported that cerebellar volume correlates with an increased total brain volume. In the majority of autistic subjects, reduced size of the cerebellar hemisphere is observed [42, 82], but this trend is not detectable in cohorts of high-functioning autistic individuals [56]. Regional hypoplasia affects the vermis in autistic individuals relatively often [20, 22, 52] and may be associated with the

deficits in attention-orienting [49, 104], stereotypic behavior and reduced exploration observed in autism [86]. In the examined autistic cohort, selective and severe hypoplasia of lobes 1–4 associated with hypoconvolution of a large portion of the dentate nucleus appears to correspond to clinically detected defects of movement coordination. These findings suggest that differences in the type, topography and severity of cerebellar developmental defects may contribute to different clinical manifestations.

In the 4–7-year-old autistic children examined in this study, the volume of the Purkinje cells was 38% smaller than that of the age-matched control group [111]. Moreover, it has been reported that Purkinje cells of the autistic subjects revealed a 40% decrease in the expression of glutamic acid decarboxylase 67 (GAD67) mRNA [114]. In autism, the basket cells provide an increased GABAergic feed-forward inhibition to Purkinje cells. The result could be disruption in the timing of Purkinje cell firings and altered inhibition of the cerebellar nuclei, which could directly affect cerebello-cortical output and contribute to the changes in motor behavior and cognition observed in autism [115]. These findings and the reduced volume (by 26%) of the neurons of the dentate nucleus seen in the 4–7-year-old autistic children [111] suggest that in autism, interactions between the Purkinje cells and dentate nucleus are modified on the structural, molecular and functional levels.

The (a) detected changes within the subependymal cell layer with subependymal nodular dysplasia, (b) subcortical and periventricular heterotopias and (c) neocortex, archicortex, dentate gyrus, cornu Ammonis and cerebellar dysplasia reflect focal modification of neurogenesis, migration and alterations of the cytoarchitecture of brain cortex, subcortical structures and cerebellum in autism. Detection of dysplastic changes only in one control brain and of the broad spectrum of focal developmental alterations in the brains of 92% of the autistic subjects indicates that focal changes are a reflection of global developmental abnormalities and that regional changes may have their own contribution to the clinical heterogeneity of autism.

Acknowledgments This study was supported in part by funds from the New York State Office of Mental Retardation and Developmental Disabilities, a grant from the Department of Defense Autism Spectrum Disorders Research Program (AS073234, Program Project; J.W., T.W., A.C.), a grant from Autism Speaks (Princeton, NJ), and grant R01 HD43960 (J.W.) from the National Institutes of Health, National Institute of Child Health and Human Development. Tissue and clinical records' acquisition was coordinated by the Autism Tissue Program (Princeton, NJ; Directors: Jane Pickett, Ph.D. and Daniel Lightfoot, Ph.D.). The tissue was obtained from the Harvard Brain Tissue Resource Center, Belmont, MA, supported in part by PHS grant number R24-MH 068855; the National Institute of Child Health and Human Development Brain and Tissue Bank for Developmental Disorders at the University of Maryland, Baltimore; and the Brain Bank for Developmental Disabilities and Aging of the New York

State Institute for Basic Research in Developmental Disabilities, Staten Island, NY. We thank Drs. Helmut Hainsen and Christoph Schmitz for help in implementation of the celloidin protocol, and Mrs. Jadwiga Wegiel, Cathy Wang and En Wu Zhang for histology. We are deeply indebted to the families of the tissue donors who have made this study possible.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. American Psychiatric Association (2000) Diagnostic and statistical manual of mental disorders DSM-IV-TR. American Psychiatric Association, Washington, DC
2. Andrade DM (2009) Genetic basis in epilepsies caused by malformations of cortical development and in those with structurally normal brain. *Hum Genet* 126:173–193
3. Avila J, Dominguez J, Diaz-Nido J (1994) Regulation of microtubule dynamics by microtubule-associated protein expression and phosphorylation during neuronal development. *Int J Dev Biol* 38:13–25
4. Bailey AP, Luthert A, Dean B et al (1998) A clinicopathological study of autism. *Brain* 121:889–905
5. Bailey AR, Giunta BN, Obregon D et al (2008) Peripheral biomarkers in autism: secreted amyloid precursor protein- α as a probable key player in early diagnosis. *Int J Clin Exp Med* 1:338–344
6. Barth PG (1987) Disorders of neuronal migration. *Can J Neurol Sci* 14:1–16
7. Bauman ML, Kemper TL (1985) Histoanatomic observations of the brain in early infantile autism. *Neurology* 35:866–867
8. Baybis M, Yu J, Lee A, Golden JA et al (2004) mTOR cascade activation distinguishes tubers from focal cortical dysplasia. *Ann Neurol* 56:478–487
9. Bobinski M, de Leon MJ, Convit A et al (1999) MRI of entorhinal cortex in mild Alzheimer's disease. *Lancet* 353:38–40
10. Boddaert N, Zilbovicius M, Philipe A et al (2009) MRI findings in 77 children with non-syndromic autistic disorder. *PLoS One* 4:e4415
11. Bruce S, Nyberg F, Melén E et al (2009) The protective effect of farm animal exposure on childhood allergy is modified by NPSR1 polymorphisms. *J Med Genet* 46:159–167
12. Carlson BA, Houser OW, Gomez MR (1999) Brain imaging in the tuberous sclerosis complex. In: Gomez M, Sampson J, Whittemore V (eds) *Tuberous sclerosis complex*, 3rd edn. Oxford University Press, New York, pp 85–93
13. Casanova MF, Buxhoeveden DP, Switala AE, Roy E (2002) Minicolumnar pathology in autism. *Neurology* 58:428–432
14. Casanova MF, van Kooten IAE, Switala EH et al (2006) Minicolumnar abnormalities in autism. *Acta Neuropathol* 112:287–303
15. Caspi M, Atlas R, Kantor A, Sapir T, Reiner O (2000) Interaction between LIS1 and doublecortin, two lissencephaly gene products. *Hum Mol Genet* 9:2205–2213
16. Chang BS, Walsh CA (2009) The genetic basis of human cerebral cortical malformations. In: Runge MS, Patterson C (eds) *Principles of molecular medicine*. Human Press Inc, Totowa, NJ, pp 1073–1079
17. Chauhan A, Chauhan V (2006) Oxidative stress in autism. *Pathophysiology* 13:171–181

18. Chen G, Rajkowska G, Du F, Seraji-Bozorgzad N, Manji NH (2000) Enhancement of hippocampal neurogenesis by lithium. *J Neurochem* 75:1729–1734
19. Chow TW, Cummings JL (1999) Frontal-subcortical circuits. In: Miller BL, Cummings JL (eds) *The human frontal lobes: functions and disorders*. Guilford Press, New York, pp 3–26
20. Ciesielski KT, Harris RJ, Hart BL, Pabst H (1997) Cerebellar hypoplasia and frontal lobe cognitive deficits in disorders of early childhood. *Neuropsychologia* 35:643–655
21. Courchesne E, Hesselink JR, Jernigan TL, Yeung-Courchesne R (1987) Abnormal neuroanatomy in a nonretarded person with autism. Unusual findings with magnetic resonance imaging. *Arch Neurol* 44:335–341
22. Courchesne E, Yeung-Courchesne R, Press GA, Hesselink JR, Jernigan TL (1988) Hypoplasia of cerebellar vermal lobules VI and VII in autism. *N Engl J Med* 318:1349–1354
23. Crino PB, Trojanowski JQ, Eberwine J (1997) Internexin, MAP1B, and nestin in cortical dysplasia as markers of developmental maturity. *Acta Neuropathol* 93:619–627
24. Crino PB, Henske EP (1999) New developments in the neurobiology of the tuberous sclerosis complex. *Neurology* 53:1384–1390
25. Crino PB, Miyata H, Vinters HV (2002) Neurodevelopmental disorders as a cause of seizures: neuropathologic, genetic, and mechanistic considerations. *Brain Pathol* 12:212–233
26. Curtis MA, Penney EB, Pearson J, Dragunow M, Connor B, Faull RL (2005) The distribution of progenitor cells in the subependymal layer of the lateral ventricle in the normal and Huntington's disease human brain. *Neuroscience* 132:777–788
27. Damasio H, Maurer RG, Damasio AR, Chui HC (1980) Computerized tomographic scan findings in patients with autistic behavior. *Arch Neurol* 37:504–510
28. Department of Health and Human Services (2007) Morbidity and mortality weekly report. In: Department of Health and Human Services, Centers for Disease Control and Prevention, pp 1–28
29. des Portes V, Francis F, Pinard JM et al (1998) Doublecortin is the major gene causing X-linked subcortical laminar heterotopia (SCLH). *Hum Mol Genet* 7:1063–1070
30. Diño MR, Willard FH, Mugnaini E (1999) Distribution of unipolar brush cells and other calretinin immunoreactive components in the mammalian cerebellar cortex. *J Neurocytol* 28:99–123
31. Dubeau F, Tampieri D, Lee N et al (1995) Periventricular and subcortical nodular heterotopia: a study of 33 patients. *Brain* 118:1273–1287
32. Duman RS, Nakagawa S, Malberg J (2001) Regulation of adult neurogenesis by antidepressant treatment. *Neuropsychopharmacology* 25:836–844
33. Eriksson PS, Perfilieva E, Bjork-Eriksson T et al (1998) Neurogenesis in the adult human hippocampus. *Nat Med* 4:1313–1317
34. Englund C, Kowalczyk T, Daza RAM et al (2006) Unipolar brush cells of the cerebellum are produced in the rhombic lip and migrate through developing white matter. *J Neurosci* 26:9184–9195
35. Ess KC, Kamp CA, Tu BP, Gutmann DH (2005) Developmental origin of subependymal giant cell astrocytoma in tuberous sclerosis complex. *Neurology* 64:1446–1449
36. Falkai P, Schneider-Axmann T, Honer WG (2000) Entorhinal cortex pre-alpha cell clusters in schizophrenia: quantitative evidence of a developmental abnormality. *Biol Psychiatry* 47:937–943
37. Fatemi SH, Kroll JL, Stary JM (2001) Altered levels of Reelin and its isoforms in schizophrenia and mood disorders. *Neuroreport* 12:3209–3215
38. Fatemi SH, Stary JM, Halth AR, Realmuto GR (2001) Dysregulation of reelin and Bcl-2 proteins in autistic cerebellum. *J Autism Dev Disord* 31:529–535
39. Fox JW, Lamperti ED, Eksioglu YZ (1998) Mutations in filamin 1 prevent migration of cerebral cortical neurons in human periventricular heterotopia. *Neuron* 21:1315–1325
40. Fuh JL, Wang SJ (1995) Caudate hemorrhage: clinical features, neuropsychological assessments and radiological findings. *Clin Neurol Neurosurg* 97:296–299
41. Gabriels RL, Cuccaro ML, Hill DE, Ivers BJ, Goldson E (2005) Repetitive behaviors in autism: relationships with associated clinical features. *Res Dev Disabil* 26:169–181
42. Gaffney GR, Tsai LY, Kuperman S, Minchin S (1987) Cerebellar structure in autism. *Am J Dis Child* 141:1330–1332
43. Gillberg C, Coleman M (1996) Autism and medical disorders: a review of the literature. *Dev Med Child Neurol* 38:191–202
44. Gleeson JG, Allen KM, Fox JW (1998) Doublecortin, a brain-specific gene mutated in human X-linked lissencephaly and double cortex syndrome, encodes a putative signaling protein. *Cell* 92:63–72
45. Gleeson JG, Lin PT, Flanagan LA, Walsh CA (1999) Doublecortin is a microtubule-associated protein and is expressed widely by migrating neurons. *Neuron* 23:257–271
46. Gorlin JB, Henske E, Warren ST (1993) Actin-binding protein (ABP-280) filamin gene (FLN) maps telomeric to the color vision locus (R/GCP) and centromeric to G6PD in Xq28. *Genomics* 17:496–498
47. Guidotti A, Auta J, Davis JM (2000) Decrease in reelin and glutamic acid decarboxylase67 (GAD67) expression in schizophrenia and bipolar disorder: a postmortem brain study. *Arch Gen Psychiatry* 57:1061–1069
48. Happe F, Ronald A, Plomin R (2006) Time to give up on a single explanation for autism. *Nat Neurosci* 9:1218–1220
49. Harris NS, Courchesne E, Townsend J, Carper RA, Lord C (1999) Neuroanatomic contributions to slowed orienting of attention in children with autism. *Brain Res Cogn Brain Res* 8:61–71
50. Hashimoto T, Tayama M, Mori K, Fujino K, Miyazaki M, Kuroda Y (1989) Magnetic resonance imaging in autism: preliminary report. *Neuropediatrics* 20:142–146
51. Hashimoto T, Tayama M, Miyazaki M, Murakawa K, Kuroda Y (1993) Brainstem and cerebellar vermis involvement in autistic children. *J Child Neurol* 8:149–153
52. Hashimoto T, Tayama M, Murakawa K et al (1995) Development of the brainstem and cerebellum in autistic patients. *J Autism Dev Disord* 25:1–18
53. Heinsen H, Arzberger T, Schmitz C (2000) Celloidin mounting (embedding without infiltration)—a new, simple and reliable method for producing serial sections of high thickness through complete human brains and its application to stereological and immunohistochemical investigations. *J Chem Neuroanat* 20:49–59
54. Herbert MR, Ziegler DA, Deutsch CK (2003) Dissociations of cerebral cortex, subcortical and cerebral white matter volumes in autistic boys. *Brain* 126:1182–1192
55. Hollander E, Anagnostou E, Chaplin W (2005) Striatal volume on magnetic resonance imaging and repetitive behaviors in autism. *Biol Psychiatry* 58:226–232
56. Holtum JR, Minshew NJ, Sanders RS, Phillips NE (1992) Magnetic resonance imaging of the posterior fossa in autism. *Biol Psychiatry* 32:1091–1101
57. Honer WG, Bassett AS, Falkai P, Beach TG, Lapointe JS (1996) A case study of temporal lobe development in familial schizophrenia. *Psychol Med* 26:191–195
58. Ito M (2008) Control of mental activities by internal models in the cerebellum. *Nat Rev Neurosci* 9:304–313

59. Jacobs BL, Fornal CA (1999) Activity of serotonergic neurons in behaving animals. *Neuropsychopharmacology* 21:9S–15S
60. Jarskog LF, Gilmore JH, Selinger ES, Lieberman JA (2000) Cortical bcl-2 protein expression and apoptotic regulation in schizophrenia. *Biol Psychiatry* 48:641–650
61. Kaplan MP, Chin SS, Fliegner KH, Liem RK (1990) Alpha-internexin, a novel neuronal intermediate filament protein, precedes the low molecular weight neurofilament protein (NF-L) in the developing rat brain. *J Neurosci* 10:2735–2748
62. Kemper TL (1988) Neuroanatomic studies of dyslexia and autism. In: *Disorders of the developing nervous system: changing views on their origins, diagnosis, and treatments*. Alan R. Liss Inc, New York, pp 125–154
63. Kemper TL, Bauman ML (1993) The contribution of neuropathologic studies to the understanding of autism. *Behav Neurol* 11:175–187
64. Kemper TL, Bauman M (1998) Neuropathology of infantile autism. *J Neuropathol Exp Neurol* 57:645–652
65. Kremer S, De Saint MA, Minotti L et al (2002) Focal cortical dysplasia possibly related to a probable prenatal ischemic injury. *J Neuroradiol* 29:200–203
66. Langen M, Durston S, Staal WG, Palmen SJ, van Engelen H (2007) Caudate nucleus is enlarged in high-functioning medication-naïve subjects with autism. *Biol Psychiatry* 62:262–266
67. Lee A, Maldonado M, Baybis M et al (2003) Markers of cellular proliferation are expressed in cortical tubers. *Ann Neurol* 53:668–673
68. Lopez-Hurtado E, Prieto JJ (2008) A microscopic study of language-related cortex in autism. *Am J Biochem Biotechnol* 4:130–145
69. Lord C, Risi S, Lambrecht L et al (2000) The autism diagnostic observation schedule-generic: a standard measure of social and communication deficits associated with the spectrum of autism. *J Autism Dev Disord* 30:205–223
70. Madsen TM, Treschow A, Bengzon J, Bolwig TG, Lindvall O, Tingstrom A (2000) Increased neurogenesis in a model of electroconvulsive therapy. *Biol Psychiatry* 47:1043–1049
71. Malberg JE, Eisch AJ, Nestler EJ, Duman RS (2000) Chronic antidepressant treatment increases neurogenesis in adult rat hippocampus. *J Neurosci* 20:9104–9110
72. Manev H, Uz T, Smalheiser NR, Manev R (2001) Antidepressants alter cell proliferation in the adult brain in vivo and in neural cultures in vitro. *Eur J Pharmacol* 411:67–70
73. Marcotte L, Crino PB (2006) The neurobiology of the tuberous sclerosis complex. *Neuromol Med* 8:531–546
74. Marin-Padilla M, Parisi JE, Armstrong DL, Sargent SK, Kaplan JA (2002) Shaken infant syndrome: developmental neuropathology, progressive cortical dysplasia, and epilepsy. *Acta Neuropathol* 103:321–332
75. Marshall CR, Noor A, Vincent JB et al (2008) Structural variation of chromosomes in autism spectrum disorder. *Am J Human Gen* 82:477–488
76. Matus A (1988) Microtubule-associated proteins: their potential role in determining neuronal morphology. *Ann Rev Neurosci* 11:29–44
77. Meencke HJ, Janz D (1985) The significance of microdysgenesis in primary generalized epilepsy: an answer to the considerations of Lyon and Gastaut. *Epilepsia* 26:368–371
78. Mendez MF, Adams NL, Lewandowski KS (1989) Neurobehavioral changes associated with caudate lesions. *Neurology* 39:349–354
79. Miles JH, Takahashi TN, Bagby S et al (2005) Essential versus complex autism: definition of fundamental prognostic subtypes. *Am J Med Genet A* 135:171–180
80. Miyata H, Chiang AC, Vinters H (2004) Insulin signaling pathways in cortical dysplasia and TSC-tubers: tissue microarray analysis. *Ann Neurol* 56:510–519
81. Muller RA (2007) The study of autism as a distributed disorder. *Ment Retard Dev Disabil Res Rev* 13:85–95
82. Murakami W, Courchesne E, Press GA, Yeung-Courchesne R, Hesselink JR (1989) Reduced cerebellar hemisphere size and its relationship to vermal hypoplasia in autism. *Arch Neurol* 46:689–694
83. Newschaffer CJ, Fallin D, Lee NL (2002) Heritable and non-heritable risk factors for autism spectrum disorders. *Epidemiol Rev* 24:137–153
84. Nunzi MG, Birnstiel S, Bhattacharyya BJ, Slater NT, Mugnaini E (2001) Unipolar brush cells form a glutamatergic projection system within the mouse cerebellar cortex. *J Comp Neurol* 434:329–341
85. Palmen SJ, van Engelen H, Hof PR, Schmitz C (2004) Neuropathological findings in autism. *Brain* 127:2572–2583
86. Pierce K, Courchesne E (2001) Evidence for a cerebellar role in reduced exploration and stereotyped behavior in autism. *Biol Psychiatry* 49:655–664
87. Piven J, Saliba K, Bailey J, Arndt S (1997) An MRI study of autism: the cerebellum revisited. *Neurology* 49:546–551
88. Poldrack RA, Prabhakaran V, Seger CA, Gabrieli JD (1999) Striatal activation during acquisition of a cognitive skill. *Neuropsychology* 13:564–574
89. Redcay E, Courchesne E (2005) When is the brain enlarged in autism? A meta-analysis of all brain size reports. *Biol Psychiatry* 58:1–9
90. Ritvo ER, Freeman BJ, Scheibel AB et al (1986) Lower Purkinje cell counts in the cerebella of four autistic subjects: initial findings of the UCLA-NSAC Autopsy Research Report. *Am J Psychiatry* 143:862–866
91. Rodier PM, Ingram JL, Tisdale B, Nelson S, Romano J (1996) Embryological origin of autism: developmental abnormalities of the cranial nerve nuclei. *J Comp Neurol* 370:247–261
92. Rorke LB, Fogelson MH, Riggs HE (1968) Cerebellar heterotopia in infancy. *Dev Med Child Neurol* 10:644–650
93. Rorke LB (1994) A perspective: the role of disordered genetic control of neurogenesis in the pathogenesis of migration disorders. *J Neuropathol Exp Neurol* 53:105–117
94. Rutter M, Bailey A, Bolton P, Le Couteur A (1994) Autism and known medical conditions: myth and substance. *J Child Psychol Psychiatry* 35:311–322
95. Saitoh O, Courchesne E (1998) Magnetic resonance imaging study of the brain in autism. *Psychiatry Clin Neurosci* 52(Suppl):S219–S222
96. Sarnat HB, Benjamin DR, Siebert JR et al (1992) Cerebral dysgenesis: embryology and clinical expression. *Cell* 69:581–595
97. Schmahmann JD, Doyon J, McDonald D et al (1999) Three-dimensional MRI atlas of the human cerebellum in proportional stereotaxic space. *Neuroimage* 10:233–260
98. Schmidtke K, Manner H, Kaufmann R, Schmolck H (2002) Cognitive procedural learning in patients with fronto-striatal lesions. *Learn Mem* 9:419–429
99. Sears LL, Vest C, Mohamed S, Bailey J, Ranson BJ, Piven J (1999) An MRI study of the basal ganglia in autism. *Prog Neuropsychopharmacol Biol Psychiatry* 23:613–624
100. Simms ML, Kemper TL, Timbie CM, Bauman ML, Blatt GJ (2009) The anterior cingulate cortex in autism: heterogeneity of qualitative and quantitative cytoarchitectonic features suggests possible subgroups. *Acta Neuropathol* 118:673–684
101. Sokol DK, Chen D, Farlow MR et al (2006) High levels of Alzheimer beta-amyloid precursor protein (APP) in children

- with severely autistic behavior and aggression. *J Child Neurol* 21:444–449
102. Tassi L, Colombo N, Garbelli R et al (2002) Focal cortical dysplasia: neuropathological subtypes, EEG, neuroimaging and surgical outcome. *Brain* 125:1719–1732
 103. The Autism Genome Project Consortium (2007) Mapping autism risk loci using genetic linkage and chromosomal rearrangements. *Nat Genet* 39:319–328
 104. Townsend J, Courchesne E, Covington J et al (1999) Spatial attention deficits in patients with acquired or developmental cerebellar abnormality. *J Neurosci* 19:5632–5643
 105. Tuchman RF, Rapin I, Shinnar S (1991) Autistic and dysphasic children. I. Clinical characteristics. *Pediatrics* 88:1211–1218
 106. Tuchman RF, Rapin I (2002) Epilepsy in autism. *Lancet Neurol* 1:352–358
 107. Usui N, Matsuda K, Mihara T et al (2001) MRI of cortical dysplasia-correlation with pathological findings. *Neuroradiology* 43:830–837
 108. van Kooten I, Palmen SJ, von Engelan CP et al (2008) Neurons in the fusiform gyrus are fewer and smaller in autism. *Brain* 131:987–999
 109. Voelbel GT, Bates ME, Buckman JF, Pandina G, Hendren RL (2006) Caudate nucleus volume and cognitive performance: are they related in childhood psychopathology? *Biol Psychiatry* 60:942–950
 110. Volkmann K, Rieger S, Babaryka A, Koster RW (2008) The zebrafish cerebellar rhombic lip is spatially patterned in producing granule cell populations of different functional compartments. *Dev Biol* 313:167–180
 111. Wegiel J, Wisniewski T, Chauhan A (2010) Type, topography and sequelae of neuropathological changes shaping clinical phenotype of autism. In: Chauhan A, Chauhan V, Brown WT et al (eds) *Autism: oxidative stress, inflammation, and immune abnormalities*. Taylor & Francis/CRC Press, Boca Raton, FL, pp 1–34
 112. Weiss LA, Arking DE, The Gene Discovery Project of Johns Hopkins & the Autism Consortium (2009) A genome-wide linkage and association scan reveals novel loci for autism. *Nature* 461:802–811
 113. Wolf HK, Normann S, Green AJ et al (1997) Tuberous sclerosis-like lesions in epileptogenic human neocortex lack allelic loss at the TSC1 and TSC2 regions. *Acta Neuropathol* 93:93–96
 114. Yip J, Soghomonian JJ, Blatt GJ (2007) Decreased GAD67 mRNA levels in cerebellar Purkinje cells in autism: pathophysiological implications. *Acta Neuropathol* 113:559–568
 115. Yip JJ, Soghomonian J, Blatt GJ (2008) Increased GAD67 mRNA expression in cerebellar interneurons in autism: implications for Purkinje cell dysfunction. *J Neurosci Res* 86:525–530

**9th ANNUAL INTERNATIONAL MEETING FOR AUTISM RESEARCH (IMFAR)
PHILADELPHIA, PA MAY 20-22, 2010**

**Section 130: Neuropathology
Saturday, May 22, 2010 9:45am- 11:45 am**

Poster Presentation 130.040 61

Neuronal Growth Delay within the Claustrum of Autistic Subjects.

S. Y. Ma*, I. Kuchna, K. Nowicki, J. Wegiel, H. Imaki, I. Cohen, E. London, M. Flory, W. T. Brown, T. Wisniewski and J. Wegiel,
New York State Institute for Basic Research in Developmental Disabilities,

Background: The claustrum ontogeny and morphology indicate a pallial and subpallial origin and explains why the Claustrum receives inputs from multiple cortical areas, integrates the multiple inputs into a new signal, and redirects sensory information throughout the striatum and thalamus. Involvement in sensorimotor integration and cognition suggests that the claustrum plays a role in higher order functionality enabling the organism to rapidly adapt to the subtleties and nuances of an ever-changing environment (Edelstein and Denaro 2004). The severely compromised ability to adapt characteristic of autism suggests that these high order functions are not processed properly and that developmental alterations of the claustrum may contribute to the autistic phenotype.

Objectives: The aim of this study was to test the hypothesis that claustrum development is modified and to characterize the type of structural changes employing unbiased morphometric methods.

Methods: Brain hemispheres of 13 autistic and 14 control subjects 4 to 64 years of age were fixed in 10% formalin, dehydrated and embedded in celloidin. The fractionator method was used to determine the number of neurons, the Cavalieri method to estimate the volume of the claustrum, and Nucleator method to determine the volume of neurons and neuronal nuclei.

Results: In subjects 4 to 23 years of age both the claustrum and the prepiriform claustrum were characterized by developmental delays. Volume of the claustrum was reduced by 30% on average, and the total number of neurons by 29% whereas the numerical density was not modified. The volume of neurons in this younger group of autistic subjects was reduced by 32% and the volume of neuronal nuclei was proportionally less (by 31%).

Conclusions: The claustrum showed a similar range of developmental delay to the structures interacting with the Claustrum including the entorhinal cortex, several striatal subdivisions and the thalamus as shown in our other studies. These results support the hypothesis that developmental delay of the claustrum may contribute to the rigidity of behaviors and the lack of ability to adapt to a changing environment seen in autism.

Poster Presentation 130.049 70

Repetitive and Stereotyped Behaviors in Autism Are Driven by Abnormal Development of the Striatum but Not of the Substantia Nigra. K. Nowicki*, T. Wisniewski, I. Kuchna, J. Wegiel, H. Imaki, S. Y. Ma, I. Cohen, E. London, M. Flory, W. T. Brown and J. Wegiel, *New York State Institute for Basic Research in Developmental Disabilities*

Background: Repetitive and stereotyped behaviors defined as recurring, nonfunctional activities or interests that occur regularly and interfere with daily functioning are defining signs of autism (Gabriels et al., 2005).

Several studies have implicated the role of basal ganglia and fronto-striatal circuitry in the pathophysiology of autism, related especially to repetitive and stereotyped behaviors.

Objectives: The aim of this morphometric study was to determine whether the developmental trajectories of neurons within the dorsal, ventral and lateral parts of the substantia nigra, the source of dopaminergic innervation of the human striatum, are similar to the pattern of development of four striatal subdivisions including the caudate, putamen, nucleus accumbens and globus pallidus.

Methods: We applied the Nucleator method to detect patterns of developmental changes of neuronal soma and nuclear volume within five subdivisions of the nigro-striatal system in the brains of 12 subjects with idiopathic autism and 12 age-matched controls.

Results: Volume of neurons within the three subdivisions of the substantia nigra, including the dorsal, ventral and lateral parts innervating four parts of the striatum was almost identical in the autistic and control subjects both 4 to 8 years of age and >8 years of age. However, in the brains of autistic children 4 to 8 years of age we found a significant delay of neuron growth with the volume of cell body smaller than in control subjects by 22% in the caudate nucleus (0.025*), by 21% in the putamen ($p < 0.042^*$), by 31% in the globus pallidus ($p < 0.008^{**}$) and by 37% in the nucleus accumbens ($p < 0.004^{**}$). The volume of neuron nucleus was also significantly reduced. No significant difference between the volume of neuronal soma and nucleus in autistic and control subjects >8 years of age suggests that a correction of neuronal growth occurs in late childhood and adulthood.

Conclusions:

The observed pattern indicates a normal development of neurons producing dopamine but a delayed development of striatal neurons. It suggests that development of the nigro-striatal system is desynchronized and that the repetitive and stereotyped behaviors in autism are driven by delayed development of neurons within the striatum but not in the substantia nigra.

Oral Presentations

Section 135: Neuropathology I, Saturday, May 22, 2010 9:45am- 11:45 am

Oral Presentation 135.005 Defects of Neurogenesis, Neuronal Migration and Dysplastic Changes in the Brains of Autistic Subjects.

J.Wegiel*, I. Kuchna, K. Nowicki, H. Imaki, J. Wegiel, E. Marchi, S. Y. Ma, A. Chauhan, V. Chauhan, I. Cohen, E. London, W. T. Brown and T. Wisniewski, *New York State Institute for Basic Research in Developmental Disabilities*

Background: An emerging theory of autism related encephalopathy integrates evidence of (a) abnormal acceleration of brain growth in early childhood (Redcay and Courchesne 2005), (b) minicolumn pathology (Casanova et al 2002, 2006), (c) curtailed neuronal development (Bauman and Kemper 1985), brain-structure-specific delays of neuronal growth in early childhood, and desynchronization of neuronal development.

Objectives: The aims of this study were to detect the patterns of focal qualitative developmental brain defects, including the type, topography and severity of changes, and to identify the structures and brain regions that are prone to developmental alterations in autism.

Methods: Formalin-fixed brain hemispheres of 13 autistic (4–60 years of age) and 14 age-matched control subjects were embedded in celloidin and cut into 200- μ m thick coronal sections, which were stained with cresyl violet and used for neuropathological evaluation.

Results: Thickening of the subependymal germinal matrix and subependymal nodular dysplasias in two autistic children were indicative of active neurogenesis. Subcortical, periventricular, hippocampal and cerebellar heterotopias were detected in the brains of four autistic subjects (31%) reflecting abnormal neuronal migration. Multifocal cerebral dysplasias seen in the neocortex (31%), in the entorhinal cortex (15%), in the cornu Ammonis (31%) and in the dentate gyrus (15%) reflects frequent developmental distortions of brain cytoarchitecture. Cerebellar flocculonodular dysplasia detected in six subjects (46%) and hypoplasia in one case indicate local failure of cerebellar development in 54% of the autistic subjects. Dystrophy with calcification and focal damage within meninges, cortex and white matter was present in all examined brains of autistic and control subjects.

Conclusions: Detection of focal dysplasia in one control subject and of a broad spectrum of focal qualitative neuropathological developmental changes in 12 of 13 (92%) examined brains of autistic subjects indicates a multi-regional dysregulation of neurogenesis, neuronal migration and maturation in autism. This may contribute to the heterogeneity of the observed clinical phenotype.



NEURONAL GROWTH DELAY WITHIN THE CLAUSTRUM OF AUTISTIC SUBJECTS

Shuang Yong Ma, Izabela Kuchna, Krzysztof Nowicki, Jarek Wegiel, Humi Imaki,
Ira L. Cohen, Eric London, Michael Flory, W.Ted Brown, Thomas Wisniewski, Jerzy Wegiel
New York State Institute for Basic Research in Developmental Disabilities, Staten Island, New York, U.S.A.

In 2005, Nobel Prize laureate Dr. Francis Crick and his co-worker Christoph Koch proposed the claustrum as the ideal candidate for the brain "consciousness" center, because it acts like a conductor in an orchestra to integrate information from all of the various modalities.


CLAUSTRUM ANATOMY

The claustrum is a thin ribbon of gray matter located between the insular cortex and the striatum, closely associated with the amygdala and built of 16 million cells of both cortical and subcortical origin. 94% of claustrum neurons are large projection spiny neurons with numerous coarse pigment granules (Braak and Braak 1982).

CLAUSTRUM CONNECTIVITY

The most fascinating feature of the claustrum is its extensive reciprocal connections to and from almost all brain regions (Fernandez-Miranda *et al.* 2008). The information from the various brain regions is segmented topographically in the claustrum and in an overlapping manner.

- The anterior claustrum projects to and receives signals from the frontal cortex.
- The middle claustrum is associated with the parietal cortex.
- The posterior/inferior claustrum is associated with the temporal and occipital cortices.
- The dorsal claustrum forms a visual loop with BA17 (Fernandez-Miranda *et al.* 2008, Morys *et al.* 1993, Pearson *et al.* 1982).
- The claustrum also projects to the amygdala (Amaral and Insausti 1992), the hippocampus (Amaral and Cowan 1980), and the striatum (Arikuni and Kubota 1985, Crick and Koch 2005).
- Social brain circuitry (orbito-frontal cortex, superior temporal gyrus and amygdala) (Brothers *et al.* 1990) projects to the claustrum and receives claustral connections (Fernandez-Miranda *et al.* 2008).



CLAUSTRUM FUNCTION IN NORMAL AND PATHOLOGICAL CONDITIONS

- Higher order cognitive functions such as fear recognition (Stein *et al.* 2007)
- Experiential dread (Berns *et al.* 2006)
- Cognitive impairment (Dubroff *et al.* 2008)
- Memory storage (Morys *et al.* 1996)
- Associative learning (Chachi and Powell 2004)
- Repetitive behaviors and addiction (Morys *et al.* 1996; Naqvi *et al.* 2007)
- Multimodal processing of olfactory, auditory, visual, and tactile information.
- Emotional and behavioral responses (Bennet and Baired 2006)
- Suppression of natural urges (Lerner *et al.* 2008)
- Seizures (Zhang *et al.* 2001)
- Contribution to psychoses (Sperner *et al.* 1996)

HYPOTHESIS

The severely compromised ability to adapt that is observed in individuals with autism, suggests that numerous high order functions are not processed properly and that developmental defects in the claustrum may contribute to the cognitive impairment, repetitive behaviors, deficits of social interactions, altered processing of sensory signals and other components of the autistic phenotype.

AIM

The aim of this morphometric study of the claustrum of autistic and control subjects was to determine whether and how the development of claustrum, known to be responsible for integration of the functions of the social and somatosensory circuits, is affected in autism.

MATERIAL AND METHODS

Nine brain hemispheres of autistic subjects from 4 to 36 years of age and nine control subjects 4 to 32 years of age were examined. In the majority of cases, the right hemisphere was examined (16 cases, 89%). Four age-matched pairs of autistic and control subjects represented the youngest group (4/4, 5/4, 7/7, and 8/8 years of age). Five age-matched pairs represented late childhood and adulthood (13/14, 17/20, 21/23, 23/28, and 36/32). Brain hemispheres were fixed in 10% formalin, dehydrated in ascending concentrations of ethyl alcohol for 19 days (dehydration results in the reduction of brain hemisphere weight by 47% on average), and embedded in celloidin. Free-floating celloidin 200- μ m-thick sections were stained with cresyl violet.

Clinical inclusion criteria.

Selection of autistic cases was based on clinical evaluation including Autism Diagnostic Interview-Revised (ADI-R) score. Subjects with atypical autism were excluded from this morphometric study.

Neuropathological exclusion criteria.

Expanded neuropathological evaluation was based on the examination of 120 coronal hemispheric CV-stained serial sections. Brains affected with premortem pathology (ischemia, anoxia-related neuronal degeneration/loss, gliosis, and neovascularization) and severe postmortem degradation (autolysis) affecting brain region volume; cytoarchitecture; and the number, shape, and size of neurons were excluded from this study.



Morphometric equipment.

Image analyzer with computer-controlled automatic stage installed on Axiophot II microscope (Zeiss) and Stereo-Investigator and Nucleator software (Microbrightfield, VA, USA).

Morphometric methods.



The Cavalieri method was used to estimate the volume of the ventral and dorsal claustrum, the disector method was used to determine the number of neurons, and the nucleator to determine the volume of neurons and neuronal nuclei. Objective lens: 1.25x (volumetry) and 40x (disector and nucleator). The final magnification used was 48 x and 1,480 x, respectively. The mean number of neurons examined per one region/case: 641 in the autism group, and 697 in the control group. The coefficient of error was less than 0.01 (Schaffer).

Statistical analysis. Claustrum volume, number of neurons, neuronal density, mean neuron size and mean nuclear size were compared in repeated-measures ANOVA.





Claustrum volume

The claustrum volume was 22% less in autistic (388 mm³) than in control (494 mm³) subjects from 4 to 8 years of age. In autistic subjects 13 to 36 years of age, the volume of claustrum increased by 18% (to 472 mm³) and was insignificantly higher than in age-matched controls (452 mm³). This observation is consistent with MRI studies of 16 high-functioning (IQ 85) autistic and 14 control males from 7 to 12 years of age which revealed a lower volume of the claustrum of autistic (555 mm³) than control subjects (701 mm³) (Davies 2008). The lower volumes detected in postmortem study than in clinical MRI study are an effect of tissue shrinkage during dehydration.





Number of neurons

The mean numerical density of neurons was almost identical in the autistic (29,156/mm³; n = 9) and control subjects (29,260/mm³; n = 9). The mean total number of neurons in the nine autistic subjects (12.5 millions) was insignificantly less (by 8%) than in the control subjects (13.6 millions).



Mean neuron volume

The mean volume of the neuronal body in 4- to 8-year old autistic children was less by 32% (1,465 µm³, n = 4) than in control group (2,154 µm³; n = 4; p < 0.05). However, the mean volume of neuronal body in 13- to 36-year-old autistic subjects was less only by 19% (1,640 µm³; n = 5) than in control group (2,010 µm³; n = 5; p < 0.02).




Mean volume of neuron nucleus

The mean volume of neuron nucleus in 4- to 8-year old autistic children (252 µm³; n = 4), was 41% less than in the control group (427 µm³; n = 4; p < 0.009). The mean volume of the neuronal nucleus in 13- to 36-year-old autistic subjects was insignificantly less (269 µm³; n = 5) than in the control group (358 µm³; n = 5).

Desynchronization of development of claustral circuitry

The striking differences in the developmental volume deficits of neurons in the claustrum and in brain structures projecting to the claustrum (shown in previous reports) that were detected in 4- to 8-year-old autistic children are signs of desynchronization of claustral circuitry development. The observed pattern of developmental delays suggests that a number of physiologic processes that collectively ensure the maintenance of undisturbed contact with the environment are altered in the early childhood of autistic subjects.



Underconnectivity

The smaller volume of the claustrum and the reduced size of neurons could be indicative of underconnectivity in the brains of autistic subjects. Underconnectivity of the claustrum may result in defects/deficits of integration of cortical and subcortical modalities and may contribute to the complex autistic phenotype.

CONCLUSIONS

1. Stereological study revealed a significant delay of neuronal growth in the claustrum, which is responsible for the integration of a number of physiological processes that collectively maintain contact with the patient environment.
2. The detected desynchronization of neuronal growth in the claustrum and interacting brain modalities, including social brain, sensorimotor system, and memory system, appears to be a major contributor to the clinical symptoms of autism.
3. Examination of older children and adults with autism shows that in the claustrum of teenagers, acceleration of neuronal growth results in partial correction of neuronal size.
4. The findings of this study suggeststst, that correction of the delay of neuronal growth in the early stage of development of children with autism may prevent functional alterations across the lifespan.

Acknowledgements

The Autism Brain Atlas Project is supported by Autism Speaks, the Department of Defense Autism Spectrum Disorders Research Program (A5073334), and the NYS Office of Mental Retardation and Developmental Disabilities.

The tissue was provided by the Harvard Tissue Resource Center, Boston, MA (R24-MH068855), and the Brain and Tissue Bank at the University of Maryland, Baltimore, MD.

The Autism Tissue Program (Princeton, NJ) coordinated the selection of material for the morphometric study of cortical and subcortical subdivisions for this project.

This study is the product of the coordinated efforts of families who donated tissue and provided information; the above-listed institutions; and many unnamed clinicians, psychiatrists, geneticists, radiologists, neuropathologists, medical examiners, and members of the Advisory Board of the Autism Tissue Program.

Conflict of interest: None



REPETITIVE AND STEREOTYPED BEHAVIORS IN AUTISM ARE DRIVEN BY ABNORMAL DEVELOPMENT OF THE STRIATUM BUT NOT OF THE SUBSTANTIA NIGRA

Krzysztof Nowicki, Thomas Wisniewski, Izabela Kuchna, Jarek Wegiel, Humi Imaki, Shuang Yong Ma, Ira L. Cohen, Eric London, W. Ted Brown, Jerzy Wegiel

New York State Institute for Basic Research in Developmental Disabilities, Staten Island, New York, U.S.A.

BACKGROUND

Anatomical substrate of repetitive and stereotyped behaviors.

Repetitive and stereotyped behaviors, defined as recurring, nonfunctional activities or interests that occur regularly and interfere with daily functioning, are diagnostic signs of autism (Gabriels et al., 2005). Several studies have implicated the role of the basal ganglia and fronto-striatal circuitry in the pathophysiology of autism, especially in repetitive and stereotyped behaviors.

The substantia nigra, together with the caudate nucleus, putamen, n. accumbens and globus pallidus, belongs to the basal ganglia. These structures form the cortico-subcortical circuits that play a key role (a) in programming and execution of movements, and (b) in the reward system.

Partial anatomical and functional overlap of motor and reward systems.

The reward system includes (a) the prefrontal cortex, (b) the limbic system components with the amygdala, hippocampus, and anterior cingulate, (c) the substantia nigra and limbic striatum with the nucleus accumbens, ventral caudate nucleus, and putamen (Lopes de Silva et al., 1990; Kotter and Stephan, 1997).

Substantia nigra and brain reward pathway. Dopamine has been associated with pleasure, and it has been called the "anti-stress molecule" and "pleasure molecule" (Bloom et al., 2000). Released dopamine stimulates a number of receptors (D₁-D₅), which results in increased feelings of well-being and stress reduction. A reduced number of dopamine receptor sites in n. accumbens and the amygdala may result in repetitive behavior in an attempt to stimulate dopamine release.

Reward deficiency syndrome is associated with abnormalities of the basal ganglia, including the substantia nigra, caudate, putamen, and nucleus accumbens, as well as the prefrontal cortex, hypothalamus, and central nucleus of the amygdala. Our studies indicate that almost all of these structures are affected by developmental neuronal delay in people with autism. The substantia nigra appears to play a key role in both the motor and reward systems, but the morphology of the substantia nigra dopaminergic neurons is not known.

Nucleus accumbens dopaminergic innervation and role in reward system.

The nucleus accumbens is at the center of network processing of reward-related information. The shell of the nucleus accumbens (Heimer et al., 1997) is a part of the extended amygdala involved in integrating emotions, whereas, the core belongs to the striato-pallidal system integrating motor responses. The nucleus accumbens projections to the motor areas, such as the ventral pallidum, turn reward information into motivated action (Mogenson et al., 1980). Dopaminergic projections from the substantia nigra to the nucleus accumbens have a critical role in stimulus-reward learning. Repetitive behaviors and stereotypies may reflect altered motor responses modulated by the altered reward system.



Substantia nigra structure. The substantia nigra pars compacta forms a shallow layer of approximately 1.4 million pigment-positive, dopamine-producing neurons. They form ventral and dorsal tiers, and the lateral part. Neurons from both the ventral and dorsal tiers project to the striatum and cerebral cortex. The dorsal tier also projects to the amygdala. Numerous reciprocal connections feedback within basal ganglia networks.

HYPOTHESIS

- Development of the dopaminergic system contributing to repetitive and stereotyped behaviors is altered in autism.
- The clinical outcome of altered brain development is a complex product of changes in different dopaminergic system components.
- The absence of clinical signs of dopamine deficit or excess suggests that the substantia nigra is not altered in autistic subjects, and that modifications of the development of striatal components are the main contributor to repetitive and stereotyped behaviors.

AIM

The aim of this morphometric study was to determine whether the developmental trajectories of neurons within the dorsal, ventral, and lateral parts of the substantia nigra, the source of dopaminergic innervation of the human striatum, are similar to the patterns of development of four striatal subdivisions, including the caudate, putamen, nucleus accumbens, and globus pallidus.

MATERIAL AND METHODS

Material. Eleven brain hemispheres of autistic subjects from 4 to 56 years of age and 11 control subjects from 4 to 64 years of age were examined. In the majority of the cases, the right hemisphere was examined (18 cases, 82 %). Four age-matched pairs of autistic and control subjects represented the youngest group (4/4, 5/4, 7/7, and 8/8 years of age). Seven pairs represented late childhood and adulthood (11/14, 13/15, 17/20, 21/23, 22/29, 23/32, and 56/64 year old).

Brain hemispheres were fixed in 10% formalin, dehydrated in ethyl alcohol for 19 days, and embedded in celloidin. Dehydration reduces brain hemisphere weight by 47% on average. Free-floating, 200-µm-thick sections were stained with cresyl violet.

Clinical and neuropathological inclusion/exclusion criteria. Clinical inclusion criteria were based on Autism Diagnostic Interview-Revised (ADI-R) score. Subjects with atypical autism were excluded from this morphometric study.

Neuropathological examination of 120 coronal hemispheric CV-stained serial sections revealed autism-associated developmental changes reflecting defects of neurogenesis, migration (heterotopias), and cytoarchitecture (dysplasia) in 92% of autistic subjects. Five brains of autistic subjects and four brains of control subjects with signs of premortem pathology (ischemia or anoxia) or severe postmortem degradation (autolysis) were excluded from this study.

Morphometric methods and statistical analysis.

The volume of neurons and neuronal nuclei within the three subdivisions of the substantia nigra, including the ventral and dorsal tier and lateral part, was evaluated in autistic and control subjects.

In our previous study of this cohort, we determined the volume of neurons and nuclei in other components of the nigro-striatal system, including the caudate nucleus, putamen, globus pallidus, nucleus accumbens, and amygdala.


The Nucleator was used to determine the volume of neurons and neuronal nuclei (objective lens: 40 x; final magnification 1,480 x). The mean number of neurons examined per case: 291 in the autism group, and 282 in the control group. The coefficient of error was less than 0.01 (Schaffer). For neuron morphometry, the image analyzer with computer-controlled automatic stage installed on Axiophot II microscope (Zeiss) and Stereo-Investigator and Nucleator software (Microbrightfield, VA, USA) was applied.

The substantia nigra subdivisions volume, number of neurons, neuronal density, and mean neuronal size were compared in autistic and control cohorts in repeated-measures ANOVA.

RESULTS

SUBSTANTIA NIGRA

No difference between the volume of pigmented neurons and their nuclei in the substantia nigra of control and autistic subjects




Neuron volume in control cohort.

The mean volume of the neurons in the ventral and dorsal tiers and the lateral part of the substantia nigra in 4- to 64-year-old control subjects was similar (9,904 µm³, 9,624 µm³, and 10,041 µm³, respectively). The lack of significant difference suggests the homogeneity of pigmented neurons within different portions of the pars compacta, with a mean volume of 9,856 µm³ in the control group. There was no statistically significant difference in neuron volume between 4- to 8-year-old and 14- to 64-year-old control subjects.

Neuron volume in autistic cohort. The study of autistic subjects from 4 to 56 years of age did not reveal any difference in the volume of the pigmented neurons in the ventral and dorsal tiers, and in the lateral part of the substantia nigra (9,949 µm³, 9,903 µm³, and 10,281 µm³, respectively). The mean volume of neurons in the substantia nigra of autistic subjects (10,044 µm³) was comparable with those observed in the control cohort (9,856 µm³).

In both the autistic and control cohorts, there was no difference in the volume of pigmented neurons between subjects younger than 8 years of age and in the teenagers/adults group.




Nucleus volume in neurons of autistic and control subjects. The nucleus volume in neurons in the ventral and dorsal tiers and in the lateral part of the substantia nigra is comparable across anatomical subdivisions, age subgroups, and autistic and control cohorts.

CAUDATE, PUTAMEN, GLOBUS PALLIDUS, NUCLEUS ACCUMBENS

Desynchronized development of nigro-striatal circuitry in autistic subjects: a potential contribution to repetitive and stereotyped behaviors

In the brains of autistic children 4 to 8 years of age, we found a significant delay of neuron growth, with the volume of cell bodies smaller than in control subjects by: 22% in the caudate nucleus ($p < 0.025^*$), 21% in the putamen ($p < 0.042^*$), 31% in the globus pallidus ($p < 0.008^{**}$) and 37% in the nucleus accumbens ($p < 0.004^{**}$). The volume of the neuronal nucleus was also significantly reduced.

Much less, or lack of, significant difference between the volume of neuronal soma and nucleus in autistic and control subjects older than 8 years of age suggests that a correction of neuronal growth occurs in late childhood and adulthood.



The unaffected size of the neurons of the substantia nigra providing dopamine to the motor and reward system, but the significant structure-specific delay of neuronal growth in striatal subdivisions indicates that development of the nigrostriatal system is desynchronized and that differences are most striking in 4- to 8-year-old autistic subjects. The observed pattern of developmental desynchronization of the nigrostriatal system might be a key contributor to the repetitive and stereotyped behaviors of autistic subjects.

CONCLUSIONS

- The observed pattern indicates the normal development of neurons within the ventral and dorsal tiers and in the lateral part of the substantia nigra.
- This pattern may suggest that dopaminergic projections to other components of the nigro-striatal system are not deficient.
- No difference in the volume of pigmented neurons in the substantia nigra of autistic and control subjects, but the significant delay of the growth of neurons in the striatal subdivisions of autistic children, indicates desynchronization of development of both the motor system and the award system.
- This study suggests that the repetitive and stereotyped behaviors seen in autism are driven by a delayed development of neurons within the striatum but not in the substantia nigra.

Acknowledgements: This project was supported by grants from the Department of Defense Autism Spectrum Disorders Research Program (AS073234) and Autism Speaks, as well as by the NYS Office of Mental Retardation and Developmental Disabilities. The Harvard Brain Tissue Resource Center (R24-MH 068855), and the Brain and Tissue Bank at the University of Maryland, Baltimore, provided tissue. The Autism Tissue Program coordinated tissue acquisition.

Dystrophy with Calcification within Brains of Autistic and Control Subjects

Izabela Kuchna, Krzysztof Nowicki, Humi Imaki, Jarek Wegiel, Shuang Yong Ma, Elaine Marchi, Ira L. Cohen,
Eric London, W. Ted Brown, Thomas Wisniewski, Jerzy Wegiel

New York State Institute for Basic Research in Developmental Disabilities, Staten Island, New York, U.S.A.

OBJECTIVES

The overall aim of our study is to detect a global pattern of developmental quantitative and qualitative alterations in the brains of autistic subjects. Application of a unique tissue preservation protocol and examination on average of 120 hemispheric sections per case resulted in detection of the three major types of pathological changes:

- (a) Developmental qualitative defects of neurogenesis, migration and cytoarchitecture
- (b) Desynchronization of neuronal growth;
- (c) A new form of meningeal and brain tissue dystrophy with calcification.

Developmental qualitative defects of neurogenesis, migration and cytoarchitecture. Developmental abnormalities were detected in 92% of examined brains of autistic subjects including:

- Alterations of neurogenesis within the subependymal layer,
- Defects of migration with heterotopias, and
- Defects of cytoarchitecture with dysplastic changes in the brain cortex, hippocampus, and cerebellum.

Role in autism. These three types of developmental changes may contribute to a broad spectrum of interindividual differences among subjects diagnosed with autism (Wegiel et al, Acta Neuropath, 2010; free access).

Desynchronization of neuronal growth. Stereological study of 13 major brain structures with 21 anatomical subdivisions revealed:

- (a) A very significant delay of neuronal growth in seven examined brain structures, including the claustrum, n. accumbens, thalamus, Purkinje cells, globus pallidus, amygdala, and entorhinal cortex, resulting in neuronal body volume deficit ranging from -41% to -28% in the brains of autistic subjects;
- (b) A moderate delay in three brain structures, including the dentate nucleus in the cerebellum, putamen and caudate nucleus (-19% to -16%), and
- (c) No signs of delay of neuronal growth in the cornu Ammonis, lateral geniculate body, and substantia nigra.

Role in autism: The detected desynchronization of neuronal growth in the brain of autistic subjects appears to be a major contributor to the clinical phenotype of autism. (Wegiel et al, 2010. In: Autism. Oxidative Stress, Inflammation and Immune Abnormalities. Eds.: A. Chauhan, V. Chauhan, W.T. Brown)

A new form of meningeal and brain tissue necrosis and dystrophy with calcification. Brain tissue lesions with intracranial calcifications are observed in many diseases including those with viral and bacterial infections, vascular pathology, toxic injury, brain tumors, teratomas, lissencephaly, Fahr's disease, and very often in parasitic infections (Rabbits et al 1969).

Our study of global patterns of neuropathological changes in the brains of autistic and age-matched control subjects has revealed the new type of meningeal, cortical and white matter necrosis and dystrophy with calcification in the brains of autistic and control individuals.

Role in autism: To be determined.

AIMS

The aims of this study are to:

- Characterize the type, topography and prevalence of a new form of brain pathology in autistic and control subjects.
- Determine the risk of this type of lesions contribution to the autistic phenotype.

MATERIAL & METHODS

Material. The brain hemispheres of 13 autistic and 14 control subjects 4 to 64 years of age were fixed in 10% formalin, dehydrated and embedded in celluloid and then cut into 200-µm- or 50-µm-thick coronal serial sections. Free-floating hemispheric sections were stained with cresyl violet.

Neuropathological evaluation. Every 1.2 mm distance section from one hemisphere was examined. On average, 120 histological slides were examined per brain hemisphere.


RESULTS

Prevalence. Dystrophy with calcification was found in all 13 autistic and 14 control brains examined, but with significant interindividual differences pertaining to the extension of meningeal/cortical and white matter damage.


Topography. Lesions were detected on the lateral surface of the brain hemisphere and usually in the frontal superior, middle and inferior gyri, temporal superior, middle and inferior gyri, and middle occipital gyrus. However, lesions were not detected on the medial surface of the brain and cerebellum.

Laterality. For three autistic cases and one control case, both hemispheres were available for macro- and microscopic evaluation. The pattern of changes in both hemispheres was comparable with minor differences in lesion topography and severity.



Macroscopic mapping of a new form of dystrophy with calcification in autism and control. Lateral view of right brain hemisphere of 7-year-old male diagnosed with autism and severe early-onset epilepsy (B6493) and 7-year-old control male (UMB4898) illustrate a topography of lesions. Linear leptomeningeal and cortical lesions (red arrowheads) cross the superior, middle and inferior frontal gyrus in the frontal (F) lobe and gyri of the temporal (T) and occipital (O) lobes.




More severe dystrophy with calcification in the brain of autistic subject with seizures. Selected serial CV-stained coronal hemispheric sections from the brain of an autistic and control subject illustrate the topography and depth of the lesions (arrowheads). Lesions penetrate deeper into the cortex and white matter of the autistic subject. The contribution of these lesions to severe epilepsy is not excluded. Much less pronounced changes in the control subject could be nonsymptomatic.



Leptomeningeal, cortical and white matter lesions. The depth of linear lesions varied locally from a few mm to more than 30 mm. Shallow lesions penetrated from leptomeninges into the cortex (Figs. a and b), whereas deep lesions penetrated from leptomeninges linearly through the cortical ribbon and white matter of one or two adjacent gyri (Fig. c). Deep lesions caused misplacement of cortical ribbon and local disruption of cortical ribbon and white matter continuity.




Focal loss of neuroarchitecture. In this new form of pathology, necrosis and dystrophy focally disrupt the continuity of the cortical ribbon and white matter. Progression of death and lysis of neurons in the cortex and axons in white matter replaced by amorphous material (asterisks) leads to the almost complete loss of neuroarchitecture.



Progression of changes within lesioned area and adjacent neuropil. Progression of dystrophic changes is associated with the condensation and mineralization of amorphous products of cell lysis in the affected area (arrowheads). However, the border of the lesioned area and adjacent cortex shows only a narrow zone with distinct alterations of neuronal architecture.



Calcification. Lesion area shows progressive accumulation of mineralized deposits, with calcium being the main component (van Kossa method of calcium staining). The mineralization increases with age. Local differences in the amount of calcium deposits may reflect different time of onset and duration of changes in a given site.



Focal loss of gliooarchitecture. Micrograph shows the total loss of gliooarchitecture in lesion (L) area and sharp border (arrowheads) between lesion and intact cortex (C) (staining for astrocytes GFAP). A striking feature of the lesion border was the lack of any reaction of astrocytes to severe neuronal damage and loss.




Vascular pathology In the lesion area, the leptomeningeal vessels and vessels penetrating the cortex and white matter become tortuous with increased cellularity of thickened vascular wall and reduced vessel lumen. This vascular pathology may suggest that the causative factor is of blood origin.



Fragmentation and degradation of vascular wall residues were a common finding in the affected areas. The lesion area was often saturated with erythrocytes and activated macrophages.



Progression of necrotic changes resulted in digestion of the cytoplasm of neurons, glial cells, and cells of the vascular wall. The sign of the unique lytic activity of the intercellular space in the lesion area was numerous distorted, thread-like cell nuclei.



Mean length (mm) of linear lesions in children and adult autistic and control subjects

Lobe	Autism 4-13 y	Autism 17-60 y	Control 4-15 y	Control 20-64 y
Frontal	30.7	26.9	29.9	25.4
Temporal	35.7	36.9	43.7	46.1
Occipital	17.7	22.9	24.0	16.8
Total length	84.1	86.7	97.6	88.3

The total length of linear lesions was comparable in autistic and control subjects in both age groups. The almost identical length of lesions in both children and adults suggests that pathology develops in early childhood (< 4 years).

CONCLUSIONS

- The similar morphology of lesions in control and autistic subjects 4 to 64 years of age suggests that this newly discovered type of dystrophy with calcification is a common form of human brain pathology with age of onset of less than 4 years.
- A similar length of lesions in children and adults indicates that the linear size of lesions undergoes relatively limited modifications with age.
- However, the presence of degenerated neurons and cells of the vascular wall in lesions in all subjects regardless of age, suggests expansion of the depth of lesion and indicates that the process of brain tissue damage is active decades after the original causative events.
- The 84 to 97-mm-long lesions of leptomeninges, cortex and white matter in all examined brains of autistic and control subjects reflect a common pathological mechanism with yet undetermined subclinical or clinical manifestations.
- Our hypothesis that deep lesions penetrating across the gray and white matter of 2-3 gyri contribute to or cause early-onset epilepsy requires further studies. Other neurological correlates cannot be excluded.

Acknowledgements

The Autism Brain Atlas Project is supported by Autism Speaks, the Department of Defense Autism Spectrum Disorders Research Program (AS07334), and the NYS Office of Mental Retardation and Developmental Disabilities. The tissue was provided by the Harvard Brain Tissue Resource Center, Boston, MA (R24-MH068855), and the Brain and Tissue Bank at the University of Maryland, Baltimore, MD. Autism Tissue Program (Princeton, NJ) coordinated the selection of material for this study. Conflict of Interest: None