

Head Waves in Asian Marginal Seas

Peter H. Dahl

phone: (206) 543-2667 fax: (206) 543-6785 email:dahl@apl.washington.edu

Jee Woong Choi

phone: (206) 616-8597 fax: (206) 543-6785 email:choijw@apl.washington.edu

Applied Physics Laboratory
College of Ocean and Fisheries Sciences
University of Washington
Seattle, Washington 98105

Award Number: N00014-03-1-0271

LONG-TERM GOALS

To develop a robust means to utilize readily-obtained precursor (or head-wave-like) arrival measurements for purposes of geoaoustic inversion. Under certain conditions these arrivals can be classified as *interference head waves*.

These arrivals are important features of shallow water propagation that are exhibited at relatively short ranges (< 1 km). They are more directly linked with seabed properties, in contrast to long-range, trapped modal arrivals.

OBJECTIVES

The objective this year was to further the understanding of interference head waves, specifically in terms of their frequency and amplitude dependence on source and acquisition geometry. This builds on our analysis of precursor arrivals measured from the Yellow Sea as part of the August 1996 joint U.S.-China Yellow Sea Experiment that was published this year [1].

APPROACH

The approach involved revisiting a 1965 work by Chekin [2], along with studies by Červený and Ravindra [3] that first utilized the parameter ζ , which embodies source-to-range, sediment sound speed gradient, and sound frequency. Combining both works, an analytical expression for the frequency dependence of the interference head wave was derived that is a function of ζ . This expression was tested against simulations using the RAM Parabolic Equation (PE) code.

WORK COMPLETED

The analysis of the frequency dependence of interference head waves was completed for which a better understanding of interference head waves, and how they can be inverted for sound speed and attenuation in the sediment, was achieved. Results were published in the July issue of the *Journal of the Acoustical Society of America*. Some results from this work are summarized below.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2007		2. REPORT TYPE Annual		3. DATES COVERED 00-00-2007 to 00-00-2007	
4. TITLE AND SUBTITLE Head Waves In Asian Marginal Seas				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Washington, College of Ocean and Fisheries Sciences, Seattle, WA, 98105				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT To develop a robust means to utilize readily-obtained precursor (or head-wave-like) arrival measurements for purposes of geoacoustic inversion. Under certain conditions these arrivals can be classified as interference head waves.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

RESULTS

Figure 1 shows the basic structure of an interference head wave, which can be thought of as the superposition of C_n waves, $n=0,1,2, \dots$




Figure 1. (a) Sound-speed profile in seabed showing constant sound speed, zero gradient (dashed line) and non-zero positive gradient (solid line). (b) Equivalent ray path of a first-order head wave corresponding to the zero gradient case (dashed line) and multiple ray paths (C_n waves) corresponding to the positive gradient case (solid lines). The superposition of the C_n waves constitutes an interference head wave.

The interference head wave propagating through seabed sediment with a linear sound speed gradient is was studied as a function of the parameter ζ , that as mentioned above, is itself a function of frequency f , sediment sound speed gradient, and geometry. For small ζ the spectrum of the interference head wave goes as $|S(f)|/f$, where $S(f)$ is the source amplitude spectrum; as ζ increases the dominant frequency of the spectrum changes to that of the source spectrum. For a range of ζ values, the spectrum may display a shift in dominant frequency. This is illustrated in Fig. 2 that shows the changing behavior of the *received spectrum* of the interference head wave as a function of ζ .

Understanding this basic nature of the interference head wave in terms of its spectral properties, is essential for utilizing such arrivals in geoacoustic inversion. In view of the importance of the parameter ζ , we propose calling it the *Chekin parameter*, to recognize the 1965 work by Chekin [2] from which the parameter can be derived.




Figure 2. (a) Comparison of the interference head wave amplitude spectrum (thick gray line) predicted from the ray theory approach with that obtained from the PE simulation (thin solid line, and falling nearly entirely within gray line.) The thick solid line is amplitude spectrum of the source, and, for reference, the dashed line represents the amplitude spectrum of a first-order head wave. (b) The PE simulated time series of the interference head wave (within circle) from which the spectrum in (a) is estimated. The two arrival groups arriving after the interference head wave are the water-borne arrivals. The first group consists of the arrivals that have one or fewer interactions with the sea surface; these include the direct, bottom, sea surface, bottom-surface, surface-bottom, and bottom-surface-bottom paths. The second group consists of the arrivals that have interacted with the sea surface twice.

IMPACT/APPLICATIONS

Results of this research will constitute an important contribution to WESTPAC data bases. The technical approach refined in this work can be exploited to obtain geoacoustic bottom properties based on logistically simpler, short range measurements.

RELATED PROJECTS

The work on this project relates to a planned experiment off the Korean coastline, for which the principal organizers are NRL and the Korean Agency for Advanced Development (ADD). The PI and Dr. Choi (now working at Hanyang University in Korea) will participate in this experiment with the goal of obtaining field measurements of interference head waves or related precursor arrivals, from which geoacoustic inversions can be made.

REFERENCES

- [1] P. H. Dahl and J. W. Choi, "Precursor arrivals in the Yellow Sea, their distinction from first-order head waves, and their geoacoustic inversion," *J. Acoust. Soc. Am.*, 120 (6), 3525-3533, December 2006.
- [2] B. S. Chekin, "The effect on a head wave of small inhomogeneities in a refracting medium," *Bull. (Izv.) Acad. Sci. USSR, Earth Phys. Ser.*, 3, 143-147 (1965).
- [3] V. Červený and R. Ravindra, *Theory of Seismic Head Waves*, (University of Toronto Press, Toronto 1971), Chap. 6.

PUBLICATIONS

- J. W. Choi and P. H. Dahl, "Spectral properties of the Interference Head Wave," *J. Acoust. Soc. Am.*, 122 (6), 146-150, July 2007.
- P. H. Dahl and J. W. Choi, "Precursor arrivals in the Yellow Sea, their distinction from first-order head waves, and their geoacoustic inversion," *J. Acoust. Soc. Am.*, 120 (6), 3525-3533, December 2006.