

Improving Formulation of Marine Stable Boundary Layers Using CBLAST Weak Wind Data

Larry Mahrt

College of Oceanic and Atmospheric Sciences

Oregon State University

Corvallis, OR 97331

phone: (541) 737-5691 fax: (541) 737-2540 email: mahrt@coas.oregonstate.edu

Grant Number: N00014-05-1-0032

<http://blg.oce.orst.edu/cblast-weakwind>

LONG-TERM GOAL

The long-term goals are to better understand air-sea exchange of momentum, heat and moisture and its impact on the marine boundary layer and to better understand the influence of SST variations on the area-averaged (grid-averaged) flux-gradient relationship. Substantial modification of the formulation of the surface moisture flux for all conditions and the boundary layer for very stable conditions are also long term goals.

OBJECTIVES

The objectives are to quantify the influences of surface heterogeneity and strong stability on the transfer coefficients for the bulk formula and to formulate the difference between the transfer coefficients for momentum, heat and moisture.

APPROACH

Analyze the LongEZ eddy-correlation data in the CBLAST Weak Wind Pilot Experiment 2001 and compare with eddy-correlation data from the CBLAST WHOI ASIT tower during the intensive period and prior to the intensive period. LongEZ data from SHOWEX will be incorporated for comparison. We will also analyze eddy-correlation data collected by the CIRPAS Pelican during the CBLAST Weak Wind Experiment. The analyzed fluxes will be used for evaluation of regional modeling and LES simulations.

WORK COMPLETED

Work during the past year concentrated on examination of the wind and stress variability in the cases of weakest winds. We have also carried out joint work with Shouping Wang. We have developed a new roughness length formulation, which Shouping has tested in COAMPS. A short paper has been submitted to Geophysical Review Letters. We have not been unable to obtain the corrected data from the ASIT tower.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2007		2. REPORT TYPE Annual		3. DATES COVERED 00-00-2007 to 00-00-2007	
4. TITLE AND SUBTITLE Improving Formulation Of Marine Stable Boundary Layers Using CBLAST Weak Wind Data			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Oregon State University, College of Oceanic and Atmospheric Sciences, Corvallis, OR, 97331			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES code 1 only					
14. ABSTRACT The long-term goals are to better understand air-sea exchange of momentum, heat and moisture and its impact on the marine boundary layer and to better understand the influence of SST variations on the area-averaged (grid-averaged) flux-gradient relationship. Substantial modification of the formulation of the surface moisture flux for all conditions and the boundary layer for very stable conditions are also long term goals.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

RESULTS

With very weak winds, the wind and stress fields become dominated by background mesoscale motions, generally of unknown origin. For weak large-scale, the wind direction meanders in an unpredictable manner. This meandering often takes the form of sudden wind direction shifts in contrast to conventional understanding. A modified bulk formula has been developed, but much more evaluation is required before offering it to the modeling community. Numerical models do not capture such motions not only because of resolution issues, but also because such motions on resolved scales are seriously damped by explicit and implicit numerical horizontal diffusion required for well behaved numerical performance. In actuality, such mesoscale motions lead to shear-generation of turbulence and enhanced surface fluxes. In CBLAST Low, such mesoscale motions become important when the large-scale flow at 10 m decreases to less than 2 m/s. While such winds occur less than 10% of the time, they often lead to fog development. The large wind direction variability (Figure 1) prevents adjustment of the wind driven wave field, which is superimposed upon the normally larger-amplitude swell.




Figure 1. The standard deviation of the one-minute wind direction within one-hour records at the ASIT Tower. The mathematical upper limit of the standard deviation of wind direction is 104 degrees. Some weak-wind cases were eliminated because of fog formation and condensation on the sensors.




Figure 2. Roughness lengths for a) momentum, b) heat and c) moisture for the new formulation (solid curves), COARE v 2.5 (dashed) and COARE v 3.0 (dotted).

New formulations of the roughness lengths for momentum, heat and moisture were developed for the standard bulk flux formula with Monin-Obukov similarity theory (Vickers et al., 2007). The new formulations are based on turbulence measurements collected by the NOAA LongEZ in CBLAST Low and SHOWEX. The proposed roughness lengths are generally smaller than existing formulations, especially for moisture transport. Significant improvement is found for the latent heat flux compared to existing algorithms used by the COAMPS surface flux scheme and in the TOGA COARE v 2.5 and v 3.0 bulk flux algorithms. The reduced latent heat flux was verified using ASIT tower data (unpublished).

The more efficient transfer of heat compared to moisture is apparently related to the dynamic nature of temperature and the fact that temperature fluctuations dominate buoyancy generation of turbulence for our datasets. The proposed momentum roughness length (and thus the momentum flux) is smaller than given by existing algorithms in conditions of modestly weak winds and weak turbulence partly because of numerous cases of wind following swell. The proposed formulations need to be tested using independent datasets. The proposed formulation did not include the very weakest wind cases where the wind field is dominated by meandering mesoscale motions and the drag coefficient (roughness length for momentum) is apparently augmented by such motions.

RELATED PROJECTS

We are currently constructing a new surface flux scheme for Eric Skillingstad's LES model with the option for a TOGA COARE 2.6a -like method that includes the Charnock term plus a smooth flow term, and LKB roughness lengths for heat and moisture.

PUBLICATIONS

Vickers, D. and L. Mahrt, 2006: Evaluation of the air-sea bulk formula and sea-surface temperature variability. *J. Geophys. Res.* 111, 5002-5016. [published, refereed]

Skillingstad, S., D. Vickers, L. Mahrt and R. Samelson, 2006: Effects of Mesoscale Sea-Surface Temperature Fronts on the Marine Boundary Layer. *Boundary-Layer Meteorol.*, 123, 219-237. [published, refereed]

Mahrt L. 2007: Weak-wind mesoscale meandering in the nocturnal boundary layer. *Env. Fluid Mech.*, 7, 331-347. [published, refereed]

Vickers, D., L. Mahrt and S. Wang, 2007: New formulations for sea-surface roughness lengths. Submitted to *Geophys. Res. Letters*. [refereed]