

Award Number: W81XWH-08-1-0104

TITLE: The role of eIF4E activity in breast cancer

PRINCIPAL INVESTIGATOR: James N McElwaine, Ph.D.

CONTRACTING ORGANIZATION: University of Cambridge
Cambridge CB2 1T2

REPORT DATE: August 2010

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT:

Approved for public release; distribution unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. **PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.**

1. REPORT DATE (DD-MM-YYYY) 31-08-2010		2. REPORT TYPE Final		3. DATES COVERED (From - To) 01 AUG 2008 - 31 JUL 2010	
4. TITLE AND SUBTITLE The role of eIF4E activity in breast cancer				5a. CONTRACT NUMBER W81XWH-08-1-0104	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) McElwaine, JN Email: jnml1@amtp.cam.ac.uk				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) AND ADDRESS(ES) University of Cambridge 16 Mill Lane Cambridge CB2 1T2 UK				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research And Materiel Command Fort Detrick, MA 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Increased eIF4E expression occurs in many breast cancers and makes fundamental contributions to carcinogenesis by stimulating expression of cancer-related genes at post-transcriptional levels. This key role is highlighted by the facts that eIF4E levels can predict prognosis and that eIF4E is an established therapeutic target. However, eIF4E activity is a complex function of expression levels and phosphorylation statuses of eIF4E and its regulatory proteins. Our hypothesis was that combined analyses of these pathway components would allow insights into eIF4E activity and its influence on cancer. We have established that mathematically combining assessments of expressions of eIF4E-regulators with assessments of expression of eIF4E in clinical tumours provides improved prognostic insights over examination of eIF4E alone. In doing so we have determined the mathematical relationships between expression of each pathway component and pathway activity. Using human cell lines, we have experimentally demonstrated that this combinatorial estimate of eIF4E activity does indeed reflect eIF4E activity in breast cells. This represents a critical validation of our technique for estimation of eIF4E activity and allows us to propose that the estimate may act as a potentially powerful therapy predictive marker for cancer therapies directed against the eIF4E pathway.					
5. SUBJECT TERMS None provided.					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT UU	18. NUMBER OF PAGES 20	19a. NAME OF RESPONSIBLE PERSON USAMRMC
a. REPORT U	b. ABSTRACT U	c. THIS PAGE U			19b. TELEPHONE NUMBER

Standard Form 298 (Rev. 8-98)
Prescribed by ANSI Std. Z39.18

Table of Contents

	<u>Page</u>
Introduction.....	1
Body.....	1
Key Research Accomplishments.....	7
Reportable Outcomes.....	7
Conclusion.....	8
References.....	9
Appendices.....	10

1 Introduction

Increased expression of eIF4E has frequently been reported in breast cancers and is thought to make fundamental contributions to disease development and progression [1]. Increased eIF4E activity acts to enhance the translation of oncogenic cancer-related transcripts that contain highly structured 5' untranslated regions (UTRs) in their mRNAs. Over-expression of eIF4E has been shown to correlate with poor prognosis in breast cancer[2] therefore the level of eIF4E over-expression has been studied extensively as a prognostic marker with some success. Furthermore, eIF4E is an established target for cancer therapy [3] and clinical trials of the efficacy and safety of cancer therapeutics that target eIF4E have been carried out, again, with some success. However, it is clear that eIF4E expression may not equate to eIF4E activity, since eIF4E activity is additionally regulated by a family of binding proteins, the 4E-BPs, that bind to and inhibit eIF4E activity[4]. Our hypothesis was that analysis of eIF4E *activity* in individual breast tumours, as opposed to eIF4E *expression*, gives improved prognostic, predictive and biological understanding of individual breast cancers, and overall insights into the mechanisms of breast carcinogenesis.

In order to test this hypothesis we aimed to investigate eIF4E activity in breast carcinogenesis using a novel cross-disciplinary approach. First, we aimed to determine expression levels of eIF4E and its regulators in breast tumours, and to estimate the individual contributions of each factor to eIF4E activity by relating each expression level to cancer survival. This approach would allow us to build, in collaboration with mathematicians, an equation relating the levels of these factors together to estimate eIF4E activity. Next we aimed to validate these mathematical relationships in tissue culture systems where we would be able to measure both expression levels and eIF4E activity. Finally, we aimed to determine whether measures of eIF4E activity in cell lines, and estimates of eIF4E activity in tumours, would act as predictive markers for cellular responses to eIF4E directed therapies.

2 Body

Tissue micro-arrays containing samples from 424 breast tumours were established, supported by detailed clinicopathological data. The cohort included a wide range of patient and tumour characteristics, with mean patient follow-up of 91.9 months. We carried out IHC for eIF4E, 4E-BP1, 4E-BP2 and p4E-BP1 on TMA sections and assessed immunoreactivity within tumour cells, taking into account the proportions of cells staining positively and average intensity, giving scores of 0 (negative) or 27 (positive). The distribution of the scores is shown in Figure 2.

KaplanMeier survival analyses were used to determine survival with respect to eIF4E. Analyses were carried out with expression divided into IHC scores, although scores of 0, 2 and 3 were combined as each individual group was small, for overall survival (OS), disease-free survival (DFS) and disease-specific survival (DSS) (Figure 2). High eIF4E scores were indicative of poor prognosis. Prognosis seemed to worsen with each increasing eIF4E score for OS, whereas patterns for DFS and DSS suggested weaker, but still detectable, influences of individual scores with an overall grouping into two classes (05 good prognosis; 6 or 7 poor prognosis). We have examined relationships between eIF4E expression and survival using Cox regressions. We have included either eIF4E scores, or eIF4E expression dichotomised arbitrarily or as suggested by the apparent bimodal distribution seen above, and modelled these with respect to OS, DFS and DSS. Models that most accurately reflected the data included eIF4E scores rather than dichotomised data, showing the value of scoring proportion and intensity of positive tumour cells. In these models each increase of 1 in eIF4E score gave increases in hazard ratios (HRs) of 1.22 ($P = 0.004$), 1.3 ($P = 0.008$) and 1.33 ($P = 0.005$) for OS, DFS and DSS, respectively. Thus individuals with scores of 7 have DFS HRs of 6.15 (95% CIs: 312, $P = 0.008$) as compared with individuals with scores of 0. We have examined independence from the Nottingham Prognostic Index (NPI), which takes account of tumour size, grade and lymph node status. In multivariate Cox regressions the prognostic value of eIF4E was independent of NPI with eIF4E remaining significant for OS (NPI $P < 0.00001$; eIF4E $P = 0.02$), DFS (NPI $P < 0.00001$; eIF4E $P = 0.045$)




Figure 1: Distribution of Scores.




Figure 2: Expression of eukaryotic translation initiation factor 4E (eIF4E), but not eIF4E-binding protein (4E-BP) 1, 4E-BP2 or p4E-BP1, is associated with prognosis. (**A-C**) KaplanMeier survival analyses for overall (**A**), disease-free (**B**) and disease-specific survival (**C**) for patient groups with tumours with differing eIF4E scores. (**D-F**) KaplanMeier survival analyses for disease-free survival for patient groups with tumours with differing scores for 4E-BP1 (**D**), 4E-BP2 (**E**) or p4E-BP1 (**F**). As scores of 0, 2 and 3 were relatively rare, these have been grouped together. Censoring ticks have been omitted for clarity.




Figure 3: Additional prognostic information is gained by combining assessment of eukaryotic translation initiation factor 4E (eIF4E)-binding proteins (4E-BPs) with eIF4E analysis. KaplanMeier survival analyses of disease-free survival for patients with either high (6 or 7) eIF4E scores (**AC**) or high (45.4) Nottingham Prognostic Index scores (**DF**). Patients were dichotomised using either eIF4E expression (**A** and **D**), the y function (which includes eIF4E and 4E-BP2 scores) (**B** and **E**), or the z function (which includes scores for all 4 markers) (**C** and **F**).

and DSS (NPI $P < 0.00001$; eIF4E $P = 0.029$).

KaplanMeier survival analyses were also used to determine survival with respect to the other markers. We present data for DFS (Figure 2 **DF**). These antigens provided little prognostic insight and we were unable to construct significant Cox equations to model their individual survival influences. In the case of 4E-BP2, there was a nonsignificant trend for high scores to associate with good prognosis.

Next we examined influences of 4E-BPs in the context of eIF4E expression. We necessarily restricted these analyses to the 282 patients for whom scores of all four antigens were available. Using this dataset, models including solely eIF4E expression gave HRs of 1.21 ($P = 0.011$), 1.24 ($P = 0.035$) 1.27 ($P=0.02$) for OS, DF DSS, respectively, for each increase of 1 in eIF4E score. We were unable to construct significant Cox models on the basis of combinations of 4E-BPs without including eIF4E, suggesting that eIF4E is their critical effector. We found models combining expression of 4E-BP1 or p4E-BP1 (as IHC scores, or dichotomised into two groups) with eIF4E, provided little additional prognostic value over that found with eIF4E alone. However, including 4E-BP2 in a model for OS enhanced the model significantly with hazard increasing by 1.28 with each point increase in eIF4E score ($P = 0.005$) and decreasing 0.11 with each increase in 4E-BP2 scor ($P=0.02$). We refined this by combining eIF4E and 4E-BP2 scores into a single non-linear variable in which high levels of eIF4E or 4E-BP2 act to increase or decrease the value respectively (achieved using $y = \max(0, X - B2/3.5)$, where X and $B2$ represent eIF4E and 4E-BP2 scores). This variable y predicted survival more accurately than examination of eIF4E alone; each increase of 1 in y carried HRs of 1.32 ($P = 0.0003$), 1.32 ($P = 0.013$) and 1.36 ($P = 0.006$) for OS, DFS and DSS, respectively. We also investigated statistically nonsignificant. Cox models including expression of all four antigens. In order to combine terms, we considered models that included each variable separately and found their relative effect on HRs using maximum likelihood estimation. Although individual components were not statistically significant and had only little effects on likelihoods (with the exception of eIF4E), a combination gave improved prognostic power. This variable defined as $z = XB1/4 + PB1/2 - B2/4$, where $B1$ and $PB1$ represent 4E-BP1 and p4E-BP1. Each increase of 1 in z gave HRs of 1.15 ($P = 0.006$), 1.26 ($P = 0.002$) and 1.28 ($P = 0.0008$) for OS, DFS and DSS, respectively. This variable has a highly significant relationship with survival but this should be treated with caution because the constants were determined using regressions for OS, and, thus to an extent, significance is self fulfilling, at least for OS. The utility of z , however, is supported by the fact that its relationship with S and survival is more significant with DFS and DSS than OS, a result not predetermined by the approach. We also examined whether y or z give prognostic insights independently of NPI using multivariate analyses. NPI and either y or z remain significant in models for DFS (NPI $P < 0.00001$; y $P = 0.04$ or z $P = 0.03$) and DSS (NPI $P < 0.00001$; y $P = 0.02$ or z $P = 0.02$). The statistical significance of relationships of y and z with survival show additional prognostic value from examining multiple eIF4E pathway components. In addition, we have shown the value of these variables using KaplanMeier analyses. First, we focused on patients with high eIF4E scores (6 or 7), as it is in this context that differential expression of 4E-BPs would be most relevant. Patients with eIF4E scores 6 or 7 have a relatively poor prognosis (Figure 2**AC**), but no difference was detected between groups scored as 6 or 7 in terms of DFS (Figure 2 **B** and **A**). When y was applied to this cohort some discrimination occurred with improved prognosis for patients whose y scores were lowered by 4E-BP2 (Figure 2**B**), although the discrimination remained When z was applied to this cohort further discrimination occurred (Figure 2**C**) showing how 4E-BPs affect patient outcome through eIF4E.

Rapamycin and derivatives have been used as cancer therapeutics with limited success. Growth of some cancers is inhibited effectively while a proportion appears relatively resistant to the drugs. Efficacy could be improved by use of predictive biomarkers to allow selection and treatment of only patients with cancers likely to respond to these drugs. The drugs inhibit the kinase mTOR, which is frequently up-regulated in cancer, and efforts to establish predictive markers have centred on determining the phosphorylation states of mTOR itself or of the primary targets of mTORs action, 4E-BP1 or S6K, as each of these may relate to mTOR activity directly. In an alternative approach, we determined the functional activity of a key effector pathway for mTOR in cell lines, and demonstrated that this




Figure 4: Translational efficiencies specified by structured 5UTRs correlate with sensitivities to rapamycin. Data from Figure 1B (y-axis) and 4B (x-axis) were plotted for each cell line. Linear regression was performed to determine the relationship in the cell lines excluding the outlier MCF7 cells; the linear model is shown as a line (black) with 95% confidence intervals (grey lines) ($P = 0.0037$).

predicts sensitivity to rapamycin.

We determined the sensitivities to rapamycin of a panel of cell lines, including lines representative of colon, lung and breast cancer; more than 3 fold variation was seen in sensitivity. Up-regulated mTOR enhances the translational efficiency of mRNAs with structured 5 untranslated regions, as a result of increased activities of the translation factors eIF4E and eIF4A. We established and validated an assay to measure these translational efficiencies. These translational efficiencies were predictive of rapamycin sensitivity ($p = 0.0037$), with more efficient translation indicative of enhanced sensitivity. Levels of phosphorylated 4E-BP1 were unrelated to rapamycin sensitivity. We concluded that assessment of the functional impact of mTOR activity gave greater predictive value than measurement of the levels of phosphorylation of its direct targets.

Relative translational efficiencies specified by this eIF4E-responsive 5'UTR were determined in the panel of cell lines. Cells were transiently transfected with vectors to allow expression of GFP mRNAs with control or the structured 5'UTR as before, and translational efficiencies were determined. A range of translational efficiencies was seen, with A549 cells determining the lowest, and HB2 cells the highest. Surprisingly, the two cells lines of non-cancer origin (HB2 and MCF10A) were found to determine relatively efficient translation from the reporter 5'UTR. Importantly, translational efficiencies specified by the eIF4E-responsive 5'UTR correlated with rapamycin sensitivity (Figure 2). Initially, we analysed this relationship using Spearmans rho; we found a strong and significant positive association (rho coefficient: 0.72; $p=0.037$). However, the correlation was particularly evident in 8 of the 9 cell lines; if MCF7 cells, which were more sensitive to rapamycin than predicted, were excluded from the analysis the strength and significance of the relationship was increased (rho coefficient: 0.83; $p = 0.015$). Similarly in linear regression, a highly significant relationship was seen when MCF7 cells were excluded from the analysis (Figure 2; $p = 0.0037$).

3 Key Research Accomplishments

1) We have established that mathematically combining assessments of expressions of eIF4E-regulators with assessments of expression of eIF4E in clinical tumours provides improved prognostic insights over examination of eIF4E alone, presumably as this combinatorial function estimates eIF4E *activity*. This work has been published [5].

2) We have demonstrated that an experimentally determined measure of eIF4E activity predicts cellular sensitivity to rapamycin (manuscript under revision), suggesting that measurement of eIF4E activity within tumours could be used as a predictive marker for therapy with RAD001 or other rapamycin-derivatives in cancer.

3) We are in the process of testing whether our combinatorial estimate of eIF4E activity in tumours can be used as a predictive marker for neo-adjuvant RAD001 therapy in breast cancer.

4 Reportable outcomes

This work has led directly to two publications, one of which is currently under review.

Coleman LJ, Peter MB, Teall TJ, Brannan RA, Hanby AM, Honarpisheh H, Shaaban AM, Smith L, Speirs V, Verghese ET, **McElwaine JN** & Hughes TA (2009) Combined analysis of eIF4E and 4E-binding protein expression predicts breast cancer survival and estimates eIF4E activity. *Br. J. Cancer.* 100, 1393–9.

Satheesha S, Coleman LJ, Ingram N, Madhok B, **McElwaine JN**, Hughes TA. Translational efficiencies defined by a structured 5 untranslated region predict cellular sensitivity to rapamycin. Under review.

5 Conclusions

We have tested and supported our main hypothesis: analysis of eIF4E activity in breast tumours gives improved prognostic insights into breast cancer over analysis of only eIF4E expression.

We have also demonstrated that measurement of eIF4E activity provides a predictive marker for response of cell lines to treatment with the drug rapamycin. Work is still underway to establish whether our clinically practicable method of estimating eIF4E activity in tumour samples provides a predictive marker for response of breast tumours to neo-adjuvant treatment with the rapamycin-derivative RAD001.

Finally, we have developed a generally-applicable cross-disciplinary methodology for combining the prognostic/predictive insights from individual biomarkers into estimates of the activities of biological pathways. This approach could be applied to estimate pathway activities and therefore predict response for any molecularly-directed cancer therapy. The most obvious example is the her2 pathway that is targeted by herceptin (trastuzumab). This therapy is assigned on the basis of over-expression of the biomarker and molecular target her2. However, a significant proportion of patients fail to respond to the therapy, while it has been reported that some her2-negative tumours do respond. It is likely that an estimate of her2 activity, based on combined assessments of expression of her2 and its direct regulators, would provide an improved predictive biomarker.

JN McElwaine received support from this grant.

References

- [1] De Benedetti A & Graff JR (2004) eIF-4E expression and its role in malignancies and metastases. *Oncogene* 23, 3189-99
- [2] Li BD, McDonald JC, Nassar R & De Benedetti A (1998) Clinical outcome in stage I to III breast carcinoma and eIF4E overexpression. *Ann Surg* 227, 756-61; discussion 61-3
- [3] Graff JR, Konicek BW, Carter JH & Marcusson EG (2008) Targeting the eukaryotic translation initiation factor 4E for cancer therapy. *Cancer Res* 68, 631-4
- [4] Richter JD & Sonenberg N (2005) Regulation of cap-dependent translation by eIF4E inhibitory proteins. *Nature* 433, 477-80
- [5] Coleman LJ, Peter MB, Teall TJ, Brannan RA, Hanby AM, Honarpisheh H, Shaaban AM, Smith L, Speirs V, Verghese ET, McElwaine JN & Hughes TA (2009) Combined analysis of eIF4E and 4E-binding protein expression predicts breast cancer survival and estimates eIF4E activity. *Br J Cancer* 100, 1393-9
- [6] Hughes TA & Brady HJM (2005) Expression of axin2 is regulated by the alternative 5' untranslated regions of its mRNA. *J Biol Chem* 280, 8581-8
- [7] Gray NK & Hentze MW (1994) Regulation of protein synthesis by mRNA structure. *Mol Biol Rep* 19, 195-200
- [8] Smith L, Coleman LJ, Cummings M, Satheesha S, Shaw SO, Speirs V & Hughes TA (2010) Expression of oestrogen receptor beta isoforms is regulated by transcriptional and post-transcriptional mechanisms. *Biochem J* 429, 283-90
- [9] Meric-Bernstam F & Gonzalez-Angulo AM (2009) Targeting the mTOR signaling network for cancer therapy. *J Clin Oncol* 27, 2278-87
- [10] Dowling RJ, Topisirovic I, Fonseca BD & Sonenberg N (2010) Dissecting the role of mTOR: Lessons from mTOR inhibitors. *Biochim Biophys Acta* 1804, 433-9

6 Appendices

Combined analysis of eIF4E and 4E-binding protein expression predicts breast cancer survival and estimates eIF4E activity

LJ Coleman^{1,5}, MB Peter^{2,5}, TJ Teall^{1,5}, RA Brannan³, AM Hanby^{1,3}, H Honarpisheh¹, AM Shaaban³, L Smith¹, V Speirs¹, ET Verghese³, JN McElwaine⁴ and TA Hughes^{*,1}

¹Leeds Institute of Molecular Medicine, Leeds University, Leeds LS9 7TF, UK; ²Department of Surgery, LGI, Leeds LS1 3EX, UK; ³Department of Histopathology, SJUH, Leeds LS9 7TF, UK; ⁴Department of Applied Mathematics and Theoretical Physics, Cambridge University, Cambridge CB2 0WA, UK

Increased eukaryotic translation initiation factor 4E (eIF4E) expression occurs in many cancers, and makes fundamental contributions to carcinogenesis by stimulating the expression of cancer-related genes at post-transcriptional levels. This key role is highlighted by the facts that eIF4E levels can predict prognosis, and that eIF4E is an established therapeutic target. However, eIF4E activity is a complex function of expression levels and phosphorylation statuses of eIF4E and eIF4E-binding proteins (4E-BPs). Our hypothesis was that the combined analyses of these pathway components would allow insights into eIF4E activity and its influence on cancer. We have determined expression levels of eIF4E, 4E-BP1, 4E-BP2 and phosphorylated 4E-BP1 within 424 breast tumours, and have carried out analyses to combine these and relate the product to patient survival, in order to estimate eIF4E activity. We show that this analysis gives greater prognostic insights than that of eIF4E alone. We show that eIF4E and 4E-BP expression are positively associated, and that 4E-BP2 has a stronger influence on cancer behaviour than 4E-BP1. Finally, we examine eIF4E, estimated eIF4E activity, and phosphorylated 4E-BP1 as potential predictive biomarkers for eIF4E-targeted therapies, and show that each determines selection of different patient groups. We conclude that eIF4E's influence on cancer survival is modulated substantially by 4E-BPs, and that combined pathway analyses can estimate functional eIF4E.

British Journal of Cancer (2009) **100**, 1393–1399. doi:10.1038/sj.bjc.6605044 www.bjcancer.com

Published online 14 April 2009

© 2009 Cancer Research UK

Keywords: translation factors; predictive biomarkers; Cox regression; pathway biomarker

The eukaryotic translation initiation factor 4E (eIF4E) has key roles in carcinogenesis (De Benedetti and Graff, 2004). eIF4E is often overexpressed in carcinoma cells as compared to equivalent normal epithelium in many tumour types including breast (Kerekatte *et al*, 1995), lung (Rosenwald *et al*, 2001) and colon (Rosenwald *et al*, 1999). The oncogenic role of this overexpression has been shown by various experimental observations; for example, forced eIF4E overexpression within many cell types leads to transformation (De Benedetti and Graff, 2004), and within transgenic mice increases incidence of multiple tumour types (Ruggero *et al*, 2004). eIF4E has at least two normal cellular functions. First, it is an essential component of the multimeric factor eIF4F, which initiates cap-dependent translation – the mechanism responsible for most protein synthesis (Gray and Wickens, 1998). eIF4E's role is to bind to mRNA caps allowing recruitment of eIF4F, and subsequently the translational machinery. The complex formed scans linearly along the 5' untranslated region (UTR) until an initiation codon in good context is encountered, at which point further elements of translational machinery are recruited and protein synthesis starts. Second,

eIF4E regulates expression of some genes by controlling nuclear export of their transcripts (Culjkovic *et al*, 2007), a function that also requires eIF4E's cap-binding activity (Culjkovic *et al*, 2005). Under most normal conditions, availability of active eIF4E is thought to be rate limiting for both functions. One might expect a general translational stimulation to result from the increased eIF4E expression in cancers, on account of enhanced mRNA cap recognition, yet effects of eIF4E overexpression are more subtle. Approximately 10% of mammalian transcripts have 5'UTRs that may form complex secondary structures that reduce the abilities of both eIF4F to bind to mRNAs and the translational machinery to scan 5'UTRs (Pesole *et al*, 2001); the result is that these transcripts are translated inefficiently (Hughes, 2007). The majority of human transcripts with these inhibitory 5'UTRs code for growth or cancer-associated proteins (Kozak, 1991). Increased eIF4E is thought to reduce the effects of 5'UTR structure by enhancing cap-recognition and scanning, therefore increasing translation of these specific oncogenic transcripts (De Benedetti and Graff, 2004). Similarly, increased eIF4E expression enhances nuclear export of a set of transcripts associated with oncogenesis (Culjkovic *et al*, 2007). As a consequence of this central role, eIF4E is an established target for cancer therapy (Smolewski, 2006; Graff *et al*, 2008).

The importance of eIF4E in cancer has been underlined by the fact that eIF4E expression levels can be used to determine prognosis. Cases in which eIF4E is highly overexpressed tend to have poor prognoses (Li *et al*, 1998). A substantial confounding factor is that eIF4E expression does not equate to eIF4E activity, thereby making interpretation of potential influences of eIF4E

*Correspondence: Dr TA Hughes, Leeds Institute of Molecular Medicine, St James's University Hospital, Leeds University, Leeds, LS9 7TF, UK; E-mail: t.hughes@leeds.ac.uk

⁵These authors contributed equally to this work.

Received 27 January 2009; revised 18 March 2009; accepted 25 March 2009; published online 14 April 2009

levels difficult. eIF4E activity is a complex function of eIF4E expression and expressions and activities of eIF4E-binding proteins (4E-BP1, 2 and 3) that bind to and inhibit eIF4E (Richter and Sonenberg, 2005) (Supplementary Figure S1). Activity is further regulated by phosphorylation of 4E-BP1 (and other 4E-BPs it is assumed), with only hypophosphorylated forms being able to inhibit eIF4E. Additional regulation occurs by differential phosphorylation of eIF4E itself, although there are conflicting reports as to how this influences activity (Scheper and Proud, 2002). The result is that high expression of eIF4E may not lead to high eIF4E activity if, for example, hypophosphorylated 4E-BP1 were also highly expressed. Many cancer-related signalling pathways, including PI3K and p38, converge to regulate eIF4E and 4E-BP phosphorylation; therefore, eIF4E activity seems to be a key cancer-signalling node (Polunovsky and Bitterman, 2006). Here, we have tested the hypothesis that combined analyses of expressions and phosphorylation states of eIF4E, and its regulators allows greater understanding of eIF4E activity and its influence on cancer than examination of eIF4E expression alone.

MATERIALS AND METHODS

Patients

Ethical approval was obtained (Leeds East 05/Q1206/136). Archival cancer tissue and data were obtained for 424 patients diagnosed at LTH NHS Trust from 1983–2006. Tissue microarrays (TMAs) were constructed containing 0.6 mm cores selected from representative tumour areas as determined by a consultant breast histopathologist (AMS) from H&E stained sections. Survival periods – overall: initial diagnosis to death; disease-free: initial diagnosis to the diagnosis of recurrence/metastasis; disease-specific: initial diagnosis to death after recurrence or metastasis (cancer-specific death confirmed in most cases).

Westerns and immunohistochemistry

MCF7 and MDA-MB-231 cells were cultured/transfected as earlier (Johnson *et al*, 2008; Maraqa *et al*, 2008). An eIF4E expression vector was obtained from John Blenis (Harvard Medical School) and Nahum Sonenberg (McGill). Western analyses were carried out as earlier (Maraqa *et al*, 2008) using the reagents in Supplementary Table S1. TMA sections of 5 μ m were dewaxed and blocked in hydrogen peroxidase block (20 min). Antigens were retrieved and stained as described in Supplementary Table S1, and as used elsewhere (Zhou *et al*, 2004; Dutton *et al*, 2005; Lee *et al*, 2005; Engelman *et al*, 2008). Envision detection was used (DAKO, Glostrup, Denmark). Negative controls (primary antibodies omitted) were included in each immunohistochemistry (IHC) batch; in addition, adjacent normal epithelium, lymphocytes and blood vessel endothelium served as internal controls. Controls were performed for p4E-BP1 antibodies in which sections were pretreated with Lambda Phosphatase (Nebraska, NE, USA). Cores were scored for immunoreactivity by two or more individuals (LJC, TJT, ETV and RAB), taking into account the average intensity and percentage of positively stained tumour cells (as used earlier for eIF4E (Zhou *et al*, 2006)). Staining intensity scores (0 no staining, 1 weak, 2 moderate and 3 strong) were added to percentages positively stained scores (1 <5%; 2, 6–25%, 3, 26–75% and 4 >75%), giving totals of 0 or 2–7. Consensus scores were determined for cores with different initial scores, and all scoring was overseen by a consultant breast histopathologist (AMH).

Mathematical analyses

Data were analysed using Kaplan–Meier survival curves. Dependence on prognostic indicators was determined using Cox proportional hazards models; significance values relate to likelihood ratio

tests of the null hypothesis that indicators do not effect hazard rates (Cox and Oakes, 1984). SPSS (SPSS, Chicago, IL, USA) and the Statistics Toolbox in MATLAB (ecdf.m and coxphfit.m) (MathWorks, Natick, MA, USA) were used. Tests were two sided and $P < 0.05$ was considered significant.

RESULTS

Antibody validation

We have used IHC to determine expression levels in breast tumours of the main regulatory molecules of the eIF4E pathway – namely, eIF4E, 4E-BP1, 4E-BP2 and phosphorylated 4E-BP1 (Thr37/46) (termed p4E-BP1). We have not examined 4E-BP3 because it is not thought to have a role in breast (Poulin *et al*, 1998), or phosphorylated forms of eIF4E, as their influences on the activity and in cancer remain uncertain (Scheper and Proud, 2002; Salehi and Mashayekhi, 2006; Buxade *et al*, 2008). First, we optimised the antibody use on archival breast tissue. We established that antibodies were specific for their antigens using western blots against lysates of breast cancer cell lines (Supplementary Figure S2A). In addition, we showed phospho-specificity of antibodies against p4E-BP1 by carrying out IHC on serial tissue sections with and without pretreatment with protein phosphatase (Supplementary Figure S2B).

Patient cohort and immunohistochemistry

Tissue micro-arrays containing samples from 424 breast tumours were established, supported by detailed clinicopathological data (Supplementary Table S2). The cohort included a wide range of patient and tumour characteristics, with mean patient follow-up of 91.9 months. We carried out IHC for eIF4E, 4E-BP1, 4E-BP2 and p4E-BP1 on TMA sections and assessed immunoreactivity within tumour cells, taking into account the proportions of cells staining positively and average intensity, giving scores of 0 (negative) or 2–7 (positive). Representative staining patterns are shown (Figure 1, and at higher magnification in Supplementary Figure S3). Tumour stroma and normal tissue were negative for eIF4E, 4E-BP1 and 4E-BP2, whereas very occasional low-intensity staining for p4E-BP1 was noted in normal epithelial cells. Staining was generally cytoplasmic, although nuclear staining was noted in a minority of cases (Supplementary Figure S4); this was separately analysed and was found not to be of prognostic value and is not discussed. As expected, data were not available for some patients because of the TMA core loss during processing, a well-recognised occurrence, therefore, data for all four antigens were available for only 282 patients. The full range of scores were observed for each antigen (Figure 1). It was notable that staining was most frequently not detectable for p4E-BP1. Others have reported more frequent expression of p4E-BP1 (Zhou *et al*, 2004), therefore we carried out IHC for an alternative p4E-BP1 species (Ser65); we found immunoreactivity with this antibody to be similarly infrequent (see discussion).

Expressions of eIF4E and 4E-BPs correlate with grade

Associations between antigen expressions and a wide range of clinicopathological parameters were examined. No correlations were found with nodal status, tumour size or histological type. Weak positive/borderline no correlations were found with oestrogen receptor α status and eIF4E expression (Spearman's ρ coefficient 0.21; $P < 0.001$) and 4E-BP2 (0.22; $P < 0.001$), but not with 4E-BP1 or p4E-BP1. Strong correlations between expression of markers and tumour grade were found. eIF4E expression (split into three classes, 0–3, 4–5 and 6–7) was positively associated with grade (χ^2 -test, $P = 0.011$), whereas expression of both 4E-BPs was negatively associated with grade




Figure 1 The full range of expression intensities and proportions for eukaryotic translation initiation factor 4E (eIF4E), eIF4E-binding protein (4E-BP) 1, 4E-BP2 and p4E-BP1 occur within breast cancers. (A–H) Representative tumour tissue microarray cores showing immunoreactivity as labelled. These cores were scored A 4, B 7, C 3, D 7, E 3, F 7, G 3 and H 7. (I) Histograms showing distributions of immunohistochemistry scores within breast cancers. Scores (x-axis) and numbers of cores assigned to each score (y-axis) are shown. Percentages of the cohort are given above the bar for each score.

(4E-BP1 $P=0.002$; 4E-BP2 $P=0.029$). p4E-BP1 was positively associated with grade ($P=0.012$). A positive association between eIF4E expression and grade has been reported earlier (Li *et al*, 2002). Correlations for other markers were consistent with their influences on carcinogenesis being through the eIF4E pathway; 4E-BPs, eIF4E inhibitors, were negatively associated with grade, whereas 4E-BP1 phosphorylation, which would relieve 4E-BP1-induced inhibition of eIF4E, was positively associated.

High expression of eIF4E correlates with poor prognosis

Kaplan–Meier survival analyses were used to determine survival with respect to eIF4E. Analyses were carried out with expression divided into IHC scores, although scores of 0, 2 and 3 were combined as each individual group was small, for overall survival (OS), disease-free survival (DFS) and disease-specific survival (DSS) (Figure 2A–C). High eIF4E scores were indicative of poor prognosis. Prognosis seemed to worsen with each increasing eIF4E score for OS, whereas patterns for DFS and DSS suggested weaker, but still detectable, influences of individual scores with an overall grouping into two classes (0–5 good prognosis; 6 or 7 poor prognosis). We have examined relationships between eIF4E expression and survival using Cox regressions. We have included either eIF4E scores, or eIF4E expression dichotomised arbitrarily

or as suggested by the apparent bimodal distribution seen above, and modelled these with respect to OS, DFS and DSS. Models that most accurately reflected the data included eIF4E scores rather than dichotomised data, showing the value of scoring proportion and intensity of positive tumour cells. In these models each increase of 1 in eIF4E score gave increases in hazard ratios (HRs) of 1.22 ($P=0.004$), 1.3 ($P=0.008$) and 1.33 ($P=0.005$) for OS, DFS and DSS, respectively. Thus individuals with scores of 7 have DFS HRs of 6.15 (95% CIs: 3–12, $P=0.008$) as compared with individuals with scores of 0. The prognostic value of eIF4E has been reported earlier as independent of grade/nodal status in breast cancer (Li *et al*, 1998); we have examined independence from the Nottingham Prognostic Index (NPI), which takes account of tumour size, grade and lymph node status (Haybittle *et al*, 1982). In multivariate Cox regressions the prognostic value of eIF4E was independent of NPI with eIF4E remaining significant for OS (NPI $P<0.00001$; eIF4E $P=0.02$), DFS (NPI $P<0.00001$; eIF4E $P=0.045$) and DSS (NPI $P<0.00001$; eIF4E $P=0.029$).

Prognostic value of 4E-BP1, 4E-BP2 and p4E-BP1

Kaplan–Meier survival analyses were also used to determine survival with respect to the other markers. We present data




Figure 2 Expression of eukaryotic translation initiation factor 4E (eIF4E), but not eIF4E-binding protein (4E-BP) 1, 4E-BP2 or p4E-BP1, is associated with prognosis. (A–C) Kaplan–Meier survival analyses for overall (A), disease-free (B) and disease-specific survival (C) for patient groups with tumours with differing eIF4E scores. (D–F) Kaplan–Meier survival analyses for disease-free survival for patient groups with tumours with differing scores for 4E-BP1 (D), 4E-BP2 (E) or p4E-BP1 (F). As scores of 0, 2 and 3 were relatively rare, these have been grouped together. Censoring ticks have been omitted for clarity.

for DFS (Figure 2D–F), and for OS and DSS (Supplementary Figure S5). These antigens provided little prognostic insight and we were unable to construct significant Cox equations to model their individual survival influences. In the case of 4E-BP2, there was a nonsignificant trend for high scores to associate with good prognosis.

Mathematical modelling of influences of 4E-BPs: 4E-BPs modify eIF4E activity and provide additional prognostic insights

Next we examined influences of 4E-BPs in the context of eIF4E expression. We necessarily restricted these analyses to the 282 patients for whom scores of all four antigens were available. Using this dataset, models including solely eIF4E expression gave HRs of 1.21 ($P=0.011$), 1.24 ($P=0.035$) 1.27 ($P=0.02$) for OS, DFS and DSS, respectively, for each increase of 1 in eIF4E score. We were unable to construct significant Cox models on the basis of combinations of 4E-BPs without including eIF4E, suggesting that eIF4E is their critical effector. We found models combining expression of 4E-BP1 or p4E-BP1 (as IHC scores, or dichotomised into two groups) with eIF4E, provided little additional prognostic value over that found with eIF4E alone. However, including 4E-BP2 in a model for OS enhanced the model significantly with hazard increasing by 1.28 with each point increase in eIF4E score ($P=0.005$) and decreasing 0.11 with each increase in 4E-BP2 score ($P=0.02$). We refined this by combining eIF4E and 4E-BP2 scores into a single non-linear variable in which high levels of eIF4E or 4E-BP2 act to increase or decrease the value respectively (achieved using $\max(0, X-B2/3.5)$, where X and B2 represent eIF4E and 4E-BP2 scores). This variable, termed 'y', predicted survival more accurately than examination of eIF4E alone; each increase of 1 in y carried HRs of 1.32 ($P=0.0003$), 1.32 ($P=0.013$) and 1.36 ($P=0.006$) for OS, DFS and DSS, respectively. We also investigated

Cox models including expression of all four antigens. In order to combine terms, we considered models that included each variable separately and found their relative effect on HRs using maximum likelihood estimation. Although individual components were not statistically significant and had only little effects on likelihoods (with the exception of eIF4E), a combination gave improved prognostic power. This variable termed 'z' can be described as $X-B1/4 + PB1/2-B2/4$, where B1 and PB1 represent 4E-BP1 and p4E-BP1. Each increase of 1 in z gave HRs of 1.15 ($P=0.006$), 1.26 ($P=0.002$) and 1.28 ($P=0.0008$) for OS, DFS and DSS, respectively. This variable has a highly significant relationship with survival but this should be treated with caution because the constants were determined using regressions for OS, and, thus to an extent, significance is self fulfilling, at least for OS. The utility of z, however, is supported by the fact that its relationship with survival is more significant with DFS and DSS than OS, a result not predetermined by the approach. We also examined whether y or z give prognostic insights independently of NPI using multivariate analyses. NPI and either y or z remain significant in models for DFS (NPI $P<0.00001$; y $P=0.04$ or z $P=0.03$) and DSS (NPI $P<0.00001$; y $P=0.02$ or z $P=0.02$).

The statistical significance of relationships of y and z with survival show additional prognostic value from examining multiple eIF4E pathway components. In addition, we have shown the value of these variables using Kaplan–Meier analyses. First, we focused on patients with high eIF4E scores (6 or 7), as it is in this context that differential expression of 4E-BPs would be most relevant. Patients with eIF4E scores 6 or 7 have a relatively poor prognosis (Figure 2A–C), but no difference was detected between groups scored as 6 or 7 in terms of DFS (Figures 2B and 3A). When y was applied to this cohort some discrimination occurred with improved prognosis for patients whose y scores were lowered by 4E-BP2 (Figure 3B), although the discrimination remained statistically nonsignificant. When z was applied to this cohort




Figure 3 Additional prognostic information is gained by combining assessment of eukaryotic translation initiation factor 4E (eIF4E)-binding proteins (4E-BPs) with eIF4E analysis. Kaplan–Meier survival analyses of disease-free survival for patients with either high (6 or 7) eIF4E scores (**A–C**) or high (>5.4) Nottingham Prognostic Index scores (**D–F**). Patients were dichotomised using either eIF4E expression (**A** and **D**), the y function (which includes eIF4E and 4E-BP2 scores) (**B** and **E**), or the z function (which includes scores for all 4 markers) (**C** and **F**).

further discrimination occurred (Figure 3C) showing how 4E-BPs affect patient outcome through eIF4E. Second, we have focused on patients with high NPI (and consequently poor DFS, Supplementary Figure S6). These patients were further stratified according to eIF4E expression (cutoff 5.5 as suggested by the distribution in Figure 2B) into separate groups (Figure 3D). As before, when z was applied (Figure 3F) further discrimination occurred allowing identification of patients with very poor (high z), or relatively good prognosis (low z). In this case, z discriminated into statistically significantly different groups (Figure 3F Log rank $P = 0.039$) when use of eIF4E alone was not significant (Figure 3D Log rank $P = 0.15$). In this case y was substantially less successful as a prognostic indicator (Figure 3E Log rank $P = 0.5$) showing the importance of combining all four components.

Expressions of eIF4E and 4E-BPs are positively associated

Although including assessments of 4E-BPs provided additional prognostic information over that from only eIF4E, we were surprised that influences of 4E-BPs, especially of 4E-BP1, were relatively weak. One explanation for this was that expression levels of eIF4E and 4E-BPs were not independent of each other. This would mean that at a given eIF4E level, differential expression of 4E-BPs – therefore differential modification of eIF4E activity – would be relatively rare, thereby minimising apparent influences of 4E-BPs in our analyses. Associations between marker expressions were examined using Spearman's correlation tests (Table 1). eIF4E expression showed moderate positive associations with expression of both 4E-BPs ($P < 0.0001$), although weak positive/no association with p4E-BP1. Expression of 4E-BPs was also moderately positively associated with each other ($P = 0.02$). Expression of p4E-BP1 was positively associated with 4E-BP1 ($P < 0.0001$) (expected as 4E-BP1 must be expressed to be phosphorylated).

Table 1 Expressions of eIF4E, 4E-BP1 and 4E-BP2 are positively associated

	eIF4E	4E-BP1	4E-BP2	p4E-BP1
eIF4E	—	0.31	0.34	0.21
4E-BP1	0.31	—	0.37*	0.36
4E-BP2	0.34	0.37*	—	0.14
p4E-BP1	0.21	0.36	0.14	—

Abbreviations: eIF4E = eukaryotic translation initiation factor 4E; 4E-BP = eIF4E-binding protein; p4E-BP1 = phosphorylated 4E-BP1. Spearman's ρ correlation figures are shown. Associations are either moderately (* $P = 0.02$) or highly significant (others $P < 0.0001$).

eIF4E activity scores are potential biomarkers for eIF4E-targeted therapies

The eIF4E pathway is a target for cancer therapy with drugs that either inhibit eIF4E activity indirectly (mTOR inhibitors that reduce 4E-BP1 phosphorylation thereby inhibiting eIF4E (Smolewski, 2006)), or directly (by binding to/reducing expression of eIF4E (Graff *et al*, 2008)). However, a concern with these agents is toxicity resulting from general translational repression. Consequently, selection of patients who are most likely to benefit from such agents using predictive biomarkers may aid their efficacy. Three selection criteria are apparent: individuals with highest eIF4E levels, highest eIF4E activities or highest levels of phosphorylated 4E-BP1 (especially relevant for mTOR inhibitors as these act by reducing 4E-BP phosphorylation). We have compared selection of potential treatment groups using these criteria from our cohort for whom scores of all four antigens were available: first (group 1), those with eIF4E scores of 7 (37/282; 13.1%); second (group 2), those with high estimated pathway activities ($z \geq 5.75$, cutoff chosen to give a similar-sized group,




Figure 4 Use of biomarkers for dividing patients into potential treatment groups for eukaryotic translation initiation factor 4E (eIF4E)-directed therapy; different markers select very different groups. A Venn diagram demonstrating relationships between potential treatment groups selected on the basis of high eIF4E expression (1), high estimated eIF4E activity ('z') (2), and high p4E-BP1 expression (3).

38/282; 13.5%); finally (group 3), those with p4E-BP1 scores of 6 or 7 (26/282; 9.2%). We have examined whether biomarker choice leads to selection of different potential treatment groups (Figure 4). Substantially different patient groups were selected using each potential predictive biomarker. Only 10 patients appeared in all groups (16% of total patients selected in any group). Group 1 contained the largest proportion of uniquely selected individuals (56% of total), whereas the majorities of groups 2 and 3 (74 and 81% respectively) overlapped with at least one other group. Group 2 (high z values) had large overlaps with both other groups (46% of group 1 and 81% of group 3). In addition, group 2 included all individuals with high eIF4E and p4E-BP1 (i.e. individuals likely to have high eIF4E activity by all measures) reflecting the fact that z successfully takes account of both eIF4E and p4E-BP1, thereby supporting its utility as a potential predictive marker.

DISCUSSION

Expression of eIF4E in cancer has been studied extensively, however, expression does not equate to activity; therefore, interpretation of its influence is more complex than simply assessing expression. Our hypothesis was that combined examination of eIF4E and its regulators would allow greater insights into eIF4E's influence on cancer. Therefore, we determined the expression levels of eIF4E and its most well-established regulatory proteins 4E-BP1, 4E-BP2 and p4E-BP1 within tumour cells of a large cohort of cancer patients, and have combined these data into an improved measure of prognosis and estimate of eIF4E activity. In common with initial publications on eIF4E's role in cancer (Kerekatte *et al*, 1995; Li *et al*, 1998) and much of the subsequent literature, we have focused on breast cancer.

There is a wealth of literature on eIF4E expression in cancer and, although patient cohorts used have been relatively small (<200 individuals), the conclusion that high eIF4E levels are associated with poor prognosis is well established (De Benedetti and Graff, 2004). We have also found this (Figure 2A–C) using the largest cohort to date. The 4E-BPs have received less attention. 4E-BP1 is more highly expressed in tumours than normal tissues (Salehi and Mashayekhi, 2006), and expression correlates inversely with tumour progression (Martin *et al*, 2000) – observations in agreement with our findings. Despite the inverse correlation with progression, we found examination of 4E-BP1 expression not to give significant prognostic insights (Figure 2D, Supplementary Figure S5A). To our knowledge, there are no published studies concerning 4E-BP2 in cancer samples. We found that 4E-BP2 showed a nonsignificant trend for association with good prognosis (Figure 2E, Supplementary Figure S5B). High p4E-BP1 levels have

earlier been shown to correlate with grade and poor prognosis (Zhou *et al*, 2004; Castellvi *et al*, 2006; Rojo *et al*, 2007). In contrast, we found detectable p4E-BP1 (Thr37/46) in only 37.7% of the patients compared with >59% in other reports (Zhou *et al*, 2004; Rojo *et al*, 2007; Akcakanat *et al*, 2008)), and to provide little prognostic power when analysed alone (Figure 2F, Supplementary Figure S5C). We have carried out additional analyses with the same antibody as reported earlier for p4E-BP1 (Ser65) (Zhou *et al*, 2004) and similarly found low expression (50% undetectable, 28% the lowest positive score). These differences may relate to the cohorts used as earlier studies had higher proportions of node positive (Zhou *et al*, 2004), subsequently metastatic (Akcakanat *et al*, 2008) or high-grade cases (Rojo *et al*, 2007).

There are very few studies where multiple eIF4E pathway components have been analysed. The ratios of p4E-BP1 to total 4E-BP1 and of eIF4E to 4E-BP1 have been shown to correlate with high tumour grade (Salehi and Mashayekhi, 2006; Armengol *et al*, 2007), but their relationships with survival were not examined. A positive correlation between expression of eIF4E and p4E-BP1 has also been noted, with the conclusion being that eIF4E was 'active' in these cells (Nathan *et al*, 2004). We have determined expression levels of eIF4E, 4E-BP1, 4E-BP2, p4E-BP1 and have undertaken analyses to relate these together and to survival. First, we found no statistically significant relationship to survival for any combination without including eIF4E – suggesting that eIF4E is the critical effector. Second, in support of our initial hypothesis, we showed that combined assessment of the four components allowed improved prognostic insights over eIF4E alone ('z' function; text, Figure 3). It is important to note that we did not predetermine relationships between components in z, rather these were defined by the best fit with the data; the fact that the relationships reflect our expectations from understanding the pathway supports the view that z is a true estimate of eIF4E activity. Third, we showed differential expression of 4E-BP2 in cancer to be more influential in terms of survival than 4E-BP1. This was shown by the observations that expression of 4E-BP2, but not 4E-BP1, showed a trend towards being a prognostic factor alone (Figure 2E, Supplementary Figure S5B), provided an improved prognostic indicator in combination with eIF4E (y), and was the most statistically significant component of z after eIF4E itself. This observation may relate to the fact that 4E-BP2 binds, and therefore inhibits eIF4E more strongly than 4E-BP1 (Abiko *et al*, 2007). Interestingly, we found that expressions of eIF4E and 4E-BPs were positively associated (Table 1): an unexpected finding as they are functionally opposed and correlate oppositely with grade. One explanation is that 4E-BP translation may be specifically derepressed by eIF4E's action on the 5'UTRs of their transcripts, representing a negative feedback loop within the eIF4E pathway.

Clinical trials of the efficacy and safety of cancer therapeutics that target eIF4E have been carried out (Graff *et al*, 2008) and some toxicity has been reported (O'Donnell *et al*, 2008; Taberero *et al*, 2008). Selection of individuals who are most likely to benefit from the agents may be appropriate in order to avoid potentially harmful and/or ineffective therapy in some patients. We show that substantially different patient groups are chosen using three potential predictive biomarkers, and therefore that use of the best biomarker is important for targeting of these therapies (Figure 4). Patients with high estimated eIF4E activity ('z') (group 2) and high p4E-BP1 levels (group 3) should provide good candidates for treatment. The former group has a particularly poor prognosis (Figure 3C and F), and therefore great potential for clinical benefit from these drugs.

ACKNOWLEDGEMENTS

We thank Professors Blenis and Sonenberg for generously supplying a plasmid, and Yorkshire Cancer Research, US DoD

BCRP, Breast Cancer Research Action Group and Breast Cancer Campaign for financial support.

Supplementary Information accompanies the paper on British Journal of Cancer website (<http://www.nature.com/bjc>)

REFERENCES

- Abiko F, Tomoo K, Mizuno A, Morino S, Imataka H, Ishida T (2007) Binding preference of eIF4E for 4E-binding protein isoform and function of eIF4E N-terminal flexible region for interaction, studied by SPR analysis. *Biochem Biophys Res Commun* **355**: 667–672
- Akccakanat A, Sahin A, Shaye AN, Velasco MA, Meric-Bernstam F (2008) Comparison of Akt/mTOR signaling in primary breast tumors and matched distant metastases. *Cancer* **112**: 2352–2358
- Armengol G, Rojo F, Castellvi J, Iglesias C, Cuatrecasas M, Pons B, Baselga J, Ramon y Cajal S (2007) 4E-binding protein 1: a key molecular 'funnel factor' in human cancer with clinical implications. *Cancer Res* **67**: 7551–7555
- Buxade M, Parra-Palau JL, Proud CG (2008) The MnkS: MAP kinase-interacting kinases (MAP kinase signal-integrating kinases). *Front Biosci* **13**: 5359–5373
- Castellvi J, Garcia A, Rojo F, Ruiz-Marcellan C, Gil A, Baselga J, Ramon y Cajal S (2006) Phosphorylated 4E binding protein 1: a hallmark of cell signaling that correlates with survival in ovarian cancer. *Cancer* **107**: 1801–1811
- Cox DR, Oakes D (1984) *Analysis of Survival Data*. Chapman & Hall: London
- Culjkovic B, Topisirovic I, Borden KL (2007) Controlling gene expression through RNA regulons: the role of the eukaryotic translation initiation factor eIF4E. *Cell Cycle* **6**: 65–69
- Culjkovic B, Topisirovic I, Skrabanek L, Ruiz-Gutierrez M, Borden KL (2005) eIF4E promotes nuclear export of cyclin D1 mRNAs via an element in the 3'UTR. *J Cell Biol* **169**: 245–256
- De Benedetti A, Graff JR (2004) eIF-4E expression and its role in malignancies and metastases. *Oncogene* **23**: 3189–3199
- Dutton A, Reynolds GM, Dawson CW, Young LS, Murray PG (2005) Constitutive activation of phosphatidylinositol 3 kinase contributes to the survival of Hodgkin's lymphoma cells through a mechanism involving Akt kinase and mTOR. *J Pathol* **205**: 498–506
- Engelman JA, Chen L, Tan X, Crosby K, Guimaraes AR, Upadhyay R, Maira M, McNamara K, Perera SA, Song Y, Chirieac LR, Kaur R, Lightbown A, Simendinger J, Li T, Padera RF, Garcia-Echeverria C, Weissleder R, Mahmood U, Cantley LC, Wong KK (2008) Effective use of PI3K and MEK inhibitors to treat mutant Kras G12D and PIK3CA H1047R murine lung cancers. *Nat Med* **14**: 1351–1356
- Graff JR, Konicek BW, Carter JH, Marcusson EG (2008) Targeting the eukaryotic translation initiation factor 4E for cancer therapy. *Cancer Res* **68**: 631–634
- Gray N, Wickens M (1998) Control of translation initiation in animals. *Annu Rev Cell Dev Biol* **14**: 399–458
- Haybittle JL, Blamey RW, Elston CW, Johnson J, Doyle PJ, Campbell FC, Nicholson RI, Griffiths K (1982) A prognostic index in primary breast cancer. *Br J Cancer* **45**: 361–366
- Hughes TA (2007) 5' untranslated regions: critical regulators of cap-dependent translation. In *Leading-Edge Messenger RNA Research Communications*, Ostrovskiy M (ed), pp 33–45. Nova Science Publishers, Inc: New York, USA
- Johnson N, Speirs V, Curtin NJ, Hall AG (2008) A comparative study of genome-wide SNP, CGH microarray and protein expression analysis to explore genotypic and phenotypic mechanisms of acquired antiestrogen resistance in breast cancer. *Breast Cancer Res Treat* **111**: 55–63
- Kerekatte V, Smiley K, Hu B, Smith A, Gelder F, De Benedetti A (1995) The proto-oncogene/translation factor eIF4E: a survey of its expression in breast carcinomas. *Int J Cancer* **64**: 27–31
- Kozak M (1991) An analysis of vertebrate mRNA sequences: intimations of translational control. *J Cell Biol* **115**: 887–903
- Lee JW, Choi JJ, Lee KM, Choi CH, Kim TJ, Lee JH, Kim BG, Ahn G, Song SY, Bae DS (2005) eIF-4E expression is associated with histopathologic grades in cervical neoplasia. *Hum Pathol* **36**: 1197–1203
- Li BD, Gruner JS, Abreo F, Johnson LW, Yu H, Nawas S, McDonald JC, DeBenedetti A (2002) Prospective study of eukaryotic initiation factor 4E protein elevation and breast cancer outcome. *Ann Surg* **235**: 732–738; discussion 738–9
- Li BD, McDonald JC, Nassar R, De Benedetti A (1998) Clinical outcome in stage I to III breast carcinoma and eIF4E overexpression. *Ann Surg* **227**: 756–761; discussion 761–3
- Maraqa L, Cummings M, Peter MB, Shaaban AM, Horgan K, Hanby AM, Speirs V (2008) Carcinoembryonic antigen cell adhesion molecule 6 predicts breast cancer recurrence following adjuvant tamoxifen. *Clin Cancer Res* **14**: 405–411
- Martin ME, Perez MI, Redondo C, Alvarez MI, Salinas M, Fando JL (2000) 4E binding protein 1 expression is inversely correlated to the progression of gastrointestinal cancers. *Int J Biochem Cell Biol* **32**: 633–642
- Nathan CA, Amirghahari N, Abreo F, Rong X, Caldito G, Jones ML, Zhou H, Smith M, Kimberly D, Glass J (2004) Overexpressed eIF4E is functionally active in surgical margins of head and neck cancer patients via activation of the Akt/mammalian target of rapamycin pathway. *Clin Cancer Res* **10**: 5820–5827
- O'Donnell A, Faivre S, Burris III HA, Rea D, Papadimitrakopoulou V, Shand N, Lane HA, Hazell K, Zoellner U, Kovarik JM, Brock C, Jones S, Raymond E, Judson I (2008) Phase I pharmacokinetic and pharmacodynamic study of the oral mammalian target of rapamycin inhibitor everolimus in patients with advanced solid tumors. *J Clin Oncol* **26**: 1588–1595
- Pesole G, Mignone F, Gissi C, Grillo G, Licciulli F, Liuni S (2001) Structural and functional features of eukaryotic mRNA untranslated regions. *Gene* **276**: 73–81
- Polunovsky VA, Bitterman PB (2006) The cap-dependent translation apparatus integrates and amplifies cancer pathways. *RNA Biol* **3**: 10–17
- Poulin F, Gingras AC, Olsen H, Chevalier S, Sonenberg N (1998) 4E-BP3, a new member of the eukaryotic initiation factor 4E-binding protein family. *J Biol Chem* **273**: 14002–14007
- Richter JD, Sonenberg N (2005) Regulation of cap-dependent translation by eIF4E inhibitory proteins. *Nature* **433**: 477–480
- Rojo F, Najera L, Lirola J, Jimenez J, Guzman M, Sabadell MD, Baselga J, Ramon y Cajal S (2007) 4E-binding protein 1, a cell signaling hallmark in breast cancer that correlates with pathologic grade and prognosis. *Clin Cancer Res* **13**: 81–89
- Rosenwald IB, Chen JJ, Wang S, Savas L, London IM, Pullman J (1999) Upregulation of protein synthesis initiation factor eIF-4E is an early event during colon carcinogenesis. *Oncogene* **18**: 2507–2517
- Rosenwald IB, Hutzler MJ, Wang S, Savas L, Fraire AE (2001) Expression of eukaryotic translation initiation factors 4E and 2alpha is increased frequently in bronchioloalveolar but not in squamous cell carcinomas of the lung. *Cancer* **92**: 2164–2171
- Ruggero D, Montanaro L, Ma L, Xu W, Londei P, Cordon-Cardo C, Pandolfi PP (2004) The translation factor eIF-4E promotes tumor formation and cooperates with c-Myc in lymphomagenesis. *Nat Med* **10**: 484–486
- Salehi Z, Mashayekhi F (2006) Expression of the eukaryotic translation initiation factor 4E (eIF4E) and 4E-BP1 in esophageal cancer. *Clin Biochem* **39**: 404–409
- Scheper GC, Proud CG (2002) Does phosphorylation of the cap-binding protein eIF4E play a role in translation initiation? *Eur J Biochem* **269**: 5350–5359
- Smolewski P (2006) Recent developments in targeting the mammalian target of rapamycin (mTOR) kinase pathway. *Anticancer Drugs* **17**: 487–494
- Taberner J, Rojo F, Calvo E, Burris H, Judson I, Hazell K, Martinelli E, Ramon y Cajal S, Jones S, Vidal L, Shand N, Macarulla T, Ramos FJ, Dimitrijevic S, Zoellner U, Tang P, Stumm M, Lane HA, Leibold D, Baselga J (2008) Dose- and schedule-dependent inhibition of the mammalian target of rapamycin pathway with everolimus: a phase I tumor pharmacodynamic study in patients with advanced solid tumors. *J Clin Oncol* **26**: 1603–1610
- Zhou S, Wang GP, Liu C, Zhou M (2006) Eukaryotic initiation factor 4E (eIF4E) and angiogenesis: prognostic markers for breast cancer. *BMC Cancer* **6**: 231
- Zhou X, Tan M, Stone Hawthorne V, Klos KS, Lan KH, Yang Y, Yang W, Smith TL, Shi D, Yu D (2004) Activation of the Akt/mammalian target of rapamycin/4E-BP1 pathway by ErbB2 overexpression predicts tumor progression in breast cancers. *Clin Cancer Res* **10**: 6779–6788